

Green tea extract for treatment of cancers

A systematic review protocol

Kai Cheng, MM^{a,*}, Nan-Nan Chi, MM^a, Jun-Dong Liu, MB^b

Abstract

Background: Previous clinical studies suggested that green tea extract (GTE) may benefit patients with a variety of cancers. However, its efficacy is still inconclusive. Thus, the objective of this study will systematically collate the clinical studies testing its efficacy and safety for cancers.

Methods: We will perform a systematic review of clinical studies assessing the efficacy of GTE in variety of cancers. We will search Cochrane Central Register of Controlled Trials (CENTRAL), EMBASE, MEDLINE, Cumulative Index to Nursing and Allied Health Literature (CINAHL), Allied and Complementary Medicine Database (AMED), and Chinese Biomedical Literature Database (CBM) using a comprehensive strategy. We will also screen the reference lists of relevant studies to identify any additional studies for potential inclusion. All databases will be searched up to February 1, 2019. All eligible case-control studies and randomized controlled trials will be included in this study. Two independent authors will review all searched literature. Upon inclusion of trials, we will extract data by using a predefined standardized form. The risk of bias assessment will be evaluated by using Cochrane risk of bias tool. We will use RevMan 5.3 software to pool the data and carry out meta-analysis.

Results: The primary outcome includes overall response rate. The secondary outcomes comprise of overall survival, progression-free survival, the disease control rate, and any adverse events.

Conclusions: The results of this study will contribute to the understanding of the efficacy of GTE in the setting of cancers and promote future research of GTE in patients with cancers.

Dissemination and ethics: The results of this systematic review are expected to be published through peer-reviewed journals. This study does not need ethic approval, because it does not utilize individual patient data.

Systematic review registration: PROSPERO CRD42019125111.

Abbreviations: AMED = Allied and Complementary Medicine Database, CENTRAL = Cochrane Central Register of Controlled Trials, CIs = confidence intervals, GTE = green tea extract.

Keywords: cancer, efficacy, green tea extract, safety, systematic review

1. Introduction

Cancers are lethal disorder with poor outcomes and increasing incidence around the world.^[1,2] There are more than 100 types of cancer, such as lung cancer, breast cancer, prostate cancer, lymphoma, gastric cancer, ovarian cancer, and so on.^[3] Symptoms often vary according to the different types of cancer. A variety of factors contribute to the cancers, including, alcohol, obesity, dietary factors, and infection.^[4,5] The cancer management may include chemotherapy, radiation, and/or surgery.^[6,7]

However, all those treatments still have limited efficacy, and also accompany lots of toxicities.^[8] Therefore, more potential therapies are still urgently needed for patients with cancers.

Green tea is a very popular and widely consumed beverage globally.^[9,10] Previous studies have reported that highest green tea intake can significantly lower cancer risk.^[9-11] It contains several catechin components, also known as green tea extracts (GTE), including Epigallocatechin gallate, Epigallocatechin, Epicatechin-3-gallate, and Epicatechin.^[12] Several preclinical studies of green tea and its components are reported to show promising efficacy for the growth inhabitation of tumors with fewer adverse events.^[13,14] Furthermore, a numerous of clinical trials have reported that GTE can be used to treat a large range of cancers effectively, including breast cancer, colorectal cancer, prostate cancer, ovarian cancer, liver cancer, lung cancer, lymphoma, gastric cancer, and pancreatic cancer.^[9,10,15-29] Presently, no systematic review study has assessed the efficacy and safety of GTE for patients with a variety of cancers. Thus, this systematic review will first investigate the efficacy and safety of GTE for cancers.

N-NC and KC contributed equally to this study.

This study was funded by the Heilongjiang Natural Science Foundation Project (H2016088). The supporter has no roles in this study.

The authors have no conflicts of interest to disclose.

^a Second Ward of Gastroenterology Department, ^b Department of Pharmacy, First Affiliated Hospital of Jiamusi University, Jiamusi, China.

** Correspondence: Kai Cheng, Second Ward of Gastroenterology Department, First Affiliated Hospital of Jiamusi University, No. 348 Dexiang Street, Jiamusi, 154002, China (e-mail: jdfychengkai@126.com).*

Copyright © 2019 the Author(s). Published by Wolters Kluwer Health, Inc.

This is an open access article distributed under the Creative Commons Attribution License 4.0 (CCBY), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Medicine (2019) 98:15(e15117)

Received: 12 March 2019 / Accepted: 13 March 2019

<http://dx.doi.org/10.1097/MD.00000000000015117>

2. Methods

2.1. Objective

This systematic review aims to assess the efficacy and safety of GTE for patients with cancers.

Table 1**Search strategy for CENTRAL database.**

Number	Search terms
1	Mesh descriptor: (neoplasms) explode all trees
2	((cancer [*]) or (cancers [*]) or (lung cancer [*]) or (breast cancer [*]) or (prostate cancer [*]) or (colorectal cancer [*]) or (ovarian cancer [*]) or (liver cancer [*]) or (lymphoma [*]) or (gastric cancer [*]) or (pancreatic cancer [*])):ti, ab, kw
3	Or 1–2
4	Mesh descriptor: (tea) explode all trees
5	((green teav) or (extractv) or (epigallocatechin gallatev) or (epigallocatechin) or (epicatechin-3-gallatev) or (epicatechin):ti, ab, kw
6	Or 4–5
7	MeSH descriptor: (randomized controlled trials) explode all trees
8	((random [*]) or (allocation [*]) or (random allocation [*]) or (placebo [*]) or (blind [*]) or (RCT [*]) or (clinical trials [*]) or (controlled trials [*])):ti, ab, kw
9	MeSH descriptor: (case-control studies) explode all trees
10	((case-control [*]) or (studies [*]) or (case-control studies [*]) or (case [*]) or (control [*]) or (study [*])):ti, ab, kw
11	Or 7–10
12	3 and 6 and 11

CENTRAL = Cochrane Central Register of Controlled Trials, RCT = randomized controlled trial.

2.2. Study registration

The protocol of this study has been registered in PROSPERO with CRD42019125111.

2.3. Inclusion criteria for study selection

2.3.1. Types of studies. All qualified randomized controlled trials (RCTs) and case-control studies will be included in this study without language or publication restrictions. However, non-clinical studies, case studies, uncontrolled studies will not be included.

2.3.2. Types of patients. Patients who are clinically diagnosed with cancers, including breast cancer, colorectal cancer, prostate cancer, ovarian cancer, liver cancer, lung cancer, lymphoma, gastric cancer, and pancreatic cancer will all be fully considered for inclusion without the restrictions of race, gender, and age. However, patients with other severe disorders that may affect the outcome measurements will be excluded, such as acute heart failure, severe stroke and so on.

2.3.3. Types of interventions. Any types of GTE therapy will be considered for inclusion, except the combination of GTE with other interventions. Control therapy can be any treatments, except the GTE, or any kinds of green tea.

2.3.4. Types of outcome measurements. The primary outcome is measured by overall response rate (ORR). The secondary outcomes are assessed by overall survival (OS), progression-free survival (PFS), the disease control rate (DCR), and any adverse events.

2.4. Search methods for the identification of studies

2.4.1. Search electronic bibliographic databases. We will search Cochrane Central Register of Controlled Trials (CENTRAL), EMBASE, MEDLINE, the Cumulative Index to Nursing and Allied Health Literature (CINAHL), the Allied and Complementary Medicine Database (AMED), and Chinese databases of Chinese Biomedical Literature Database (CBM) up to February 1, 2019 using a comprehensive strategy. Reference lists of relevant studies will also be identified for potential consideration. The sample of comprehensive search strategy for CENTRAL is presented in Table 1. Similar comprehensive search strategies will also be applied to any other electronic databases.

2.4.2. Search for other resources. References lists of relevant trials will also be searched to avoid missing any potential studies.

2.5. Data collection and extraction

2.5.1. Study selection. Two authors will independently screen the title and abstract of all identified studies for eligibility. Then, full-texts will be read for further selection in accordance with the predefined eligibility criteria. Any disagreements between 2 authors will be solved by consulting a third author. The whole process of study selection will be presented in Figure 1.

2.5.2. Data extraction and management. Two authors will independently extract data according to the predefined and standardized data extraction form. This form comprises of the data of study characteristics, such as title, first author, published year, location; patient populations, such as race, age, gender, eligibility criteria; study design, such as sample size, randomization, concealment, and blinding; interventions, such as dosage, frequency, and duration; and outcomes, such as primary, secondary, and safety outcome measurements. All divergences regarding the data extraction between 2 authors will be solved by a third author through discussion.

2.5.3. Dealing with missing data. Any missing data will be inquired by contacting the primary authors. If we can not receive those data, then we will only analyze the available data and will discuss its potential impacts.

2.6. Risk of bias assessment

We will use Cochrane Risk of Bias Tool to evaluate the risk of bias for each included study. This tool assesses the risk of bias in 7 domains, and each domain will be determined as high, unclear, or low risk of bias. Any dissimilarity regarding the risk of bias assessment between 2 authors will be settled down by consensus with a third author.

2.7. Statistical analysis

RevMan 5.3 software will be utilized to pool the data and conduct meta-analysis in this study.

2.7.1. Measurement of treatment effect. Dichotomous data will be shown as risk ratio and 95% confidence intervals (CIs). Continuous data will be demonstrated as mean difference or standardized mean difference and 95% CIs.




Figure 1. Flowchart of study selection.

2.7.2. Assessment of heterogeneity. Cochrane I^2 test will be used to assess the heterogeneity. If the value of $I^2 > 50\%$, significant heterogeneity is considered. Otherwise, if the value of $I^2 \leq 50\%$, acceptable heterogeneity is considered.

2.7.3. Data synthesis. If acceptable heterogeneity will be detected, a fixed-effect model will be used to pool and meta-analysis will be performed. Otherwise, if significant heterogeneity will be identified, a random-effect model will be applied, and subgroup analysis will be carried out. Meta-analysis will be conducted according to the results of subgroup analysis. If there is still significant heterogeneity after the subgroup analysis, meta-analysis will not be conducted. However, a narrative summary will be described.

2.7.4. Subgroup analysis. Subgroup analysis will be carried out to identify any reasons that may cause the substantial heterogeneity. It will be performed based on the different treatments, controls, and outcome measurements.

2.7.5. Sensitivity analysis. Sensitivity analysis will be conducted to check the stability of the pooled results by removing the low quality of included studies.

2.7.6. Publication biases. If there will be more than 10 qualified studies are included, Funnel plot and Egger linear regression test will be operated to identify any potential publication biases.

3. Discussion

Currently, no systematic review has been performed regarding the efficacy and safety of GTE for the treatment of patients with cancers. In this systematic review, we will first investigate the efficacy and safety of GTE for a variety of cancers. The results of this study will summarize the latest evidence for assessing the efficacy and safety of GTE for the treatment of cancers. Its findings may provide helpful evidence for both patients and clinicians.

Author contributions

Conceptualization: Kai Cheng, Nan-Nan Chi.
Data curation: Kai Cheng, Nan-Nan Chi, Jun-Dong Liu.
Formal analysis: Nan-Nan Chi, Jun-Dong Liu.
Funding acquisition: Kai Cheng.
Investigation: Kai Cheng.
Methodology: Nan-Nan Chi, Jun-Dong Liu.
Project administration: Kai Cheng.
Resources: Nan-Nan Chi, Jun-Dong Liu.
Software: Nan-Nan Chi, Jun-Dong Liu.
Supervision: Kai Cheng.
Validation: Kai Cheng, Nan-Nan Chi.
Visualization: Kai Cheng.
Writing – Original Draft: Kai Cheng, Nan-Nan Chi, Jun-Dong Liu.

Writing – Review & Editing: Kai Cheng, Nan-Nan Chi, Jun-Dong Liu.

References

- [1] De Flora S, La Maestra S. Epidemiology of cancers of infectious origin and prevention strategies. *J Prev Med Hyg* 2015;56:E15–20.
- [2] Koubkova L, Hrstka R, Dobes P, et al. Second primary cancers - causes, incidence and the future. *Klin Onkol* 2014;27:11–7.
- [3] Denis H, Davoine C, Bermudez E, et al. Specific immunotherapies in the treatment of cancers. *Bull Cancer* 2019;106:37–47.
- [4] Yang L, Drake BF, Colditz GA. Obesity and other cancers. *J Clin Oncol* 2016;34:4231–7.
- [5] Lee PN, Thornton AJ, Hamling JS. Epidemiological evidence on environmental tobacco smoke and cancers other than lung or breast. *Regul Toxicol Pharmacol* 2016;80:134–63.
- [6] Mohiuddin M, Regine WF, Stevens J, et al. Combined intraoperative radiation and perioperative chemotherapy for unresectable cancers of the pancreas. *J Clin Oncol* 1995;13:2764–8.
- [7] Bernier J, Poortmans PM. Surgery and radiation therapy of triple-negative breast cancers: from biology to clinics. *Breast* 2016;28:148–55.
- [8] Cetina-Pérez L, Serrano-Olvera A, Flores-Cisneros L, et al. Epidemiological profile, gastrointestinal toxicity, and treatment of pelvic cancers in patients managed with radiotherapy to the abdominal pelvic area. *Rev Invest Clin* 2018;70:112–6.
- [9] Hu Y, McIntosh GH, Le Leu RK, et al. Supplementation with Brazil nuts and green tea extract regulates targeted biomarkers related to colorectal cancer risk in humans. *Br J Nutr* 2016;116:1901–11.
- [10] Samavat H, Dostal AM, Wang R, et al. The Minnesota Green Tea Trial (MGTT), a randomized controlled trial of the efficacy of green tea extract on biomarkers of breast cancer risk: study rationale, design, methods, and participant characteristics. *Cancer Causes Control* 2015;26:1405–19.
- [11] Imai K, Suga K, Nakachi K. Cancer-preventive effects of drinking green tea among a Japanese population. *Prev Med* 1997;26:769–75.
- [12] Mattoli L, Mercati V, Burico M, et al. Experimental evidence of the presence of bimolecular caffeine/catechin complexes in green tea extracts. *J Nat Prod* 2018;81:2338–47.
- [13] Fujiki H, Suganuma M, Okabe S, et al. Japanese green tea as a cancer preventive in humans. *Nutr Rev* 1996;54:S67–70.
- [14] Yoshizawa S, Horiuchi T, Fujiki H, et al. Antitumor promoting activity of (–)-epigallocatechin gallate, the main constituent of “tannin” in green tea. *Phytother Res* 1987;1:44–7.
- [15] Samavat H, Ursin G, Emory TH, et al. A randomized controlled trial of green tea extract supplementation and mammographic density in postmenopausal women at increased risk of breast cancer. *Cancer Prev Res (Phila)* 2017;10:710–8.
- [16] Lazzeroni M, Guerrieri-Gonzaga A, Gandini S, et al. A presurgical study of lecithin formulation of green tea extract in women with early breast cancer. *Cancer Prev Res (Phila)* 2017;10:363–70.
- [17] Dostal AM, Samavat H, Bedell S, et al. The safety of green tea extract supplementation in postmenopausal women at risk for breast cancer: results of the Minnesota Green Tea Trial. *Food Chem Toxicol* 2015;83:26–35.
- [18] Kumar NB, Pow-Sang J, Egan KM, et al. Randomized, placebo-controlled trial of green tea catechins for prostate cancer prevention. *Cancer Prev Res (Phila)* 2015;8:879–87.
- [19] Crew KD, Ho KA, Brown P, et al. Effects of a green tea extract, Polyphenon E, on systemic biomarkers of growth factor signalling in women with hormone receptor-negative breast cancer. *J Hum Nutr Diet* 2015;28:272–82.
- [20] Trudel D, Labbé DP, Araya-Farias M, et al. A two-stage, single-arm, phase II study of EGCG-enriched green tea drink as a maintenance therapy in women with advanced stage ovarian cancer. *Gynecol Oncol* 2013;131:357–61.
- [21] Stendell-Hollis NR, Thomson CA, Thompson PA, et al. Green tea improves metabolic biomarkers, not weight or body composition: a pilot study in overweight breast cancer survivors. *J Hum Nutr Diet* 2010;23:590–600.
- [22] Wang P, Aronson WJ, Huang M, et al. Green tea polyphenols and metabolites in prostatectomy tissue: implications for cancer prevention. *Cancer Prev Res (Phila)* 2010;3:985–93.
- [23] Brausi M, Rizzi F, Bettuzzi S. Chemoprevention of human prostate cancer by green tea catechins: two years later. A follow-up update. *Eur Urol* 2008;54:472–3.
- [24] Bettuzzi S, Brausi M, Rizzi F, et al. Chemoprevention of human prostate cancer by oral administration of green tea catechins in volunteers with high-grade prostate intraepithelial neoplasia: a preliminary report from a one-year proof-of-principle study. *Cancer Res* 2006;66:1234–40.
- [25] Luo H, Tang L, Tang M, et al. Phase IIa chemoprevention trial of green tea polyphenols in high-risk individuals of liver cancer: modulation of urinary excretion of green tea polyphenols and 8-hydroxydeoxyguanosine. *Carcinogenesis* 2006;27:262–8.
- [26] Choan E, Segal R, Jonker D, et al. A prospective clinical trial of green tea for hormone refractory prostate cancer: an evaluation of the complementary/alternative therapy approach. *Urol Oncol* 2005;23:108–13.
- [27] Laurie SA, Miller VA, Grant SC, et al. Phase I study of green tea extract in patients with advanced lung cancer. *Cancer Chemother Pharmacol* 2005;55:33–8.
- [28] The Indian-US Head and Neck Cancer Cooperative Group. Green tea and leukoplakia. *Am J Surg* 1997;174:552–5.
- [29] Dai Q, Ji BT, Jin F, et al. Case-control study of pancreatic cancer in Shanghai population—relationship between pancreatic cancer and tobacco, alcohol and green tea. *Tumor* 1996;1:5–10.