


PRIMARY RESEARCH

Open Access



TACR2 is associated with the immune microenvironment and inhibits migration and proliferation via the Wnt/ β -catenin signaling pathway in prostate cancer

Wang Jianfeng, Wang Yutao and Bi Jianbin 

Abstract

Background: The tachykinin receptor 2 (TACR2) is encoded by the tachykinin receptor correlation gene. Recent microarray analysis for prostate cancer suggests that TACR2 expression is associated with clinical phenotype and disease-free survival among patients with prostate cancer.

Results: TACR2 protein levels were lower in prostate cancer tissues than in adjacent normal prostate tissue. TACR2 expression significantly correlated with clinical stage, Gleason scores, and survival outcomes. TACR2 expression positively correlated with mast cells and negatively correlated with M2 macrophages. Overexpression of TACR2 promoted the migration and proliferation of prostate cancer cells by regulating the Wnt signaling pathway.

Conclusions: The TACR2-Wnt/ β -catenin signaling pathway is critical in prostate cancer. TACR2 may affect tumor cells' occurrence and development by changing the content of immune cells in the tumor microenvironment. These findings suggest that TACR2 may be a candidate molecular biomarker for prostate cancer therapy.

Keywords: Tachykinin receptor 2 (TACR2), Immune microenvironment, Wnt/ β -catenin signaling pathway, Prostate cancer

Background

Prostate cancer is the second most common malignant tumor in men and the fifth most common cause of cancer death worldwide. Although its mortality has decreased in recent years, more than 300,000 men still die of prostate cancer every year; especially in developed countries, prostate cancer prevalence remains very high [1–3]. Prostate-specific antigen (PSA) is a kallikrein-like serine protease secreted by prostatic epithelial cells. Its serum expression levels can be used to screen for prostate cancer. Nevertheless, the utility of PSA is limited and controversial [4–6]. Therefore, new biomarkers are urgently

needed to measure the progression of prostate cancer more accurately.

In a previous study, we established and validated a prognostic risk score formula called CMU5 based on the expression of five genes: *FAM72D*, *ARHGAP33*, *TACR2*, *PLEK2*, and *FA2H*; the signature reliably predicted prostate cancer outcome [7]. Among these five genes, we focused on *TACR2*, which encodes tachykinin receptor NK2, one of the three tachykinin receptors that belong to the G-protein-coupled receptor superfamily [8]. It is not only expressed in the peripheral and central nervous system but also in some peripheral tissues [8, 9]. Tachykinins interact with tachykinin receptors and participate in critical physiological processes of the respiratory, cardiovascular, immune, endocrine, gastrointestinal, and genitourinary systems, as well as other peripheral

*Correspondence: jianbinbi@cmu.edu.cn
Department of Urology, The First Hospital of China Medical University, Shenyang, Liaoning, People's Republic of China



© The Author(s) 2021. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

organ systems [10, 11]. Tachykinins are involved in the development of many diseases, including cancer. Abnormal expression of tachykinin receptor coding genes was found in breast cancer, colon cancer, and other tumors. Nevertheless, the role of TACR2 in cancer progression remains unclear [11–13].

The Wnt signaling pathway includes the canonical Wnt/ β -catenin signaling pathway, the non-canonical Wnt/ Ca^{2+} signaling pathway, and the Wnt/planar cell polarity signaling pathway [14]. The classic Wnt/ β -catenin signaling pathway is the most characterized Wnt pathway [15]; it is involved in various cellular functions, including cell proliferation, stem cell self-renewal, and organogenesis [15–17]. The Wnt/ β -catenin signaling pathway is closely related to the occurrence and development of breast cancer, colon cancer, ovarian cancer, prostate cancer, and other malignant tumors. It is essential for the migration, proliferation, angiogenesis, and invasion of cancer cells [16–20]. In this study, we determined the effect of TACR2 on the immune microenvironment of prostate cancer by examining the content of immune cells. In addition, we found that TACR2 may affect the migration and proliferation of prostate cancer cells by regulating the Wnt/ β -catenin signaling pathway. These findings suggest that TACR2 may be used as a marker of prostate cancer development and may become a potential therapeutic target.

Results

TACR2 was downregulated in tumor tissue

To explore the expression pattern of TACR2, we used the TIMER database to determine that TACR2 expression was significantly downregulated in a variety of cancer tissues. Of these, TACR2 expression was significantly downregulated in prostate cancer, clear cell renal cell carcinoma, chromophobe tumor, and papillary renal cell carcinoma (Fig. 1A). We performed a paired t-test and clinical correlation analysis in TCGA-PRAD. We found that TACR2 expression levels were low in tissues adjacent to prostate cancer and were higher in patients with higher clinical stages and higher Gleason scores (Fig. 1B, $P < 0.001$). In the survival analysis with disease-free recurrence as the follow-up endpoint, TACR2 high-expressing prostate cancer patients had better survival outcomes than low-expressing patients (Fig. 1C, $P < 0.001$).

TACR2 immune correlation

In GSE46602, we divided the expression matrix into two groups according to the median expression of TACR2. We then compared the differential pathways in the two sets of samples according to the GSEA analysis strategy. The T cell receptor and Wnt pathways were significantly enriched (Fig. 1D). These findings suggest that TACR2

may be closely related to these two pathways. Therefore, to conduct a detailed analysis of these two pathways, we measured the immune cell content in TCGA-PRAD (Fig. 2A) and then analyzed the correlation between TACR2 expression levels and immune cell content (Fig. 2B). We found that TACR2 expression levels positively correlated with MAST cells (Fig. 2C: $\text{COR} = 0.22$, $P < 0.001$). TACR2 expression levels negatively correlated with M2 macrophages (Fig. 2D: $\text{COR} = -0.17$, $P < 0.001$). These trends suggest that TACR2 expression is a prognostic protective factor for prostate cancer.

TACR2 co-expression network


We attempted to construct a co-expression network of TACR2 by analyzing its protein-coding genes and exploring the biological processes involved in TACR2 by enriching the co-expression network's functions. According to the cutting height = 120,000, 495 samples were included, and a tree diagram of 265 samples was obtained (Fig. 3A). We built scale-free co-expression networks. The best soft threshold was 5, and R-squared was 0.9 (Fig. 3B). A hierarchical cluster tree was constructed, with each leaf representing a gene and each branch representing a co-expression module. A total of eight co-expression modules were generated (Fig. 3C). The Brown module ($\text{COR} = 0.6$; $P < 0.001$) and gray module ($\text{COR} = 0.48$; $P < 0.001$) had high correlations with TACR2 (Fig. 3D). There was a correlation between Brown module membership and the gene significance of TACR2 (Fig. 3E: $\text{COR} = 0.69$; $P < 0.001$).

TACR2 pathway analysis

We found that TACR2 is associated with the Wnt pathway in the GSE46602 cohort. We obtained the same result in TCGA-PRAD (Fig. 4A). Then, we enriched the co-expression network of TACR2. The enrichment results showed that TACR2 is involved in biological processes such as muscle contraction (Fig. 4B). In the pathway analysis, GSE46602 and the TCGA-PRAD cohort confirmed that TACR2 might regulate the Wnt pathway. This pathway affects the progression of prostate cancer. Therefore, we performed a Pearson pathway analysis on the expression of TACR2 and the genes in the Wnt pathway. We found that TACR2 and gene expression levels in the Wnt pathway were also closely related (Fig. 4C).

Expression of TACR2 and β -catenin in prostate cancer and adjacent normal tissue


TCGA database analysis showed that transcription levels of TACR2 were significantly different between prostate cancer and adjacent normal tissues. Western blot was used to measure expression levels of TACR2 protein in 30 pairs of prostate cancer and adjacent normal tissues.



Consistent with the TCGA database, expression levels of TACR2 protein in prostate cancer tissues were significantly lower than in adjacent tissues (Fig. 5A, B, $P < 0.01$). Next, we found that the expression of β -catenin in prostate cancer tissues was significantly higher than in adjacent tissues (Fig. 5A, C, $P < 0.01$). The clinicopathological data of these 30 patients with prostate cancer was shown in Additional file 1: Table S1.

Overexpression of TACR2 deactivates the Wnt/ β -catenin signaling pathway

To verify the effect of TACR2 on Wnt/ β -catenin signaling transduction, we selected hormone-resistant prostate cancer cells (DU145 and PC3) and hormone-sensitive prostate cancer cells (LNCaP) and transfected them with TACR2 overexpression lentiviral vector and corresponding negative control lentiviral vector. Western blot test



showed that protein levels of TACR2 in the TA-OE cell lines were significantly higher than in the NC cells (Fig. 5D, E, P < 0.01). β -Catenin is the core component of the Wnt/ β -catenin signaling pathway; its increased levels suggest pathway activation [21, 22]. Compared with the NC cells, expression levels of β -catenin and Cyclin D1 proteins in the TA-OE cell lines were significantly higher (Fig. 5D, E, P < 0.01). After isolating nuclear proteins, we found that the nuclear protein levels of β -catenin in TA-OE cells were significantly lower than that in NC cells (Fig. 5F, G, P < 0.01). This evidence suggests that TACR2 overexpression significantly blocks activation of the Wnt/ β -catenin signaling pathway.

Overexpression of TACR2 inhibited cell activity, proliferation, and migration

To further study the biological function of TACR2 in prostate cancer, the EdU test was used to measure the proliferation of TACR2-overexpressing cells. We found that the cells' proliferation ability was lower than that of NC cells (Fig. 6A, P < 0.01). The CCK-8 was used to analyze the activity of TACR2-overexpressing cells. The TA-OE cell lines activity was significantly lower

than that of NC cell lines (Fig. 6B, P < 0.05). We used a wound-healing migration assay and a Transwell assay to determine the role of TACR2 in prostate cancer cell migration. Transwell assays showed that the cell migration decreased with overexpression of TACR2 (Fig. 6C, P < 0.01). Consistently, the wound-healing migration assay showed that the migration ability of PC3 and DU145 decreased significantly after TACR2 overexpression (Fig. 6D, P < 0.01).

Discussion

In recent years, with the popularity of PSA screening and advances in prostate cancer treatment, prostate cancer incidence has been declining or remaining stable; nevertheless, the mortality rate of prostate cancer has not declined [23]. PSA's limitation as a biomarker for prostate cancer has been widely reported [24, 25]. Therefore, more specific and useful new biomarkers are needed to surpass the limitations of traditional biomarkers. Identifying new biomarkers and clarifying their signal transduction process will reveal the molecular mechanisms of the occurrence and development of prostate cancer. Doing so will





Fig. 3 Results of WGCNA analysis in TCGA-PRAD. **A** According to the cutting height = 120,000, 495 samples were included, and the tree diagram of 265 samples was obtained. **B** We built scale-free co-expression networks. The best soft threshold was 5, R-squared was 0.9. **C** A hierarchical cluster tree was constructed, with each leaf representing a gene and each branch representing a co-expression module, and a total of eight co-expression modules were generated. **D** Correlation coefficients between different factors and co-expression modules, where the Brown module (COR = 0.6; $P = 4E-41$) and grey module (COR = 0.48; $P = 1E-24$) had a high correlation with TACR2. The correlation between Brown module membership and gene significance for TACR2 was COR = 0.69; $P = 9.5e-74$

help to predict disease outcome and identify better treatment decisions.

TACR2 is involved in a series of biological processes, including controlling intestinal motility, hormone secretion, and visceral sensitivity [9]. The relationship between

TACR2 and tumor remains unclear. Using the TIMER database, we found that the expression of TACR2 was significantly downregulated in a variety of cancer tissues, including prostate cancer, clear cell renal cell carcinoma, chromophobe tumor, and papillary renal cell carcinoma.



Expression levels of TACR2 significantly correlated with clinical stage, Gleason score, and survival outcomes. Using GSEA, we analyzed the differential pathways between TACR2 high- and low-expression groups of gse46602 and found that the T cell receptor and Wnt

pathways were enriched. Therefore, we performed a detailed analysis of these two pathways.

Recent studies showed that tachykinins and their receptors regulate the development, activation, and function of critical immune cells and the immune system.

(See figure on next page.)

Fig. 5 Expression of TACR2 and β -catenin in prostate cancer and prostate cancer cell lines. **A, B** The expression of TACR2 protein in prostate cancer tissues (T) and adjacent tissues (N) ($P < 0.01$). **A, C** The expression of β -catenin in prostate cancer tissues and adjacent tissues ($P < 0.01$). **D, E** After transfection with TACR2 overexpression lentiviral vector and corresponding negative control lentiviral vector in DU145, PC3, and LNCaP, protein levels of TACR2, β -catenin and Cyclin D1 were measured using western blot. **F, G** After isolating nuclear proteins, the nuclear protein levels of β -catenin in TA-OE and NC cells were measured using western blot. The samples derive from the same experiment and those blots were processed in parallel




Fig. 5 (See legend on previous page.)

Tachykinins are also involved in immune cell migration, immune cell proliferation, and inflammation regulation with cytokines [26, 27]. We analyzed the correlation between expression levels of TACR2 and the content of immune cells and found that expression levels of TACR2 positively correlated with mast cells and negatively correlated with M2 macrophages. The number of infiltrating mast cells in tumor tissue is closely related to tumor angiogenesis, differentiation, and metastasis. In the prostate cancer microenvironment, mast cells drive adverse events, including tumor invasion, angiogenesis, biochemical recurrence, and metastasis after radical prostatectomy [28–30]. M2 macrophages cannot present antigens. The higher their content, the weaker the patient's immune response and the worse the outcome [31–33]. These results suggest that TACR2 may regulate the immune cell receptor pathway by changing the immune cell tumor microenvironment and then affecting the occurrence and development of tumor cells. TACR2 is expected to be a target for immunotherapy of prostate cancer.

Next, we analyzed the protein-coding genes of TACR2 to construct its co-expression network and explored the biological process of TACR2 by enriching the function of the co-expression network. The enrichment results showed that TACR2 was involved in muscle contraction and other biological processes. In the pathway analysis, the TCGA-PRAD cohort confirmed that TACR2 might regulate the Wnt signaling pathway. We also found that TACR2 is closely related to the expression levels of critical genes in the Wnt pathway. We measured protein expression in 30 pairs of prostate cancer and adjacent normal tissues. Consistent with the data analysis, the expression of TACR2 protein in prostate cancer tissues was significantly lower, while the expression of β -catenin was significantly higher. Three cell types (hormone-sensitive LNCaP cells, non-hormone-sensitive PC3, and DU145 cells) were selected for TACR2 overexpression. Compared with NC cells, the total protein and nuclear protein levels of β -catenin in TA-OE cells was significantly greater. The protein expression level of Cyclin D1, the downstream target gene of β -catenin, was also consistent with the trend of β -catenin in NC and TA-OE cells. This evidence suggests that TACR2 overexpression significantly blocks the activation of the Wnt/ β -catenin signaling pathway. Further cell function experiments

showed that overexpression of TACR2 inhibited the activity, proliferation, and migration of prostate cancer cells.

In conclusion, TACR2 is expressed at low levels in prostate cancer tissue and inhibits prostate cancer cells' migration and proliferation via the Wnt/ β -catenin signaling pathway. These findings suggest TACR2 may affect tumor cells' occurrence and development by changing the content of immune cells in the tumor microenvironment. Nevertheless, the specific mechanisms of TACR2 and the Wnt/ β -catenin signaling pathway and the mechanism of regulating the tumor microenvironment require further study. These findings suggest that TACR2 may be a new biomarker and may provide a theoretical basis for prostate cancer treatment.

Methods

Statement

The collection of tissue samples was approved by the Ethics Committee of the First Hospital of China Medical University according to the Declaration of Helsinki, and all participants provided written informed consent (approval number, [2018] 2018-190-2). All methods were performed in accordance with the relevant guidelines and regulations.

Matrix source

The prostate cancer mRNA matrix and clinical information were obtained from TCGA (The Cancer Genome Atlas) database (<https://www.cancer.gov/>) [34]. We considered 547 prostate samples, including 52 normal prostate samples and 495 tumor samples. We downloaded 36 prostate cancer tissues matrix and the clinical information from GSE46602, the platform for which is the GPL570 Affymetrix Human Genome U133 Plus 2.0 Array [35].

Pathway analysis

There are few studies on TACR2 at present. To explore the pathways related to TACR2 in prostate cancer, we conducted Gene Set Enrichment Analysis (GSEA) [36]. The GSEA method is used to determine whether a gene set of interest differs significantly in two biological states. We used GSEA to explore functional pathways related to TACR2 expression, grouping according to the median of TACR2 expression. Using a single data set that causes

(See figure on next page.)

Fig. 6 Overexpression of TACR2 inhibited cell activity, proliferation, and migration. **A** The effect of TACR2 overexpression on the proliferation of DU145, PC3, and LNCaP cells were determined using the EdU assay (magnification $\times 200$). **B** Overexpression of TACR2 suppressed cell viability according to the CCK-8 assay. **C** Overexpression of TACR2 inhibited cell migration according to the Transwell assay (magnification $\times 20$). **D** According to the wound-healing assay, overexpression of TACR2 inhibited cell migration in 5637 and T24 cells (magnification $\times 20$)




Fig. 6 (See legend on previous page.)

high false-positive results, we performed cross-validation analysis between TCGA-PRAD, and GSE46602.

TACR2 co-expression mRNA analysis

We found that TACR2 was significantly related to the Wnt pathway. We know that the Wnt pathway is involved in a variety of biological processes in the body. To determine which biological processes TACR2 relates to, we used Gene Ontology (GO) (<http://geneontology.org/>) analysis to illustrate the biological function of TACR2 co-expressed protein-coding genes [37]. We used the weighted gene co-expression network (WGCNA) method to explore the co-expression factors of TACR2 by constructing a scale-free network [38]. The number of genes in the minimum module was defined as 30. By drawing a correlation heat map between the module and TACR2, we determined the module with the strongest correlation with TACR2. We extracted protein-coding genes with Pearson correlation coefficients >0.4. GO analysis of these genes identified the most relevant biological functions and molecular results of TACR2. We did not directly screen based on the Pearson correlation coefficient of TACR2 and other protein-coding genes; instead, we first determined its co-expression module before the screening. Such a screening strategy assumes that co-expression modules usually have similar biological behaviors. The most relevant co-expression module of TACR2 can be obtained using this method, determining the most relevant biological function.

Immune cell proportion correlation

There is tachykinin-mediated modulation of the immune response [39]. Therefore, we attempted to determine associated immune processes by measuring prostate cancer tissues' immune cell content. We ran the CIBERSORT algorithm and determined the immune cell proportion based on the bulk tissue gene expression matrices. LM22 is a gene signature matrix that defines 22 immune cell subtypes obtained from a website (<https://cibersort.stanford.edu/>).

Clinical specimen collection

From January 2020 to December 2020, we performed radical prostatectomy on prostate cancer patients according to EAU GUIDELINES ON PROSTATE CANCER, and 30 pairs of prostate cancer tissue samples and adjacent normal prostate tissue samples (>1 cm from the tumor) were collected at the First Hospital of China Medical University (Shenyang, China). No patients received endocrine therapy, chemotherapy, or radiotherapy before radical prostatectomy. All tissues were examined histologically. Prostate cancer tissue samples had high-density cancer foci, and no cancer foci were found in the adjacent

normal prostate tissue samples. Tissue samples were stored at -80°C before use.

Cell lines and reagents

The human prostate cancer cell lines, PC3, DU145, and LNCaP Clone FGC, were obtained from the National Collection of Authenticated Cell Cultures (Shanghai, China) and were STR certified. PCR was used to detect Mycoplasma in the culture medium, and the passage time of the cells was not more than 6 months. The cell lines were cultured in RPMI 1640 (HyClone, USA) supplemented with 10% fetal bovine serum (FBS, Gibco) and grown at 37°C , 5% CO_2 . The antibodies included TACR2 (Proteintech, 25270-1-AP), beta-Catenin (Proteintech, 51067-2-AP), GAPDH (Cell Signaling Technology, 5174S), Cyclin D1 (Cell Signaling Technology, 55506S), Lamin B1 (Cell Signaling Technology, 13435S).

Construction of cell lines with TACR2 overexpression

PC3, DU145, and LNCaP cells were seeded in 6-well or other plates and were transfected 24 h later. Genechem (Shanghai, Genechem) was commissioned to design and synthesize the TACR2 overexpression lentiviral vector and the corresponding negative control lentiviral vector. Three cell lines were transfected according to the manufacturer's protocol. The stable negative control (NC) and TACR2 overexpression (TA-OE) cell lines were established in a puromycin-containing medium.

Western blot analysis

The total proteins were extracted from tissues and cell lines using RIPA lysis buffer (containing protease and phosphatase inhibitors). Nuclear protein was extracted using the Nuclear and Cytoplasmic Protein Extraction kit (Beyotime). Protein concentration was determined with bicinchoninic acid assay (Beyotime Institute of Biotechnology). The denatured protein sample (40 $\mu\text{g}/\text{lane}$) was added to the 10% sodium dodecyl sulfate–polyacrylamide gel for electrophoresis (140 V, 60 min), then separated and transferred to the polyvinylidene fluoride membrane (340 mA, 90 min). Subsequently, membranes were blocked in a sealed container with 5% non-fat milk (37°C , 1 h) and incubated with anti-TACR2 (1:500), anti- β -catenin (1:5000) or anti-GAPDH (1:1000) primary antibodies overnight in 5% fat-free milk at 4°C .

EdU assay

The stable cells were seeded on 24-well plates. According to the manufacturer's instructions, EdU (BeyoClick™, EDU-488, China) reagent was added to the medium at the ratio of 1:1000. According to the manufacturer's protocol, the culture was continued at 37°C for 2 h; after labeling, complete the following experimental steps.

Finally, the number of proliferating cells was counted under the fluorescence microscope (Olympus Corporation, Japan).

Cell viability assay

The stable PC3, DU145, and LNCaP cells were seeded into 96-well plates at the density of 5×10^3 cells/well. Cell Counting Kit-8 (CCK-8) (Bimake, USA) solution was added to each well to a final concentration of 0.5 mg/ml and incubated at 37 °C for 1 h. The absorbance was measured at 450 nm using a plate reader (Model 680; Bio-Rad Laboratories).

Transwell assay

Transwell chambers without matrix were used for cell migration tests. We plated 200 μ l of serum-free medium containing stable cells (1×10^5) into the upper chambers (Corning, NY, USA) of 24-well plates, and 600 μ l medium (10% FBS) was plated into the lower chamber. After 48 h of incubation, the cells under the membrane were fixed with 4% paraformaldehyde for 10 min and stained with crystal violet stain for 10 min. The remaining cells in the upper chamber were removed with cotton swabs. The migration of cells on the surface under the membrane was measured using an inverted light microscope, and the migration efficiency was calculated using ImageJ.

Wound-healing migration assay

Wound-healing tests were used to measure cell migration. Stable PC3 and DU145 cells were seeded on 24-well plates. When 90–100% confluence was achieved, a 200- μ l sterile pipette was used to create a linear scratch. After washing with a serum-free medium, the cells were cultured for 48 h. Cell images at 0 and 48 h after injury were recorded under a microscope.

Statistical analysis

GraphPad Prism version 8.0 (GraphPad Software Inc., USA) was used for statistical analyses. The data of at least three independent experiments were expressed as mean \pm standard deviation. The differences between groups were analyzed using Student's *t*-test. Statistically, significance was set at $P < 0.05$. Pearson coefficients > 0.4 were considered significant. We used R-version 3.6.3 for all the R package statistical analyses.

Abbreviations

TACR2: Tachykinin receptor 2; PSA: Prostate-specific antigen; TCGA: The Cancer Genome Atlas; WGCNA: Weighted gene co-expression network; GO: Gene Ontology; NC: Negative control cell lines; TA-OE: TACR2 overexpression cell lines; CCK-8: Cell Counting Kit-8.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12935-021-02126-0>.

Additional file 1: Table S1. The clinicopathological data of 30 patients with prostate cancer.

Acknowledgements

We gratefully thank the Urology Department at the First Hospital of China Medical University. We want to thank TCGA and GEO for their free use.

Authors' contributions

JW contributed significantly to study design, execution, and acquisition of data; YW performed the data analyses and wrote the manuscript; JB contributed to the conception of the study. All authors read and approved the final manuscript.

Funding

This study was supported by the Natural Science Foundation of Liaoning Province (Grant No. 20180551219).

Availability of data and materials

Publicly available datasets were analyzed. These data can be found in the following locations: the datasets TCGA-PRAD for this study can be found at <http://cancergenome.nih.gov/>. The datasets GSE46602 and for this study can be found at <http://www.ncbi.nlm.nih.gov/geo/>.

Declarations

Ethics approval and consent to participate

The Ethics Committee of the First Hospital of China Medical University approved this study according to the Declaration of Helsinki, and all participants provided written informed consent.

Consent for publication

Not applicable.

Competing interests

The authors declare that there are no conflicts of interest.

Received: 17 May 2021 Accepted: 30 July 2021

Published online: 07 August 2021

References

- Cioni B, Zaalberg A, van Beijnum JR, et al. Androgen receptor signaling in macrophages promotes TREM-1-mediated prostate cancer cell line migration and invasion. *Nat Commun*. 2020;11(1):4498.
- Rouyer M, Oudard S, Joly F, et al. Overall and progression-free survival with cabazitaxel in metastatic castration-resistant prostate cancer in routine clinical practice: the FUJI cohort. *Br J Cancer*. 2019;121(12):1001–8.
- Loiseau A, Boudon J, Oudot A, et al. Titanate nanotubes engineered with gold nanoparticles and docetaxel to enhance radiotherapy on xenografted prostate tumors. *Cancers*. 2019;11(12):1962.
- Ferrer-Batallé M, Llop E, Ramírez M, et al. Comparative study of blood-based biomarkers, α 2,3-sialic acid PSA and PHI, for high-risk prostate cancer detection. *Int J Mol Sci*. 2017;18(4):845.
- Totten SM, Adusumilli R, Kullolli M, et al. Multi-lectin affinity chromatography and quantitative proteomic analysis reveal differential glycoform levels between prostate cancer and benign prostatic hyperplasia sera. *Sci Rep*. 2018;8(1):6509.
- Cucchiara V, Cooperberg MR, Dall'Era M, et al. Genomic markers in prostate cancer decision making. *Eur Urol*. 2018;73(4):572–82.
- Wang Y, Lin J, Yan K, Wang J. Identification of a robust five-gene risk model in prostate cancer: a robust likelihood-based survival analysis. *Int J Genom*. 2020;2020:1097602.

8. Xiang D, Xiao J, Fu L, et al. DNA methylation of the Tacr2 gene in a CUMS model of depression. *Behav Brain Res*. 2019;365:103–9.
9. Mao YL, Shen CL, Zhou T, et al. Ablation of Tacr2 in mice leads to gastric emptying disturbance. *Neurogastroenterol Motil*. 2017;29(11):e13117.
10. Steinhoff MS, von Mentzer B, Geppetti P, Pothoulakis C, Bunnett NW. Tachykinins and their receptors: contributions to physiological control and the mechanisms of disease. *Physiol Rev*. 2014;94(1):265–301.
11. Palma C. Tachykinins and their receptors in human malignancies. *Curr Drug Targets*. 2006;7(8):1043–52.
12. Nizam E, Erin N. Differential consequences of neurokinin receptor 1 and 2 antagonists in metastatic breast carcinoma cells; effects independent of substance P. *Biomed Pharmacother*. 2018;108:263–70.
13. Fang W, Fu C, Chen X, et al. Neurokinin-2 receptor polymorphism predicts lymph node metastasis in colorectal cancer patients. *Oncol Lett*. 2015;9(5):2003–6.
14. Wang Z, Shan W, Li H, et al. The PACAP-derived peptide MPAPO facilitates corneal wound healing by promoting corneal epithelial cell proliferation and trigeminal ganglion cell axon regeneration. *Int J Biol Sci*. 2019;15(12):2676–91.
15. Li Y, Gong H, Ding J, et al. Inhibition of GSK3 represses the expression of retinoic acid synthetic enzyme ALDH1A2 via Wnt/ β -catenin signaling in WtT49 cells. *Front Cell Dev Biol*. 2020;8:94.
16. Cheng X, Xu X, Chen D, Zhao F, Wang W. Therapeutic potential of targeting the Wnt/ β -catenin signaling pathway in colorectal cancer. *Biomed Pharmacother*. 2019;110:473–81.
17. Krishnamurthy N, Kurzock R. Targeting the Wnt/ β -catenin pathway in cancer: update on effectors and inhibitors. *Cancer Treat Rev*. 2018;62:50–60.
18. Tian T, Guo T, Zhen W, Zou J, Li F. BET degrader inhibits tumor progression and stem-like cell growth via Wnt/ β -catenin signaling repression in glioma cells. *Cell Death Dis*. 2020;11(10):900.
19. Arend RC, Londoño-Joshi AI, Straughn JM Jr, Buchsbaum DJ. The Wnt/ β -catenin pathway in ovarian cancer: a review. *Gynecol Oncol*. 2013;131(3):772–9.
20. Ren L, Chen H, Song J, et al. MiR-454-3p-mediated Wnt/ β -catenin signaling antagonists suppression promotes breast cancer metastasis. *Theranostics*. 2019;9(2):449–65.
21. Zhao Y, Zhu H, Yang Y, et al. AQP1 suppression by ATF4 triggers trabecular meshwork tissue remodeling in ET-1-induced POAG. *J Cell Mol Med*. 2020;24(6):3469–80.
22. Silva AK, Yi H, Hayes SH, Seigel GM, Hackam AS. Lithium chloride regulates the proliferation of stem-like cells in retinoblastoma cell lines: a potential role for the canonical Wnt signaling pathway. *Mol Vis*. 2010;16:36–45.
23. Culp MB, Soerjomataram I, Efstathiou JA, Bray F, Jemal A. Recent global patterns in prostate cancer incidence and mortality rates. *Eur Urol*. 2020;77(1):38–52.
24. Maolake A, Izumi K, Natsagdorj A, et al. Tumor necrosis factor- α induces prostate cancer cell migration in lymphatic metastasis through CCR7 upregulation. *Cancer Sci*. 2018;109(5):1524–31.
25. Saini S. PSA and beyond: alternative prostate cancer biomarkers. *Cell Oncol*. 2016;39(2):97–106.
26. Khorasani S, Boroumand N, Lavi Arab F, Hashemy SI. The immunomodulatory effects of tachykinins and their receptors. *J Cell Biochem*. 2020;121(5–6):3031–41.
27. Green DP, Limjunyawong N, Gour N, Pundir P, Dong X. A mast-cell-specific receptor mediates neurogenic inflammation and pain. *Neuron*. 2019;101(3):412–420.e3.
28. Hempel Sullivan H, Maynard JP, Heaphy CM, et al. Differential mast cell phenotypes in benign versus cancer tissues and prostate cancer oncologic outcomes. *J Pathol*. 2020;253(4):415–26.
29. Xu W, Qian J, Zeng F, et al. Protein kinase Ds promote tumor angiogenesis through mast cell recruitment and expression of angiogenic factors in prostate cancer microenvironment. *J Exp Clin Cancer Res*. 2019;38(1):114.
30. Hempel Sullivan H, Heaphy CM, Kulac I, et al. High extratumoral mast cell counts are associated with a higher risk of adverse prostate cancer outcomes. *Cancer Epidemiol Biomark Prev*. 2020;29(3):668–75.
31. Xue Y, Tong L, LiuAnwei Liu F, et al. Tumor-infiltrating M2 macrophages driven by specific genomic alterations are associated with prognosis in bladder cancer. *Oncol Rep*. 2019;42(2):581–94.
32. Yamaguchi T, Fushida S, Yamamoto Y, et al. Tumor-associated macrophages of the M2 phenotype contribute to progression in gastric cancer with peritoneal dissemination. *Gastric Cancer*. 2016;19(4):1052–65.
33. Lan J, Sun L, Xu F, et al. M2 macrophage-derived exosomes promote cell migration and invasion in colon cancer. *Cancer Res*. 2019;79(1):146–58.
34. Wang Z, Jensen MA, Zenklusen JC. A practical guide to the cancer genome atlas (TCGA). *Methods Mol Biol*. 2016;1418:111–41.
35. Mortensen MM, Høyer S, Lynnerup AS, et al. Expression profiling of prostate cancer tissue delineates genes associated with recurrence after prostatectomy. *Sci Rep*. 2015;5:16018.
36. Subramanian A, Tamayo P, Mootha VK, et al. Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci USA*. 2005;102(43):15545–50.
37. Ashburner M, Ball CA, Blake JA, et al. Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat Genet*. 2000;25(1):25–9.
38. Langfelder P, Horvath S. WGCNA: an R package for weighted correlation network analysis. *BMC Bioinform*. 2008;9:559.
39. Bost KL. Tachykinin-mediated modulation of the immune response. *Front Biosci*. 2004;9:3331–2.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

