

Author disclosures are available with the text of this letter at www.atsjournals.org.

Francesca Innocenti, M.D.*
Azienda Ospedaliero-Universitaria Careggi
Firenze, Italy

Vittorio Palmieri, M.D.
Ospedale dei Colli Monaldi-Cotugno-CTO
Naples, Italy

Irene Tassinari, M.D.
Elisa Capretti, M.D.
Anna De Paris, M.D.
Adriana Gianni, M.D.
Anna Marchesini, M.D.
Michele Montuori, M.D.
Riccardo Pini, M.D.
Azienda Ospedaliero-Universitaria Careggi
Firenze, Italy

ORCID ID: 0000-0002-7805-7619 (F.I.).

*Corresponding author (e-mail: innocenti.fra66@gmail.com).

References

- Rhodes A, Evans LE, Alhazzani W, Levy MM, Antonelli M, Ferrer R, et al. Surviving Sepsis Campaign: International guidelines for management of sepsis and septic shock: 2016. *Intensive Care Med* 2017;43:304–377.
- Stolk RF, van der Pasch E, Naumann F, Schouwstra J, Bressers S, van Herwaarden AE, et al. Norepinephrine dysregulates the immune response and compromises host defense during sepsis. *Am J Respir Crit Care Med* 2020;202:830–842.
- Uhel F, van der Poll T. Norepinephrine in septic shock: a mixed blessing. *Am J Respir Crit Care Med* 2020;202:788–789.
- Capretti E, De Paris A, Gianni A, Marchesini A, D'Argenzio F, Montuori M, et al. Variation of the myocardial contractility in response to the administration of norepinephrine during septic shock. EUSEM Annual Meeting. 2020, Virtual.
- Palmieri V, Innocenti F, Guzzo A, Guerrini E, Vignaroli D, Pini R. Left ventricular systolic longitudinal function as predictor of outcome in patients with sepsis. *Circ Cardiovasc Imaging* 2015;8:e003865, discussion e003865.
- Innocenti F, Palmieri V, Stefanone VT, Donnini C, D'Argenzio F, Cigana M, et al. Epidemiology of right ventricular systolic dysfunction in patients with sepsis and septic shock in the emergency department. *Intern Emerg Med* 2020;15:1281–1289.
- Innocenti F, Palmieri V, Guzzo A, Stefanone VT, Donnini C, Pini R. SOFA score and left ventricular systolic function as predictors of short-term outcome in patients with sepsis. *Intern Emerg Med* 2018;13:51–58.
- Rudski LG, Lai WW, Afilalo J, Hua L, Handschumacher MD, Chandrasekaran K, et al. Guidelines for the echocardiographic assessment of the right heart in adults: a report from the American Society of Echocardiography endorsed by the European Association of Echocardiography, and the Canadian Society of Echocardiography. *J Am Soc Echocardiogr* 2010;23:685–713, quiz 786–788.
- Sacha GL, Bauer SR, Lat I. Vasoactive agent use in septic shock: beyond first-line recommendations. *Pharmacotherapy* 2019;39:369–381.
- Hernández G, Teboul JL, Bakker J. Norepinephrine in septic shock. *Intensive Care Med* 2019;45:687–689.
- Hamzaoui O, Jozwiak M, Geffriaud T, Sztrymf B, Prat D, Jacobs F, et al. Norepinephrine exerts an inotropic effect during the early phase of human septic shock. *Br J Anaesth* 2018;120:517–524.
- Hamzaoui O, Georger JF, Monnet X, Ksouri H, Maizel J, Richard C, et al. Early administration of norepinephrine increases cardiac preload and cardiac output in septic patients with life-threatening hypotension. *Crit Care* 2010;14:R142.
- Thooft A, Favory R, Salgado DR, Taccone FS, Donadello K, De Backer D, et al. Effects of changes in arterial pressure on organ perfusion during septic shock. *Crit Care* 2011;15:R222.

Copyright © 2021 by the American Thoracic Society



Adverse Health Effects in People with and without Preexisting Respiratory Conditions during Bushfire Smoke Exposure in the 2019/2020 Australian Summer

To the Editor:

Australia had unprecedented bushfires affecting multiple states in the summer of 2019/2020. Prolonged exposure to bushfire smoke over December 2019 and January 2020 is estimated to have resulted in over 400 excess deaths and over 3,000 additional hospitalizations (1). Some evidence has suggested that people with asthma are at higher risk for adverse health effects after bushfire or wildfire smoke exposure (2–5), although others suggest individuals with asthma are more inclined to take protective measures (6). We aimed to understand the impact of the 2019/2020 bushfire season on the health and behavior of people with and without preexisting respiratory conditions in affected Australian states.

A cross-sectional study was conducted to compare health effects of the 2019/2020 bushfires in people with and without respiratory conditions. Respiratory conditions were defined as self-reported asthma, emphysema, chronic bronchitis, chronic obstructive pulmonary disease, bronchiectasis, and any other chronic lung conditions. Participants 18 years or over with and without respiratory conditions were recruited from postcodes affected by bushfire smoke in the summer of 2019/2020 in six states (New South Wales, Victoria, South Australia, Tasmania, Australian Capital Territory, and Queensland) in Australia during August 2020. *A priori* power analysis was conducted using large sample approximation in G*Power 3.1.9.7 (7). To detect at least 20% difference in risk of adverse health effects after smoke exposure among people with and without respiratory conditions (i.e., odds ratio of 1.2), the sample size necessary to achieve in a two-sided test with $\alpha = 0.05$ and power of 80% is 961.

A market research company, Dynata, distributed the survey link by email to a randomly selected sample of their panel members in the affected areas in selected states. We aimed to recruit 500 people with preexisting (self-reported) respiratory conditions and 500 without preexisting respiratory conditions, on the basis of responses to the survey. Surveys were deidentified and no identifying information was provided or collected. Eligible participants were asked to provide

Ⓜ This article is open access and distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives License 4.0 (<https://creativecommons.org/licenses/by-nc-nd/4.0/>). For commercial usage and reprints, please contact Diane Gern (dgern@thoracic.org).

Supported by Medical Research Futures Fund Bushfire Impact Research grant APP1201320.

Author Contributions: Conception and design of study: C.R.M. and G.B.M. Drafting of manuscript: C.R.M., P.-Y.N., and G.B.M. Data analysis and interpretation: C.R.M., P.-Y.N., and G.B.M. Input for analysis and interpretation: C.R.M., P.-Y.N., M.T., H.S., A.A.C., S.S., and G.B.M.

Originally Published in Press as DOI: 10.1164/rccm.202012-4471LE on May 11, 2021

Table 1. Differences in Mask Use, Health System Access, and Health Effects during the 2019/2020 Bushfire Season between People with and without Respiratory Conditions

Factor	With Respiratory Conditions [n (%)] (N = 490)	Without Respiratory Conditions [n (%)] (N = 527)	Odds Ratio		
			Estimate	95% CI	P Value
Exposed to bushfire smoke in the last 12 mo	343 (70.0)	337 (63.9)	1.32	1.0–1.7	0.04*
Wore a mask or P2 respirator during fires	98 (20.0)	51 (9.7)	2.3	1.6–3.4	<0.0001*
Experienced breathing difficulties	172 (35.1)	53 (10.1)	5.2	3.5–7.5	<0.0001*
Developed chest infection within 1 wk of BFSE	65 (13.3)	21 (4.0)	3.7	2.2–6.2	<0.0001*
Urgent visit to GP for breathing difficulties	51 (10.4)	4 (0.8)	15.2	5.5–42.2	<0.0001*
Visit to ED for breathing difficulties	21 (4.3)	6 (1.1)	3.9	1.6–9.7	0.0040*
Admitted to hospital for breathing difficulties	12 (2.4)	3 (0.6)	4.4	1.2–15.8	0.0200*
Increased use of reliever inhalers	207 (42.2)	13 (2.5)	28.9	15.6–53.7	<0.0001*
Increased use of controller or controller/reliever combination inhalers	149 (30.4)	11 (2.1)	20.5	10.2–41.1	<0.0001*
Increased use of corticosteroids	39 (8.0)	6 (1.1)	7.5	3.2–17.6	<0.0001*
Prescribed antibiotics	30 (6.1)	6 (1.1)	5.7	2.3–13.9	0.0002*
Any adverse health effect	294 (60.0)	80 (15.2)	8.4	6.1–11.5	<0.0001*

Definition of abbreviations: BFSE = bushfire smoke exposure; CI = confidence interval; ED = emergency department; GP = general practitioner. *Significant at $\alpha = 0.05$.

informed consent before completing the survey. The survey was launched on August 3, 2020, and closed on August 21, 2020. The study was approved by the Human Research Ethics Committee of the University of New South Wales (project number HC200477).

The online survey collected data on sociodemographic characteristics, preexisting conditions, self-reported bushfire smoke exposure (BFSE) within the last 12 months, use of masks, outdoor air avoidance, and other nonpharmaceutical interventions undertaken. An adverse health effect is defined as any of the following occurring between December 2019 and February 2020: increased use of reliever, controller, or combination inhalers; increased use of corticosteroids; receiving antibiotics; an urgent visit to a general practitioner; admission to an emergency department (ED) or a hospital for breathing difficulties; or developing a chest infection, all within a week of BFSE. State of residence was determined on the basis of reported postcode. Age was recorded as a continuous variable and converted to a binary variable (<65 yr old) for analysis. Similarly, smoking status and household income were recorded as categorical and converted to binary variables (“current smoker” and “income <\$18,200 per annum”). All other factors in regression analysis are binary variables.

We calculated cluster (postcode)-level sampling weights using data for age group, sex, area socioeconomic status, remoteness, and the prevalence of chronic disease using data from the 2017–2018 Australian National Health Survey (8). Each participant was assigned a sampling weight on the basis of their unique combination of attributes, calculated as the ratio of the number of participants with that combination of attributes in the surveyed sample to the expected number of people with the same combination of attributes in the Australian Health Survey. These factors were selected for adjustment on the basis of previous evidence, which shows that older adults and people of low socioeconomic status may be at an increased risk of mortality and ED/hospital admission after short-term exposure to bushfire smoke (9, 10).

We used descriptive statistics to summarize sociodemographic characteristics, preexisting conditions, avoidance behaviors, health effects, and smoke exposure. Logistic regression (with and without sampling weights) was performed to quantify the association of potential risk factors with respiratory conditions and with the occurrence of adverse health effects. Stratified analysis was conducted for subpopulations with and without preexisting respiratory conditions for the association between BFSE and adverse health

Table 2. Logistic Regression Analysis for Predictors of Adverse Health Effects during the 2019/2020 Bushfire Season, Adjusted for the Australian Population

Factor	Reporting Adverse Effects [n (%)] (N = 374)	Reporting No Adverse Effects [n (%)] (N = 643)	Odds Ratio (Unweighted)			Adjusted Odds Ratio (Weighted) [*]		
			Estimate	95% CI	P Value	Estimate	95% CI	P Value
Residing in NSW	228 (61.0)	267 (41.5)	2.20	1.7–2.9	<0.0001 [†]	1.27	0.6–2.6	0.51
Age <65	278 (74.3)	454 (70.6)	1.21	0.9–1.6	0.17	2.88	1.2–7.2	0.02 [†]
Annual household income <\$18,200	17 (4.5)	32 (5.0)	0.91	0.5–1.7	0.75	0.68	0.3–1.8	0.45
Current smoker	88 (23.5)	129 (20.1)	1.23	0.9–1.7	0.24	2.37	0.9–6.2	0.08 [†]
History of cardiovascular disorders	129 (34.5)	208 (32.3)	1.10	0.8–1.5	0.51	1.53	0.7–3.4	0.29
Avoided outdoor air during the bushfire	209 (55.9)	262 (40.7)	1.84	1.4–2.4	<0.0001 [†]	0.74	0.4–1.4	0.36
Reduced physical activity during the bushfire	186 (49.7)	179 (27.8)	2.56	1.9–3.5	<0.0001 [†]	2.11	1.0–4.7	0.07 [†]
Used masks or face covering during the bushfire	79 (21.1)	17 (2.6)	9.86	5.7–16.9	<0.0001 [†]	34.77	13.0–93.4	<0.0001 [†]
Used P2 respirators during the bushfire	56 (15.0)	20 (3.1)	5.49	3.2–9.5	<0.0001 [†]	7.28	2.5–21.5	0.0004 [†]
Exposed to bushfire smoke in the last 12 mo	305 (81.6)	375 (58.3)	3.16	2.3–4.3	<0.0001 [†]	2.05	0.7–6.1	0.19
Preexisting respiratory conditions	294 (78.6)	196 (30.5)	8.38	6.1–11.5	<0.0001 [†]	34.59	11.3–106.3	<0.0001 [†]
Participants with ≥1 preexisting respiratory conditions								
Exposed to bushfire smoke in the last 12 mo	238 (81.0) [‡]	105 (41.5) [§]	3.68	2.5–5.4	<0.0001 [†]	4.12	1.8–9.6	0.001 [†]
Participants without preexisting respiratory conditions								
Exposed to bushfire smoke in the last 12 mo	67 (83.4)	270 (60.4) [¶]	3.38	1.7–6.6	0.0004 [†]	0.93	0.3–3.5	0.92

Definition of abbreviations: CI = confidence interval; NSW = New South Wales.

*All analyses are clustered at postcode level. Adjusted odds ratios are calculated using the weighted sample, standardized to the 2017–2018 Australian National Health Survey data for age group, sex, area socioeconomic status, remoteness, and the prevalence of chronic diseases.

[†]Significant at $\alpha = 0.05$.

[‡]N = 294.

[§]N = 196.

^{||}N = 80.

[¶]N = 447.

effects. Interaction between BSFE and preexisting respiratory conditions was assessed using SAS's joint test. Statistical significance was defined at α -level of 0.05. Analysis was completed using SAS software (version 9.4; SAS Institute).

Of 1,017 participants, 490 (48.2%) had preexisting respiratory conditions, 680 (66.8%) were exposed to bushfire smoke during the preceding summer, and 553 (54.4%) reported a bushfire within 50 km of their home. The median age was 54 years (range 18–89 yr) and sex distribution was close to 50%. Nonrespiratory medical conditions were present in 686 (67.5%) participants, most commonly hypertension ($n = 284$, 27.9%), allergies ($n = 274$, 26.9%), diabetes ($n = 132$, 13.0%), and dermatitis ($n = 124$, 12.2%).

Participants reported that, after bushfire smoke exposure, they experienced "chest infection" ($n = 86$, 8.5%), breathing difficulties ($n = 225$, 22.1%), urgent primary care visit for breathing difficulties ($n = 55$, 5.4%), and ED visit or admission to hospital ($n = 15$, 1.5%). Increased use of oral corticosteroids occurred in 45 (4.4%), and 36 (3.5%) were prescribed antibiotics; 221 (21.7%) reported increased use of reliever medication and 161 (15.8%) stated that they increased their use of controller or combined controller/reliever medication.

Table 1 shows that participants with respiratory conditions were significantly more likely to experience adverse health effects, wear a mask or P2 respirator during fires, or be prescribed antibiotics or oral corticosteroids. People older than 65 years reported fewer visits to health facilities than people younger than 65 years (6.0% vs. 7.4%) and were more likely to report outdoor air avoidance (50.9% vs. 44.6%).

Table 2 shows the predictors of adverse health effects. After adjusting for sociodemographic characteristics of the Australian population, the effect of BFSE on the risk of adverse effects was greater in people with preexisting respiratory disease than in people without preexisting respiratory disease (P for interaction = 0.0427). Younger age (<65 yr) was associated with higher probability of adverse effects. Cardiovascular disease was not a risk factor. Adverse health effects were more common among people who use respirators and masks. This may be attributable to confounding by indication, in that those with preexisting conditions, or who were experiencing symptoms due to the smoke, were more likely to use respiratory protection. In stratified analysis among people without preexisting respiratory conditions, the rate of adverse health effects was higher in people exposed to bushfire smoke (19.9%) than in people not exposed (6.8%) ($P = 0.0001$). Among

people with preexisting respiratory conditions, a similar trend was observed (69.4% for BFSE and 38.1% for nonexposure, $P < 0.0001$).

The primary limitation of our study is its cross-sectional design, which did not allow us to measure temporal changes in outcome. Participants with preexisting respiratory conditions are more likely to remember adverse events than participants without these conditions, which may lead to recall bias and overestimation of the risk of adverse events. Future studies would benefit from a cohort design, which would overcome these limitations. Nonetheless, the chance of participants without respiratory conditions not recalling these events is partially mitigated by the fact that the 2019–2020 bushfire was a major natural disaster and the adverse effects surveyed were relatively uncommon.

In conclusion, smoke exposure was significantly associated with adverse health effects during the Australian bushfire season in 2019/2020 not only among people with respiratory conditions but also among healthy people. Surprisingly, older age (65 yr and above) was associated with a significantly lower risk of adverse health effects. Our data suggest older people may be more cautious and less mobile in outdoor settings than younger people during bushfires. Younger people (<65 yr) may benefit from public health messaging about outdoor air avoidance and respirator use. Adverse health effects due to smoke exposure also impacted people without respiratory conditions. However, people with respiratory conditions are at greater risk and should be a priority for mitigation measures into the future. ■

Author disclosures are available with the text of this letter at www.atsjournals.org.

C. Raina MacIntyre, Ph.D.*
The Kirby Institute at University of New South Wales
Sydney, New South Wales, Australia
and

Arizona State University College of Public Affairs
and Community Solutions
Phoenix, Arizona

Phi-Yen Nguyen, M.P.H.
The Kirby Institute at University of New South Wales
Sydney, New South Wales, Australia
and

University of New South Wales School of Population Health
Sydney, New South Wales, Australia

Mallory Trent, M.S.P.H.
The Kirby Institute at University of New South Wales
Sydney, New South Wales, Australia

Holly Seale, Ph.D.
Abrar Ahmad Chughtai, Ph.D.
University of New South Wales School of Population Health
Sydney, New South Wales, Australia

Smita Shah, M.B. Ch.B., M.C.H.
Western Sydney Local Health District
Prevention Education and Research Unit
Westmead, New South Wales, Australia
and
The University of Sydney Faculty of Medicine and Health
Sydney, New South Wales, Australia

Guy B. Marks, Ph.D.
Woolcock Institute of Medical Research
Glebe, New South Wales, Australia

and

University of New South Wales South Western Sydney Clinical School
Liverpool, New South Wales, Australia

ORCID IDs: 0000-0002-3060-0555 (C.R.M.); 0000-0002-0476-3385 (P.-Y.N.); 0000-0001-5264-2660 (M.T.); 0000-0002-1877-5395 (H.S.); 0000-0003-4203-7891 (A.A.C.); 0000-0002-8976-8053 (G.B.M.).

*Corresponding author (e-mail: rainam@protonmail.com).

References

- Borchers Arriagada N, Palmer AJ, Bowman DM, Morgan GG, Jalaludin BB, Johnston FH. Unprecedented smoke-related health burden associated with the 2019-20 bushfires in eastern Australia. *Med J Aust* 2020;213:282–283.
- Kiser D, Metcalf WJ, Elhanan G, Schnieder B, Schlauch K, Joros A, *et al*. Particulate matter and emergency visits for asthma: a time-series study of their association in the presence and absence of wildfire smoke in Reno, Nevada, 2013-2018. *Environ Health* 2020;19:92.
- Gan RW, Liu J, Ford B, O'Dell K, Vaidyanathan A, Wilson A, *et al*. The association between wildfire smoke exposure and asthma-specific medical care utilization in Oregon during the 2013 wildfire season. *J Expo Sci Environ Epidemiol* 2020;30:618–628.
- Borchers Arriagada N, Horsley JA, Palmer AJ, Morgan GG, Tham R, Johnston FH. Association between fire smoke fine particulate matter and asthma-related outcomes: Systematic review and meta-analysis. *Environ Res* 2019;179:108777.
- Martin KL, Hanigan IC, Morgan GG, Henderson SB, Johnston FH. Air pollution from bushfires and their association with hospital admissions in Sydney, Newcastle and Wollongong, Australia 1994-2007. *Aust N Z J Public Health* 2013;37:238–243.
- Künzli N, Avol E, Wu J, Gauderman WJ, Rappaport E, Millstein J, *et al*. Health effects of the 2003 Southern California wildfires on children. *Am J Respir Crit Care Med* 2006;174:1221–1228.
- Faul F, Erdfelder E, Buchner A, Lang A-G. Statistical power analyses using G*Power 3.1: tests for correlation and regression analyses. *Behav Res Methods* 2009;41:1149–1160.
- Kalisch DW. National Health Survey: First Results (2017-18 Financial Year). Canberra: Australian Bureau of Statistics. 2015.
- US EPA. Integrated Science Assessment (ISA) for Particulate Matter (Final Report, Dec 2009). US Environmental Protection Agency. Washington, DC, EPA/600/R-08/139F, 2009.
- Rappold AG, Cascio WE, Kilaru VJ, Stone SL, Neas LM, Devlin RB, *et al*. Cardio-respiratory outcomes associated with exposure to wildfire smoke are modified by measures of community health. *Environ Health* 2012;11:71.

Copyright © 2021 by the American Thoracic Society



Optimism with Caution: Elexacaftor–Tezacaftor–Ivacaftor in Patients with Advanced Pulmonary Disease

To the Editor:

We read with great interest the article by Burgel and colleagues, which described significant and rapid improvements in outcomes of patients with severe cystic fibrosis (CF)-related lung disease after commencing

Ⓐ This article is open access and distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives License 4.0 (<https://creativecommons.org/licenses/by-nc-nd/4.0/>). For commercial usage and reprints, please contact Diane Gern (dgern@thoracic.org).

Originally Published in Press as DOI: 10.1164/rccm.202103-0682LE on May 5, 2021