

SHORT COMMUNICATION

Exploration of anthelmintic activity of *Cassia* spp. extracts on gastrointestinal nematodes of sheep

Sri Wahyuni^{1,2}, Sunarso Sunarso¹, Bambang Waluyo Hadi Eko Prasetyono¹, Fadjar Satrija³

¹Faculty of Animal and Agricultural Sciences, University of Diponegoro (UNDIP), Jalan Prof. Soedarto, UNDIP Tembalang Campus, Semarang 50275, Indonesia

²Faculty of Animal Husbandry, University of Darul Ulum Islamic Center Sudirman (UNDARIS), Jalan Tentara Pelajar 13 Ungaran 50514, Indonesia

³Faculty of Veterinary Medicine, IPB University, Jalan Agatis IPB Dramaga Campus, Bogor 16680, Indonesia

ABSTRACT

Objective: This study aimed to explore the phytochemical constituents and anthelmintic activities of four *Cassia* spp. leaves against *Haemonchus contortus*.

Materials and Methods: The extracts were prepared from four species of *Cassia* spp. (*C. siamea*, *C. fistula*, *C. surattensis*, and *C. spectabilis*). Phytochemical screening of the extract was done based on the Harborne method. Evaluation of the anthelmintic activities against *H. contortus* was done *in vitro* using infective larvae (L₃) migration inhibition assay (LMIA). Measurement of larvae migrating was conducted through a nylon filter with a pore size of 20 µm. The doses of *Cassia* spp. extract implemented were 25, 50, 100, and 200 mg/ml.

Results: Tannins, alkaloids, phenol hydroquinone, flavonoids, steroids, triterpenoids, and saponins were present in all the extracts, whereas alkaloids were absent in *C. fistula*. No triterpenoids were found in *C. surattensis* and *C. spectabilis*. Movement of *H. contortus* larvae was significantly inhibited after exposure to *Cassia* extracts at various dosage levels ($p < 0.05$). The test results using LMIA on L₃ *H. contortus* showed the lowest inhibition in the negative control. Among the species of *Cassia*, the *C. surattensis* (at 200 mg/ml) showed the highest ($p < 0.05$) inhibition level on the larvae. The latter result corresponded to the effect of albendazole.

Conclusion: Compared to other *Cassia* spp., *C. surattensis* exhibited the highest inhibition against L₃ *H. contortus*. However, the inhibition effect of *C. surattensis* was still lower as compared to albendazole.

ARTICLE HISTORY

Received December 03, 2018

Revised December 31, 2018

Accepted January 12, 2019

Published May 11, 2019

KEYWORDS

Anthelmintic effect; *C. siamea*; *C. fistula*; *C. surattensis*; *C. spectabilis*; *H. contortus*



This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 Licence (<http://creativecommons.org/licenses/by/4.0>)

Introduction

The gastrointestinal nematode is one of the animal health risks related to the use of pasture for small as well as large ruminant productions. The most dominant gastrointestinal nematodes in small ruminants in the tropics, including Indonesia, are *Haemonchus contortus* [1,2]. Control of gastrointestinal nematodes of ruminants mainly relies on the use of synthetic chemical anthelmintics. However, internal parasitic disease control using synthetic chemicals has raised big concern, such as chemical residues in the animal products, especially when the animals on the extensive livestock management system. In this respect,

any alternatives substitute to the role of synthetic chemicals is, therefore, of important to be developed. Among the alternatives, tannin-rich plants seemed to be good candidates to control the internal parasitic disease in animals particularly the small ruminants on the pasture rotation system [3].

The research on anthelmintic effect of the plants containing tannin to nematodes has been carried out [4,5]. *Cassia* spp. is the plants with high condensed tannin content [6]. Kundu et al. [5] showed the broad spectrum anthelmintic effects of *C. angustifolia*, *C. alata*, and *C. occidentalis* crude ethanol extracts against helminth parasite

Correspondence Sri Wahyuni ✉ swahyuniundaris@gmail.com 📧 Faculty of Animal Husbandry, University of Darul Ulum Islamic Center Sudirman (UNDARIS), Jalan Tentara Pelajar 13 Ungaran 50514, Indonesia.

How to cite: Wahyuni S, Sunarso S, Prasetyono BWHE, Satrija F. Exploration of anthelmintic activity of *Cassia* spp. extracts on gastrointestinal nematodes of sheep. J Adv Vet Anim Res 2019; 6(2):236–40.

of domestic fowls (*Heterakis gallinarum*, *Catatropis* sp., and *Raillietina tetragona*) *in vitro* at concentration levels of 10–40 mg/ml. Likewise, *Cassia tora* leaves extracts showed anthelmintic activity against *Pherethima posthuma* [7].

In respect, particularly to *C. fistula* Linn, the methanolic extract from pods, pulp, and seeds from such plant (at the concentration 100 mg/ml) showed an anthelmintic activity against *P. posthuma* [8,9]. Other *Cassia* spp., such as *C. siamea*, *C. surattensis*, and *C. spectabilis*, had reported having antibacterial activity, antibiofilm, antifungal, and antioxidant [10–13]. However, anthelmintic activities above mentioned *Cassia* spp. have never been documented. Anthelmintic activity can be affected by phytochemical compounds. Some phytochemical compounds, such as tannin, anthraquinone glycoside, naphthopyrone glycoside, phenolic compounds, flavonoids, and many others, had isolated from *Cassia* plants and suggested to the biological activity of the plants [14]. Hence, this study aimed to explore the phytochemical compound and anthelmintic activities against *H. contortus* of *C. siamea*, *C. fistula*, *C. surattensis*, and *C. spectabilis* leaves extracts.

Materials and Methods

Ethical approval

All the procedures in the study accordance were done after the approval from the Ethics Committee for Animal Use at the IPB University (approval number 44-2017 IPB).

Plants material

The plants were collected in the area of Diponegoro University, Semarang, Central Java, Indonesia. The collected plants were identified by Herbarium Bogoriense of Indonesian Institute of Sciences as *C. siamea*, *C. fistula*, *C. surattensis*, and *C. spectabilis* by the reference number 335/IPH.1.01/H.07/II/2016.

Extracts preparation

The leaves of *Cassia* spp. were separately picked and sorted to remove undesired plant parts. Drying of leaves in the room temperature was done for 7–10 days and then milled to produce homogenous flour particles. The *Cassia* leaf flours were then turned into extracts using maceration technique with n-hexane and ethanol solvent. *Cassia* spp. leaf flour (1,000 gm) was initially incubated in n-hexane (5,000 ml) and later in ethanol, 96% (5,000 ml) in the room temperature for 72 h. After incubation, extracting the extract was carried out with Whatman No. 1 filter paper. Filtering results were collected and evaporated using a vacuum rotary evaporator at 50 rpm, 40°C–50°C until the extract was obtained. Storage of the extracts before analysis was used at 4°C analysis.

Phytochemical screening

Phytochemical screening of *Cassia* spp. extract using the [15] method was done to detect tannins, saponins, alkaloids, flavonoids, steroids, triterpenoids, and hydroquinone phenolics.

Preparation *H. contortus* infective larvae (L₃)

Haemonchus contortus for this study was cultured from stool of a *H. contortus* purely infected sheep donor. Feces of the donor sheep was collected every morning using an apron. The feces were then mixed with vermiculite to obtain moist and airy medium *H. contortus* egg hatching. Incubation of the mixture was carried out in the room temperature for 7–8 days so that *H. contortus* eggs can hatch and develop into infective larvae (L₃). The L₃ were then harvested by a modification of the Baermann method [16]. In order to remove fecal impurities, the infective larvae harvested are screened. The collected larvae were stored at 4°C before being used for the assay.

Infective larvae assay

An *in vitro* larval migration inhibition assay (LMIA) was performed to determine anthelmintic effect of the four *Cassia* spp. Extracts according to Rabel et al. [17] method. The experiment used factorial (4 × 6) design with of *Cassia* spp. as the first factor and dose levels of the extracts as the second factor [18]. The LMIA started with the mixing of 100 L₃ (s) in each *Cassia* extracts that were prepared into 1.5-ml solution with the dosage levels of 25, 50, 100, and 200 mg/ml diluted in phosphate buffer saline (PBS).

These mixtures were then incubated at the room temperature for 3 h. After incubation, the mixtures were then washed thrice by using PBS, centrifuged at 3,500 rpm for 5 min, before the supernatant was removed. The remaining sediments (the larvae) were then sieved using 20-µm grid sieve mounted on PBS-filled microplate well. Number of the larvae migrated through the sieve to the microplate well was then counted by observing the process under a stereoscopic microscope at 20× magnification. The percentage of larval migration inhibition was then calculated using the formula of $[(A - B)/A] \times 100$ where *A* is the number of larvae that were prepared (100) and *B* is the number of larvae that migrated through the sieve at each treatment [17].

Statistical analysis

Results of the phytochemical analysis were presented descriptively, while the results of the LMIA were analyzed based on analysis of variance using SAS v 9.0 [19]. The significant effect of LMIA was further tested with DMRT. The difference in treatment was stated to be significant at $p < 0.05$.

Results

Percentage yield of extract

The highest percentage of *Cassia* spp. extract yield was found in *C. spectabilis*, followed by *C. surattensis*, *C. siamea*, and *C. fistula*. *Cassia surattensis* extract had the highest tannin content followed by the extracts *C. siamea*, *C. fistula*, and *C. spectabilis*, respectively (Table 1).

Phytochemical screening of four *Cassia* spp. extract

Phytochemical compounds identified in the four *Cassia* spp. extracts are shown in Table 2. Tannins, alkaloids, phenol hydroquinone, flavonoids, steroids, triterpenoids, and saponins were present in all the extracts. Alkaloids were absent in *C. fistula*, whereas no triterpenoids was found in *C. surattensis* and *C. spectabilis* extracts.

Larval migration assay of extracts *Cassia* spp.

Table 3 showed that all the *Cassia* spp. extracts at various doses significantly affected the percentage of inhibited L₃ *H. contortus* ($p < 0.05$). The *H. contortus* larval migration inhibition was lowest in the negative control (PBS). Among the various dose and species of *Cassia*, the *C. surattensis* at the dose level 200 mg/ml resulted in the highest inhibited of L₃ *H. contortus* ($p < 0.05$). The later result corresponded to the effect of albendazole.

Discussion

Our data showed that all *Cassia* spp. used in the present study contained most of the phytochemical compounds, i.e., alkaloids, phenol hydroquinone, flavonoid, steroid, triterpenoid, tannin, and saponin. Unlike *C. siamea*, *C. fistula* contained no alkaloids, while *C. surattensis* and *C. spectabilis* contained no triterpenoids. Corresponding to our result, Mohammed et al. [20] found that *C. siamea* contained anthraquinones, alkaloids, tannins, saponins, flavonoids, polyphenols, and glycosides. Likewise, Kamagaté et al. [21] reported that the ethanol extract of *C. siamea* leaf contained flavonoid (D-pinitol, luteolin), dihydronaphtalen one [(4-trans)-acetyl-3,6,8-trihydroxy-3-methyldihydronaphtalenone], triterpenoid (lupeol), and [4-(cis)-acetyl-3,6,8-trihydroxy-3-methyldihydronaphtalenone].

Our finding showed that *C. fistula* contains phenol hydroquinone, flavonoids, steroids, triterpenoids, tannins, and saponins. However, it contained no alkaloid. In contrast, Panda et al. [22] found alkaloids, flavonoids, tannins and phenolic compound, glycosides, protein, amino acids, saponins, and triterpenoids. In Another study revealed that *C. fistula* leaves mainly contain oxalic acid, tannin, oxy anthraquinone, and anthraquinone derivative [23]. In general, the composition and concentration of secondary

metabolites in particular plant is influenced by several factors, including genetic factors, climate, soil, harvest time, and solar radiation [24].

Compared to PBS group, all *Cassia* spp. were capable of increasing the percentage inhibition of *H. contortus*, of the doses, applied. Yet, the values were still lower as compared to the albendazole group. In this respect, *C. surattensis* had anthelmintic activity, especially against *H. contortus*. Several studies used else species of the genus *Cassia*, such as the *C. tora*, *C. auriculata*, *C. angustifolia*, *C. occidentalis*, and *C. alata* which contain active compounds as tannins, flavonoids, saponins, and alkaloids that shown to be anthelmintic [5,7,25]. The anthelmintic activities may be contributed by tannins, alkaloids, flavonoids, steroids, phenol hydroquinone, triterpenoids, and saponins.

Very limited study on the mechanisms of phytochemical compounds of *Cassia* spp. to destruct *H. contortus*. However, tannin has been demonstrated to block through uncoupling the oxidative phosphorylation leading to the death of parasites. Another possible mechanism could be through binding of tannins to free proteins in the gastrointestinal tract of the animal, or glycoprotein on the cuticle of the helminth's body surface causing a paralysis, and thus death [26]. Cuticle and digestive tissue of larvae are significantly damaged due to the presence of tannins. The size of the polymer and the tannin molecule has a relationship in influencing the strength of the

Table 1. Percentage yield and tannin content from all four extracts of *Cassia* spp.

Extract	Percentage yield (W/W)	Tannin (%)
<i>C. siamea</i>	4.49	4.92
<i>C. fistula</i>	2.36	3.98
<i>C. surattensis</i>	5.23	7.82
<i>C. spectabilis</i>	11.37	2.67

Table 2. The phytochemical constituent of extracts of *Cassia* spp.

Phytochemical	Extracts			
	<i>C. siamea</i>	<i>C. fistula</i>	<i>C. surattensis</i>	<i>C. spectabilis</i>
Alkaloids	+	-	+	+
Phenol Hydroquinone	+	+	+	+
Flavonoids	+	+	+	+
Steroids	+	+	+	+
Triterpenoids	+	+	-	-
Tannins	+	+	+	+
Saponins	+	+	+	+

+: present; -: absent.

Table 3. Percentage *L*₃ *H. contortus* inhibited after exposure to 3 h on the species and dose of extract of *Cassia* spp. different*.

Extracts	PBS (negative control)	C-25	C-50	C-100	C-200	Albendazole (positive control)
<i>C. siamea</i>	22.11 ± 2.84 ^e	59.89 ± 1.30 ^f	63.18 ± 1.09 ^{def}	63.34 ± 1.22 ^{def}	65.31 ± 1.25 ^{cd}	82.71 ± 1.19 ^a
<i>C. fistula</i>	24.89 ± 7.32 ^e	59.88 ± 1.54 ^{bc}	62.30 ± 4.42 ^{def}	60.90 ± 3.13 ^f	63.46 ± 1.49 ^{def}	81.82 ± 1.19 ^a
<i>C. surattensis</i>	23.81 ± 0.85 ^e	63.56 ± 2.41 ^{def}	65.90 ± 2.56 ^{cd}	67.99 ± 2.06 ^c	71.62 ± 2.49 ^b	84.94 ± 0.78 ^a
<i>C. spectabilis</i>	22.56 ± 2.64 ^e	59.95 ± 1.25 ^f	61.10 ± 1.71 ^{ef}	62.27 ± 1.71 ^{def}	64.98 ± 1.17 ^{cde}	85.36 ± 0.94 ^a

Values are mean ± SD of six replicates.

^{a-e}Means within a row and column with different superscripts differ ($p < 0.05$).

C- means dose of species *Cassia* extracts (mg/ml).

anthelmintic effect. In addition, there are effects of the structural units of monomeric tannins because gallocatechin and epigallocatechin monomers have significant anthelmintic activity on the contrary monomer catechin and epicatechin [27]. In addition to tannin, other bioactive compounds in *Cassia* spp. may contribute the anthelmintic activity against *H. contortus*. Indeed [28–30], suggested flavonoid, terpenoid, saponin, and alkaloids may exert an anthelmintic effect on *H. contortus*.

Previous studies in laboratory animals have demonstrated the safety of *Cassia* spp. as medicinal plants. Thus, no clinical signs of toxicity and mortality were found in mice receiving single oral doses of 2,000–5,000 mg/kg body weight extracts of *C. spectabilis* [13,31], *C. siamea* [21], and *C. surattensis* [32], respectively. However, long-term administration of *Cassia* spp. extracts may cause reversible hepatotoxicity in rats and mice [21].

Conclusion

Compared to other *Cassia* spp., *C. surattensis* exhibited the highest migration inhibition against *L*₃ *H. contortus*. However, the inhibition effect *C. surattensis* was still lower as compared to albendazole. Owing to the latter fact, it could be inferred that *C. surattensis* may be used to substitute the role albendazole as anthelmintic provided that dose is increased. Hence, the future study is necessary to confirm optimal doses of *C. surattensis* as substitute for albendazole.

Acknowledgments

This research was financed by the Doctoral Dissertation Research Grant of the Ministry of Research, Technology and Higher Education of Republic of Indonesian through Domestic Graduate Scholarship (“BPP-DN”) program. The authors would like to thank Mr. Sulaiman and Department Animal Infectious Diseases, and Veterinary Public Health, Faculty Veterinary Medicine, IPB University, Bogor for providing laboratory facilities during the study.

Conflict of Interests

The authors declare that they have no conflict of interest.

Authors’ contribution

Sri Wahyuni and Fadjar Satrija designed the study, conducted the experiments, analyzed the data, and prepared the article. Sunarso Sunarso and Bambang Waluyo Hadi Eko Prasetyono corrected the article.

References

- [1] Satrija F, Beriajaya. The epidemiology and control of gastrointestinal nematodes of ruminants in Indonesia, with spec I:1 reference of small ruminants in West Java. In: Biological of gastrointestinal nematodes of ruminants using predacious fungi. FAO, Roma, Italia, pp. 66–71, 1998.
- [2] Nurhidayah N, Satrija F, Retnani EBR. Gastrointestinal parasitic infection of swamp buffalo in Sentra Peternakan Rakyat (SPR) of Banten Province Indonesia: prevalence, risk factor and its impact to production performance. Trop Anim Sci J 2019; 41(1):6–12; <https://doi.org/10.5398/tasj.2019.42.1.6>
- [3] Robertson HA, Niezen JH, Waghorn GC, Charleston WAG, Jinlong M. The effect of six herbages on liveweight gain, wool growth and faecal egg count of parasitised ewe lambs. Proc N Zeal Soc Anim Prod 1995; 55:199–201.
- [4] Githiori JB, Athanasiadou S, Thamsborg SM. Use of plants in novel approaches for control of gastrointestinal helminths in livestock with emphasis on small ruminants. Vet Parasitol 2006; 139(4):308–20; <https://doi.org/10.1016/j.vetpar.2006.04.021>
- [5] Kundu S, Roy S, Lyndem LM. Broad spectrum anthelmintic potential of *Cassia* plants. Asian Pac J Trop Biomed 2014; 4(1):5436–41; <https://doi.org/10.12980/APJTB.4.2014C1252>
- [6] Alvarez AC, de Ugaz OL. Taninos. Rev Química VI 1992; 47–63.
- [7] John J, Mehta A, Shukla S, Mehta P. A report on anthelmintic activity of *Cassia tora* leaves. Songklanakarin J Sci Technol 2009; 31(3):269–71.
- [8] Sumi S, Saj OP. Antibacterial, anthelmintic and phytochemical investigations on the pod extracts of *Cassia fistula* Linn. Int J Med Pharm Sci 2012; 2(1):6–15.
- [9] Irshad M, Sing M, Rizvi MA. Assesment of anthelmintic activity of *Cassia fistula* L. Middle East J Sci Res 2010; 5(5):346–9.
- [10] Bhadauria S, Singh H. Bioactive nature of flavonoids from *Cassia siamea* and *Lantana camara*. Indian J Fund Appl Life Sci 2011; 1(2):107–10.
- [11] Jothy SL, Torey A, Darah I, Choong YS, Saravanan D, Chen Y, et al. *Cassia spectabilis* (DC) Irwin et Barn: a promising traditional herb in health improvement. Molecules 2012; 10292–305; <https://doi.org/10.3390/molecules170910292>

- [12] Motina BK. Antimicrobial activities of crude extract from *Cassia surattensis*. NU Sci J 2013; 10(1):10–7.
- [13] Suhasini K, Shridhar NB, Sanganal JS, Pattar J, Rao S. Evaluation of acute and subacute toxicity studies of *Cassia spectabilis* Leaf extract in Wistar Albino Rats. Int J Pharmacol Toxicol Sci 2013; 3(4):24–9.
- [14] Singh S, Singh SK, Yadav A. A review on *Cassia* species: pharmacological, traditional and medicinal aspects in various countries. Am J Phytomed Clin Therap 2013; 3:291–312.
- [15] Harborne JB. Phytochemical methods: a guide to modern techniques on plant analysis. 3rd edition, Kluwer Academic Publishers, UK, pp. 299–312, 1998.
- [16] Hansen J, Perry B. The epidemiology, diagnosis, and control of helminth parasites of ruminants: a Handbook. ILRAD, Nairobi, pp. 80–92, 1994.
- [17] Rabel B, Mcgregor R, Douch PGC. Improved bioassay for estimation of inhibitory effects of ovine gastrointestinal mucus and anthelmintics on nematode larval migration. Int J Parasitol 1994; 24(5):671–6; [https://doi.org/10.1016/0020-7519\(94\)90119-8](https://doi.org/10.1016/0020-7519(94)90119-8)
- [18] Gomez AA, Gomez KA. Statistical procedures for agricultural research. 1st edition, John Wiley & Sons, Inc., Canada, Vol. 6, p. 680, 1984.
- [19] SAS Institute. The SAS system for windows. Ver 9.0. SAS Institute Inc, Cary, NC, 2002.
- [20] Mohammed A, Liman ML, Atiku MK. Chemical composition of the methanolic leaf and stem bark extracts of *Senna siamea* Lam. J Pharmacogn Phytother 2013; 5(5):98–100.
- [21] Kamagaté M, Koffi C, Kouamé NM, Akoubet A, Alain N, Yao R, et al. Ethnobotany, phytochemistry, pharmacology and toxicology profiles of *Cassia siamea* Lam. J Phytopharm 2014; 3(1): 57–76.
- [22] Panda SK, Padhi LP, Mohanty G. Antibacterial activities and phytochemical analysis of *Cassia fistula* (Linn.) leaf. J Adv Pharm Technol Res 2016; 2(1):62–7; <https://doi.org/10.4103/2231-4040.79814>
- [23] Danish M, Singh P, Mishra G, Srivastava S, Jha KK, Khosa RL. *Cassia fistula* Linn. (Amulthus)—an important medicinal plant: a review of its traditional uses, phytochemistry and pharmacological properties. J Nat Prod Plant Resour 2011; 1(1):101–8; <https://doi.org/10.1080/01652176.1983.9693876>
- [24] Gobbo-Neto L, Lopes NP. Plantas medicinais: fatores de influência no conteúdo de metabólitos secundários. Química Nova 2007; 30(2):374–81; <https://doi.org/10.1590/S0100-40422007000200026>
- [25] Chaudhary S, Kumar A. Phytochemical analysis and assessment of *in vitro* anthelmintic activity of *Cassia auriculata* Linn leaves. Am J Phytomed Clin Therap 2014; 2(2):161–7.
- [26] Tiwari P, Kumar B, Kumar M, Kaur M, Debnath J, Sharma P. Comparative anthelmintic activity of aqueous and ethanolic stem extract of *Tinospora cardifolia*. Int J Drug Dev Res 2011; 3(1):70–83.
- [27] Williams AR, Fryganas C, Ramsay A, Mueller-Harvey I, Thamsborg SM. Direct anthelmintic effects of condensed tannins from diverse plant sources against *Ascaris suum*. PLoS One 2014; 9(5):e97053; <https://doi.org/10.1371/journal.pone.0097053>
- [28] Patel J, Kumar GS, Quesha MdS, Jena PK. Anthelmintic activity of ethanolic extract of whole plant of *Eupatorium odoratum* L. Int J Phytomed 2010; 2:127–32; <https://doi.org/10.5138/ijpm.2010.0975.0185.02020>
- [29] Botura MB, dos Santos JDG, da Silva GD, de Lima HG, de Oliveira JVA, de Almeida MAO, et al. *In vitro* ovicidal and larvicidal activity of *Agave sisalana* Perr. (sisal) on gastrointestinal nematodes of goats. Vet Parasitol 2013; 192:211–217; <https://doi.org/10.1016/j.vetpar.2012.10.012>
- [30] Jain P, Singh S, Singh SK, Verma SK, Kharya MD, Solanki S. Anthelmintic potential of herbal drugs. Int J Res Dev Pharm Life Sci 2013; 2(3):412–27.
- [31] Sangetha S, Zuraini Z, Sasidharan, Suryani S. Fungicidal effect and oral acute toxicity of *Cassia spectabilis* leaf extract. Jpn J Med Mycol 2008; 49:299–304; <https://doi.org/10.3314/jjmm.49.299>
- [32] Sumathy V, Zuraini Z, Sasidharan S. *In vivo* toxicity study of *Cassia surattensis* flower extract. Res J Pharm Biol Chem Sci 2011; 2(3):607–17.