



Cite this: *RSC Adv.*, 2019, 9, 40072

Received 21st September 2019
Accepted 19th November 2019

DOI: 10.1039/c9ra07653k

rsc.li/rsc-advances

Synthesis of novel polycarbonyl Schiff bases by ring-opening reaction of ethyl 5-acyl-4-pyrone-2-carboxylates with primary mono- and diamines†

D. L. Obydenov, * L. R. Khammatova, V. D. Steben'kov  and V. Y. Sosnovskikh 

An approach for the introduction of the tricarbonyl moiety into aromatic, heterocyclic, and aliphatic amines with the use of acylpyrones has been developed for the synthesis and the design of novel polycarbonyl Schiff base ligands, including salphen structures. This Michael addition–ring-opening reaction proceeds under mild conditions (stirring at 0–20 °C) *via* the attack at the C-6 position of the pyrone ring in good to high yields (up to 99%) with excellent selectivity. The products can be easily isolated by crystallization without the use of chromatography. The scope of the reaction, tautomeric equilibrium of open-chain products, and their cyclization into pyridone structures were investigated.

Introduction

Polycarbonyl compounds are widely used in various fields of materials chemistry, analytical and inorganic chemistry due to their multifarious chemical properties¹ and complex-formation ability.² Functional derivatives of 1,3-diketones, such as Schiff bases, play an especially important role as *N,N',O,O'*-ligands and can give stable metal-chelate complexes with various metal ions.³ These coordination compounds are actively used as catalytic systems for diverse organic reactions (synthesis of heterocycles^{4a–c} and polymerization^{4f}), as well as chemosensors,⁵ materials for OLEDs,⁶ OFEDs,⁷ and solar cells.⁸

To date, although a large number of metal salen complexes have been studied, methods for further functionalization of the ligand structure remain important. The introduction of additional coordinating centers for the synthesis of multinucleating salphen complexes makes it possible to improve their catalytic, magnetic, optical properties and ion sensing.⁹

It is well known that the main method for the preparation of Schiff bases involves the condensation of carbonyl compounds with aromatic or aliphatic amines,¹⁰ whereas the use of Michael reaction as an effective synthetic tool remains limited for the construction of ligands.¹¹ Unlike the 1,3-dicarbonyl motif, the introduction of the polycarbonyl moiety into the molecule based on 1,3,5-triketones and other open-chain polyketones has some drawback, such as side reactions (the retro-Claisen cleavage, the intramolecular cyclization), a low selectivity, and low chemical reactivity of these substrates due to the presence of the acidic protons involved in the formation of the

intramolecular hydrogen bonds.¹² In the literature there are only a few examples of salen ligand synthesis based on 2,4,6-hexanetriones.¹³ Using acyl-4-pyrones as highly reactive synthetic equivalents of 1,3,5-triones, which can undergo Michael reaction under the action of amines as a key stage to circumvent the above difficulties. In this case, the reaction proceeds under mild conditions to avoid the formation of heterocyclization products and with high chemoselectivity. The ability of 4-pyrones to react with various N-nucleophiles can allow the introduction of the polycarbonyl moiety in wide scope of substrates.

5-Acylcomanoates **1** (ref. 14) are convenient building blocks for the construction of heterocyclic systems *via* a nucleophilic attack at the C-6 position due to the presence of the electron-withdrawing substituent at the C-5 position followed by pyrone ring-opening (Fig. 1). Usually, 4-pyrones react with N-nucleophiles *via* the ANRORC (Addition of the Nucleophile, Ring Opening, and Ring Closure) mechanism and give more stable azaheterocycles.^{14a,b} Earlier,¹⁵ for the reaction of 2,5-dicarbethoxy-4-pyrone and ethyl 5-benzoylcomanoate with *o*-phenylenediamine, the formation of polyfunctional open-chain polycarbonyl substrates only as intermediates in the synthesis of polycyclic pyridones have been shown.

In addition, in organic chemistry close attention is paid to the development of highly efficient synthetic methods, such as “Click Chemistry”. This term describes reactions that are high yielding, wide in scope, stereospecific, simple to perform, high atom-economic, include facile the isolation of a product without the use of chromatography and can be carried out in easily removable or benign solvents.¹⁶ The search for such transformations is actively conducted, and Michael reaction and ring-opening reaction belong to a number of click-reactions.¹⁶ Pyrones can undergo the tandem of Michael reaction/pyrone ring-opening, therefore, they are of interest for the

Institute of Natural Sciences and Mathematics, Ural Federal University, 51 Lenina Ave., 620000 Ekaterinburg, Russian Federation. E-mail: dobydenov@mail.ru

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c9ra07653k






Fig. 1 Reactions of acylpyrones **1** with primary amines.

development of new synthetic click-approaches for the construction of polycarbonyl structures.

This work describes the general approach for the introduction of the 1,3,5-triketone moiety into the molecules bearing the amino group with the use of 5-acyl-4-pyrone-2-carboxylic acid esters to obtain and design new polycarbonyl Schiff base ligands, including salphen structures, as well as to study their tautomeric equilibrium and the substrate scope of this method.

Results and discussion

We started the study with the reaction of 5-acylcomanoates **1** with various aromatic mono- and diamines **2** in EtOH at 0 °C or 20 °C (Scheme 1 and Table 1). It was found that pyrones **1** gave tetracarbonyl enaminones **3a–o** in medium to high yields (47–96%). The nature of the substituents in the aromatic amine has a slight effect on the outcome of the reaction though unreactive amines bearing the electron-withdrawing group, such as 2-aminopyridine and 2-aminopyrimidine, gave enaminones **3k** and **3m** at 20 °C in good yields (74% and 47%, respectively). The transformation was found to be tolerant to steric hindrance of a bulky acyl group and, for pivaloylpyrone **1f** bearing the bulky *tert*-butyl group, product **3i** was obtained in 61% yield.

It was found that the polycarbonyl moiety can be introduced into the molecule of aliphatic amine, such as benzylamine (**2h**). Indeed, enaminones **3n,o** were obtained in lower yield (43–54%) compared to aromatic amines. This result can be explained by easier intramolecular cyclization of enaminone **3** into stable pyridones because of higher nucleophilicity of the alkylamine.







Products **3** are multifunctionalized structures bearing the enaminodione and diketone moieties, which can undergo keto-enol tautomerism and *Z/E* isomerization as the result of the push-pull nature of the double bond. Two geometric isomers *Z-3* and *E-3* and also tautomer *keto-E-3*, which exist in equilibrium in the solution of CDCl₃, were detected by NMR spectroscopy (Scheme 1 and Table 1).

A clear influence of the substituents on the tautomeric equilibrium is observed, which can be explained by the stability of the hydrogen bond formed and steric interactions. The introduction of the electron-donating MeO group into the aryl substituent of *N*-phenyl-substituted enaminones **3a** increases the content of the most stable isomer *E-3* from 82% to 95%. Although a change in the structure of the amine has an effect on the observed equilibrium, the introduction of the *ortho* substituent reduces the content of major tautomer *E-3* to 76–79%. Enaminones **3h,j,l,m**, which bear the thenoyl substituent, are especially distinguished in this series because the content of the major isomer *E-3* reaches 87–93%.

The structure of tautomer *E-3* was confirmed on the basis of the coupling constants extracted from the ¹H-coupled ¹³C NMR spectrum of **3b**, along with the analysis of ¹H–¹³C HMBC spectrum of **3b**. It is known¹⁷ that, in the case of enones, coupling constant ³J_{H,C} between the β-olefinic protons and the carbonyl group strongly depends on the dihedral angle and is a good criterion for establishing the configuration of the double bond when other experiments cannot be applied. For NC-substituted enaminones, the CH proton of the N–CH= moiety has ³J_{H,CN} coupling with the cyano group in the range of 4.6–5.8 Hz and 9.6–10.8 Hz for their *cis*- and *trans*-orientation, respectively.¹⁸ Also in the literature the ³J_{H,C=O} coupling has been measured¹⁹ for similar structures, alkoxymethylidenediones (*cis*-³J_{H,C=O} = 2.6–4.0 Hz, *trans*-³J_{H,C=O} = 7.7–10.1 Hz) and the magnitudes of these ³J_{H,C=O} coupling


Table 1 Reaction scope study of pyrones 1 with amines 2^a

Adduct	Pyrone	Amine	R	X	Yield, %	<i>E</i> -3 : <i>Z</i> -3 : keto- <i>E</i> -3
3a	1a	2a	Ph	H	87	82 : 13 : 5
3b	1b	2a	4-MeOC ₆ H ₄	H	85	95 : 2 : 3
3c	1a	2b	Ph	OH	68	76 : 17 : 7
3d	1c	2b	4-ClC ₆ H ₄	OH	81	76 : 17 : 7
3e	1b	2c	4-MeOC ₆ H ₄	NH ₂	86	79 : 15 : 6
3f	1c	2c	4-ClC ₆ H ₄	NH ₂	78	78 : 16 : 6
3g	1d	2c	2-Naph	NH ₂	90	79 : 13 : 8
3h	1e	2c	2-Th	NH ₂	90	89 : 8 : 3
3i	1f	2c	<i>t</i> -Bu	NH ₂	61	79 : 14 : 7



^a Pyrone **1** (1.11 mmol) and amine **2** (1.11 mmol) were stirred in EtOH (6 mL) at 0 °C for 1 h. ^b The reaction was performed at 20 °C for 2 h. ^c The reaction was performed at 20 °C for 4 days, 2.22 mmol of the amine was used.

constants are convenient parameters for establishment of enaminodione configuration in our case. It was found that for enaminodione **3b** the carbonyl group of the aryl substituent (C-4') is a quartet at δ 193.0 ppm with $^3J_{\text{H,C}} = 3.7$ Hz, which indicates the *cis*-arrangement of the H-6 proton relative to C-4' (Fig. 2). The signal for the carbonyl group of the diketone moiety (C-4) in the spectrum of **3b** appears as a doublet of doublets and exhibits a $^3J_{\text{C-H}}$ coupling of 7.8 Hz with the H-6 proton and a small $^2J_{\text{C-H}}$ coupling of 3.7 Hz, which is consistent with the magnitude of the coupling constant $^2J_{\text{C-H}}$ in the =CH-C=O

Fig. 2 Selected ¹³C NMR signals of *E*-3b.

moiety (2.3–3.3 Hz).²⁰ The value of the first (7.8 Hz) is more than $^3J_{\text{H,C}}$ for C-4' (3.7 Hz) and is in agreement with the *trans*-arrangement of H-6 and C-4, which corresponds structure *E*-3b.

The ¹H NMR spectra of polycarbonyl compounds (het)aryl substituted **3a–m** feature a main set of signals of tautomer *E*-3 with the 5*E*-configuration of the double bond which consists of a doublet of the NH proton at δ 12.07–12.60 ppm ($^3J = 12.4$ –13.6 Hz) and a downfield singlet for the OH group (δ 14.27–14.50 ppm), apparently involved in a strong intramolecular hydrogen bonds, a doublet of the =CHN proton at δ 7.86–8.81 ppm ($^3J = 12.4$ –13.6 Hz), and a vinylic proton singlet at δ 6.59–7.15 ppm. For Bn-substituted **3n,o**, the signals of both the =CHN proton and the NH proton are shifted upfield compared with those of (het)aryl substituted **3a–m** and appeared at δ 7.49–7.69 and 10.95 ppm, respectively.

The characteristic signals of minor tautomer *Z*-3 with 5*Z*-configuration include a doublet of the =CHN proton (δ 8.27–9.30 ppm, $^3J = 12.8$ –14.0 Hz), which is shifted downfield by *ca.* 0.5 ppm compared with the same proton of *E*-3, a singlet for the =CHCO proton (δ 5.84–6.34 ppm) is shifted upfield by 0.7–1.2 ppm compared to *E*-3 isomer due to a shielding influence of the aromatic ring of the acyl moiety (Fig. 3). Such a difference in




Fig. 3 Fragment of the ^1H NMR spectrum of **3a** in CDCl_3 .

the chemical shifts of these protons was not observed for pivaloyl-substituted enaminones **3i,n** (only 0.01–0.2 ppm). The OH proton and the NH proton appear as broadened signals at δ 15.0–15.4 ppm and δ 11.19–12.60 ppm (for aryl substituted **3a–m**, δ 10.04–10.95 ppm for benzyl substituted **3n,o**), respectively, as a result of the formation of the intramolecular hydrogen bonds. It is important to note that the OH proton signal is shifted downfield by *ca.* 0.8 ppm in comparison with the same proton of *E*-3 isomer, which indicates the stronger intramolecular hydrogen bond ($\text{OH}\cdots\text{O}=\text{C}$) in the *Z*-3 isomer and is consistent with the proposed structures.

It is well known^{14b,d} that diketobutanoates predominantly exist as the enol form and contain some quantity of the diketo form (0–10%). Therefore, the structure of keto-*E*-3 was established on the basis of the relative content of the tautomer (0–13%). In ^1H NMR spectra the feature signals of the tautomer are a doublet ($^3J \approx 13.0$ Hz) of the downfield NH proton at δ 12.22–12.60 ppm (for keto-*E*-**3a–m**) or δ 11.23–11.25 ppm (for keto-*E*-**3n–o**) and a singlet of the CH_2 group at δ 4.07–4.40 ppm. The


signal of the NH proton of the tautomer is shifted downfield by 0.2–0.6 ppm compared to the same proton of *E*-3 isomer because of stronger electron-withdrawing properties of the non-conjugated carbonyl group of the diketone moiety.

Enaminones **3** are of interest as chemically active substances for further transformations. When substrates **3** bearing the amino group react with pyrone **1** in EtOH at room temperature for 24 h, they give polycarbonyl ligands **4a–e** in 64–79% yields (Method A, Scheme 2 and Table 2), which are yellow or red precipitates and stable during storage in the solid state. Polycarbonyl ligands **4** can also be obtained directly from *o*-phenylenediamines **2** by the reaction with two equivalents of 4-pyrones **1** in 31–63% yields (Method B).

Theoretically, compounds **4** can be represented as ten open-chain tautomeric structures, but in the ^1H NMR spectra it is possible to detect and assign some signals only for *5E,5'E-4* and *5E,5'Z-4* tautomers by comparing NMR signals of parent polycarbonyl compounds **3**.

The structure of compounds **4** was confirmed on the basis of elemental analysis data, IR and NMR spectra. In ^1H NMR spectra in CDCl_3 major isomer *5E,5'E-4* appeared as a singlet of the H-3 proton at δ 6.82–6.95 ppm, doublets of the H-6 proton and the NH group at δ 7.92–8.09 ppm and 12.41–12.58 ppm ($^3J = 12.4$ – 12.6 Hz), respectively. The characteristic signals of minor isomer *5E,5'Z-4* include two singlets at δ 5.90–6.36 and 6.41–6.45 ppm attributed to the vinylic protons of the diketone motifs and two doublets of the $=\text{CHN}$ protons at δ 7.99–8.12 ppm and 8.36–8.53 ppm.

It was found that the content of the most stable *5E,5'E-4* tautomer is equal to 39–74%. The nature of the substituents in the aryl group has a strong influence on the content of this tautomer. In the case of electron-donating substituents, the content of *5E,5'E*-isomer is the highest (69–74% for **4b,d,e**), while the introduction of the electron-withdrawing substituent ($\text{Ar} = 4\text{-ClC}_6\text{H}_4$) leads to a decrease in the content of the main isomer to 39%. These data are also consistent with the amount




Scheme 2 Reactions of pyrones **1** with diamines **2**.

Table 2 Scope of the pyrone transformation with diamines 2^a

Bis-adduct	Ar	R	Yield from enaminone 3, % (Method A)	Yield from diamine, % (Method B)	5 <i>E</i> ,5' <i>E</i> -4 : 5 <i>E</i> ,5' <i>Z</i> -4
4a	Ph	H	64	57	50 : 17
4b	4-MeOC ₆ H ₄	H	73	38	69 : 17
4c	4-ClC ₆ H ₄	H	65	75	39 : 26
4d	2-Th	H	75	31	71 : 14
4e	2-Th	Me	79	63	74 : 10

^a Method A: pyrone 1 (0.26 mmol) and enaminone 3 (0.26 mmol) were stirred in EtOH (2 mL) at 20 °C for 24 h. Method B: pyrone 1 (0.72 mmol) and diamine 2 (0.36 mmol) were stirred in EtOH (2 mL) at 0 °C for 1 h and then at 20 °C for 24 h.

Scheme 3 Reaction of pyrone 1a with *p*-phenylenediamine.

of minor 5*E*,5'*Z*-isomer (10–26%), the largest content (26%) was found for 4c (Ar = 4-ClC₆H₄).


Compounds 4 are polydentate salphen structures, which makes them promising ligands for the formation of binuclear complex compounds with various metal ions, which have been recently received much attention for various application from catalysis to materials chemistry.⁹ It was found that compound 4c upon treatment with copper(II) acetate gives binuclear chelate complex 5, whose structure is in agreement with the literature data¹³ and is confirmed by elemental analysis, ESI-MS data, NMR and IR²¹ spectroscopies. ¹H NMR spectra for Cu(II)-coordinated compounds are usually not informative as the result of strong signal peak broadening and unpredictable chemical shifts due to paramagnetic properties of Cu(II) compounds.²² In contrast, binuclear complexes can produce narrow ¹H NMR resonance because of magnetic interaction of two paramagnetic copper(II) centers.²² The ¹H NMR spectrum of compound 5 appears to be useful and features a single set of signals, which consists of a broadened triplet for the Me group at δ 1.10 ppm, broadened quartet at δ 3.76 ppm, a broadened singlet for =CH proton at δ 5.74 ppm, and a multiplet for the aromatic protons at δ 7.24–7.80 ppm.

It is interesting to note that the reaction of benzoylpyrone 1a (2 equiv.) with *p*-phenylenediamine (2i, 1 equiv.) proceeds very readily to form compound 6 in almost quantitative yield (99%) (Scheme 3). Such a large difference in reactivity compared to *o*-phenylenediamine can be explained by less steric hindrances

during the formation of the polycarbonyl ligands. In contrast to aromatic diamines, the reaction of pyrone 1a with ethylenediamine led to a product, which exists in solution as a complex mixture of tautomers.

The main drawback of the polycarbonyl Schiff base ligands bearing the triketone moiety is their ability to cyclize into pyridones, which may limit their application. Therefore, further we decided to study the chemical properties of the polycarbonyl compounds 3, which are connected with their stability and the possibility of further modification for the design of new ligands.


Enaminones 3e,f,h,j bearing the amino group in the *ortho* position can undergo intramolecular cyclization to form more



Scheme 4 Cyclization reactions of polycarbonyl compounds 3.

Table 3 Synthesis of pyrido[1,2-*a*]quinoxaline-6,8-diones 7,8

Starting enaminone	Ar	R	Product 7, 8	Yield 7, %	Yield 8, %	7 : 8 (in DMSO- <i>d</i> ₆)
3e	4-MeOC ₆ H ₄	H	a	72	89	81 : 19
3f	4-ClC ₆ H ₄	H	b	89	89	76 : 24
3h	2-Th	H	c	86	92	93 : 7
3j	2-Th	Me	d	85	80	97 : 3

Scheme 5 Reaction of pyrone **1e** with 2,3-diaminopyridine.

stable polycyclic pyridones **7** and **8** (Scheme 4 and Table 3). It was found that reflux of compounds **3** in EtOH without the addition of acid leads to the formation of dihydropyridones **7** in 72–89%. Although quinoxalines **8** were detected in the ¹H NMR spectra of products **7** (the content reached 24%), structures **7** are proved by the data of elemental analysis and the IR spectra of the obtained precipitates **7** exhibit the absence of the characteristic bands of the carbonyl groups of pyridones **8** at 1683–1710 cm⁻¹. The obtained result probably shows that the formation of compounds **8** occurs upon the dissolution of dihydropyridone **7** in DMSO-*d*₆. Recrystallization of compounds **7** from acetic acid or boiling in EtOH with the addition of hydrochloric acid lead to the formation of dehydration products **8** in 80–92%.

Enaminones **3a,d**, which were obtained from aniline and *o*-aminophenol, undergo cyclization in the presence of acids gave thermodynamically more stable pyridones **9** in 78–79% (Scheme 4). On the other hand, it was important to find ways for the modification of such polycarbonyl ligands into structures containing heterocyclic substituents by reactions on the active diketobutanoic moiety. Using enaminone **3c** as a representative example, it was shown that this compound is able to react with *o*-aminophenol (**2b**) under reflux in benzene to form benzoxazine **10** in 85% yield.

The reaction of thenoylpyrone **1e** with 2,3-diaminopyridine (**2j**) currently led to dihydropyridone **11** (the content of dehydration product **12** is 8% in DMSO-*d*₆) in 91% yield without

the isolation of polycarbonyl intermediate **3** (Scheme 5). Compound **11** gave pyrido[1,2-*a*]quinoxaline-6,8-dione **12** in 81% yield under heating in AcOH. The structure of products **11,12** was supported by the literature data²³ regarding reactions of 2,3-diaminopyridones with polycarbonyl compounds.

Conclusions

In summary, we have found a convenient, convergent, and effective approach for the introduction of a triketone moiety into various amines using ethyl esters of 5-acyl-4-pyrone-2-carboxylic acids, which allows the synthesis of a wide range of new polycarbonyl Schiff base ligands and salphen structures. Important advantages of this reaction include its wide substrate scope, catalyst free, atom economy, high yields, mild conditions and easy performing and isolation of products without the use of chromatography. Obtained polycarbonyl compounds in the solution of CDCl₃ exist as several tautomers, the major tautomer has 5*E*-configuration of the double bond. The found strategy for the introduction of the triketone moiety can be used further for the deep functionalization of polymers and biopolymers containing a free amino group for the creation of novel catalysts or materials.

Experimental

General

NMR spectra were recorded on Bruker DRX-400 (¹H – 400 MHz and ¹³C – 100 MHz) and Bruker Avance III-500 (¹H – 500 MHz and ¹³C – 126 MHz) spectrometers in DMSO-*d*₆ or CDCl₃. Chemical shifts are reported relative to TMS, CHCl₃ (δ = 7.26 ppm, ¹H NMR), DMSO-*d*₆ (δ = 2.50 ppm, ¹H NMR), CDCl₃ (δ = 77.06 ppm, ¹³C NMR) and DMSO-*d*₆ (δ = 39.52 ppm, ¹³C NMR) as internal standards. IR spectra were recorded on a Shimadzu IRSpirit-T (QATR-S) instrument (FTIR mode, ZnSe crystal). Elemental analysis was performed on a PerkinElmer PE 2400 automatic analyzer. The mass spectrum (ESI-MS) of compound **5** was measured with a Waters Xevo QToF instrument. All solvents used were dried and distilled per standard procedures. Melting points were determined on a Stuart SMP40 apparatus. Pyrones **1** were prepared according the literature procedure.²⁴

General approach for the preparation of enaminodiones **3**

Pyrone **1** (1.11 mmol) was added to EtOH (6 mL) containing amine **2** (1.11 mmol) (for **3m** – 2.22 mmol) under cooling on an

ice bath. The resulted mixture was stirred at 0 °C for 1 h (for **3k** – at 20 °C for 2 h, or **3m** – at 20 °C for 1 day). The precipitate was filtered and washed with EtOH (for **3i**, the reaction mixture was additionally maintained at –20 °C for 12 h).

(2Z,5E)-Ethyl-5-benzoyl-2-hydroxy-4-oxo-6-(phenylamino)hexa-2,5-dienoate (3a). Yield 87% (0.353 g), yellow powder, mp 128–129 °C. IR (ATR): 2987, 1728, 1606, 1575, 1252, 847 cm⁻¹. This compound exists in CDCl₃ as a mixture of tautomers *E-3a* : *Z-3a* : keto-**3a** = 82 : 13 : 5. ¹H NMR (CDCl₃, ppm) δ: *E-3a* (82%) – 1.31 (t, *J* = 7.1 Hz, 3H, Me), 4.30 (q, *J* = 7.1 Hz, 2H, OCH₂), 7.01 (s, 1H, =CH), 7.10 (d, *J* = 7.7 Hz, 2H, H-2, H-6 Ph), 7.22 (tt, *J* = 7.4 Hz, *J* = 1.1 Hz, 1H, H-4 Ph), 7.38 (dd, *J* = 7.7 Hz, *J* = 7.4 Hz, 2H, H-3, H-5 Ph), 7.48 (dd, *J* = 7.8 Hz, *J* = 7.2 Hz, 2H, H-3, H-5 Ph'), 7.57 (tt, *J* = 7.8 Hz, *J* = 1.2 Hz, 1H, H-4 Ph'), 7.73 (dd, *J* = 7.2 Hz, *J* = 1.4 Hz, 1H, H-2, H-6 Ph'), 8.08 (d, *J* = 13.4 Hz, 1H, =CHN), 12.41 (d, *J* = 13.4 Hz, 1H, NH), 14.36 (s, 1H, OH); ¹³C NMR (CDCl₃, ppm) δ 14.1, 62.1, 103.9, 111.5, 117.9, 126.2, 128.6, 129.3, 130.1, 132.2, 138.5, 139.8, 153.3, 160.1, 162.6, 194.08, 194.12. Anal. calcd for C₂₁H₁₉NO₅: C, 69.03; H, 5.24; N, 3.83. Found: C, 69.08; H, 5.13; N, 4.04.

(2Z,5E)-Ethyl-2-hydroxy-5-(4-methoxybenzoyl)-4-oxo-6-(phenylamino)hexa-2,5-dienoate (3b). Yield 85% (0.373 g), yellow powder, mp 131–133 °C. IR (ATR): 3205, 3005, 2849, 1715, 1580, 1245, 1108, 1023 cm⁻¹. This compound exists in CDCl₃ as a mixture of tautomers *E-3b* : *Z-3b* : keto-**3b** = 95 : 2 : 3. ¹H NMR (CDCl₃, ppm) δ: *E-3b* (95%) – 1.31 (t, *J* = 7.1 Hz, 3H, Me), 3.89 (s, 3H, MeO), 4.29 (q, *J* = 7.1 Hz, 2H, OCH₂), 6.90 (s, 1H, =CH), 6.97 (d, AA'BB', *J* = 8.9 Hz, 2H, H-3 H-5 Ar), 7.12 (d, *J* = 7.7 Hz, 2H, H-2, H-6 Ph), 7.21 (tt, *J* = 7.4 Hz, *J* = 0.9 Hz, 1H, H-4 Ph), 7.39 (dd, *J* = 7.9 Hz, *J* = 7.4 Hz, 2H, H-3, H-5 Ph), 7.74 (d, AA'BB', *J* = 8.9 Hz, 2H, H-2, H-6 Ar), 8.05 (d, *J* = 13.3 Hz, 1H, =CHN), 12.34 (d, *J* = 13.3 Hz, 1H, NH), 14.38 (s, 1H, OH); ¹³C NMR (CDCl₃, ppm) δ 14.0 (Me), 55.5 (MeO), 62.1 (CH₂), 103.9 (C-3), 111.7 (C-5), 113.8 (C-3, C-5 Ar), 117.8 (C-2, C-6 Ph), 126.0 (C-4 Ph), 130.0 (C-3, C-5 Ph), 131.8 (C-2, C-6 Ar), 132.1 (C-1 Ar), 138.6 (C-1 Ph), 152.4 (C-6), 159.9 (C-2), 162.6 (C-1), 163.1 (C-4 Ar), 193.0 (Ar-C-4=O), 193.9 (C-4=O). Anal. calcd for C₂₂H₂₁NO₆: C, 66.83; H, 5.35; N, 3.54. Found: C, 66.61; H, 5.27; N, 3.86.

(2Z,5E)-Ethyl-5-benzoyl-2-hydroxy-6-((2-hydroxyphenyl)amino)-4-oxohexa-2,5-dienoate (3c). Yield 68% (0.288 g), yellow powder, mp 172–174 °C. IR (ATR): 3057, 1733, 1640, 1608, 1562, 1446, 1251, 1157, 1099, 847 cm⁻¹. This compound exists in CDCl₃ as a mixture of tautomers *E-3c* : *Z-3c* : keto-**3c** = 76 : 17 : 7. ¹H NMR (CDCl₃, ppm) δ: *E-3c* (76%) – 1.31 (t, *J* = 7.1 Hz, 3H, Me), 4.30 (q, *J* = 7.1 Hz, 2H, OCH₂), 6.79 (br, 1H, HOAr), 6.86–6.93 (m, 1H, Ar), 6.95–7.10 (m, 4H, Ar, =CH), 7.48 (t, *J* = 7.4 Hz, 2H, H-3, H-5 Ph), 7.57 (t, *J* = 7.2 Hz, 1H, H-4 Ph), 7.73 (d, *J* = 7.3 Hz, 2H, H-2, H-6 Ph), 8.13 (d, *J* = 13.3 Hz, 1H, =CHN), 12.60 (d, *J* = 13.3 Hz, 1H, NH), 14.80 (s, 1H, OH). Anal. calcd for C₂₁H₁₉NO₆: C, 66.14; H, 5.02; N, 3.67. Found: C, 66.32; H, 4.91; N, 3.38.

(2Z,5E)-Ethyl-5-(4-chlorobenzoyl)-2-hydroxy-6-(2-hydroxyphenylamino)-4-oxohexa-2,5-dienoate (3d). Yield 81% (0.374 g), yellow powder, mp 180–182 °C. IR (ATR): 3190, 1734, 1609, 1576, 1248, 744 cm⁻¹. This compound exists in CDCl₃ as a mixture of tautomers *E-3d* : *Z-3d* : keto-**3d** = 76 : 17 : 7. ¹H NMR (CDCl₃, ppm) δ: *E-3d* (76%) – 1.31 (t, *J* = 7.1 Hz, 3H, Me),

4.30 (q, *J* = 7.1 Hz, 2H, OCH₂), 6.10 (br, 1H, HOAr), 6.88–7.12 (m, 5H, Ar, =CH), 7.45 (d, AA'BB', *J* = 8.4 Hz, 2H, H-3, H-5 Ar'), 7.68 (d, AA'BB', *J* = 8.4 Hz, 2H, H-2, H-6 Ar'), 8.10 (d, *J* = 13.6 Hz, 1H, =CHN), 12.59 (d, *J* = 13.6 Hz, 1H, NH), 14.49 (s, 1H, OH). Anal. calcd for C₂₁H₁₈ClNO₆: C, 60.66; H, 4.36; N, 3.37. Found: C, 60.61; H, 4.10; N, 3.30.

(2Z,5E)-Ethyl-6-(2-aminophenylamino)-2-hydroxy-5-(4-methoxybenzoyl)-4-oxohexa-2,5-dienoate (3e). Yield 83% (0.378 g), red powder, mp 131–133 °C. IR (ATR): 3394, 1731, 1611, 1595, 1246, 749 cm⁻¹. This compound exists in CDCl₃ as a mixture of tautomers *E-3e* : *Z-3e* : keto-**3e** = 79 : 15 : 6. ¹H NMR (CDCl₃, ppm) δ: *E-3e* (79%) – 1.31 (t, *J* = 7.1 Hz, 3H, Me), 3.77 (br, 2H, NH₂), 3.88 (s, 3H, MeO), 4.29 (q, *J* = 7.1 Hz, 2H, OCH₂), 6.78–6.87 (m, 2H, Ar), 6.92–7.00 (m, 4H, =CH, Ar, Ar'), 7.07 (td, *J* = 7.6 Hz, *J* = 0.9 Hz, 1H, H-4 Ar), 7.74 (d, AA'BB', *J* = 8.7 Hz, 2H, H-2, H-6 Ar'), 7.95 (d, *J* = 13.1 Hz, 1H, =CHN), 12.24 (d, *J* = 13.1 Hz, 1H, NH), 14.33 (s, 1H, OH). Anal. calcd for C₂₂H₂₂N₂O₆: C, 64.38; H, 5.40; N, 6.83. Found: C, 64.59; H, 5.60; N, 6.64.

(2Z,5E)-Ethyl-6-(2-aminophenylamino)-5-(4-chlorobenzoyl)-2-hydroxy-4-oxohexa-2,5-dienoate (3f). Yield 78% (0.359 g), orange powder, mp 138–140 °C. IR (ATR): 3404, 3351, 3068, 1723, 1597, 117, 841 cm⁻¹. This compound exists in CDCl₃ as a mixture of tautomers *E-3f* : *Z-3f* : keto-**3f** = 78 : 16 : 6. ¹H NMR (CDCl₃, ppm) δ: *E-3f* (78%) – 1.34 (t, *J* = 7.1 Hz, 3H, Me), 3.79 (br, 2H, NH₂), 4.32 (q, *J* = 7.1 Hz, 2H, OCH₂), 6.81–6.90 (m, 2H, Ar), 6.98 (d, *J* = 7.9 Hz, 1H, H-6 Ar), 7.00 (s, 1H, =CH), 7.07 (td, *J* = 7.7 Hz, *J* = 0.9 Hz, 1H, H-4 Ar), 7.47 (d, AA'BB', *J* = 8.4 Hz, 2H, H-3, H-5 Ar'), 7.69 (d, AA'BB', *J* = 8.4 Hz, 2H, H-2, H-6 Ar'), 7.97 (d, *J* = 13.2 Hz, 1H, =CHN), 12.33 (d, *J* = 13.2 Hz, 1H, NH), 14.28 (s, 1H, OH). Anal. calcd for C₂₁H₁₉ClN₂O₅: C, 60.80; H, 4.62; N, 6.72. Found: C, 60.78; H, 4.62; N, 6.67.

(2Z,5E)-Ethyl-5-(2-naphthoyl)-6-(2-aminophenylamino)-2-hydroxy-4-oxohexa-2,5-dienoate (3g). Yield 90% (0.430 g), orange powder, mp 169–171 °C. IR (ATR): 3401, 3183, 2987, 1728, 1532, 1246, 1101, 827 cm⁻¹. This compound exists in CDCl₃ as a mixture of tautomers *E-3g* : *Z-3g* : keto-**3g** = 79 : 13 : 8. ¹H NMR (CDCl₃, ppm) δ: *E-3g* (79%) – 1.25 (t, *J* = 7.1 Hz, 3H, OCH₂Me), 3.79 (br, 2H, NH₂), 4.27 (q, *J* = 7.1 Hz, 2H, OCH₂), 6.79 (td, *J* = 7.4 Hz, *J* = 0.9 Hz, 1H, H-5 Ar), 6.85 (dd, *J* = 8.0 Hz, *J* = 0.9 Hz, 1H, H-3 Ar), 6.95 (d, *J* = 7.5 Hz, 1H, H-6 Ar), 7.08 (td, *J* = 7.7 Hz, *J* = 0.9 Hz, 1H, H-4 Ar), 7.10 (s, 1H, =CH), 7.51–7.68 (m, 2H, Naph), 7.84 (dd, *J* = 8.5 Hz, *J* = 1.5 Hz, 1H, Naph), 7.89–8.00 (m, 3H, Naph), 8.07 (d, *J* = 13.1 Hz, 1H, =CHN), 8.24 (d, *J* = 0.7 Hz, 1H, H-1 Naph), 12.38 (d, *J* = 13.1 Hz, 1H, NH), 14.34 (s, 1H, OH). Anal. calcd for C₂₅H₂₂N₂O₅: C, 69.76; H, 5.15; N, 6.51. Found: C, 69.88; H, 5.14; N, 6.30.

(2Z,5E)-Ethyl-6-(2-aminophenylamino)-2-hydroxy-4-oxo-5-(thiophene-2-carbonyl)hexa-2,5-dienoate (3h). Yield 90% (0.386 g), orange powder, mp 143–145 °C. IR (ATR): 3416, 3345, 3135, 1722, 1635, 1578, 1248, 836 cm⁻¹. This compound exists in CDCl₃ as a mixture of tautomers *E-3h* : *Z-3h* : keto-**3h** = 89 : 8 : 3. ¹H NMR (CDCl₃, ppm) δ: *E-3h* (89%) – 1.33 (t, *J* = 7.1 Hz, 3H, Me), 3.79 (br, 2H, NH₂), 4.31 (q, *J* = 7.1 Hz, 2H, OCH₂), 6.82–6.88 (m, 2H, H-5, H-3 Ar), 7.04 (dd, *J* = 7.2 Hz, *J* = 1.0 Hz, 1H, H-6 Ar), 7.04 (s, 1H, =CH), 7.09 (td, *J* = 7.9 Hz, *J* = 1.1 Hz, 1H, H-4 Ar), 7.14 (dd, *J* = 5.0 Hz, *J* = 3.8 Hz, 1H, H-4 Th), 7.55 (dd, *J* = 3.8 Hz, *J* = 1.0 Hz, 1H, H-3 Th), 7.67 (dd, *J* = 5.0 Hz, *J*

=1.0 Hz, 1H, H-5 Th), 8.14 (d, $J = 13.1$ Hz, 1H, =CHN), 12.22 (d, $J = 13.1$ Hz, 1H, NH), 14.28 (s, 1H, OH). Anal. calcd for $C_{19}H_{18}N_2O_5S$: C, 59.06; H, 4.70; N, 7.25. Found: C, 58.88; H, 4.88; N, 7.49.

(2Z,5E)-Ethyl-5-((2-aminophenylamino)methylene)-2-hydroxy-7,7-dimethyl-4,6-dioxooct-2-enoate (3i). Yield 61% (0.244 g), yellow powder, mp 104–106 °C. IR (ATR): 3223, 3100, 2959, 2921, 1695, 1677, 1637, 1504, 1108, 882 cm^{-1} . This compound exists in $CDCl_3$ as a mixture of tautomers *E-3i* : *Z-3i* : keto-**3i** = 79 : 14 : 7. 1H NMR ($CDCl_3$, ppm) δ : *E-3i* (79%) – 1.35 (s, 9H, *t*-Bu), 1.38 (t, $J = 7.1$ Hz, 3H, Me), 3.75 (br, 2H, NH_2), 4.36 (q, $J = 7.1$ Hz, 2H, OCH_2), 6.59 (s, 1H, =CH), 6.83–6.92 (m, 2H, Ar), 7.05 (dd, $J = 7.8$ Hz, $J = 0.9$ Hz, 1H, H-6 Ar), 7.11 (td, $J = 7.6$ Hz, $J = 1.1$ Hz, 1H, H-4 Ar), 7.86 (d, $J = 12.4$ Hz, 1H, =CHN), 11.95 (d, $J = 12.4$ Hz, 1H, NH), 14.36 (s, 1H, OH). Anal. calcd for $C_{19}H_{24}N_2O_5S$: C, 63.22; H, 6.71; N, 7.77. Found: C, 63.53; H, 6.58; N, 7.89.

(2Z,5E)-Ethyl-6-(2-amino-4,5-dimethylphenylamino)-2-hydroxy-4-oxo-5-(thiophene-2-carbonyl)hexa-2,5-dienoate (3j). Yield 96% (0.442 g), red powder, mp 157–159 °C. IR (ATR): 3411, 3370, 2992, 1717, 1590, 1256, 794 cm^{-1} . This compound exists in $CDCl_3$ as a mixture of tautomers *E-3j* : *Z-3j* : keto-**3j** = 87 : 9 : 4. 1H NMR ($CDCl_3$, ppm) δ : *E-3j* (87%) – 1.33 (t, $J = 7.1$ Hz, 3H, OCH_2Me), 2.16 (s, 3H, Me), 2.19 (s, 3H, Me), 3.59 (br, 2H, NH_2), 4.31 (q, $J = 7.1$ Hz, 2H, OCH_2), 6.65 (s, 1H, Ar), 6.80 (s, 1H, Ar), 7.02 (s, 1H, =CH), 7.14 (dd, 1H, $J = 5.0$ Hz, $J = 3.8$ Hz, H-4 Th), 7.54 (dd, 1H, $J = 3.8$ Hz, $J = 1.1$ Hz, H-3 Th), 7.67 (dd, 1H, $J = 5.0$ Hz, $J = 1.1$ Hz, H-5 Th), 8.11 (d, $J = 13.2$ Hz, 1H, =CHN), 12.23 (d, $J = 13.2$ Hz, 1H, NH), 14.32 (s, 1H, OH); ^{13}C NMR ($CDCl_3$, ppm) δ 14.1, 19.0, 19.4, 62.1, 103.5, 111.6, 119.5, 120.1, 124.8, 127.8, 128.7, 132.9, 133.3, 135.7, 136.3, 145.3, 153.9, 159.7, 162.7, 185.3, 193.0. Anal. calcd for $C_{21}H_{22}N_2O_5S$: C, 60.85; H, 5.35; N, 6.76. Found: C, 60.51; H, 5.65; N, 6.65.

(2Z,5E)-Ethyl-5-benzoyl-2-hydroxy-4-oxo-6-(pyridin-2-ylamino)hexa-2,5-dienoate (3k). Yield 74% (0.301 g), yellow powder, mp 159–160 °C. IR (ATR): 3219, 2995, 1741, 1622, 1590, 1563, 1253, 887 cm^{-1} . This compound exists in $CDCl_3$ as a mixture of tautomers *E-3k* : *Z-3k* : keto-**3k** = 80 : 13 : 7. 1H NMR ($CDCl_3$, ppm) δ : *E-3k* (80%) – 1.35 (t, $J = 7.1$ Hz, 3H, Me), 4.34 (q, $J = 7.1$ Hz, 2H, OCH_2), 7.00 (d, $J = 8.1$ Hz, 1H, Py), 7.10 (dd, $J = 7.3$ Hz, $J = 4.8$ Hz, 1H, Py), 7.12 (s, 1H, =CH), 7.51 (dd, $J = 7.7$ Hz, $J = 7.3$ Hz, 2H, H-3, H-5 Ph), 7.61 (tt, $J = 7.7$ Hz, $J = 1.4$ Hz, 1H, H-4 Ph), 7.73 (td, $J = 7.8$ Hz, $J = 1.3$ Hz, 1H, Py), 7.79 (2H, dd, $J = 7.9$ Hz, $J = 1.4$ Hz, H-2, H-6 Ph), 8.30 (dd, $J = 4.8$ Hz, $J = 1.3$ Hz, 1H, Py), 8.81 (d, $J = 12.5$ Hz, 1H, =CHN), 12.41 (d, $J = 12.5$ Hz, 1H, NH), 14.44 (s, 1H, OH); ^{13}C NMR ($CDCl_3$, ppm) δ 14.1, 62.2, 103.8, 112.3, 112.9, 120.7, 128.6, 129.6, 132.3, 138.8, 139.4, 149.0, 149.9, 151.2, 160.8, 162.5, 194.4, 194.8. Anal. calcd for $C_{20}H_{18}N_2O_5S$: C, 65.57; H, 4.95; N, 7.65. Found: C, 65.62; H, 4.76; N, 7.94.

(2Z,5E)-Ethyl-6-(1,5-dimethyl-3-oxo-2-phenyl-2,3-dihydro-1H-pyrazol-4-ylamino)-2-hydroxy-4-oxo-5-(thiophene-2-carbonyl)hexa-2,5-dienoate (3l). Yield 83% (0.444 g), yellow powder, mp 170–172 °C. IR (ATR): 3101, 1715, 1668, 1587, 1493, 1246, 1020, 910, 846 cm^{-1} . This compound exists in $CDCl_3$ as a mixture of tautomers *E-3l* : *Z-3l* : keto-**3l** = 92 : 4 : 4. 1H NMR ($CDCl_3$, ppm) δ : *E-3l* (92%) – 1.34 (t, $J = 7.1$ Hz, 3H, OCH_2Me), 2.38 (s, 3H, Me),

3.11 (s, 3H, Me), 4.32 (q, $J = 7.1$ Hz, 2H, OCH_2), 7.12 (dd, $J = 5.0$ Hz, $J = 3.9$ Hz, 1H, H-4 Th), 7.15 (s, 1H, =CH), 7.33–7.38 (m, 3H, Ph), 7.48 (dd, $J = 8.0$ Hz, $J = 7.4$ Hz, 2H, H-3, H-5 Ph), 7.60 (dd, $J = 5.0$ Hz, $J = 0.8$ Hz, 1H, H-5 Th), 7.64 (dd, $J = 3.9$ Hz, $J = 0.8$ Hz, 1H, H-3, Th), 9.03 (d, $J = 12.6$ Hz, 1H, =CHN), 12.52 (d, $J = 12.6$ Hz, 1H, NH), 14.27 (s, 1H, OH). Anal. calcd for $C_{24}H_{23}N_3O_6S$: C, 59.86; H, 4.81; N, 8.73. Found: C, 59.94; H, 4.70; N, 8.72.

(2Z,5E)-Ethyl-2-hydroxy-4-oxo-6-(pyrimidin-2-ylamino)-5-(thiophene-2-carbonyl)hexa-2,5-dienoate (3m). Yield 47% (0.195 g), light yellow powder, mp 204–206 °C. IR (ATR): 3272, 3096, 3083, 1736, 1616, 1595, 1562, 1434, 1248, 987, 843 cm^{-1} . This compound exists in $CDCl_3$ as a mixture of tautomers *E-3m* : *Z-3m* : keto-**3m** = 93 : 2 : 5. 1H NMR ($CDCl_3$, ppm) δ : *E-3m* (93%) – 1.35 (t, 3H, $J = 7.1$ Hz, Me), 4.34 (q, $J = 7.1$ Hz, 2H, OCH_2), 6.99 (s, 1H, =CH), 7.07 (t, $J = 4.8$ Hz, 1H, H-5 pyrimidine), 7.18 (dd, $J = 4.9$ Hz, $J = 3.8$ Hz, 1H, H-4 Th), 7.60 (dd, $J = 3.8$ Hz, $J = 0.8$ Hz, 1H, H-3 Th), 7.73 (dd, $J = 4.9$ Hz, $J = 0.8$ Hz, 1H, H-5 Th), 8.57 (d, $J = 4.8$ Hz, 2H, H-4, H-6 pyrimidine), 8.88 (d, $J = 12.6$ Hz, 1H, =CHN), 12.07 (d, $J = 12.6$ Hz, 1H, NH), 14.50 (s, 1H, OH). Anal. calcd for $C_{17}H_{15}N_3O_5S$: C, 54.68; H, 4.05; N, 11.25. Found: C, 54.79; H, 4.06; N, 11.04.

(2Z,5E)-Ethyl-5-((benzylamino)methylene)-2-hydroxy-7,7-dimethyl-4,6-dioxooct-2-enoate (3n). Yield 54% (0.215 g), yellow powder, mp 120–122 °C. IR (ATR): 3281, 2967, 1730, 1622, 1595, 1249, 1172, 862 cm^{-1} . This compound exists in $CDCl_3$ as a mixture of tautomers *E-3n* : *Z-3n* : keto-**3n** = 79 : 11 : 10. 1H NMR ($CDCl_3$, ppm) δ : *E-3n* (79%) – 1.20 (s, 9H, *t*-Bu), 1.34 (t, $J = 7.1$ Hz, 3H, Me), 4.32 (q, $J = 7.1$ Hz, 2H, OCH_2), 4.54 (d, $J = 6.0$ Hz, 2H, CH_2Ph), 6.63 (s, 1H, =CH), 7.28 (d, $J = 7.1$ Hz, 2H, H-2, H-6 Ph), 7.32–7.44 (3H, m, Ph), 7.69 (d, $J = 13.3$ Hz, 1H, =CHN), 10.67 (dt, $J = 13.3$ Hz, $J = 6.0$ Hz, 1H, NH), 14.43 (s, 1H, OH). Anal. calcd for $C_{20}H_{25}NO_5$: C, 66.84; H, 7.01; N, 3.90. Found: C, 67.00; H, 6.91; N, 4.11.

(2Z,5E)-Ethyl-5-benzoyl-6-(benzylamino)-2-hydroxy-4-oxohexa-2,5-dienoate (3o). Yield 43% (0.181 g), yellow powder, mp 137–139 °C. IR (ATR): 3241, 3032, 2987, 1729, 1593, 1258, 1021, 907 cm^{-1} . This compound exists in $CDCl_3$ as a mixture of tautomers *E-3o* : *Z-3o* : keto-**3o** = 68 : 19 : 13. 1H NMR ($CDCl_3$, ppm) δ : *E-3o* (68%) – 1.31 (t, $J = 7.1$ Hz, 3H, Me), 4.29 (q, $J = 7.1$ Hz, 2H, OCH_2), 4.54 (d, $J = 6.0$ Hz, 2H, CH_2Ph), 7.01 (s, 1H, =CH), 7.27 (dd, $J = 6.9$ Hz, $J = 1.3$ Hz, 2H, H-2, H-6 $PhCH_2$), 7.31–7.52 (5H, m, $PhCH_2$, Ph), 7.53 (t, $J = 7.5$ Hz, 1H, H-4 Ph), 7.63 (dd, $J = 7.8$ Hz, $J = 1.4$ Hz, 1H, H-2, H-6 Ph), 7.69 (d, $J = 13.7$ Hz, 1H, =CHN), 10.95 (dt, $J = 13.7$ Hz, $J = 6.0$ Hz, 1H, NH), 14.43 (s, 1H, OH). Anal. calcd for $C_{22}H_{21}NO_5$: C, 69.64; H, 5.58; N, 3.69. Found: C, 69.48; H, 5.61; N, 3.97.

General method for the preparation of polycarbonyl compounds 4 and 6

Method A. Pyrone **1** (0.26 mmol) was added to EtOH (2 mL) containing compound **3** (0.26 mmol) under cooling on an ice bath. The resulted mixture was stirred at 0 °C for 1 h and then at 20 °C for 24 h. The precipitate was filtered and washed with EtOH.

Method B. Pyrone **1** (0.72 mmol) was added to EtOH (4 mL) containing diamine **2** (0.36 mmol) under cooling on an ice bath. The resulted mixture was stirred at 0 °C for 1 h and then

at 20 °C for 24 h. The precipitate was filtered and washed with EtOH.

(2*Z*,2'*Z*,5*E*,5'*E*)-Diethyl-6,6'-(1,2-phenylenebis(azanediyl)) bis(5-benzoyl-2-hydroxy-4-oxohexa-2,5-dienoate) (4a). Yield 64% (0.109 g, Method A), 57% (0.134 g, Method B), yellow powder, mp 148–150 °C. IR (ATR): 2987, 1728, 1716, 1609, 1571, 1247 cm⁻¹; ¹H NMR (CDCl₃, ppm) δ: (5*E*,5'*E*)-4a (50%) – 1.34 (t, *J* = 7.1 Hz, 3H, Me), 4.31 (q, *J* = 7.1 Hz, 2H, OCH₂), 6.94 (s, 1H, =CH), 7.13–7.22 (m, 1H, Ar), 7.24–7.33 (m, 1H, Ar), 7.50 (2H, dd, *J* = 7.9 Hz, *J* = 7.2 Hz, H-3, H-5 Ph), 7.55–7.66 (m, 1H, H-4 Ph), 7.75–7.81 (m, 2H, H-2, H-6 Ph), 7.98 (d, *J* = 12.5 Hz, 1H, =CHN), 12.58 (d, *J* = 12.5 Hz, 1H, NH), 14.18 (s, 1H, OH); ¹³C NMR (DMSO-*d*₆, ppm) δ 14.0, 62.1, 103.7, 113.2, 120.3, 127.2, 129.1, 129.4, 131.5, 132.4, 139.4, 154.6, 160.8, 162.3, 193.9, 194.4. Anal. calcd for C₃₆H₃₂N₂O₁₀: C, 66.25; H, 4.94; N, 4.29. Found: C, 66.38; H, 4.86; N, 4.29.

(2*Z*,2'*Z*,5*E*,5'*E*)-Diethyl-6,6'-(1,2-phenylenebis(azanediyl)) bis(2-hydroxy-5-(4-methoxybenzoyl)-4-oxohexa-2,5-dienoate) (4b). Yield 73% (0.135 g, Method A), 38% (0.098 g, Method B), yellow powder, mp 139–141 °C. IR (ATR): 3134, 2988, 2839, 1735, 1725, 1594, 1564, 1248, 1032, 839 cm⁻¹; ¹H NMR (CDCl₃, ppm) δ: (5*E*,5'*E*)-4b (69%) – 1.31 (t, *J* = 7.1 Hz, 3H, Me), 3.88 (s, 3H, OMe), 4.29 (q, *J* = 7.1 Hz, 2H, OCH₂), 6.82 (s, 1H, =CH), 6.96 (d, AA'BB', *J* = 8.8 Hz, 2H, H-3, H-5 Ar'), 7.14–7.30 (m, 2H, Ar), 7.77 (d, *J* = 8.8 Hz, AA'BB', 2H, H-2, H-6 Ar'), 7.92 (d, *J* = 12.5 Hz, 1H, =CHN), 12.49 (d, *J* = 12.5 Hz, 1H, NH), 14.18 (s, 1H, OH); ¹³C NMR (DMSO-*d*₆, ppm) δ 14.0, 55.5, 62.1, 103.7, 113.9, 120.2, 127.5, 131.5, 131.9, 132.2, 153.6, 156.8, 160.6, 162.4, 163.3, 192.7, 194.2. Anal. calcd for C₃₀H₃₆N₂O₁₂: C, 64.04; H, 5.09; N, 3.93. Found: C, 63.78; H, 4.92; N, 4.05.

(2*Z*,2'*Z*,5*E*,5'*E*)-Diethyl-6,6'-(1,2-phenylenebis(azanediyl)) bis(5-(4-chlorobenzoyl)-2-hydroxy-4-oxohexa-2,5-dienoate) (4c). Yield 65% (0.122 g, Method A), 75% (0.195 g, Method B), yellow powder, mp 140–141 °C. IR (ATR): 3147, 3063, 2987, 2924, 1737, 1724, 1613, 1253 cm⁻¹; ¹H NMR (CDCl₃, ppm) δ: (5*E*,5'*E*)-4c (39%) – 1.32 (t, *J* = 7.1 Hz, 3H, Me), 4.29 (q, *J* = 7.1 Hz, 2H, OCH₂), 6.85 (s, 1H, =CH), 7.15–7.40 (m, 2H, Ar), 7.39–7.50 (m, 2H, Ar'), 7.70 (d, *J* = 8.5 Hz, AA'BB', 2H, H-3, H-5 Ar'), 7.93 (d, *J* = 12.6 Hz, 1H, =CHN), 12.55 (d, *J* = 12.6 Hz, 1H, NH), 14.09 (s, 1H, OH). ¹³C NMR (CDCl₃, ppm) δ 14.0, 62.2, 103.6, 113.0, 120.3, 127.9, 129.0, 129.8, 130.8, 137.6, 138.8, 154.5, 160.9, 162.2, 192.5, 194.1. Anal. calcd for C₃₆H₃₀Cl₂N₂O₁₀: C, 59.93; H, 4.19; N, 3.88. Found: C, 59.64; H, 3.82; N, 4.15.

(2*Z*,2'*Z*,5*E*,5'*E*)-Diethyl-6,6'-(1,2-phenylenebis(azanediyl)) bis(2-hydroxy-4-oxo-5-(thiophene-2-carbonyl)hexa-2,5-dienoate) (4d). Yield 75% (0.130 g, Method A), 31% (0.074 g, Method B), yellow powder, mp 134–136 °C. IR (ATR): 3097, 2983, 1718, 1591, 1569, 1243, 842 cm⁻¹; ¹H NMR (CDCl₃, ppm) δ: (5*E*,5'*E*)-4d (71%) – 1.34 (t, *J* = 7.1 Hz, 3H, Me), 4.31 (q, *J* = 7.1 Hz, 2H, OCH₂), 6.95 (s, 1H, =CH), 7.15 (dd, *J* = 4.8 Hz, *J* = 3.8 Hz, 1H, H-4 Th), 7.23–7.33 (m, 2H, Ar), 7.59 (dd, *J* = 3.8 Hz, *J* = 0.8 Hz, 1H, H-3 Th), 7.70 (dd, *J* = 4.8 Hz, *J* = 0.8 Hz, 1H, H-5 Th), 8.09 (d, *J* = 12.4 Hz, 1H, =CHN), 12.48 (d, *J* = 12.4 Hz, 1H, NH), 14.12 (s, 1H, OH); ¹³C NMR (CDCl₃, ppm) δ 14.1, 62.2, 103.2, 113.5, 120.4, 127.6, 128.1, 131.5, 133.4, 134.0, 144.7, 153.0, 160.8, 162.3, 185.0, 193.7. Anal. calcd for C₃₂H₂₈N₂O₁₀S₂: C, 57.82; H, 4.25; N, 4.21. Found: C, 57.81; H, 4.20; N, 4.12.

(2*Z*,2'*Z*,5*E*,5'*E*)-Diethyl-6,6'-(4,5-dimethyl-1,2-phenylene) bis(azanediyl)bis(2-hydroxy-4-oxo-5-(thiophene-2-carbonyl)hexa-2,5-dienoate) (4e). Yield 79% (0.142 g, Method A), 63% (0.157 g, Method B), red powder, mp 174–176 °C. IR (ATR): 3148, 2984, 1742, 1726, 1610, 1566, 1411, 1238, 842 cm⁻¹; ¹H NMR (CDCl₃, ppm) δ: (5*E*,5'*E*)-4e (74%) – 1.33 (t, *J* = 7.1 Hz, 3H, OCH₂Me), 2.26 (s, 3H, Me), 4.31 (q, *J* = 7.1 Hz, 2H, OCH₂), 6.93 (s, 1H, =CH), 7.00 (br, 1H, Ar), 7.14 (dd, *J* = 5.0 Hz, *J* = 3.8 Hz, 1H, H-4 Th), 7.57 (dd, *J* = 3.8 Hz, *J* = 0.9 Hz, 1H, H-3 Th), 7.69 (dd, *J* = 5.0 Hz, *J* = 0.9 Hz, 1H, H-5 Th), 8.05 (d, *J* = 12.6 Hz, 1H, =CHN), 12.41 (d, *J* = 12.6 Hz, 1H, NH), 14.14 (s, 1H, OH); ¹³C NMR (CDCl₃, ppm) δ 14.0, 19.5, 62.1, 103.3, 113.0, 121.5, 128.0, 129.1, 133.4, 133.9, 136.9, 144.9, 153.4, 160.5, 162.4, 185.0, 193.5. Anal. calcd for C₃₄H₃₂N₂O₁₀S₂: C, 58.95; H, 4.66; N, 4.04. Found: C, 58.90; H, 4.77; N, 4.05.

Binuclear copper(II) complex 5

Cu(OAc)₂ (0.028 g, 0.15 mmol) was added to solution of polycarbonyl compounds **4c** (0.050 g, 0.069 mmol) in DMF (1 mL). The reaction mixture was stirred for 10 min at room temperature. The precipitate that formed was filtered and washed with EtOH. Yield 87% (0.051 g), red-brown powder, mp > 330 °C. ¹H NMR (DMSO-*d*₆, ppm) (selected signals) δ 1.10 (br, 3H, Me), 3.76 (br, 2H, CH₂), 5.74 (br, 1H, =CH), 7.24–7.80 (m, 6H, Ar). IR (ATR): 2982, 1733, 1723, 1713, 1631, 1569, 1338, 1630, 1393, 1340, 1247, 995, 911, 816, 749 cm⁻¹. Anal. calcd for C₃₆H₂₆Cl₂Cu₂N₂O₁₀: C, 51.19; H, 3.10; N, 3.32. Found: C, 51.06; H, 3.11; N, 3.62. HRMS (ESI) calcd for C₃₆H₂₇N₂O₁₀Cl₂Cu₂ [M + H]⁺ 842.9635, found 842.9622.

(2*Z*,2'*Z*,5*E*,5'*E*)-Diethyl-6,6'-(1,4-phenylenebis(azanediyl)) bis(5-benzoyl-2-hydroxy-4-oxohexa-2,5-dienoate) (6). Yield 99% (0.233 g, Method B), yellow powder, mp 178–180 °C. IR (ATR): 3138, 1729, 1572, 1454, 1246, 1102, 834 cm⁻¹; this compound exists in CDCl₃ as a mixture of tautomers (5*E*,5'*E*)-**6** : (5*E*,5'*Z*)-**6** : keto-(5*E*,5'*E*)-**6** = 84 : 8 : 8. ¹H NMR (CDCl₃, ppm) δ: (5*E*,5'*E*)-**6** (84%) – 1.33 (3H, t, *J* = 7.1 Hz, Me), 4.31 (2H, q, *J* = 7.1 Hz, OCH₂), 6.94 (1H, s, =CH), 7.15 (2H, s, Ar), 7.50 (dd, *J* = 7.8 Hz, *J* = 7.3 Hz, 2H, H-3, H-5 Ph), 7.57–7.60 (m, 1H, H-4 Ph), 7.75 (d, 2H, *J* = 7.3 Hz, H-2, H-6 Ph), 8.02 (d, *J* = 13.2 Hz, 1H, =CHN), 12.46 (d, *J* = 13.2 Hz, 1H, NH), 14.27 (s, 1H, OH). Anal. calcd for C₃₆H₃₂N₂O₁₀: C, 66.25; H, 4.94; N, 4.29. Found: C, 66.14; H, 4.81; N, 4.45.

General approach for the preparation of 6*a*-hydroxy-9-acyl-6*a*,7-dihydro-5*H*-pyrido[1,2-*a*]quinoxaline-6,8-diones 7

Enaminone **3e,f,h,j** (0.259 mmol) was refluxed in EtOH (2 mL) for 20 min. The precipitate that formed was filtered and washed with EtOH.

6*a*-Hydroxy-9-(4-methoxybenzoyl)-6*a*,7-dihydro-5*H*-pyrido[1,2-*a*]quinoxaline-6,8-dione (7a). Yield 72% (0.0679 g), light yellow powder, mp > 300 °C. IR (ATR): 3067, 2915, 1692, 1677, 1598, 1541, 1248, 1174, 758 cm⁻¹; ¹H NMR (DMSO-*d*₆, ppm) δ 2.95 (d, *J* = 16.6 Hz, 1H, CHH), 3.40 (d, *J* = 16.7 Hz, 1H, CHH), 3.84 (s, 3H, MeO), 6.99 (d, AA'BB', *J* = 8.9 Hz, 2H, H-3, H-5 Ar), 7.00–7.13 (m, 2H), 7.18 (td, *J* = 7.9 Hz, 1H), 7.54 (d, *J* = 7.9 Hz, 1H), 7.71 (d, AA'BB', *J* = 8.6 Hz, 2H, H-2, H-6 Ar), 7.76 (s, 1H,

OH), 8.41 (s, 1H, H-10), 11.08 (s, 1H, NH). Anal. calcd for $C_{20}H_{16}N_2O_5$: C, 66.01; H, 7.36; N, 4.41. Found: C, 65.93; H, 7.69; N, 4.43.

9-(4-Chlorobenzoyl)-6 α -hydroxy-6 α ,7-dihydro-5H-pyrido[1,2-*a*]quinoxaline-6,8-dione (7b). Yield 89% (0.0850 g), yellow powder, mp > 320 °C. IR (ATR): 3132, 3061, 3988, 2906, 2870, 2763, 1696, 1679, 1621, 1543, 1250, 749 cm^{-1} ; 1H NMR (DMSO- d_6 , ppm) δ 2.97 (d, $J = 16.7$ Hz, 1H, CHH), 3.30 (d, $J = 16.7$ Hz, 1H, CHH), 7.04–7.12 (m, 2H, Ar), 7.14–7.19 (m, 1H), 7.45 (d, AA'BB', $J = 8.5$ Hz, 2H, H-2, H-6 Ar), 7.50 (d, $J = 7.9$ Hz, 1H), 7.65 (2H, d, AA'BB', $J = 8.6$ Hz, H-3, H-5 Ar), 7.75 (s, 1H, OH), 8.46 (s, 1H, H-10), 11.02 (s, 1H, NH); ^{13}C NMR (DMSO- d_6 , ppm) δ 45.0, 82.5, 114.0, 116.3, 117.3, 123.5, 125.2, 125.7, 127.9, 128.3, 130.9, 136.7, 137.8, 149.4, 163.3, 187.2, 190.1. Anal. calcd for $C_{19}H_{13}ClN_2O_4$: C, 61.88; H, 3.55; N, 7.60. Found: C, 61.65; H, 3.25; N, 7.77.

6 α -Hydroxy-9-(thiophene-2-carbonyl)-6 α ,7-dihydro-5H-pyrido[1,2-*a*]quinoxaline-6,8-dione (7c). Yield 86% (0.0758 g), beige powder, mp > 320 °C. 1H NMR (DMSO- d_6 , ppm) δ 3.01 (d, $J = 16.6$ Hz, 1H, CHH), 3.38 (d, $J = 16.6$ Hz, 1H, CHH), 7.05–7.14 (m, 2H, Ar), 7.15–7.23 (m, 2H, Th, Ar), 7.54 (d, $J = 8.0$ Hz, 1H), 7.50 (d, $J = 7.9$ Hz, 1H), 7.71–7.80 (m, 2H, OH, Th), 7.95 (d, 1H, $J = 4.8$ Hz, H-5 Th), 8.49 (s, 1H, H-10), 11.07 (s, 1H, NH); ^{13}C NMR (DMSO- d_6 , ppm) δ 45.1, 82.3, 114.5, 116.3, 117.2, 123.5, 125.35, 125.44, 128.2 (2C), 134.2, 134.6, 144.5, 148.1, 163.3, 182.5, 187.0. Anal. calcd for $C_{17}H_{10}N_2O_3S$: C, 59.99; H, 3.55; N, 8.23. Found: C, 59.73; H, 3.58; N, 8.24.

6 α -Hydroxy-2,3-dimethyl-9-(thiophene-2-carbonyl)-6 α ,7-dihydro-5H-pyrido[1,2-*a*]quinoxaline-6,8-dione (7d). Yield 85% (0.0811 g), orange powder, mp > 300 °C. IR (ATR): 3179, 3082, 2976, 1722, 1674, 1591, 1539, 1514, 1312, 1044, 841 cm^{-1} ; 1H NMR (DMSO- d_6 , ppm) δ 2.18 (s, 3H, Me), 2.20 (s, 3H, Me), 2.96 (d, $J = 16.7$ Hz, 1H, CHH), 3.36 (d, $J = 16.7$ Hz, 1H, CHH), 6.81 (s, 1H), 7.20 (dd, $J = 4.9$ Hz, $J = 3.9$ Hz, 1H, H-4 Th), 7.37 (s, 1H), 7.70 (s, 1H, OH), 7.75 (dd, $J = 3.9$ Hz, $J = 1.0$ Hz, 1H, H-3 Th), 7.95 (dd, $J = 4.9$ Hz, $J = 1.0$ Hz, 1H, H-5 Th), 8.47 (s, 1H, H-10), 10.93 (s, 1H, NH); ^{13}C NMR (DMSO- d_6 , ppm) δ 18.8 (2Me), 45.1, 82.4, 114.2, 117.0, 117.9, 123.0, 125.8, 128.1, 131.6, 133.6, 134.1, 134.7, 144.6, 148.1, 163.3, 182.5, 186.8. Anal. calcd for $C_{19}H_{16}N_2O_4S$: C, 61.94; H, 4.38; N, 7.60. Found: C, 61.95; H, 4.35; N, 7.74.

General approach for the preparation of 9-acyl-5H-pyrido[1,2-*a*]quinoxaline-6,8-diones 8

Compound 7 (0.147 mmol) was refluxed in AcOH (2 mL) for 20 min. After cooling, the reaction mixture was diluted with H_2O (3 mL). The precipitate that formed was filtered and washed with cold EtOH. The product was dried at 100 °C.

9-(4-Methoxybenzoyl)-5H-pyrido[1,2-*a*]quinoxaline-6,8-dione (8a). Yield 89% (0.0453 g), grey powder, mp > 300 °C. IR (ATR): 3677, 2973, 2901, 1687, 1683, 1605, 1591, 1575, 1242, 1017, 850 cm^{-1} ; 1H NMR (DMSO- d_6 , ppm) δ 3.85 (s, 3H, Me), 7.05 (d, AA'BB', $J = 8.9$ Hz, 2H, H-3, H-5 Ar), 7.12 (s, 1H, H-7), 7.24 (ddd, $J = 8.0$ Hz, $J = 7.6$ Hz, $J = 1.4$ Hz, 1H, H-2), 7.29 (dd, $J = 8.0$ Hz, $J = 1.4$ Hz, 1H, H-4), 7.38 (td, $J = 7.6$ Hz, $J = 0.6$ Hz, 1H, H-3), 7.84 (d, AA'BB', $J = 8.6$ Hz, 2H, H-2, H-6 Ar), 8.20 (dd, $J = 8.6$ Hz, $J = 0.8$ Hz, 1H, H-1), 9.12 (s, 1H, H-10), 12.06 (s, 1H, NH); ^{13}C NMR

(DMSO- d_6 , ppm) δ 56.0, 113.9, 115.5, 117.0, 117.2, 123.1, 123.7, 127.1, 127.5, 129.5, 131.9, 132.6, 133.4, 137.5, 154.7, 163.6, 174.6, 191.8. Anal. calcd for $C_{20}H_{14}N_2O_5$: C, 69.36; H, 4.07; N, 8.09. Found: C, 69.51; H, 4.41; N, 8.35.

9-(4-Chlorobenzoyl)-5H-pyrido[1,2-*a*]quinoxaline-6,8-dione (8b). Yield 89% (0.0459 g), beige powder, mp > 300 °C. IR (ATR): 3063, 3036, 2987, 2905, 2841, 1710, 1241, 838 cm^{-1} ; 1H NMR (DMSO- d_6 , ppm) δ 7.15 (s, 1H, H-7), 7.23 (td, $J = 7.8$ Hz, $J = 1.6$ Hz, 1H, H-2), 7.30 (dd, $J = 7.8$ Hz, $J = 1.6$ Hz, 1H, H-4), 7.36 (td, $J = 7.5$ Hz, $J = 0.6$ Hz, 1H, H-3), 7.55 (d, AA'BB', $J = 8.6$ Hz, 2H, H-3, H-5 Ar), 7.84 (d, AA'BB', $J = 8.6$ Hz, 2H, H-2, H-6 Ar), 8.16 (d, $J = 8.5$ Hz, 1H, H-1), 9.14 (s, 1H, H-10), 12.04 (s, 1H, NH). Anal. calcd for $C_{19}H_{11}ClN_2O_3$: C, 65.06; H, 3.16; N, 7.99. Found: C, 64.99; H, 3.38; N, 7.93.

9-(Thiophene-2-carbonyl)-5H-pyrido[1,2-*a*]quinoxaline-6,8-dione (8c). Yield 92% (0.0436 g), beige powder, mp > 300 °C. IR (ATR): 3236, 3099, 2922, 1705, 1701, 1561, 1380, 900 cm^{-1} ; 1H NMR (DMSO- d_6 , ppm) δ 7.14 (s, 1H, H-7), 7.22–7.26 (m, 2H, H-4 Th, H-2), 7.29 (1H, dd, $J = 8.0$ Hz, $J = 1.3$ Hz, H-4), 7.38 (1H, td, $J = 7.5$ Hz, $J = 0.5$ Hz, H-3), 7.74 (dd, $J = 3.8$ Hz, $J = 1.0$ Hz, 1H, H-3 Th), 8.11 (dd, $J = 4.5$ Hz, $J = 1.0$ Hz, 1H, H-5 Th), 8.19 (d, 1H, $J = 8.5$ Hz, H-1), 9.20 (s, 1H, H-10), 12.07 (s, 1H, NH); ^{13}C NMR (DMSO- d_6 , ppm) δ 115.5, 117.0, 117.5, 123.1, 123.7, 127.2, 127.5, 128.8, 131.7, 133.7, 136.1, 136.2, 137.5, 143.7, 154.6, 174.2, 185.0. Anal. calcd for $C_{17}H_{10}N_2O_3S$: C, 63.34; H, 3.13; N, 8.69. Found: C, 63.32; H, 3.38; N, 8.57.

2,3-Dimethyl-9-(thiophene-2-carbonyl)-5H-pyrido[1,2-*a*]quinoxaline-6,8-dione (8d). Yield 80% (0.0412 g), light yellow powder, mp > 300 °C. IR (ATR): 3241, 3177, 3073, 2943, 1693, 1642, 1605, 1351, 1250, 848 cm^{-1} ; 1H NMR (DMSO- d_6 , ppm) δ 2.23 (s, 3H, Me), 2.27 (s, 3H, Me), 7.02 (s, 1H), 7.16 (s, 1H), 7.24 (dd, $J = 4.5$ Hz, $J = 3.2$ Hz, 1H, H-4 Th), 7.76 (d, $J = 3.2$ Hz, 1H, H-3 Th), 7.97 (s, 1H, H-1), 8.07 (d, $J = 4.5$ Hz, 1H, H-5 Th), 9.10 (s, 1H, H-10), 11.80 (s, 1H, NH). Anal. calcd for $C_{19}H_{14}N_2O_3S$: C, 65.13; H, 4.03; N, 7.99; S, 4.93. Found: C, 65.17; H, 3.94; N, 8.08.

Ethyl-5-benzoyl-4-oxo-1-phenyl-1,4-dihydropyridine-2-carboxylate (9a). Enaminone 3a (0.100 g, 0.274 mmol) and $MeSO_3H$ (0.0526 g, 0.547 mmol) were stirred in CH_2Cl_2 (2 mL) for 24 h. The solvent was evaporated, and the residue was diluted with H_2O (5 mL). The precipitate that formed was filtered and was recrystallized from toluene-hexane. Yield 78% (0.074 g), white powder, mp 112–114 °C. IR (ATR): 3055, 2970, 1722, 1640, 1592, 1491, 1241, 912 cm^{-1} ; 1H NMR ($CDCl_3$, ppm) δ 1.09 (t, $J = 7.1$ Hz, 3H, Me), 4.13 (q, $J = 7.1$ Hz, 2H, OCH_2), 7.06 (s, 1H, H-3), 7.30–7.36 (m, 2H, H-2, H-6 Ph), 7.45 (t, $J = 7.6$ Hz, 2H, H-3, H-5 Ph), 7.48–7.54 (m, 3H, Ph, Ph'), 7.57 (tt, $J = 7.4$ Hz, $J = 1.9$ Hz, 1H, H-4 Ph'), 7.89 (dd, $J = 8.5$ Hz, $J = 1.4$ Hz, 2H, H-2, H-6 Ph'), 7.91 (s, 1H, H-6). Anal. calcd for $C_{21}H_{17}NO_4$: C, 72.61; H, 4.93; N, 4.03. Found: C, 72.33; H, 4.81; N, 4.32.

Ethyl-5-(4-chlorobenzoyl)-1-(2-hydroxyphenyl)-4-oxo-1,4-dihydropyridine-2-carboxylate (9b). Compound 3d (0.300 g, 0.721 mmol) was stirred in EtOH (9 mL) containing concd HCl (0.2 mL) for 12 h at room temperature. The precipitate that formed was filtered. Yield 79% (0.227 g), light white powder, mp 245–246 °C. IR (ATR): 3060, 2983, 2902, 2740, 2576, 1728, 1665, 1632, 1585, 1552, 1528 cm^{-1} ; 1H NMR (DMSO- d_6 , ppm) δ 1.05 (t, $J = 7.1$ Hz, 3H, Me), 4.09 (q, $J = 7.1$ Hz, 2H, OCH_2), 6.77 (s, 1H,

H-3), 6.91 (td, $J = 7.7$ Hz, $J = 1.2$ Hz, 1H, H-5 Ar), 6.97 (dd, $J = 8.2$ Hz, $J = 1.1$ Hz, 1H, H-3 Ar), 7.26 (td, $J = 8.0$ Hz, $J = 1.1$ Hz, 1H, H-4 Ar), 7.33 (dd, $J = 7.8$ Hz, $J = 1.4$ Hz, 1H, H-6 Ar), 7.49 (d, AA'BB', $J = 8.4$ Hz, 2H, H-3, H-5 Ar'), 7.77 (d, AA'BB', $J = 8.4$ Hz, 2H, H-2, H-6 Ar'), 7.89 (s, 1H, H-6), 10.35 (s, 1H, OH). Anal. calcd for C₂₁H₁₆ClNO₅: C, 63.40; H, 4.05; N, 3.52. Found: C, 63.12; H, 4.04; N, 3.73.

(2E,4Z)-2-((2-Hydroxyphenylamino)methylene)-4-(2-oxo-2H-benzo[*b*]1,4-oxazin-3(4H)-ylidene)-1-phenylbutane-1,3-dione (10). Enaminone **3c** (0.100 g, 0.262 mmol) and *o*-aminophenol (0.030 g, 0.275 mmol) were refluxed in C₆H₆ (3 mL) for 9 h. The precipitate that formed was filtered. Yield 85% (0.095 mg), light white powder, mp 188–189 °C. IR (ATR): 3169, 3151, 3115, 3069, 2744, 2703, 1671, 1628, 1543, 1382 cm⁻¹; ¹H NMR (DMSO-*d*₆, ppm) δ 6.63 (s, 1H, =CH), 6.68 (t, $J = 7.6$ Hz, 1H, Ar), 6.80–6.87 (m, 2H, Ar), 6.93 (t, $J = 6.9$ Hz, 1H, Ar), 6.97 (d, $J = 8.1$ Hz, 1H, Ar), 7.22 (t, $J = 7.7$ Hz, 1H, Ar), 7.42 (d, $J = 7.8$ Hz, 1H, Ar), 7.44–7.51 (m, 3H, Ar, Ph), 7.59 (t, $J = 7.3$ Hz, 1H, H-4 Ph), 7.77 (s, 1H, =CHN), 7.83 (2H, d, $J = 7.7$ Hz, H-2, H-6 Ph), 9.56 (br, 1H, NH), 9.70 (s, 1H, OH), 10.25 (br, 1H, NH). Anal. calcd for C₂₅H₁₈N₂O₅: C, 70.42; H, 4.25; N, 6.57. Found: C, 70.09; H, 4.09; N, 6.57.

6 α -Hydroxy-9-(thiophene-2-carbonyl)-6 α ,7-dihydro-5H-dipyr-ido[1,2-*a*:2',3'-*e*]pyrazine-6,8-dione (11). General approach for the preparation of enaminodiones **3** from ethyl 4-oxo-5-(thiophene-2-carbonyl)-4H-pyran-2-carboxylate (**1e**) and 2,3-diaminopyridine. Yield 91% (0.345 g), dark grey powder, mp > 300 °C. IR (ATR): 3153, 1703, 1657, 1608, 1557, 1406, 1328, 1038, 842 cm⁻¹; ¹H NMR (DMSO-*d*₆, ppm) δ 3.00 (d, $J = 16.6$ Hz, 1H, CHH), 3.42 (d, $J = 16.6$ Hz, 1H, CHH), 7.15 (dd, $J = 8.1$ Hz, $J = 4.8$ Hz, 1H, H-2), 7.21 (dd, $J = 4.9$ Hz, $J = 3.8$ Hz, 1H, H-4 Th), 7.74 (dd, $J = 3.8$ Hz, $J = 1.1$ Hz, 1H, H-3 Th), 7.80–7.98 (s, 1H, OH), 7.98 (dd, 1H, $J = 4.9$, $J = 1.1$, H-5 Th), 8.03 (dd, $J = 8.5$ Hz, $J = 0.8$ Hz, 1H, H-1), 8.12 (1H, dd, $J = 4.9$, $J = 1.1$, H-3), 8.50 (1H, s, H-10), 11.60 (1H, s, NH); ¹³C NMR (DMSO-*d*₆, ppm) δ 44.9, 82.3, 115.6, 119.2, 121.6, 124.6, 128.3, 134.5, 134.9, 141.5, 143.7, 144.4, 147.2, 164.1, 182.5, 187.0. Anal. calcd for C₁₆H₁₁N₃O₄S: C, 56.30; H, 3.25; N, 12.31. Found: C, 56.12; H, 3.31; N, 12.08.

9-(Thiophene-2-carbonyl)-5H-dipyr-ido[1,2-*a*:2',3'-*e*]pyrazine-6,8-dione (12). General approach for the preparation of compounds **8**. Yield 81% (0.0406 g), grey powder, mp > 300 °C. IR (ATR): 3075, 2744, 1704, 1623, 1547, 1256, 802 cm⁻¹; ¹H NMR (DMSO-*d*₆, ppm) δ 7.12 (s, 1H, H-7), 7.25 (dd, $J = 4.7$ Hz, $J = 3.7$ Hz, 1H, H-4 Th), 7.31 (dd, $J = 8.4$ Hz, $J = 4.7$ Hz, 1H, H-2), 7.73 (dd, $J = 3.7$ Hz, $J = 0.7$ Hz, 1H, H-3 Th), 8.12 (dd, $J = 4.7$ Hz, $J = 0.7$ Hz, 1H, H-5 Th), 8.33 (dd, $J = 4.7$ Hz, $J = 0.7$ Hz, 1H, H-3), 8.59 (dd, $J = 8.4$ Hz, $J = 0.7$ Hz, 1H, H-1), 9.18 (s, 1H, H-10), 12.52 (s, 1H, NH), 12.26 (s, 1H, OH); ¹³C NMR (DMSO-*d*₆, ppm) δ 117.4, 119.3, 120.0, 123.6, 128.9, 131.8, 133.8, 136.31, 136.34, 137.3, 140.6, 143.6, 146.0, 155.9, 174.4, 184.9. Anal. calcd for C₁₆H₉N₃O₃S·H₂O: C, 56.30; H, 3.25; N, 12.31. Found: C, 56.31; H, 2.95; N, 12.12.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was financially supported by the Russian Science Foundation (Grant 18-13-00186). Analytical studies were carried out using equipment of the Center for Joint Use “Spectroscopy and Analysis of Organic Compounds” at the Postovsky Institute of Organic Synthesis of the Russian Academy of Sciences (Ural Branch) and the Laboratory of Complex Investigations and Expert Evaluation of Organic Materials of the Center for Joint Use at the Ural Federal University. We also thank both reviewers for helpful suggestions.

Notes and references

- (a) P. R. Christensen, A. M. Scheuermann, K. E. Loeffler and B. A. Helms, *Nat. Chem.*, 2019, **11**, 442; (b) E. A. Shokova, J. K. Kim and V. V. Kovalev, *Russ. J. Org. Chem.*, 2015, **51**, 755.
- (a) N. V. Zolotareva and V. V. Semenov, *Russ. Chem. Rev.*, 2013, **82**, 964; (b) P. A. Vigato, V. Peruzzo and S. Tamburini, *Coord. Chem. Rev.*, 2009, **253**, 1099; (c) S. N. Podyachev, S. N. Sudakova, G. S. Gimazetdinova, N. A. Shamsutdinova, V. V. Syakaev, T. A. Barsukova, N. Iki, D. V. Lapaev and A. R. Mustafina, *New J. Chem.*, 2017, **41**, 1526; (d) A. J. Brock, J. K. Clegg, F. Li and L. F. Lindoy, *Coord. Chem. Rev.*, 2018, **375**, 106; (e) D. N. Bazhin, Y. S. Kudyakova, A. S. Bogomyakov, P. A. Slepukhin, G. A. Kim, Y. V. Burgart and V. I. Saloutin, *Inorg. Chem. Front.*, 2019, **6**, 40; (f) R. Pettinari, F. Marchetti, C. Di Nicola and C. Pettinari, *Eur. J. Inorg. Chem.*, 2018, **31**, 3521.
- See reviews: (a) K. C. Gupta and A. K. Sutar, *Coord. Chem. Rev.*, 2008, **252**, 1420; (b) P. A. Vigato, V. Peruzzo and S. Tamburini, *Coord. Chem. Rev.*, 2012, **256**, 953; (c) C. J. Whiteoak, G. Salassa and A. W. Kleij, *Chem. Soc. Rev.*, 2012, **41**, 622; (d) C. Freire, M. Nunes, C. Pereira, D. M. Fernandes, A. F. Peixoto and M. Rocha, *Coord. Chem. Rev.*, 2019, **394**, 104.
- (a) Y. Ren, J. Chen, C. Qi and H. Jiang, *ChemCatChem*, 2015, **7**, 1535; (b) S. He, F. Wang, W.-L. Tong, S.-M. Yiu and M. C. W. Chan, *Chem. Commun.*, 2016, **52**, 1017; (c) J. A. Castro-Osma, M. North and X. Wu, *Chem.–Eur. J.*, 2016, **22**, 2100; (d) X. Wu, J. Mason and M. North, *Chem.–Eur. J.*, 2017, **23**, 12937; (e) S. Vagin, M. Winnacker, A. Kronast, P. T. Altenbuchner, P. Deglmann, C. Sinkel, R. Loos and B. Rieger, *ChemCatChem*, 2015, **7**, 3963; (f) W. Luo, T. Shi, S. Liu, W. Zuo and Z. Li, *Organometallics*, 2017, **369**, 1736.
- (a) S. J. Wezenberg, E. C. Escudero-Adán, J. Benet-Buchholz and A. W. Kleij, *Org. Lett.*, 2008, **10**, 3311; (b) J. Cheng, X. Ma, Y. Zhang, J. Liu, X. Zhou and H. Xiang, *Inorg. Chem.*, 2014, **53**, 3210.
- C. Bizzarri, E. Spuling, D. M. Knoll, D. Volz and S. Bräse, *Coord. Chem. Rev.*, 2018, **373**, 49.
- A. K. Asatkar, S. P. Senanayak, A. Bedi, S. Panda, K. S. Narayan and S. S. Zade, *Chem. Commun.*, 2014, **50**, 7036.
- G. Salassa, J. W. Ryan, E. C. Escudero-Adána and A. W. Kleij, *Dalton Trans.*, 2014, **43**, 210.

- 9 (a) R. M. Clarke and T. Storr, *Dalton Trans.*, 2014, **43**, 9380; (b) M. Andruh, *Chem. Commun.*, 2011, **47**, 302; (c) P. Fuentealba, V. Paredes-García, D. Venegas-Yazigi, I. D. A. Silva, C. J. Magon, R. C. de Santana, N. Audebrand, J. Manzurh and E. Spodine, *RSC Adv.*, 2017, **7**, 33305.
- 10 K. T. Mahmudov and A. J. L. Pombeiro, *Chem.–Eur. J.*, 2016, **22**, 16356.
- 11 (a) Y. S. Kudyakova, D. N. Bazhin, M. V. Goryaeva, Y. V. Burgart and V. I. Saloutin, *Russ. Chem. Rev.*, 2014, **83**, 120; (b) P. Subramaniam, C. Ramasubbu and S. Athiramu, *Green Chem.*, 2017, **19**, 2541.
- 12 (a) P. P. Mukovoz, P. A. Slepukhin, E. A. Danilova, O. P. Aysuvakova and A. P. Glinushkin, *Russ. J. Gen. Chem.*, 2018, **88**, 1363; (b) N. A. Bailey, K. C. Cox, C. P. Falshaw, D. E. Fenton, S. E. Grundy, P. Haigh, C. A. Phillips and T. J. King, *Dalton Trans.*, 1983, 2241.
- 13 (a) D. E. Fenton and S. E. Gayda, *J. Chem. Soc., Dalton Trans.*, 1977, 2095; (b) D. E. Fenton and S. E. Gayda, *Dalton Trans.*, 1977, 2109; (c) C. K. Lai, A. G. Serrette and T. M. Swager, *J. Am. Chem. Soc.*, 1992, **114**, 7948; (d) R. P. Kreh, A. E. Rodriguez and S. P. Fox, *J. Chem. Soc., Chem. Commun.*, 1984, 130.
- 14 (a) D. L. Obydenov, L. R. Khammatova, O. S. Eltsov and V. Y. Sosnovskikh, *Org. Biomol. Chem.*, 2018, **16**, 1692; (b) D. L. Obydenov, L. R. Khammatova and V. Y. Sosnovskikh, *Mendeleev Commun.*, 2017, **27**, 172; (c) D. L. Obydenov, E. O. Pan'kina and V. Y. Sosnovskikh, *Chem. Heterocycl. Compd.*, 2017, **53**, 924; (d) D. L. Obydenov, E. O. Pan'kina and V. Y. Sosnovskikh, *J. Org. Chem.*, 2016, **81**, 12532.
- 15 D. L. Obydenov and V. Y. Sosnovskikh, *Chem. Heterocycl. Compd.*, 2014, **50**, 579.
- 16 H. C. Kolb, M. G. Finn and K. B. Sharpless, *Angew. Chem., Int. Ed.*, 2001, **40**, 2004.
- 17 (a) V. S. Berseneva, A. V. Tkachev, Y. Yu. Morzherin, W. Dehaen, I. Luyten, S. Toppet and V. A. Bakulev, *J. Chem. Soc., Perkin Trans. 1*, 1998, 2133; (b) V. S. Berseneva, Y. Y. Morzherin, W. Dehaen, I. Luyten and V. A. Bakulev, *Tetrahedron*, 2001, **57**, 2179; (c) A. Ranise, F. Lucchesini, M. Caviglia, S. Alfei, A. Spallarossa and C. Caneva, *Tetrahedron*, 2013, **69**, 10858; (d) J. Schreurs, C. A. H. van Noorden-Mudde, L. J. M. van de Ven and J. W. de Ham, *Org. Magn. Reson.*, 1980, **13**, 354.
- 18 (a) J. Saloň, V. Milata, A. Gatial, N. Prónayová, J. Leško, P. Černuchová, Z. Rappoport, G. Vo-Thanh and A. Loupy, *Eur. J. Org. Chem.*, 2005, 4870; (b) J. Pigošová, A. Gatial, V. Milata, P. Černuchová, N. Prónayová, T. Liptaj and P. Matějka, *J. Mol. Struct.*, 2005, **744–747**, 315; (c) A. Gatial, S. Dorotíková, K. Plevová, V. Milata, N. Prónayová and P. Matějka, *J. Mol. Struct.*, 2015, **1090**, 112.
- 19 C. Hametner, P. Černuchová, V. Milata, G. Vo-Thanh and A. Loupy, *Org. Magn. Reson.*, 2005, **43**, 171.
- 20 A. Brbot-Šaranović, G. Pavlović and M. Cindrić, *Struct. Chem.*, 2000, **1**, 65.
- 21 (a) A. V. Pestov, L. A. Khamidullina, V. Y. Sosnovskikh, P. A. Slepukhin and I. S. Pusyrev, *Polyhedron*, 2016, **106**, 75; (b) Y. Yu, J. Chen, S. Meng, C. Li, M. Lan and Z. Zhang, *J. Mol. Catal. A: Chem.*, 2013, **380**, 104; (c) T. Rogez-Florent, S. Meignan, C. Foulon, P. Six, A. Gros, C. Bal-Mahieu, C. T. Supuran, A. Scozzafava, R. Frédérick, B. Masereel, P. Depreux, A. Lansiaux, J.-F. Goossens, S. Gluszok and L. Goossens, *Bioorg. Med. Chem.*, 2013, **21**, 1451.
- 22 (a) J. H. Satcher and A. L. Balch, *Inorg. Chem.*, 1995, **34**, 3371; (b) D. L. Reger, A. E. Pascui, P. J. Pellechia and A. Ozarowski, *Inorg. Chem.*, 2013, **52**, 12741.
- 23 (a) V. Y. Sosnovskikh, A. V. Safrygin, R. A. Irgashev, M. A. Ezhikova and M. I. Kodess, *RSC Adv.*, 2016, **6**, 30056; (b) R. A. Irgashev, A. V. Safrygin, M. A. Ezhikova, M. I. Kodess, G.-V. Röschenhaler and V. Y. Sosnovskikh, *Tetrahedron*, 2015, **71**, 1822.
- 24 D. L. Obydenov, A. O. Goncharov and V. Y. Sosnovskikh, *Russ. Chem. Bull.*, 2016, **65**, 2233.