

# Approach to epigenetic analysis in language disorders

Shelley D. Smith

Received: 17 September 2011 / Accepted: 30 October 2011 / Published online: 24 November 2011  
© Springer Science+Business Media, LLC 2011

**Abstract** Language and learning disorders such as reading disability and language impairment are recognized to be subject to substantial genetic influences, but few causal mutations have been identified in the coding regions of candidate genes. Association analyses of single nucleotide polymorphisms have suggested the involvement of regulatory regions of these genes, and a few mutations affecting gene expression levels have been identified, indicating that the quantity rather than the quality of the gene product may be most relevant for these disorders. In addition, several of the candidate genes appear to be involved in neuronal migration, confirming the importance of early developmental processes. Accordingly, alterations in epigenetic processes such as DNA methylation and histone modification are likely to be important in the causes of language and learning disorders based on their functions in gene regulation. Epigenetic processes direct the differentiation of cells in early development when neurological pathways are set down, and mutations in genes involved in epigenetic regulation are known to cause cognitive disorders in humans. Epigenetic processes also regulate the changes in gene expression in response to learning, and alterations in histone modification are associated with learning and memory deficits in animals. Genetic defects in histone modification have been reversed in animals through therapeutic interventions resulting in rescue of these deficits, making it particularly important to investigate their potential contribution to learning disorders in humans.

**Keywords** Epigenetics · Dyslexia · Reading disability · Language impairment

## Introduction

Despite the complexity of language and learning disorders, individual genes are being defined which appear to influence the development of abilities that are necessary in speech, language, and reading. Most of the identified candidate genes involve reading disability, and although the evidence supporting some of these genes is still somewhat tenuous due to small sample sizes and limited replication, most are known to be involved in early development, particularly neuronal migration (Galaburda 2005; Gabel et al. 2010; Poelmans et al. 2011). As will be discussed below, most of these candidate genes have been associated with several learning and language phenotypes, suggesting that they facilitate learning processes which are basic to learning reading and language. Similar pleiotropic effects are seen for several genes that primarily affect autism or language but have also shown effects on reading, including CNTNAP2 and ATP2C2 (Vernes et al. 2008; Newbury et al. 2011). However, despite replicated evidence for association of single nucleotide polymorphisms within and around the genes, very few coding mutations have been reported to account for their influence on these disorders. This has led to the hypothesis that mutations affecting reading and related disorders are likely to be in regulatory regions, controlling the quantity rather than quality of the gene product (Bates et al. 2011). Alterations of gene expression can be caused by mutations in gene promoters and enhancers located near the gene, but mutations in genes that mediate epigenetic controls of gene expression have been found that affect developmental learning disorders.

---

Supported by NIH-NICHD 5P50HD027802 to R.K. Olson and NIH: NIDCD R01 NIH0064386 to M.L. Rice

---

S. D. Smith (✉)  
University of Nebraska Medical Center,  
Omaha, NE, USA  
e-mail: ssmith@unmc.edu

These mutations may be in regions located further from the target gene, making it more difficult to recognize their significance.

#### Regulatory regions of genes influencing language and learning disorders

Of all of the genes that have been proposed as candidates for reading disability and language impairment, six genes have been well characterized with respect to their influence on reading and language disorders, the regions within and around the genes that appear to contain causal mutations, and the effects of the putative mutations or risk alleles on gene transcription: *DYX1C1*, *DCDC2*, *KIAA0319*, *ROBO1*, and the co-regulated genes *MRPL1* and *C2ORF3*.

***DYX1C1*** The 15q21 region was identified as a candidate region for a gene or genes influencing reading disability (RD) through linkage studies (Fulker et al. 1991; Grigorenko et al. 1997), defining the DYX1 (DYsleXia-1st reported) locus. The *DYX1C1* (DYX1-Candidate 1) gene was specifically targeted after a translocation t(2;15) (q11;q21) disrupting the previously uncharacterized gene was observed in a family with RD (Taipale et al. 2003). Since then, the DYX1C1 protein has been found to contain estrogen-receptor-binding sites (Massinen et al. 2009) and knockdown of the gene in embryonic rat brain produces delays in neuronal migration (Wang et al. 2006).

Sequence analysis of the *DYX1C1* coding regions identified a missense mutation in some RD families: rs57809907, 1249G>T, which results in Glu417X and truncates the protein by four amino acids (Taipale et al. 2003), but this variant has not been consistently associated with reading disability in other studies (Scerri et al. 2004; Wigg et al. 2004; Marino et al. 2005; Meng et al. 2005a; Dahdouh et al. 2009; Bates et al. 2010). Another missense mutation, rs17819126 (271G>A, Val91Ile) has been associated with reading ability (Bates et al. 2010), but analyses of putative effect on protein function by the SIFT (Ng and Henikoff 2003) and PolyPhen (Adzhubei et al. 2010) algorithms indicate that this should be a benign change for the protein (Ensembl release 63: [www.ensembl.org](http://www.ensembl.org)). Since studies have found association of RD with other SNPs within the gene, it seems likely that mutations affecting RD are in the regulatory rather than coding regions. A possible candidate is the -3G>A SNP rs3743205 in the 5' untranslated region (UTR). In the original report by Taipale et al. (2005), the A allele was associated with RD, but subsequent reports found association with the common G allele (Scerri et al. 2004; Wigg et al. 2004; Dahdouh et al. 2009). Interestingly, a study of RD in Chinese children also found strong association with the G allele (Lim et al. 2011). Studies of transcription factor binding in the promoter region have shown that the A allele

shows decreased binding to a repressive transcription factor, resulting in increased *DYX1C1* expression (Tapia-Paez et al. 2008), leading Lim et al. to hypothesize that the A allele is actually protective compared to the downregulating G allele, and that instances where the A allele appeared to be associated with RD could be secondary to linkage disequilibrium with a second causal variant nearby. Two other SNPs, rs12899331 and rs16787, in the promoter region were also found to be involved in transcription factor binding (Tapia-Paez et al. 2008), but these were not found to be associated with RD in later studies (Dahdouh et al. 2009). Thus, further studies are needed to define the role of particular regulatory regions of *DYX1C1* in the cause of RD.

There is also evidence for pleiotropic effects of *DYX1C1* on short-term memory and mental calculation, a mathematics measure (Marino et al. 2011a). Linkage to the *DYX1C1* region was found with speech sound disorder phenotypes (SSD) in one study (Smith et al. 2005) but not in a subsequent study (Stein et al. 2006), which located the SSD region more centromerically. Further studies are needed to determine whether the linkage signal for SSD was actually related to *DYX1C1*.

***DCDC2*** Initial linkage and association analysis defined the *DYX2* locus on chromosome 6p22 (Grigorenko et al. 1997; Cardon et al. 1994; Cardon et al. 1995; Fisher et al. 1999; Gayan et al. 1999; Kaplan et al. 2002; Deffenbacher et al. 2004), and several subsequent studies have reported associations with the *DCDC2* (doublecortin-2) gene within the region (Newbury et al. 2011; Meng et al. 2005b; Schumacher et al. 2006; Scerri et al. 2011). The structure of the gene is analogous to the X-linked *DCX* gene which is known to be involved in microtubular structure and influences neuronal migration. Mutation of the *DCX* gene produces lissencephaly in males and cortical abnormalities in females (des Portes et al. 1998; Sossey-Alaoui et al. 1998); accordingly, knockdown of *DCDC2* produces delays in neuronal migration in embryonic rat brain (Meng et al. 2005b).

Sequence analysis of coding regions of *DCDC2* in RD families has not identified causal mutations; however, association of RD was reported with a deletion in intron 2, termed BV677278, which appeared to contain transcription factor binding sites (Meng et al. 2005b). One study failed to replicate this association (Ludwig et al. 2008), but other studies have replicated it (Brkanac et al. 2007; Harold et al. 2006; Wilcke et al. 2009; Marino et al. 2011b), although the statistical significances in some of these studies were weak. Still, the importance of this region in gene regulation has been demonstrated by in vitro studies showing that sequences in the region act as enhancers for *DCDC2* expression (Meng et al. 2011). Furthermore, these differences in gene expression may have a measurable

phenotypic effect on brain structure in that BV677278 variants have been associated with differences in gray matter volume in unselected individuals (Meda et al. 2008), so this deletion appears to be an example of a mutation in a regulatory region that affects RD.

In addition to influencing RD, SNPs in *DCDC2* have been associated with both hyperactive and inattentive forms of ADHD, indicating that this gene can affect both disorders (Couto et al. 2009). More recently, evidence has been presented that *DCDC2* contributes to the risk for autism in families with both dyslexia and autism (Cuccaro et al. 2011).

*KIAA0319 DCDC2* and *KIAA0319* coding regions are separated by only 160 kb and were both included in the candidate region defined by linkage and association analyses that identified the *DYX2* locus. Association of *KIAA0319* with RD phenotypes as well as reading in the normal range has been supported by numerous studies (Newbury et al. 2011; Dennis et al. 2009; Scerri et al. 2011; Harold et al. 2006; Cope et al. 2005; Paracchini et al. 2006; Luciano et al. 2007; Paracchini et al. 2008). Although the function of the gene is not clear, knockdown of expression in embryonic rat brain results in delayed neuronal migration (Paracchini et al. 2006), similar to the knockdowns of *DYX1C1* and *DCDC2* noted above.

The SNPs showing association with RD tend to be located in the 5' UTR, the first untranslated exon, and the first intron (Elbert et al. 2011), suggesting regulatory functions. Expression of the allele containing the RD-associated SNP haplotype in this region of *KIAA0319* was shown to be decreased in cell lines from individuals with RD (Paracchini et al. 2006). Moreover, one associated SNP, rs9461045, has been shown to have a regulatory function. Reporter assays showed that the risk allele, which was hypothesized to create a binding site for the repressor OCT-1, resulted in decreased expression of *KIAA0319* in vitro, and knockdown of OCT-1 restored expression (Dennis et al. 2009).

In recognition of the likely influence of epigenetic mechanisms on *KIAA0319* expression in the etiology of RD, regions of acetylated histones were mapped in and around the gene in a neuroblastoma cell line to identify promoter regions (Couto et al. 2010). A 2.7-kb acetylated region was found spanning the 5' UTR, first exon and first intron of *KIAA0319* which corresponded to the location of five SNPs that had been associated with RD phenotypes in other studies. In addition, SNPs within or very near the acetylated region have been associated with language impairment phenotypes (Newbury et al. 2011; Rice et al. 2009) and linkage to the *DCDC2/KIAA0319* region has been reported for SSD (Smith et al. 2005). Studies in an unselected population did not show effects on language in the normal range, suggesting that this gene has more of an effect on language impairment (Scerri et al. 2011).

*ROBO1* Linkage analysis of a large family localized RD to a region on chromosome 3 (3p12–q13) which was designated *DYX5* (Nopola-Hemmi et al. 2001). A translocation within this region, t(3;8)(p12;q11), was found in an individual with RD and it was determined that this disrupted the *ROBO1* gene (Hannula-Jouppi et al. 2005), making it a candidate for RD. This gene is the human homologue of the roundabout gene in *Drosophila* and mice and is known to affect axonal guidance through the midline of the CNS and spinal cord. The coding regions of the *ROBO1* gene were sequenced in the original *DYX5* family, but no causal mutations were found. Association was found with a haplotype of SNPs in the gene in this family, and transcription of the allele containing the risk haplotype was decreased in lymphoblasts from individuals with RD. The individuals SNPs were not felt to have a regulatory function since they were also noted in unaffected individuals, pointing to an unknown regulatory mutation in the individuals with the risk haplotype. Although subsequent studies have not replicated the association of *ROBO1* SNPs with RD, linkage has been found with SSD (Stein and Schick 2004) and SNP association has been found with phonological buffer deficits in an unselected population (Bates et al. 2011) indicating that the gene's primary effects could be on language abilities related to RD.

*MRPL19 and C2ORF3* The designation of these two genes as candidates in influencing RD rests on the assumption that the causal mutation is in a regulatory region that is about 34 kb from the genes. The 2p16–p12 region was first highlighted by a genome-wide microsatellite linkage study in an extended family (Fagerheim et al. 1999), and subsequent linkage studies replicated these results across the region (Petryshen et al. 2002; Fisher et al. 2002; Francks et al. 2002; Kaminen et al. 2003). SNP association studies focused on the 2p12 region, with results indicating a region that did not contain recognizable genes (Peyrard-Janvid et al. 2004; Anthoni et al. 2007). The transcription products of three nearby genes, *FLJ13391*, *MRPL19*, and *C2ORF3*, were examined to determine if the risk haplotype of SNPs in that region had an effect on gene expression. There was no effect on the transcription of *FLJ13391*, but transcripts of one allele from *MRPL19* and *C2ORF3* were decreased in individuals who carried the risk haplotypes in the adjacent region (as determined by heterozygous SNPs within the coding regions of the two genes). This suggested that an unknown mutation in the region of SNP association has an effect on gene expression of both genes. The *MRPL19* protein is a component of the mitochondrial ribosome, but the function of the *C2ORF3* gene is unknown.

Overall, there is substantial evidence for involvement of mutations in regulatory regions of the primary candidate genes influencing RD, and several of these genes also affect

related language and learning disorders. Further investigation of epigenetic mechanisms of gene regulation is likely to be profitable, including elements that may be quite distant from the genes they affect, or factors that regulate more than one gene.

### Mechanisms of epigenetic gene regulation

The term “epigenetics” refers to the controls of gene expression that are maintained through somatic cell division (and occasionally in germline cells) but do not involve change in the DNA code itself. Stable epigenetic controls are applied and subsequently maintained in cell lineages during differentiation and cell proliferation, and reversible epigenetic changes in gene expression can occur in differentiated cells in response to external signals (Jaenisch and Bird 2003; Ptashne 2007; Day and Sweatt 2011). The two major methods of epigenetic regulation involve changes in methylation of cytosines in regulatory regions of DNA or modification of histone proteins, primarily through acetylation and methylation.

Epigenetic modifications act to control the accessibility of DNA to transcription. Methylation of cytosines in regulatory elements or the complexing of DNA around nucleosomes can block gene expression, while removal of DNA methylation or relaxing of histone complexing can make DNA more accessible. Methylation often acts on CpG islands or shores, which are regions of cytosine–guanine dinucleotide sequences in promoters of genes, thus inhibiting the binding of transcription factors. One or both strands may be methylated, which can fine-tune the degree of expression. In addition, methylation of these regions can recruit histone modifications that also block transcription machinery. In contrast, methylation within the gene exons and introns is correlated with gene expression. DNA methylation is mediated by a family of DNMT enzymes which apply and maintain methylation tags (Day and Sweatt 2011; Portela and Esteller 2010; Gropman and Batshaw 2010).

Histone modification affects the wrapping of DNA around nucleosomes, which are octomers composed of two each of four different histone proteins: H2A, H2B, H3, H4. DNA complexing with nucleosomes is part of chromatin compaction into heterochromatin, which generally is less transcriptionally active. Each histone protein has multiple sites that are subject to modification (methylation, acetylation, phosphorylation, ubiquitination, ADP ribosylation, or sumoylation) (Kouzarides 2007). Specific sites are designated by the histone type and the amino acid number, such that H4K12 designates the 14th amino acid in an H4 protein, which is a lysine (K). These modifications are reversible, mediated by families of enzymes such as histone acetylases (HATs), histone deacetylases (HDACs), histone

methylases (HMTs), histone demethylases (HDMs), and so on. The combination of histone modifications at different histone sites appears to constitute a “code” or “language” that determines when, where, and how much a particular gene is expressed (Day and Sweatt 2011; Portela and Esteller 2010; Lee et al. 2010).

### Mutations of genes affecting epigenetic mechanisms in humans and animal models

Since epigenetic mechanisms regulate the differential expression of genes in developing tissues, gene mutations that interfere with DNA methylation or histone modification may disrupt multiple organ systems. Table 1 gives several examples of developmental cognitive disorders caused by mutations in genes that disrupt epigenetic processes resulting in varying degrees of motor, craniofacial, and skeletal problems in addition to their effects on cognitive abilities. Other cognitive disorders such as Alzheimer Disease and Huntington Disease develop in adulthood through gradual neurodegeneration secondary to deregulated genes.

While the effects of mutations of genes that affect epigenetic processes can be severe and disrupt multiple systems, other genetic effects on epigenetic modification can be much more circumscribed. The “language” of methylated DNA and specific histone modifications can precisely control gene expression to produce and maintain tissue-specific and region-specific cellular differentiation. Once differentiation is completed, the same regulatory mechanisms appear to be involved in the changes in gene expression that result from learning and memory in the hippocampus (Day and Sweatt 2011). For example, certain types of learning are correlated with specific patterns of histone modification in chromatin of hippocampal cells, e.g., the learning of contextual fear responses in mice is associated with acetylation at H3K9, H3K14, H4K5, H4K8, and H4K12, as well as changes in methylation and phosphorylation at other sites. Moreover, loss of acetylation at H4K12 interferes with learning, which is normalized by introduction of an HDAC inhibitor which restores acetylation at that site (Day and Sweatt 2011; Peleg et al. 2010). Similarly, interference with the machinery that applies histone modifications or DNA methylation such as HATs, HDACs, HDMs, or DNMTs also cause learning problems; for example, mutation of the *Cbp* gene in mice, or blockage of DNA methylation through inhibition of *Dnmts* will both interfere with memory and long-term potentiation in the hippocampus (Day and Sweatt 2011; Alarcon et al. 2004; Lubin et al. 2008; Levenson et al. 2006; Miller and Sweatt 2007). The *EHMT* gene in humans encodes a histone demethylase and heterozygous deletion of this telomeric gene causes Kleefstra syndrome, a condition with severe intellectual disability, dysmorphic features, and behavioral problems such as autistic features, aggression,



**Table 1** Developmental disorders resulting from disruption of epigenetic mechanisms (Galaburda 2005)

Mechanism	Disease	Gene	Effect	Consequences
DNA methylation	Rett syndrome	<i>MeCP2</i>	Hypermethylation, abnormal mRNA splicing	Transcription repression or activation
	Fragile X syndrome	<i>FMR1</i>	Promoter hypermethylation	Transcription repression
	Prader Willi syndrome/ Angelman syndrome	del15q11-q13, <i>UBE3A</i>	Aberrant methylation in imprint control region	Transcription repression or activation
	Immunodeficiency, centromere instability, facial dysmorphism	<i>DNMT3B</i>	Hypomethylation	Transcription activation
	Alzheimer disease	<i>NEP</i>	CpG island hypomethylation	Transcription activation
Histone acetylation	Rubenstein-Taybi syndrome	<i>CBP (HAT)</i>	Reduced histone acetylation, hypertrimethylation of DNA	Transcription repression
	Coffin-Lowry syndrome	<i>RSK32</i>	Hypophosphorylation of site H3S10	Increased transcription of MAP kinase genes
Histone methylation	Oculofaciocardio-dental	<i>BCOR</i>	Disruption of HDACs	Transcription activation
	Sotos syndrome	<i>NSD1</i>	Decreased methylation of sites H4K20, H3K36	Transcription activation of multiple genes
	Kleefstra syndrome	<i>EHMT1</i>	Decreased histone methylation	Transcription activation
	Huntington Disease	<i>HTT</i>	Increased methylation at site H3K9 and possibly H3K27	Transcription activation

Galaburda (2005) adapted from Portela and Esteller (2010); Day and Sweatt (2011); Kelly et al. (2010), Lagali et al. (2010); and Gropman and Batshaw (2010)

and bipolar disorder that can change in expression and severity over time (Kleefstra et al. 2009). In *Drosophila*, mutation of the EHMT homologue results in disruption of a jumping reflex and courtship memory. These deficits were also rescued by expression of EHMT in adult flies (Kramer et al. 2011). Additional studies of mouse models of Alzheimer disease and other neurodegenerative diseases have also shown rescue of learning deficits with treatment by HDACs (Fischer et al. 2007; Guan et al. 2009).

Most recently, there have been several reports of alterations of methylation in autism spectrum disorders. Alterations in methylation of CpG islands associated with the *OXTTR* oxytocin receptor gene have been reported in brain tissues of individuals with autism spectrum disorders (Gregory et al. 2009). Abstracts at the International Congress of Human Genetics/American Society of Human Genetics meeting in Montreal in October 2011 reported that identical twins discordant for autism had significantly different genome-wide methylation patterns (Wong et al. 2011), and siblings discordant for autism had differences in 5-hydroxymethylcytosine across exonic sequences. Finally, DNA methylation was altered in CpG islands associated with the candidate gene *SHANK3* in brain samples from individuals with autism spectrum disorders, resulting in an altered pattern of isoform expression (Zhu et al. 2011). The influence of more remote regulatory regions was noted in the downregulation of the *CHRNA7* gene in autism by the Prader Willi imprinting center at 15q11.2–13.3 (Yasui et al. 2011).

Mouse models of human epigenetic syndromes, such as those listed in Table 1, can show severe phenotypic effects similar to their human counterparts (unless the models are constructed such that the mutations are only expressed in selected tissues); however, many of these disorders are caused by null mutations that have a significant effect on function. Other models of mutations of genes affecting epigenetic regulation can show much milder changes in hippocampal neurons or dendritic spines (Lagali et al. 2010). It seems possible, then, that less disruptive mutations or mutations of other genes may have much more focused effects on development and thus may be much more analogous to deficits that affect reading and language disorders. Thus, while mutations affecting epigenetic mechanisms have not been described in reading disability or language impairments, the role of epigenetic changes in learning and autism and the hints of potential therapy make it especially worthwhile to look for mutations in such genes in individuals with language and learning problems.

Approaches to the identification of epigenetic mechanisms in humans

Although candidate genes have been identified for reading disability and language impairment, the SNPs in these genes appear to account for a small portion of the phenotypic variability. In contrast, fairly substantial heritabilities have been claimed for these disorders, between

0.45–0.85 depending on population and definitions (Gayan and Olson 2001; Hawke et al. 2006; Astrom et al. 2007; Spinath et al. 2004; Tomblin and Buckwalter 1998; Dale et al. 1998; Bishop and Hayiou-Thomas 2008). There are several possible explanations for this “missing heritability,” but one of the primary reasons appears to be inherent in the current studies of SNPs, particularly in the large panels that are used for genome-wide studies. The SNPs selected for such panels are generally common in the population, which makes them more informative in comparisons between affected and unaffected individuals, but assessment of individual common SNPs ignores rare variants which are likely to have more impact, and also ignores epistatic interactions between loci (Manolio et al. 2009). There are approaches that enhance the identification of causal genes from genome-wide association studies (GWAS) data such as the simultaneous analysis of multiple variants associated with a gene (Neale and Sham 2004; Huang et al. 2011; Li et al. 2011) or focus on SNPs associated with loci which show phenotype-based differences in expression (eQTLs or eSNPs) (Innocenti et al. 2011; Majewski and Pastinen 2011) and “next generation sequencing” allows the analysis of rare as well as common variants around a gene; however, sites involved in epigenetic control of gene expression may not be included in the set of loci in gene-based analysis, and the variation in expression of eQTLs may be due in part to mutations in epigenetic regions which may be somewhat distant from the gene itself (Ernst et al. 2011). The influence of remote regulatory elements is likely to be missing in targeted screening approaches which focus on candidate genes, whether through SNP analysis or sequencing. This is due in part to the lack of information on where these regions are located, and initiatives such as the NIH Epigenomics Roadmap Program (<http://nihroadmap.nih.gov/epigenomics/initiatives.asp>) and the International Human Epigenome Consortium (<http://www.ihec-epigenomes.org/>). These are large collaborative efforts to map regions in the genome that are involved in epigenetic regulation, and the results will assist investigators in identifying regions for evaluation.

Heritable mutations that influence reading and language disorders could be in the genes that regulate epigenetic processes, analogous to the mutations in HDACs or DNMTs, or in genes such as MeCP2 or in the MAPK signaling pathway (Day and Sweatt 2011). Alternatively, mutations could be in the DNA binding regions themselves. Genome-wide association studies or even targeted SNP analysis might be able to detect such mutations, given that the sample size is large enough, the variants are not rare, and the adjacent SNPs are in linkage disequilibrium. Knowledge of the location of epigenetic regions could help prioritize the follow-up of SNPs in a GWAS that otherwise might be ignored because of lack of apparent functional

relevance (Ernst et al. 2011), and location information would also guide the placement of SNPs in a targeted array. Sequence analysis would detect rare variants, but until whole genome sequencing of large populations is financially feasible, targeted sequencing studies are also dependent upon the selection of candidate genes and regulatory regions. Studies of epigenetic mechanisms in animal models should produce additional candidate genes for examination in cognitive disorders in humans.

Another approach would be to look for genomic regions of abnormal methylation or histone modification in individuals with specific forms of language or learning disorders. However, epigenetic patterns are likely to be different in different tissues, and histone modifications especially may change over time. Fortunately, there are studies which indicate that methylation patterns affecting disorders can be consistent across tissues, such as lymphocyte and brain methylation patterns in individuals with psychiatric disorders, suggesting that lymphocyte tissues can be a good proxy for brain (Dempster et al. 2011). An abnormal methylation pattern in a region of DNA from human tissues such as lymphocytes or fibroblasts could indicate an epigenetic process that could be pursued further by determination of the effects of that abnormality on gene expression and the impact on learning in animal models. Such studies could be valuable in identifying important genes and signaling pathways involved in learning. Further genetic studies such as association and sequencing could assess the influence of these new candidates at the population level. Conversely, though, lack of a methylation abnormality in “proxy” tissues would not rule out the involvement of an epigenetic mechanism that is confined to a region of the brain.

There are many approaches to the identification of genes that affect quantitative traits such as language and learning disorders, and the most effective will take advantage of simultaneous analysis of genomic and expression analyses (Charlesworth et al. 2009). The inclusion of information on epigenetic mechanisms of gene regulation may turn out to be an important consideration in gene identification and possibly even in therapy.

## References

- Adzhubei IA, Schmidt S, Peshkin L, Ramensky VE, Gerasimova A, Bork P, et al. A method and server for predicting damaging missense mutations. *Nat Methods*. 2010;7(4):248–9.
- Alarcon JM, Malleret G, Touzani K, Vronskaya S, Ishii S, Kandel ER, et al. Chromatin acetylation, memory, and LTP are impaired in CBP<sup>+/-</sup> mice: a model for the cognitive deficit in Rubinstein-Taybi syndrome and its amelioration. *Neuron*. 2004;42(6):947–59.
- Anthoni H, Zucchelli M, Matsson H, Muller-Myhsok B, Fransson I, Schumacher J, et al. A locus on 2p12 containing the co-regulated MRPL19 and C2ORF3 genes is associated to dyslexia. *Hum Mol Genet*. 2007;16(6):667–77.

- Astrom RL, Wadsworth SJ, DeFries JC. Etiology of the stability of reading difficulties: the longitudinal twin study of reading disabilities. *Twin Res Hum Genet.* 2007;10(3):434–9.
- Bates TC, Lind PA, Luciano M, Montgomery GW, Martin NG, Wright MJ. Dyslexia and *DYX1C1*: deficits in reading and spelling associated with a missense mutation. *Mol Psychiatry.* 2010;15(12):1190–6.
- Bates TC, Luciano M, Medland SE, Montgomery GW, Wright MJ, Martin NG. Genetic variance in a component of the language acquisition device: *ROBO1* polymorphisms associated with phonological buffer deficits. *Behav Genet.* 2011;41(1):50–7.
- Bishop DV, Hayiou-Thomas ME. Heritability of specific language impairment depends on diagnostic criteria. *Genes Brain Behav.* 2008;7(3):365–72.
- Brkanac Z, Chapman NH, Matsushita MM, Chun L, Nielsen K, Cochrane E, et al. Evaluation of candidate genes for *DYX1* and *DYX2* in families with dyslexia. *Am J Med Genet B Neuropsychiatr Genet.* 2007;144B(4):556–60.
- Cardon LR, Smith SD, Fulker DW, Kimberling WJ, Pennington BF, DeFries JC. Quantitative trait locus for reading disability on chromosome 6. *Science.* 1994;266(5183):276–9.
- Cardon LR, Smith SD, Fulker DW, Kimberling WJ, Pennington BF, DeFries JC. Quantitative trait locus for reading disability: correction. *Science.* 1995;268(5217):1553.
- Charlesworth JC, Peralta JM, Drigalenko E, Goring HH, Almsay L, Dyer TD, Blangero J. Toward the identification of causal genes in complex diseases: a gene-centric joint test of significance combining genomic and transcriptomic data. *BMC Proc* 2009, 3 Suppl 7:S92.
- Cope N, Harold D, Hill G, Moskvina V, Stevenson J, Holmans P, et al. Strong evidence that *KIAA0319* on chromosome 6p is a susceptibility gene for developmental dyslexia. *Am J Hum Genet.* 2005;76(4):581–91.
- Couto JM, Gomez L, Wigg K, Ickowicz A, Pathare T, Malone M, et al. Association of attention-deficit/hyperactivity disorder with a candidate region for reading disabilities on chromosome 6p. *Biol Psychiatry.* 2009;66:368–75.
- Couto JM, Livne-Bar I, Huang K, Xu Z, Cate-Carter T, Feng Y, et al. Association of reading disabilities with regions marked by acetylated H3 histones in *KIAA0319*. *Am J Med Genet B Neuropsychiatr Genet.* 2010;153B(2):447–62.
- Cuccaro ML, Ma D, Martin ER, Haines JL, Gilbert JR, Pericak-Vance MA: Common genetic susceptibility to autism spectrum disorder (ASD) and dyslexia (DYS). Presented at the 12th International Congress of Human Genetics/61st Annual Meeting of The American Society of Human Genetics, October 12, 2011, Montreal, Canada.
- Dahdouh F, Anthoni H, Tapia-Paez I, Peyrard-Janvid M, Schulte-Korne G, Warnke A, et al. Further evidence for *DYX1C1* as a susceptibility factor for dyslexia. *Psychiatr Genet.* 2009;19:59–63.
- Dale PS, Simonoff E, Bishop DV, Eley TC, Oliver B, Price TS, et al. Genetic influence on language delay in two-year-old children. *Nat Neurosci.* 1998;1(4):324–8.
- Day JJ, Sweatt JD. Epigenetic mechanisms in cognition. *Neuron.* 2011;70(5):813–29.
- Deffenbacher KE, Kenyon JB, Hoover DM, Olson RK, Pennington BF, DeFries JC, et al. Refinement of the 6p21.3 quantitative trait locus influencing dyslexia: linkage and association analyses. *Hum Genet.* 2004;115(2):128–38.
- Dempster EL, Pidsley R, Schalkwyk LC, Owens S, Georgiades A, Kane F, Kalidindi S, Picchioni M, Kravariti E, Touloupoulou T et al: Disease-associated epigenetic changes in monozygotic twins discordant for schizophrenia and bipolar disorder. *Hum Mol Genet* 2011.
- Dennis MY, Paracchini S, Scerri TS, Prokunina-Olsson L, Knight JC, Wade-Martins R, et al. A common variant associated with dyslexia reduces expression of the *KIAA0319* gene. *PLoS Genet.* 2009;5(3):e1000436.
- des Portes V, Francis F, Pinard JM, Desguerre I, Moutard ML, Snoeck I, et al. doublecortin is the major gene causing X-linked subcortical laminar heterotopia (SCLH). *Hum Mol Genet.* 1998;7(7):1063–70.
- Elbert A, Lovett MW, Cate-Carter T, Pitch A, Kerr EN, Barr CL. Genetic variation in the *KIAA0319* 5' region as a possible contributor to dyslexia. *Behav Genet.* 2011;41(1):77–89.
- Ernst J, Kheradpour P, Mikkelsen TS, Shores N, Ward LD, Epstein CB, et al. Mapping and analysis of chromatin state dynamics in nine human cell types. *Nature.* 2011;473(7345):43–9.
- Fagerheim T, Raeymaekers P, Tonnessen FE, Pedersen M, Tranebjaerg L, Lubs HA. A new gene (*DYX3*) for dyslexia is located on chromosome 2. *J Med Genet.* 1999;36(9):664–9.
- Fischer A, Sananbenesi F, Wang X, Dobbin M, Tsai LH. Recovery of learning and memory is associated with chromatin remodelling. *Nature.* 2007;447(7141):178–82.
- Fisher SE, Marlow AJ, Lamb J, Maestrini E, Williams DF, Richardson AJ, et al. A quantitative-trait locus on chromosome 6p influences different aspects of developmental dyslexia. *Am J Hum Genet.* 1999;64(1):146–56.
- Fisher SE, Francks C, Marlow AJ, MacPhie IL, Newbury DF, Cardon LR, et al. Independent genome-wide scans identify a chromosome 18 quantitative-trait locus influencing dyslexia. *Nat Genet.* 2002;30(1):86–91.
- Francks C, Fisher SE, Olson RK, Pennington BF, Smith SD, DeFries JC, et al. Fine mapping of the chromosome 2p12-16 dyslexia susceptibility locus: quantitative association analysis and positional candidate genes *SEMA4F* and *OTX1*. *Psychiatr Genet.* 2002;12(1):35–41.
- Fulker DW, Cardon LR, DeFries JC, Kimberling WJ, Pennington BF, Smith SD. Multiple regression analysis of sib-pair data on reading to detect quantitative trait loci. *Read Writ Interdisc J.* 1991;3:299–313.
- Gabel LA, Gibson CJ, Gruen JR, LoTurco JJ. Progress towards a cellular neurobiology of reading disability. *Neurobiol Dis.* 2010;38(2):173–80.
- Galaburda AM. Dyslexia—a molecular disorder of neuronal migration: the 2004 Norman Geschwind Memorial Lecture. *Ann Dyslexia.* 2005;55(2):151–65.
- Gayan J, Olson RK. Genetic and environmental influences on orthographic and phonological skills in children with reading disabilities. *Dev Neuropsychol.* 2001;20(2):483–507.
- Gayan J, Smith SD, Cherny SS, Cardon LR, Fulker DW, Brower AM, et al. Quantitative-trait locus for specific language and reading deficits on chromosome 6p. *Am J Hum Genet.* 1999;64(1):157–64.
- Gregory SG, Connelly JJ, Towers AJ, Johnson J, Biscocho D, Markunas CA, et al. Genomic and epigenetic evidence for oxytocin receptor deficiency in autism. *BMC Med.* 2009;7:62.
- Grigorenko EL, Wood FB, Meyer MS, Hart LA, Speed WC, Shuster A, et al. Susceptibility loci for distinct components of developmental dyslexia on chromosomes 6 and 15. *Am J Hum Genet.* 1997;60(1):27–39.
- Gropman AL, Batshaw ML. Epigenetics, copy number variation, and other molecular mechanisms underlying neurodevelopmental disabilities: new insights and diagnostic approaches. *J Dev Behav Pediatr.* 2010;31(7):582–91.
- Guan JS, Haggarty SJ, Giacometti E, Dannenberg JH, Joseph N, Gao J, et al. HDAC2 negatively regulates memory formation and synaptic plasticity. *Nature.* 2009;459(7243):55–60.
- Hannula-Jouppi K, Kaminen-Ahola N, Taipale M, Eklund R, Nopola-Hemmi J, Kaariainen H, et al. The axon guidance receptor gene *ROBO1* is a candidate gene for developmental dyslexia. *PLoS Genet.* 2005;1(4):e50.
- Harold D, Paracchini S, Scerri T, Dennis M, Cope N, Hill G, et al. Further evidence that the *KIAA0319* gene confers susceptibility

- to developmental dyslexia. *Mol Psychiatry*. 2006;11(12):1085–91. 1061.
- Hawke JL, Wadsworth SJ, DeFries JC. Genetic influences on reading difficulties in boys and girls: the Colorado twin study. *Dyslexia*. 2006;12(1):21–9.
- Huang H, Chanda P, Alonso A, Bader JS, Arking DE. Gene-based tests of association. *PLoS Genet*. 2011;7(7):e1002177.
- Innocenti F, Cox NJ, Dolan ME. The use of genomic information to optimize cancer chemotherapy. *Semin Oncol*. 2011;38(2):186–95.
- Jaenisch R, Bird A. Epigenetic regulation of gene expression: how the genome integrates intrinsic and environmental signals. *Nat Genet*. 2003;33(Suppl):245–54.
- Kaminen N, Hannula-Jouppi K, Kestila M, Lahermo P, Muller K, Kaaranen M, et al. A genome scan for developmental dyslexia confirms linkage to chromosome 2p11 and suggests a new locus on 7q32. *J Med Genet*. 2003;40(5):340–5.
- Kaplan DE, Gayan J, Ahn J, Won TW, Pauls D, Olson RK, et al. Evidence for linkage and association with reading disability on 6p21.3-22. *Am J Hum Genet*. 2002;70(5):1287–98.
- Kelly TK, De Carvalho DD, Jones PA. Epigenetic modifications as therapeutic targets. *Nat Biotechnol*. 2010;28(10):1069–78.
- Kleefstra T, van Zelst-Stams WA, Nillesen WM, Cormier-Daire V, Houge G, Foulds N, et al. Further clinical and molecular delineation of the 9q subtelomeric deletion syndrome supports a major contribution of EHMT1 haploinsufficiency to the core phenotype. *J Med Genet*. 2009;46(9):598–606.
- Kouzarides T. Chromatin modifications and their function. *Cell*. 2007;128(4):693–705.
- Kramer JM, Kochinke K, Oortveld MA, Marks H, Kramer D, de Jong EK, et al. Epigenetic regulation of learning and memory by *Drosophila* EHMT/G9a. *PLoS Biol*. 2011;9(1):e1000569.
- Lagali PS, Corcoran CP, Picketts DJ. Hippocampus development and function: role of epigenetic factors and implications for cognitive disease. *Clin Genet*. 2010;78(4):321–33.
- Lee JS, Smith E, Shilatifard A. The language of histone crosstalk. *Cell*. 2010;142(5):682–5.
- Levenson JM, Roth TL, Lubin FD, Miller CA, Huang IC, Desai P, et al. Evidence that DNA (cytosine-5) methyltransferase regulates synaptic plasticity in the hippocampus. *J Biol Chem*. 2006;281(23):15763–73.
- Li MX, Gui HS, Kwan JS, Sham PC. GATES: a rapid and powerful gene-based association test using extended Simes procedure. *Am J Hum Genet*. 2011;88(3):283–93.
- Lim CK, Ho CS, Chou CH, Wayne MM. Association of the rs3743205 variant of DYX1C1 with dyslexia in Chinese children. *Behav Brain Funct*. 2011;7:16.
- Lubin FD, Roth TL, Sweatt JD. Epigenetic regulation of BDNF gene transcription in the consolidation of fear memory. *J Neurosci*. 2008;28(42):10576–86.
- Luciano M, Lind PA, Duffy DL, Castles A, Wright MJ, Montgomery GW, et al. A haplotype spanning KIAA0319 and TTRAP is associated with normal variation in reading and spelling ability. *Biol Psychiatry*. 2007;62(7):811–7.
- Ludwig KU, Schumacher J, Schulte-Korne G, Konig IR, Warnke A, Plume E, et al. Investigation of the DCDC2 intron 2 deletion/compound short tandem repeat polymorphism in a large German dyslexia sample. *Psychiatr Genet*. 2008;18(6):310–2.
- Majewski J, Pastinen T. The study of eQTL variations by RNA-seq: from SNPs to phenotypes. *Trends Genet*. 2011;27(2):72–9.
- Manolio TA, Collins FS, Cox NJ, Goldstein DB, Hindorf LA, Hunter DJ, et al. Finding the missing heritability of complex diseases. *Nature*. 2009;461(7265):747–53.
- Marino C, Giorda R, Luisa Lorusso M, Vanzin L, Salandi N, Nobile M, et al. A family-based association study does not support DYX1C1 on 15q21.3 as a candidate gene in developmental dyslexia. *Eur J Hum Genet*. 2005;13(4):491–9.
- Marino C, Mascheretti S, Riva V, Cattaneo F, Rigoletto C, Rusconi M, et al. Pleiotropic effects of DCDC2 and DYX1C1 genes on language and mathematics traits in nuclear families of developmental dyslexia. *Behav Genet*. 2011a;41(1):67–76.
- Marino C, Meng H, Mascheretti S, Rusconi M, Cope N, Giorda R, Molteni M, Gruen JR: DCDC2 genetic variants and susceptibility to developmental dyslexia. *Psychiatr Genet* 2011b epub Aug. 30, 2011.
- Massinen S, Tammimies K, Tapia-Paez I, Matsson H, Hokkanen ME, Soderberg O, et al. Functional interaction of DYX1C1 with estrogen receptors suggests involvement of hormonal pathways in dyslexia. *Hum Mol Genet*. 2009;18(15):2802–12.
- Meda SA, Gelernter J, Gruen JR, Calhoun VD, Meng H, Cope NA, et al. Polymorphism of DCDC2 reveals differences in cortical morphology of healthy individuals—a preliminary voxel based morphometry study. *Brain Imaging Behav*. 2008;2(1):21–6.
- Meng H, Hager K, Held M, Page GP, Olson RK, Pennington BF, et al. TDT-association analysis of EKN1 and dyslexia in a Colorado twin cohort. *Hum Genet*. 2005a;118(1):87–90.
- Meng H, Smith SD, Hager K, Held M, Liu J, Olson RK, et al. DCDC2 is associated with reading disability and modulates neuronal development in the brain. *Proc Natl Acad Sci U S A*. 2005b;102(47):17053–8.
- Meng H, Powers NR, Tang L, Cope NA, Zhang PX, Fuleihan R, et al. A dyslexia-associated variant in DCDC2 changes gene expression. *Behav Genet*. 2011;41(1):58–66.
- Miller CA, Sweatt JD. Covalent modification of DNA regulates memory formation. *Neuron*. 2007;53(6):857–69.
- Neale BM, Sham PC. The future of association studies: gene-based analysis and replication. *Am J Hum Genet*. 2004;75(3):353–62.
- Newbury DF, Paracchini S, Scerri TS, Winchester L, Addis L, Richardson AJ, et al. Investigation of dyslexia and SLI risk variants in reading- and language-impaired subjects. *Behav Genet*. 2011;41(1):90–104.
- Ng PC, Henikoff S. SIFT: predicting amino acid changes that affect protein function. *Nucleic Acids Res*. 2003;31(13):3812–4.
- Nopola-Hemmi J, Myllyluoma B, Haltia T, Taipale M, Ollikainen V, Ahonen T, et al. A dominant gene for developmental dyslexia on chromosome 3. *J Med Genet*. 2001;38(10):658–64.
- Paracchini S, Thomas A, Castro S, Lai C, Paramasivam M, Wang Y, et al. The chromosome 6p22 haplotype associated with dyslexia reduces the expression of KIAA0319, a novel gene involved in neuronal migration. *Hum Mol Genet*. 2006;15(10):1659–66.
- Paracchini S, Steer CD, Buckingham LL, Morris AP, Ring S, Scerri T, et al. Association of the KIAA0319 dyslexia susceptibility gene with reading skills in the general population. *Am J Psychiatry*. 2008;165(12):1576–84.
- Peleg S, Sananbenesi F, Zovoilis A, Burkhardt S, Bahari-Javan S, Agis-Balboa RC, et al. Altered histone acetylation is associated with age-dependent memory impairment in mice. *Science*. 2010;328(5979):753–6.
- Petryshen TL, Kaplan BJ, Hughes ML, Tzenova J, Field LL. Supportive evidence for the DYX3 dyslexia susceptibility gene in Canadian families. *J Med Genet*. 2002;39(2):125–6.
- Peyrard-Janvid M, Anthoni H, Onkamo P, Lahermo P, Zucchelli M, Kaminen N, et al. Fine mapping of the 2p11 dyslexia locus and exclusion of TACR1 as a candidate gene. *Hum Genet*. 2004;114(5):510–6.
- Poelmans G, Buitelaar JK, Pauls DL, Franke B. A theoretical molecular network for dyslexia: integrating available genetic findings. *Mol Psychiatry*. 2011;16(4):365–82.
- Portella A, Esteller M. Epigenetic modifications and human disease. *Nat Biotechnol*. 2010;28(10):1057–68.
- Ptashne M. On the use of the word 'epigenetic'. *Curr Biol*. 2007;17(7):R233–6.
- Rice ML, Smith SD, Gayan J. Convergent genetic linkage and associations to language, speech and reading measures in families



- of probands with Specific Language Impairment. *J Neurodev Disord.* 2009;1(4):264–82.
- Scerri TS, Fisher SE, Francks C, MacPhie IL, Paracchini S, Richardson AJ, et al. Putative functional alleles of *DYX1C1* are not associated with dyslexia susceptibility in a large sample of sibling pairs from the UK. *J Med Genet.* 2004;41(11):853–7.
- Scerri TS, Morris AP, Buckingham LL, Newbury DF, Miller LL, Monaco AP, et al. *DCDC2*, *KIAA0319* and *CMIP* are associated with reading-related traits. *Biol Psychiatry.* 2011;70(3):237–45.
- Schumacher J, Anthoni H, Dahdouh F, Konig IR, Hillmer AM, Kluck N, et al. Strong genetic evidence of *DCDC2* as a susceptibility gene for dyslexia. *Am J Hum Genet.* 2006;78(1):52–62.
- Smith SD, Pennington BF, Boada R, Shriberg LD. Linkage of speech sound disorder to reading disability loci. *J Child Psychol Psychiatry.* 2005;46(10):1057–66.
- Sossey-Alaoui K, Hartung AJ, Guerrini R, Manchester DK, Posar A, Puche-Mira A, et al. Human doublecortin (*DCX*) and the homologous gene in mouse encode a putative  $Ca^{2+}$ -dependent signaling protein which is mutated in human X-linked neuronal migration defects. *Hum Mol Genet.* 1998;7(8):1327–32.
- Spinath FM, Price TS, Dale PS, Plomin R. The genetic and environmental origins of language disability and ability. *Child Dev.* 2004;75(2):445–54.
- Stein CM, Schick JH, Taylor HG, Shriberg LD, Millard C, Kundtz-Kluge A, et al. Pleiotropic effects of a chromosome 3 locus on speech-sound disorder and reading. *Am J Hum Genet.* 2004;74(2):283–97.
- Stein CM, Millard C, Kluge A, Miscimarra LE, Cartier KC, Freebairn LA, et al. Speech sound disorder influenced by a locus in 15q14 region. *Behav Genet.* 2006;36(6):858–68.
- Taipale M, Kaminen N, Nopola-Hemmi J, Haltia T, Myllyluoma B, Lyytinen H, et al. A candidate gene for developmental dyslexia encodes a nuclear tetratricopeptide repeat domain protein dynamically regulated in brain. *Proc Natl Acad Sci U S A.* 2003;100(20):11553–8.
- Tapia-Paez I, Tammimies K, Massinen S, Roy AL, Kere J. The complex of *TFII-I*, *PARP1*, and *SFPQ* proteins regulates the *DYX1C1* gene implicated in neuronal migration and dyslexia. *FASEB J.* 2008;22(8):3001–9.
- Tomblin JB, Buckwalter PR. Heritability of poor language achievement among twins. *J Speech Lang Hear Res.* 1998;41(1):188–99.
- Vernes SC, Newbury DF, Abrahams BS, Winchester L, Nicod J, Groszer M, et al. A functional genetic link between distinct developmental language disorders. *N Engl J Med.* 2008;359(22):2337–45.
- Wang Y, Paramasivam M, Thomas A, Bai J, Kaminen-Ahola N, Kere J, et al. *DYX1C1* functions in neuronal migration in developing neocortex. *Neuroscience.* 2006;143(2):515–22.
- Wigg KG, Couto JM, Feng Y, Anderson B, Cate-Carter TD, Macciardi F, et al. Support for *EKN1* as the susceptibility locus for dyslexia on 15q21. *Mol Psychiatry.* 2004;9(12):1111–21.
- Wilcke A, Weissfuss J, Kirsten H, Wolfram G, Boltze J, Ahnert P. The role of gene *DCDC2* in German dyslexics. *Ann Dyslexia.* 2009;59:1–11.
- Wong CCY, Schalkwyk LC, Meaburn EL, Ronald A, Price TS, Plomin R, Mill J: Genome-wide DNA Methylation Profiling of Monozygotic Twins Discordant for Autism Spectrum Disorder 2011 Presented at the 12th International Congress of Human Genetics/61st Annual Meeting of The American Society of Human Genetics, October 12, 2011, Montreal, Canada.
- Yasui DH, Scoles HA, Horike S, Meguro-Horike M, Dunaway KW, Schroeder DI, et al. 15q11.2-13.3 chromatin analysis reveals epigenetic regulation of *CHRNA7* with deficiencies in Rett and autism brain. *Hum Mol Genet.* 2011;20(22):4311–23.
- Zhu L, Li X, Towers A, Wang X, Goldstein J, Yang H, Jiang Y: Epimutations and transcriptional dysregulation of *SHANK3* in brains associated with autism spectrum disorder 2011 Presented at the 12th International Congress of Human Genetics/61st Annual Meeting of The American Society of Human Genetics, October 14, 2011, Montreal, Canada.