

Erratum: Lidocaine-Induced Systemic Toxicity: A Case Report and Review of Literature

Badar Hasan ¹, Talal Asif ², Maryam Hasan ³

1. Department of Internal Medicine, University of Missouri Kansas City (UMKC) 2. Cardiology, John H. Stroger Hospital of Cook County, Chicago, USA 3. Internal Medicine, NYU Langone Medical Center, New York

✉ **Corresponding author:** Talal Asif, asift@umkc.edu

Disclosures can be found in Additional Information at the end of the article

Corresponding author: Talal Asif

1. Department of Internal Medicine, University of Missouri Kansas City (UMKC) 2. Cardiology, John H. Stroger Hospital of Cook County, Chicago, USA 3. Internal Medicine, NYU Langone Medical Center, New York

How to cite this erratum

Hasan B, Asif T, Hasan M (August 23, 2019) Correction: Lidocaine-Induced Systemic Toxicity: A Case Report and Review of Literature. Cureus 11(8): c24. doi:10.7759/cureus.c24

Correction Notice

The discussion section originally contained the following statement:

"American Society of Regional Anesthesia and Pain Medicine recommends an initial bolus of 1.5 **mg**/kg of 20% intravenous lipids (intralipids) followed by a continuous infusion of 0.25 mL/kg/minute. "

The unit of measurement (in bold above) was incorrectly labeled as "mg" when it should be "mL" - the text of the article has been updated to reflect this correction.