




## Article

# 7-Hydroxycoumarin Induces Vasorelaxation in Animals with Essential Hypertension: Focus on Potassium Channels and Intracellular Ca<sup>2+</sup> Mobilization

Rafael L. C. Jesus <sup>1</sup>, Isnar L. P. Silva <sup>1</sup>, Fênix A. Araújo <sup>2</sup>, Raiana A. Moraes <sup>2</sup>, Liliane B. Silva <sup>1</sup>, Daniele S. Brito <sup>1</sup>, Gabriela B. C. Lima <sup>1</sup>, Quiara L. Alves <sup>1</sup> and Darizy F. Silva <sup>1,2,\*</sup>

<sup>1</sup> Laboratory of Cardiovascular Physiology and Pharmacology, Federal University of Bahia, Salvador 40110-902, Brazil

<sup>2</sup> Gonçalo Moniz Institute, Oswaldo Cruz Foundation—FIOCRUZ, Salvador 40296-710, Brazil

\* Correspondence: darizy@gmail.com; Tel.: +55-719-88-52-34-36

**Abstract:** Cardiovascular diseases (CVD) are the deadliest noncommunicable disease worldwide. Hypertension is the most prevalent risk factor for the development of CVD. Although there is a wide range of antihypertensive drugs, there still remains a lack of blood pressure control options for hypertensive patients. Additionally, natural products remain crucial to the design of new drugs. The natural product 7-hydroxycoumarin (7-HC) exhibits pharmacological properties linked to antihypertensive mechanisms of action. This study aimed to evaluate the vascular effects of 7-HC in an experimental model of essential hypertension. The isometric tension measurements assessed the relaxant effect induced by 7-HC (0.001 μM–300 μM) in superior mesenteric arteries isolated from hypertensive rats (SHR, 200–300 g). Our results suggest that the relaxant effect induced by 7-HC rely on K<sup>+</sup>-channels (K<sub>ATP</sub>, BK<sub>Ca</sub>, and, to a lesser extent, K<sub>v</sub>) activation and also on Ca<sup>2+</sup> influx from sarcolemma and sarcoplasmic reticulum mobilization (inositol 1,4,5-triphosphate (IP<sub>3</sub>) and ryanodine receptors). Moreover, 7-HC diminishes the mesenteric artery's responsiveness to α<sub>1</sub>-adrenergic agonist challenge and improves the actions of the muscarinic agonist and NO donor. The present work demonstrated that the relaxant mechanism of 7-HC in SHR involves endothelium-independent vasorelaxant factors. Additionally, 7-HC reduced vasoconstriction of the sympathetic agonist while improving vascular endothelium-dependent and independent relaxation.

**Keywords:** 7-hydroxycoumarin; umbelliferone; hypertensive rats; isometric records; vasorelaxation; superior mesenteric artery superior



**Citation:** Jesus, R.L.C.; Silva, I.L.P.; Araújo, F.A.; Moraes, R.A.; Silva, L.B.; Brito, D.S.; Lima, G.B.C.; Alves, Q.L.; Silva, D.F. 7-Hydroxycoumarin Induces Vasorelaxation in Animals with Essential Hypertension: Focus on Potassium Channels and Intracellular Ca<sup>2+</sup> Mobilization. *Molecules* **2022**, *27*, 7324. <https://doi.org/10.3390/molecules27217324>

Academic Editor: Petko Petkov

Received: 1 October 2022

Accepted: 24 October 2022

Published: 28 October 2022

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

Cardiovascular diseases (CVD) are and will continue to be among the most significant health problems worldwide [1]. Hypertension remains the most prevalent common chronic disease risk factor for CVD. According to the Global Hypertension Practice Guideline 2020 from the International Society of Hypertension, hypertension accounts for 10.4 million deaths annually [2]. Additionally, hypertension comprises about 9% of global disability-adjusted life years [1,3]. However, due to the wide range of treatment options for hypertension, many patients do not achieve optimal blood pressure (BP) control. Thus, hypertensive patients who have not achieved treatment-mediated BP control using regularly available antihypertensive drugs at inadequate doses may present a condition known as resistant hypertension.

Resistant hypertension was defined by the European Guidelines (2018) [4] as a persistent increase in BP that remains above healthy levels, despite using three or more different classes of antihypertensive medications, one of which is a diuretic administered at the best-tolerated doses. As much as possible, the initial three-drug regimen should be standardized, including a blocker of the renin–angiotensin system, specifically an angiotensin-converting

enzyme inhibitor or an angiotensin receptor blocker; a long-acting calcium channel blocker, most commonly amlodipine; and a long-acting thiazide-like diuretic, preferably chlorthalidone or indapamide [5]. In addition, the American Heart Association (2018) [6] extended this definition to patients with four or more antihypertensive medications, irrespective of their BP level. However, despite the fact that these definitions are currently well accepted, several observations have challenged these definitions, since patients represent a heterogeneous population with highly variable morbid risks, which depending on genetic, environmental, cultural, and social characteristics [7]. As the number of prescribed drugs increases and, consequently, the dosage schedule becomes more complex, adherence to treatment decreases. Therefore, the search for drugs with different therapeutic targets for the treatment of hypertension seems to be interesting [5]. Thus, searching for new therapeutic strategies to reduce the morbidity and mortality induced by hypertension and CVD is necessary.

Coumarins are a natural product and exhibit various biological effects, such as anti-HIV, antimicrobial, anti-inflammatory, anticancer, anticonvulsant, and MAO-inhibiting properties [8]. Additionally, several coumarins have known vasorelaxant activities [9–11]. 7-Hydroxycoumarin (7-HC) is a seven-hydroxylated coumarin with pharmacological properties, such as antifungal, antinociceptive, anti-inflammatory, and hepatoprotective effects [12–15]. However, few studies have demonstrated the action of 7-HC in the cardiovascular system. Baccard and colleagues (2000) [16] showed that 7-HC was described as promoting increased coronary blood flow and causing positive inotropic effects in the rat heart and had a direct vasorelaxant effect on the coronary arteries. Furthermore, Jagadeesh and colleagues (2016) [17] showed that 7-HC protects against isoproterenol-induced myocardial infarction in rats through free radical scavenging.


Previous studies in our laboratory, developed by Alves and colleagues (2020) [18], demonstrated, for the first time, that 7-HC was found to be nontoxic in H9 c2 cardiomyocytes cells and exerts a vasorelaxant effect in the superior mesenteric artery of Wistar rats. This effect would involve the inhibition of intracellular  $Ca^{2+}$  mobilization and the participation of potassium channels, which appear to be activated by 7-HC in mesenteric artery rings, particularly  $K_v$ ,  $BK_{Ca}$ ,  $K_{ATP}$ , and  $K_{ir}$ , leading to vasodilation.

Despite our work elucidating 7-HC's mechanisms of action in the cardiovascular system, the main limitation of the study was that we could not infer the effects of 7-HC in a model of essential hypertension, since the hypertensive condition alters numerous signaling pathways. Therefore, we aimed to investigate the vascular effect of 7-HC and its cardiovascular activities in spontaneously hypertensive rats (SHR), specifically to describe the mechanisms of action involved in the observed responses.

## 2. Results

### 2.1. 7-HC Induces Endothelium-Independent-Relaxation in the Mesenteric Artery in Hypertensive Rats


7-HC (0.001  $\mu$ M to 300  $\mu$ M) induced similar concentration-dependent relaxation in pre-contracted (Phe, 1  $\mu$ M) mesenteric artery rings from SHR, with or without the endothelium (effect (300  $\mu$ M) =  $83.2 \pm 4.4\%$ , ( $n = 6$ ); effect (300  $\mu$ M) =  $100.4 \pm 9.8\%$ , ( $n = 6$ )) (Figure 1A,B). Furthermore, we also demonstrated that the contraction induced by Tyrode's solution containing 80 mM KCl before ( $0.78 \pm 0.07\%$  ( $n = 5$ )) and after ( $1.01 \pm 0.08\%$  ( $n = 5$ )) administration of 7-HC did not show a statistically significant difference, suggesting that 7-HC can reversibly interact with cell receptors and also does not induce tissue damage. Interestingly, the vasorelaxation induced by 7-HC in mesenteric artery rings without endothelium from SHR (effect (10  $\mu$ M) =  $20.7 \pm 4.7$  ( $n = 6$ )) was significantly attenuated compared with Wistar rats (effect (10  $\mu$ M) =  $43.1 \pm 5.7$  ( $n = 7$ )) at the 10  $\mu$ M concentration only (Figure S1). This suggests that complications arising from hypertension may influence 7-HC-induced vasorelaxation.



**Figure 1.** Vasorelaxation effects of 7-HC in the superior mesenteric artery from SHR. (A) Representative original recordings of the effects of 7-HC in endothelium-intact and endothelium-denuded mesenteric arteries of rats. (B) Relaxation responses induced by 7-HC (0.001–300  $\mu\text{M}$ ) in endothelium-intact ( $\circ$ ,  $n = 6$ ) and endothelium-denuded ( $\bullet$ ,  $n = 6$ ) rat mesenteric arterial rings, pre-contracted with Phe (1  $\mu\text{M}$ ). (C) Bar graphs showing the contraction induced by Tyrode's solution containing 80 mM KCl before and after the 7-HC concentration–response curve. Results are expressed as means  $\pm$  S.E.M. Statistical analysis was performed using unpaired Student's  $t$ -tests.

## 2.2. The Vasorelaxation Induced by 7-HC Involves $\text{K}^+$ Channel Activation in Hypertensive Rats

In endothelium-denuded rings, 7-HC (0.001  $\mu\text{M}$  to 300  $\mu\text{M}$ ) inhibited the sustained tonic contraction induced by Tyrode's solution containing 80 mM KCl (effect (300  $\mu\text{M}$ ) =  $63.8 \pm 3.9\%$ , ( $n = 6$ )) in a concentration-dependent manner. The response induced by 7-HC was significantly reduced ( $p < 0.01$ ) compared with rings contracted with Phe (1  $\mu\text{M}$ ) (effect (300  $\mu\text{M}$ ) =  $100.4 \pm 9.8\%$ , ( $n = 6$ )) (Figure 2A). However, arteries pre-contracted with phenylephrine were pre-treated with 20 mM KCl. This resulted in the partial blocking of  $\text{K}^+$  efflux by increasing the extracellular  $\text{K}^+$  concentration to 20 mM and did not change the vasorelaxant effect after cumulative administration of 7-HC (0.001  $\mu\text{M}$  to 300  $\mu\text{M}$ ) (effect (300  $\mu\text{M}$ ) =  $98.4 \pm 15.5\%$ , ( $n = 7$ )) compared with the control (Figure 2B). These results may suggest that the effect of 7-HC could depend, at least in part, on the participation of additional  $\text{K}^+$  channels. Interestingly, the 7-HC-induced vasorelaxation was shifted to the right of the concentration–response curve in the presence of TEA (1 mM), glibenclamide (10  $\mu\text{M}$ ), and 4-AP (1 mM) but not  $\text{BaCl}_2$  (30  $\mu\text{M}$ ) (Figure 2C–F) compared with the control. These results suggest that the activation of  $\text{K}_{\text{ATP}}$ ,  $\text{BK}_{\text{Ca}}$ , and, to a lesser extent,  $\text{K}_{\text{V}}$ , but not  $\text{K}_{\text{ir}}$ , plays an important role in the relaxant effect of 7-HC in mesenteric arteries of SHR.




**Figure 2.**  $K^+$  channels' involvement in 7-HC-induced relaxation in the superior mesenteric artery of SHR. Vasorelaxation responses induced by 7-HC (0.001  $\mu$ M–300  $\mu$ M) in denuded mesenteric artery rings of SHR pre-contracted with Phe (1  $\mu$ M) (●,  $n = 6$ ), 80 mM KCl (■,  $n = 5$ ) (A), 20 mM KCl + Phe (▲,  $n = 5$ ) (B), 1 mM TEA (▼,  $n = 5$ ) (C), 10  $\mu$ M glibenclamide (◆,  $n = 5$ ) (D), 1 mM 4-AP (□,  $n = 5$ ) (E), and 30  $\mu$ M BaCl<sub>2</sub> (Δ,  $n = 5$ ) (F). Results are expressed as means  $\pm$  S.E.M. Statistical analysis was performed using unpaired Student's  $t$ -tests. \*  $p < 0.05$  or \*\*  $p < 0.01$  versus Phe.

### 2.3. Effect of 7-HC on the CaCl<sub>2</sub>-Induced Concentration–Response Curves

As shown in Figure 3A, the concentration–response curve of CaCl<sub>2</sub> in a depolarizing medium was shifted to the right in the presence of 7-HC (0.3, 3, 30, and 300  $\mu$ M) compared with the control. The maximal contraction of CaCl<sub>2</sub> was significantly (\*\*\*) attenuated by all concentrations of 7-HC (effect (3  $\mu$ M) =  $31.1 \pm 7.3\%$ , ( $n = 5$ ); effect


(30  $\mu\text{M}$ ) =  $36.6 \pm 8.3\%$ , ( $n = 5$ ); effect (300  $\mu\text{M}$ ) =  $43.9 \pm 5.8\%$ , ( $n = 5$ ). These data indicate that the mechanism of action of 7-HC involves reducing the  $\text{Ca}^{2+}$  influx.



**Figure 3.** Effect of 7-HC on  $\text{Ca}^{2+}$  influx. (A) Concentration–response curve of  $\text{CaCl}_2$  on mesenteric artery segments from SHR without endothelium, in the absence ( $\bullet$ ,  $n = 15$ ) or in the presence of 7-HC ( $\blacksquare$ , 3  $\mu\text{M}$ ,  $n = 5$ ;  $\blacktriangle$ , 30  $\mu\text{M}$ ,  $n = 5$ ;  $\blacktriangledown$ , 300  $\mu\text{M}$ ,  $n = 5$ ). Representative original recordings of the effects of 3  $\mu\text{M}$  7-HC (B), 30  $\mu\text{M}$  7-HC (C), and 300  $\mu\text{M}$  7-HC (D) on isolated mesenteric artery rings pre-incubated with 80 mM calcium-free KCl. The results are expressed as means  $\pm$  S.E.M. Statistical analysis was performed using one-way ANOVA followed by Bonferroni’s post-test. \*\*\*  $p < 0.001$  versus the control.

#### 2.4. Effect of Store-Operated Calcium Entry (SOCE) Inhibition on the Relaxation Induced by 7-HC in Hypertensive Rats


The vasorelaxation induced by 7-HC (0.001  $\mu\text{M}$  to 300  $\mu\text{M}$ ) was reduced in the presence of the SOCE inhibitor SKF-96365 (effect (300  $\mu\text{M}$ ) =  $52.4 \pm 9.3\%$  ( $n = 6$ )) compared with the control (effect (300  $\mu\text{M}$ ) =  $100.4 \pm 9.8\%$  ( $n = 6$ )) (Figure 4). This result suggests that the inhibition of  $\text{Ca}^{2+}$  influx by SOCE is involved in the vasorelaxation mechanism of 7-HC in mesenteric arteries from SHR.



**Figure 4.** The vasorelaxant effect of 7-HC in the presence of SOCE inhibition in mesenteric artery of SHR. (A) Relaxation responses induced by 7-HC (0.001–300  $\mu\text{M}$ ) on the isolated endothelium-denuded mesenteric artery rings, pre-contracted with Phe, in the presence or absence of SKF-96365 (10  $\mu\text{M}$ ) (■,  $n = 5$ ) compared with the control (●,  $n = 6$ ). Representative original recordings of the effects of 7-HC in isolated mesenteric artery rings, pre-contracted with phenylephrine (B) or SKF-96365 (10  $\mu\text{M}$ ) (C). Results are expressed as means  $\pm$  S.E.M. Statistical analysis was performed using unpaired Student's *t*-tests. \*  $p < 0.05$  or \*\*  $p < 0.01$  versus Phe.

### 2.5. Effect of 7-HC on the Mobilization of Calcium from Intracellular Stores

7-HC (30  $\mu\text{M}$  and 300  $\mu\text{M}$ ) markedly reduced the Phe-induced contraction in a  $\text{Ca}^{2+}$ -free Tyrode solution ( $52.8 \pm 14.5\%$  ( $n = 5$ );  $29.1 \pm 10.7\%$  ( $n = 5$ ), respectively) compared with the control ( $100 \pm 0.0$ ) (Figure 5A), suggesting that 7-HC may inhibit the release of  $\text{Ca}^{2+}$  from intracellular stores through the IP<sub>3</sub> receptors. Moreover, 7-HC (30  $\mu\text{M}$  and 300  $\mu\text{M}$ ) significantly decreased the caffeine-induced contraction in a  $\text{Ca}^{2+}$ -free Tyrode solution ( $17.3 \pm 3.2\%$  ( $n = 5$ ),  $11.5 \pm 4.3\%$  ( $n = 5$ ), respectively) compared with the control ( $100 \pm 0$ ) (Figure 5D). These data suggest that the mechanism of action of 7-HC involves reducing the release of  $\text{Ca}^{2+}$  from the intracellular stores, through both the IP<sub>3</sub> receptors and ryanodine receptors.




**Figure 5.** Effect of 7-HC on the mobilization of intracellular  $\text{Ca}^{2+}$ . Bar graphs showing the effect of 7-HC (30  $\mu\text{M}$  and 300  $\mu\text{M}$ ) on the  $\text{Ca}^{2+}$  mobilization from phenylephrine-sensitive (A) and caffeine-sensitive (D) intracellular stocks in a  $\text{Ca}^{2+}$ -free Tyrode solution. Representative recordings of the effects of 30  $\mu\text{M}$  and 300  $\mu\text{M}$  7-HC on isolated mesenteric artery rings incubated with 1  $\mu\text{M}$  phenylephrine (B,C) or 20 mM caffeine (E,F) in a calcium-free solution. The results are expressed as means  $\pm$  S.E.M. Statistical analysis was performed using one-way ANOVA followed by Bonferroni's post-test. \*  $p < 0.05$ , \*\*  $p < 0.01$ , or \*\*\*  $p < 0.001$  versus Phe (A) or caffeine (D).

### 2.6. Influence of 7-HC Pre-Incubation on Vascular Reactivity in Hypertensive Rats

The pharmacological efficacy of Phe was significantly increased in superior mesenteric arteries from SHR compared with wistar rats. In contrast, relaxation with the endothelium-independent vasodilator and nitric oxide (NO) donor SNP or acetylcholine was decreased in SHR compared with Wistar rats (Figure S2). These results suggest that the high blood pressure established in SHR promotes vascular dysfunction compared with wistar rats. Phenylephrine's concentration–response curve was shifted to the right after pre-incubation with 7-HC (100  $\mu\text{M}$ ), with a reduction in the  $\text{pD}_2$  values ( $p < 0.001$ ) compared with the results obtained in the absence of 7-HC ( $\text{pD}_2 = 6.14 \pm 0.05$  ( $n = 5$ , presence of 7-HC);  $\text{pD}_2 = 6.64 \pm 0.07$  ( $n = 5$ , absence of 7-HC)) (Figure 6A,B). However, the maximal response did not change ( $E_{\text{max}} = 176.8 \pm 6.1$  ( $n = 5$ );  $E_{\text{max}} = 178.6 \pm 4.0$  ( $n = 5$ ), respectively, for the presence and absence of 7-HC). In addition, SNP's concentration–response curve was shifted to the left after pre-incubation with 7-HC (100  $\mu\text{M}$ ), with an increase in the  $\text{pD}_2$  values ( $p < 0.05$ ) compared with the results obtained in the absence of 7-HC ( $\text{pD}_2 = 8.66 \pm 0.09$  ( $n = 6$ , presence of 7-HC);  $\text{pD}_2 = 9.44 \pm 0.31$  ( $n = 6$ , absence of 7-HC)), but no change the maximal response ( $E_{\text{max}} = 129.2 \pm 9.0$  ( $n = 6$ );  $E_{\text{max}} = 137.3 \pm 19.7$  ( $n = 6$ ), respectively, for the presence and absence of 7-HC) (Figure 6C,D). Acetylcholine's concentration–response curve was shifted to the left after pre-incubation with 7-HC (100  $\mu\text{M}$ ), without a significant change in the maximal response and  $\text{pD}_2$  values compared with the results obtained in the absence of 7-HC ( $E_{\text{max}} = 119.9 \pm 4.2$  ( $n = 5$ ),  $\text{pD}_2 = 7.31 \pm 0.28$  ( $n = 5$ , presence of 7-HC);  $E_{\text{max}} = 109.9 \pm 14.4$ ,  $\text{pD}_2 = 6.77 \pm 0.35$  ( $n = 5$ , absence of 7-HC)) (Figure 6E,F). Taken together, these results suggest that 7-HC can reduce the responsiveness of the

mesenteric arteries of SHR to  $\alpha$ 1-adrenergic agonists and increase the responsiveness to muscarinic agonists and an NO donor, improving vascular endothelium-dependent and independent relaxation.



**Figure 6.** 7-HC modulates the vascular reactivity of the mesenteric artery in SHR. (A) Contraction responses induced by Phe ( $10^{-10}$  to  $3 \times 10^{-5}$  M, cumulatively) in the absence of 7-HC ( $\bullet$ ,  $n = 5$ ) and the presence of 7-HC ( $\circ$ ,  $n = 5$ ). (B) Bar graph with the values of pD<sub>2</sub> from Phe. (C) Relaxation responses induced by SNP ( $10^{-11}$  to  $3 \times 10^{-6}$  M, cumulatively) in the absence of 7-HC ( $\blacksquare$ ,  $n = 5$ ) and the presence of 7-HC ( $\square$ ,  $n = 5$ ). (D) Bar graph with the values of pD<sub>2</sub> from SNP. (E) Relaxation responses induced by Ach ( $10^{-11}$  to  $3 \times 10^{-6}$  M, cumulatively) in the absence of 7-HC ( $\blacktriangle$ ,  $n = 5$ ) and the presence of 7-HC ( $\triangle$ ,  $n = 5$ ). (F) Bar graph with the values of pD<sub>2</sub> from ACh. The results are expressed as means  $\pm$  S.E.M. Statistical analysis was performed using unpaired Student's *t*-tests. \*  $p < 0.05$ , \*\*  $p < 0.01$  or \*\*\*  $p < 0.001$  versus the absence of 7-HC.



### 3. Discussion

The present study indicated that the vasorelaxant effect of 7-HC in SHR occurs via an endothelium-independent pathway, which likely consists of the activation of potassium channels ( $K_{ATP}$ ,  $BK_{Ca}$ , and  $K_v$ ), and reductions in both calcium influx and intracellular calcium mobilization from the sarcoplasmic reticulum. Additionally, 7-HC reduced the pharmacological potency of  $\alpha$ 1-adrenergic agonists while increasing the potency of muscarinic agonists and NO donors in mesenteric arteries from SHR, improving vascular endothelium-dependent and independent relaxation. Interestingly, even with the pathophysiological changes generated in the hypertensive condition, 7-HC maintained its antihypertensive vascular effects, supporting its therapeutic potential.

Peripheral vascular resistance plays an essential role in the regulation of blood pressure, and several substances can act on this parameter, contributing to the control of arterial pressure [19]. Vascular endothelial and smooth muscle cells play critical roles in regulating vascular tension [20]. Vascular smooth muscle cells (VSMCs) are exposed to mechanical stress and the pressure of blood flow, and play an important role in maintaining vascular tone and resistance [21]. In addition, the endothelium regulates vascular tone by synthesizing and releasing an array of endothelium-derived relaxing factors, such as nitric oxide and endothelium-dependent hyperpolarization factors, as well as endothelium-derived contracting factors [22,23]. In the current study, when the endothelium of mesenteric artery rings was removed, the vasorelaxant effect of 7-HC was not significantly impaired, indicating that the vasorelaxant activity of 7-HC does not rely on the integrity of the endothelial layer. Thus, subsequent experiments were carried out to investigate 7-HC's endothelium-independent vasorelaxation mechanisms.

Our research group has previously demonstrated that 7-HC is nontoxic for cardiomyocytes [18,24] and did not have an irreversible action on the arterial contractile machinery in normotensive rats [18]. Our observations corroborate what is shown in the literature, since we also observed that tissue contractility of mesenteric artery rings from SHR was maintained after removing 7-HC from the tissue bath solution, reinforcing the notion that 7-HC has reversible actions on the contractile machinery and probably has no toxic effects on the vascular myocyte in SHR.

Subsequently, we designed experiments to investigate the 7-HC-mediated vasorelaxant mechanisms of action in the arteries of hypertensive animals. It has been reported in the literature that the opening of  $K^+$  channels promotes hyperpolarization of the membrane potential and closure of voltage-gated channels, which decreases the entry of  $Ca^{2+}$  and causes vasorelaxation [25–27]. In our study, the increase in extracellular  $K^+$  concentrations (from 4 to 20 mM) did not significantly attenuate 7-HC induced vasorelaxation, which diverges from the results obtained by Alves and colleagues (2020) [18] in the mesenteric artery from normotensive rats. Such differences in the tissue response can be attributed to the altered expression or activity of  $K^+$  channels, as in the experimental model of hypertension [28]. For example, an attenuation of the  $K_v$  channel's activity during hypertension has been described in VSMCs from rat mesenteric arteries [29–31]. However, 7-HC vasorelaxation was attenuated considerably when the rings were exposed to Tyrode's solution containing 80 mM KCl, suggesting the participation of  $K^+$  channels in 7-HC-induced vasorelaxation. Furthermore, the higher concentration of  $K^+$  in the solution (>20 mM) could influence the response to 7-HC in the superior mesenteric artery rings of SHR. Thus, the involvement and activation of distinct types of  $K^+$  channels in 7-HC-induced vasorelaxation were investigated.

$K^+$  channels play an important role in the regulation of vascular tone, since the activation of  $K^+$  channels is known to promote blood vessel relaxation through membrane hyperpolarization, leading to a decreased probability of voltage sensitive opening of the calcium channels, thereby lowering intracellular calcium levels [32,33]. The regional vasodilator profile of  $K^+$  channels' activation is heterogeneous and, in patients, only pinacidil, minoxidil, and diazoxide are currently used as antihypertensive drugs.

Minoxidil is an arteriolar vasodilator, and the antihypertensive activity of minoxidil is due to its sulfate metabolite. Minoxidil sulfate opens the adenosine tri-phosphate-sensitive potassium channels in vascular smooth muscle cells [34]. Pinacidil, another known potassium channel opener, also produces vasodilation comparable with that of minoxidil [35].

Disorders in the structure and function of  $K^+$  channels contribute to the pathophysiological conditions of some vascular diseases, including hypertension [31].  $K^+$  currents are essential for preserving membrane stability by suppressing the membranes' excitability. However, in the hypertensive condition, the  $K^+$  currents are lower than in the normotensive state [36]. Thus, a reduction in the potassium efflux current can affect the excitability of vascular cells, causing membrane depolarization and resulting in hypertension [37].

There are different subtypes of  $K^+$  channels. Calcium-activated  $K^+$  ( $K_{Ca}$ ) channels are the most abundant  $K^+$  channels found in the VSMC and are activated by increases in the intracellular  $Ca^{2+}$  concentration [25]. In particular,  $BK_{Ca}$  channels in the VSMC are especially important for regulating vascular tone and are responsible for a significant proportion of potassium efflux [38]. In our study, to evaluate the participation of  $BK_{Ca}$  channels in the 7-HC-mediated relaxation of SHR mesenteric rings, assays were performed utilizing 1 mM TEA, as TEA blocks  $BK_{Ca}$  channels at this concentration [39], and the vasorelaxant effect of 7-HC was significantly reduced, suggesting the participation of these channels in the mechanism of 7-HC.

A similar response was observed in experiments with mesenteric rings pre-incubated with glibenclamide (10  $\mu$ M) and 4-AP (1 mM), which are blockers of the ATP-sensitive  $K^+$  channels and the voltage-sensitive  $K^+$  channels, respectively. In both cases, the concentration-response curve was shifted to the right, indicating a decrease in the pharmacological potency of 7-HC when challenged with the respective blockers, suggesting the participation of ATP-sensitive  $K^+$  channels, or voltage-sensitive  $K^+$  channels.

The same response pattern was not observed when the mesenteric arteries were preincubated with  $BaCl_2$  (30  $\mu$ M), an inward-rectifier blocker of  $K^+$  channels. In this case, the vasorelaxant effect of 7-HC remained unaltered in the presence or absence of the blocker, suggesting that these channels may not participate in 7-HC's mechanism of action in the mesenteric arteries of SHR. By contrast, Alves and colleagues (2020) [18] proposed that 7-HC caused the opening of  $BaCl_2$ -sensitive inward rectifying  $K^+$  channels in the mesenteric rings from normotensive rats. Although there is no difference in the expression of inward rectifying  $K^+$  channels in SHR and normotensive Wistar-Kyoto rats, the inward rectifying  $K^+$  channels in SHR are less involved in the spread of Ach-induced hyperpolarization compared with normotensive rats [40]. Therefore, the different responses observed following the application of 7-HC in both strains may rely on the different physiological  $K_{ir}$  responses in normotensive rats and SHR. In addition to the elevated vascular tone observed in humans and in experimental hypertension models, such as SHR [41], there have also been reports that these cases involve alterations of various families of  $Ca^{2+}$  and/or  $K^+$  channels in the expression or functionality in the VSMC, consisting of a variety of subtypes including voltage-gated  $Ca^{2+}$  channels ( $Ca_v$ ) [42], voltage-dependent  $K^+$  channels ( $K_v$ ) [43], and the broad-conductance  $Ca^{2+}$ -activated  $K^+$  channels ( $BK_{Ca}$ ) [44,45]. This demonstrates the relevance of this study, as we observed the effect of 7-HC on the  $BK_{Ca}$  and  $K_v$  channels, which are known to participate in the pathophysiology of hypertension.

Calcium channel blockers (CCBs) were shown to lower elevated blood pressure more than 50 years ago. The drug now known as verapamil was described in 1962 as increasing renal blood flow and lowering blood pressure in hypertensive but not in normotensive subjects [46], and is still used in the treatment of hypertension. CCBs are recommended in the Eighth Joint National Committee (JNC8) guidelines to be used as a first-line treatment alone or in combination with other antihypertensives in all patients with hypertension regardless of age and race, except for patients with chronic kidney disease where angiotensin-converting enzyme inhibitors and angiotensin receptor blockers inhibitors are the recommended first-line treatment [47].

The scientific literature has reported significant abnormalities in calcium signaling in the VSMCs of SHR compared with normotensive rats [48–52]. Both extracellular  $\text{Ca}^{2+}$  influx through  $\text{Ca}_V$  or store-operated calcium entry (SOCE) can increase intracellular calcium. Previous studies have shown that the current densities of  $\text{Ca}_V$  are elevated in mesenteric VSMCs of SHR compared with WKY rats [53–55]. This elevation in the  $\text{Ca}^{2+}$  currents may be due, at least in part, to upregulation of the expression of the pore-forming  $\alpha_1C$  subunit of the  $\text{Ca}_V$  channel in the mesenteric artery from SHR [50]. SOCE increases the intracellular  $\text{Ca}^{2+}$  concentration, which is essential for producing the intracellular calcium signals needed to regulate vascular tone [56,57]. 7-HC (0.3, 3, 30, and 300  $\mu\text{M}$ ) inhibits  $\text{Ca}^{2+}$  influx-mediated vasoconstriction in mesenteric rings pre-incubated with a depolarizing solution, suggesting that 7-HC may inhibit  $\text{Ca}_V$ .

In addition, experiments were performed to assess the involvement of SOCE in the relaxation induced by 7-HC, using the nonspecific channel blocker SKF 96365. The results indicated that the 7-HC-mediated relaxation was reduced in the presence of a SOCE inhibitor, suggesting that 7-HC-induced vasorelaxation may be due to a reduction in the influx of  $\text{Ca}^{2+}$  through the SOC.

Inositol 1,4,5-trisphosphate receptors (IP3 R) and ryanodine receptors are the channels that most commonly mediate the release of  $\text{Ca}^{2+}$  from intracellular stores [58,59]. When activated by caffeine, the ryanodine receptor works through a  $\text{Ca}^{2+}$ -induced  $\text{Ca}^{2+}$  release mechanism (CICR). Alternatively, Phe, an  $\alpha_1$ -adrenergic agonist, enhances extracellular  $\text{Ca}^{2+}$  entry by activating both ROCCs and  $\text{Ca}_V$  [60], as well as inducing the release of  $\text{Ca}^{2+}$  from IP3 R. However, the vasoconstriction induced by Phe in the  $\text{Ca}^{2+}$ -free medium is due to endo/sarcoplasmic reticulum (ER/SR)  $\text{Ca}^{2+}$  release through the IP3-sensitive  $\text{Ca}^{2+}$  channels. In our experiments, this vasoconstriction was inhibited by 7-HC (30, 300  $\mu\text{M}$ ) in a concentration-dependent manner. This suggests that the endothelium-independent relaxation evoked by 7-HC may inhibit the mobilization of  $\text{Ca}^{2+}$  from intracellular stores through the IP3-sensitive  $\text{Ca}^{2+}$  channels. Furthermore, 7-HC (30, 300  $\mu\text{M}$ ) also decreases caffeine-induced contraction in endothelium-denuded arteries, suggesting the potential involvement of ryanodine receptors induced by 7-HC. These results indicate that the vasorelaxant response of 7-HC may also involve, at least in part, the inhibition of extracellular  $\text{Ca}^{2+}$  entry by  $\text{Ca}_V$  and the inhibition of  $\text{Ca}^{2+}$  released from the internal store by the IP3 receptors and ryanodine receptors, leading to a decrease in intracellular  $\text{Ca}^{2+}$ .

Hypertension is directly related to endothelial dysfunction and low NO bioavailability due to the decreased NO production and/or increased NO degradation caused by oxidative stress [61]. Therefore, this may lead to increased contractile and/or decreased vasodilatory responses. In the present study, pre-incubation with 7-HC affected contractile responses in the mesenteric arteries of hypertensive rats. We observed that the pharmacological potency of the  $\alpha_1$ -adrenergic agonist was reduced in the presence of 7-HC, reinforcing that in a hypertensive state, where adrenergic responsiveness is increasing, 7-HC may be a therapeutic alternative.

The ACh-induced relaxation in isolated arteries relies on muscarinic receptors in the endothelial cells to improve several endothelial factors [62]. At the same time, SNP, a NO donor, directly induces vasorelaxation in the vascular smooth muscle, mainly by activating soluble guanylate cyclase [63]. It has been established that the reduced relaxation response to ACh or SNP in SHR, due to endothelial dysfunction, is caused by oxidative stress [64,65], decreased production and bioavailability of NO [66–68], eNOS uncoupling [64,69], and hyper-responsiveness to contracting agents such as angiotensin II [70–72] and endothelin-1 [73]. Additionally, ACh or SNP-mediated vasorelaxation was diminished in SHR compared with normotensive rats [67,74]. Our study showed that 7-HC improved the endothelium-dependent and independent vasorelaxation evoked by ACh and SNP, respectively, suggesting that 7-HC increased the arterial sensitivity to vasorelaxant signaling pathways, which makes this coumarin a potential drug for treating diseases with endothelial dysfunction. However, further studies are needed to assess whether 7-HC

could prevent chronic endothelial dysfunction, improve NO bioavailability, and prevent vascular remodeling in hypertensive conditions.

#### 4. Material and Methods

##### 4.1. Animals

Male spontaneously hypertensive rats (SHR) at the age of 12–15 weeks, weighing 200 to 300 g, were used in all experiments. The animals were supplied by the animal facility of the Neuroscience Laboratory at the Institute of Health Sciences at the Federal University of Bahia and kept under a controlled temperature ( $23 \pm 1.0$  °C) and a light–dark cycle of 12 h (6:00 a.m. to 6:00 p.m.), with free access to food and water. The tests were performed in accordance with the guidelines for the care and use of laboratory rats adopted by the National Council for Animal Experiments Control (CONCEA–Brazil). It was approved by the Ethics Committee on Animal Use from the Institute of Health Sciences, Federal University of Bahia (CEUA/UFBA no. 130/2017).

##### 4.2. Drugs and Solutions

The drugs used in this study were: L-phenylephrine hydrochloride (Phe), acetylcholine chloride (ACh), sodium nitroprusside (SNP), barium chloride ( $\text{BaCl}_2$ ), 4-aminopyridine (4-AP), tetraethylammonium (TEA), glibenclamide (Glib), caffeine, and cremophor, all acquired from Sigma-Aldrich (Sigma Chemical Co., Saint Louis, MO, USA). All drugs were dissolved in distilled water.  $\text{K}^+$ -depolarizing solutions (20 and 80 mM KCl) were prepared by replacing equimolar NaCl with 20 or 80 mM of KCl. The 7-hydroxycoumarin (7-HC) was solubilized in cremophor and diluted to the desired concentrations with distilled water, as described by Alves and colleagues (2020) [18]. Tyrode's physiological solution was used in all arterial experiments, with the following composition at 37 °C (in mM): NaCl, 158.3; KCl, 4.0;  $\text{CaCl}_2$ , 2.0;  $\text{MgCl}_2$ , 1.05;  $\text{NaH}_2\text{PO}_4$ , 0.42;  $\text{NaHCO}_3$ , 10.0; glucose, 5.6. The reagents were acquired from Sigma (Sigma-Aldrich, St. Louis, MO, USA) or Vetec (Vetec, Rio de Janeiro, Brazil).

##### 4.3. Isolation of the Superior Mesenteric Arteries

SHR were euthanized in a  $\text{CO}_2$  chamber, and the middle portion of the superior mesenteric arteries was removed, as previously described by Silva and colleagues (2011) [75]. The mesenteric rings (2.0 mm) were stabilized with an optimal resting tension of 7.5 mN for 60 min in an organ bath with physiological Tyrode's solution (at 37 °C and pH 7.4) and gassed with a carbogenic mixture (95%  $\text{O}_2$  and 5%  $\text{CO}_2$ ). The isometric tension was recorded by a force transducer (Insight, Ribeirão Preto, Sao Paulo, Brazil) coupled with an amplifier-recorder (Insight, Ribeirão Preto, Sao Paulo, Brazil) and a personal computer equipped with data acquisition software. The presence of a functional endothelium was assessed by the ability of ACh (1  $\mu\text{M}$ ) to induce at least a 90% relaxation response in mesenteric artery rings pre-contracted with Phe (1  $\mu\text{M}$ ). Rings that relaxed by less than 10% were considered endothelium-denuded/damaged.

##### 4.4. Effect of 7-HC on the Vasculature

After the stabilization period, two successive contractions of similar magnitude were induced with Phe (1  $\mu\text{M}$ ), an  $\alpha$ 1-adrenergic agonist, in endothelium-intact and endothelium-denuded rings. In the tonic phase of the second contraction, increasing concentrations of 7-HC (0.001–300  $\mu\text{M}$ ) were cumulatively added to the organ bath, and the effects were compared. Another set of experiments investigated the reversibility of the effects of 7-HC. In these assays, a contraction was performed with Tyrode's solution, with 80 mM KCl, in absence of 7-HC; after the cumulative addition of 7-HC, the tissue was washed, stabilized, and then contracted again with Tyrode's solution with 80 mM KCl.

#### 4.5. Evaluation of K<sup>+</sup> Channel Activity in the 7-HC-Induced Vasorelaxation Response

Endothelium-denuded ring preparations were pre-incubated with Tyrode's solution containing 80 mM KCl, or Phe (1  $\mu$ M) in Tyrode's solution with 20 mM KCl, and the relaxing effect induced by 7-HC (0.001–300  $\mu$ M) was obtained. In addition, the involvement of K<sup>+</sup> channels in the vasorelaxant response induced by 7-HC was evaluated following pre-incubation of the endothelium-denuded rings for 30 min with the following different pharmacological agents: TEA (1 mM), a nonselective blocker of large-conductance Ca<sup>2+</sup>-activated K<sup>+</sup> (BK<sub>Ca</sub>) channels [76]; 4-AP (1 mM), a voltage-operated K<sup>+</sup> channel (K<sub>v</sub>) blocker [77]; BaCl<sub>2</sub> (30  $\mu$ M), a blocker of inward rectifying K<sup>+</sup> channels (K<sub>ir</sub>) [78]; and glibenclamide (10  $\mu$ M), a specific ATP-sensitive K<sup>+</sup> (K<sub>ATP</sub>) channel subtype-selective blocker [79].

#### 4.6. Investigation of the Effects Induced by 7-HC in Ca<sup>2+</sup> Influx

The method was adapted from that described by Alves et al. 2020 [18]. To investigate the influence of 7-HC on the influx of Ca<sup>2+</sup> in endothelium-denuded rings, cumulative concentrations of CaCl<sub>2</sub> (100–10,000  $\mu$ M) were added to a depolarization medium of Ca<sup>2+</sup>-free Tyrode's solution containing 80 mM KCl (for 15 min) in the absence (control) or presence of different concentrations of 7-HC (3, 30 and 300  $\mu$ M). Another experiment was performed to investigate the influx of Ca<sup>2+</sup> through SOCE (store-operated calcium entry), where experiments were performed using SKF 96365 (10  $\mu$ M, a SOCE blocker). Endothelium-denuded rings were pre-incubated with SKF for 30 min; subsequently, Phe (1  $\mu$ M) was added.

#### 4.7. Investigation of 7-HC-Induced Mobilization of Calcium from Intracellular Stores

After the stabilization period, vessels were pre-contracted with Tyrode's solution containing 80 mM KCl for 3 min, then washed with a Ca<sup>2+</sup>-free Tyrode solution containing EGTA 1 mM, followed by the addition of 1  $\mu$ M Phe (in an organ bath at 37 °C) or 20 mM caffeine (in an organ bath at 23 °C). Rings were rinsed with Tyrode's physiological solution, and Tyrode's solution containing 80 mM KCl was added for 3 min to load the Ca<sup>2+</sup> stores within the vascular smooth muscle. This procedure was repeated until two transient contractions of similar magnitude had been obtained. Next, this experimental protocol was repeated by incubating 7-HC (10 and 100  $\mu$ M) in Ca<sup>2+</sup>-free Tyrode's solution for 5 min, then applying Phe or caffeine, as previously described [80]. The role of 7-HC on the mobilization of calcium from the intracellular stores was assessed by comparing Phe or caffeine-mediated contraction in the absence or presence of 7-HC via the inositol 1,4,5-triphosphate (IP3) and ryanodine receptors, respectively.

#### 4.8. Influence of 7-HC on the Vascular Reactivity in Hypertensive Rats

To investigate the influence of 7-HC on arterial reactivity in SHR, cumulative concentration-response curves were obtained for  $\alpha$ 1- adrenergic (Phe), muscarinic (ACh), and nitric oxide (NO) donor (SNP) agonists. After the stabilization period, the arterial rings were contracted with Phe (1  $\mu$ M) and then assessed to determine the relaxant effect induced by ACh (10<sup>-11</sup> to 3  $\times$  10<sup>-6</sup> M) or SNP (10<sup>-11</sup> to 3  $\times$  10<sup>-6</sup> M). The contractile response to Phe (10<sup>-10</sup> to 3  $\times$  10<sup>-5</sup> M) was also assessed. Next, the tissues were washed and incubated for 30 min with 7-HC (100  $\mu$ M) and then challenged again with the same cumulative curves in the presence of 7-HC.

#### 4.9. Data Analysis

Data are presented as the mean  $\pm$  standard error of the mean (S.E.M.), and *n* represents the number of mesenteric artery rings obtained from different animals. All calculations were performed using Prism software version 5.0 (GraphPad Software Inc., CA, USA). The pharmacological potency, expressed as pD<sub>2</sub> (pD<sub>2</sub> =  $-\log EC_{50}$ ), was defined in the experiments where E<sub>max</sub> (the maximum effect generated by the agonist) could be determined. Data were analyzed with unpaired *t*-tests, two-tailed tests, and 95% confidence intervals.

One-way ANOVA followed by Bonferroni's post-test was used when data were compared among more than two groups. Statistical significance was set at  $p < 0.05$ .

## 5. Conclusions

The results demonstrated that 7-HC induces vasorelaxation in the mesenteric artery from animals with essential hypertension involving the activation of  $K^+$  channels, inhibition of the  $Ca^{2+}$  influx through  $Cav$  and  $SOC$ , and a reduction in the release of calcium from ryanodine- and  $IP_3$ -sensitive intracellular stores. Furthermore, these results demonstrated that the vasorelaxant effect of 7-HC on the vascular smooth muscle cells from the superior mesenteric artery of SHR is independent of endothelium-derived relaxing factors, most likely through the activation of the  $K_{ATP}$ ,  $BK_{Ca}$ , and  $K_v$  channels. Additionally, 7-HC can attenuate calcium influx, which leads to a reduction in the mobilization of intracellular calcium. Furthermore, 7-HC also seems to be able to improve endothelial function and the relaxing ability of vascular smooth muscle, attenuating the vascular dysfunctions associated with hypertension. Thus, 7-HC is a promising compound with direct vascular activity and may be considered a potential strategy for the management of hypertension.

**Supplementary Materials:** The following supporting information can be downloaded at <https://www.mdpi.com/article/10.3390/molecules27217324/s1>. Figure S1: Vasorelaxation effects of 7-HC in the superior mesenteric artery from Wistar rats and SHR. Figure S2: Vascular abnormalities in the superior mesenteric artery from hypertensive rats compared with normotensive rats.

**Author Contributions:** Conceptualization, R.L.C.J. and D.F.S.; methodology, R.L.C.J., I.L.P.S., D.S.B., G.B.C.L. and L.B.S.; formal analysis, R.L.C.J., F.A.A. and R.A.M.; data curation, R.L.C.J., R.A.M. and F.A.A.; writing—original draft preparation, R.L.C.J., F.A.A. and R.A.M.; writing—review and editing, R.L.C.J., Q.L.A. and D.F.S.; supervision, Q.L.A. and D.F.S.; project administration, R.L.C.J. and D.F.S.; funding acquisition, D.F.S. All authors have read and agreed to the published version of the manuscript.

**Funding:** This study was funded by grants from National Council for Scientific and Technological Development (CNPq), process number 233867/2014–7 and 306106/2017–5; the Coordination for the Improvement of Higher Education Personnel–Brazil (CAPES), finance code 001; the Foundation for Research Support in the State of Bahia (FAPESB). D.F.S. is a CNPq Research Fellow.

**Institutional Review Board Statement:** The tests were performed in accordance with the guidelines for the care and use of laboratory rats adopted by the National Council for Animal Experiments Control (CONCEA–Brazil). It was approved by the Ethics Committee on Animal Use of the Institute of Health Sciences, Federal University of Bahia (CEUA/UFBA no. 130/2017).

**Data Availability Statement:** Not applicable.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. GBD 2017 Risk Factor Collaborators. Global, Regional, and National Comparative Risk Assessment of 84 Behavioural, Environmental and Occupational, and Metabolic Risks or Clusters of Risks for 195 Countries and Territories, 1990–2017: A Systematic Analysis for the Global Burden of Disease Study. *Lancet* **2018**, *392*, 1923–1994. [[CrossRef](#)]
2. Unger, T.; Borghi, C.; Charchar, F.; Khan, N.A.; Poulter, N.R.; Prabhakaran, D.; Ramirez, A.; Schlaich, M.; Stergiou, G.S.; Tomaszewski, M.; et al. 2020 International Society of Hypertension Global Hypertension Practice Guidelines. *Hypertension* **2020**, *75*, 1334–1357. [[CrossRef](#)] [[PubMed](#)]
3. Al Ghorani, H.; Götzinger, F.; Böhm, M.; Mahfoud, F. Arterial Hypertension—Clinical Trials Update 2021. *Nutr. Metab. Cardiovasc. Dis.* **2022**, *32*, 21–31. [[CrossRef](#)] [[PubMed](#)]
4. Williams, B.; Mancia, G.; Spiering, W.; Agabiti Rosei, E.; Azizi, M.; Burnier, M.; Clement, D.L.; Coca, A.; de Simone, G.; Dominiczak, A.; et al. 2018 ESC/ESH Guidelines for the Management of Arterial Hypertension. *Eur. Heart J.* **2018**, *39*, 3021–3104. [[CrossRef](#)] [[PubMed](#)]
5. Acelajado, M.C.; Hughes, Z.H.; Oparil, S.; Calhoun, D.A. Treatment of Resistant and Refractory Hypertension. *Circ. Res.* **2019**, *124*, 1061–1070. [[CrossRef](#)] [[PubMed](#)]

6. Carey, R.M.; Calhoun, D.A.; Bakris, G.L.; Brook, R.D.; Daugherty, S.L.; Dennison-Himmelfarb, C.R.; Egan, B.M.; Flack, J.M.; Gidding, S.S.; Judd, E.; et al. Resistant Hypertension: Detection, Evaluation, and Management: A Scientific Statement From the American Heart Association. *Hypertension* **2018**, *72*, E53–E90. [[CrossRef](#)]
7. Lamirault, G.; Artifoni, M.; Daniel, M.; Barber-Chamoux, N. Nantes University Hospital Working Group On Hypertension Resistant Hypertension: Novel Insights. *Curr. Hypertens. Rev.* **2020**, *16*, 61–72. [[CrossRef](#)]
8. Srikrishna, D.; Godugu, C.; Dubey, P.K. A Review on Pharmacological Properties of Coumarins. *Mini Rev. Med. Chem.* **2018**, *18*, 113–141. [[CrossRef](#)]
9. He, J.-Y.; Zhang, W.; He, L.-C.; Cao, Y.-X. Imperatorin Induces Vasodilatation Possibly via Inhibiting Voltage Dependent Calcium Channel and Receptor-Mediated  $\text{Ca}^{2+}$  Influx and Release. *Eur. J. Pharmacol.* **2007**, *573*, 170–175. [[CrossRef](#)] [[PubMed](#)]
10. Najmanová, I.; Doseděl, M.; Hrdina, R.; Anzenbacher, P.; Filipický, T.; Říha, M.; Mladěnka, P. Cardiovascular Effects of Coumarins besides Their Antioxidant Activity. *Curr. Top. Med. Chem.* **2015**, *15*, 830–849. [[CrossRef](#)]
11. Oliveira, E.J.; Romero, M.A.; Silva, M.S.; Silva, B.A.; Medeiros, I.A. Intracellular Calcium Mobilization as a Target for the Spasmolytic Action of Scopoletin. *Planta Med.* **2001**, *67*, 605–608. [[CrossRef](#)] [[PubMed](#)]
12. De Almeida Barros, T.A.; de Freitas, L.A.R.; Filho, J.M.B.; Nunes, X.P.; Giulietti, A.M.; de Souza, G.E.; dos Santos, R.R.; Soares, M.B.P.; Villarreal, C.F. Antinociceptive and Anti-Inflammatory Properties of 7-Hydroxycoumarin in Experimental Animal Models: Potential Therapeutic for the Control of Inflammatory Chronic Pain. *J. Pharm. Pharmacol.* **2010**, *62*, 205–213. [[CrossRef](#)] [[PubMed](#)]
13. de Araújo, R.S.A.; Guerra, F.Q.S.; de Lima, E.O.; de Simone, C.A.; Tavares, J.F.; Scotti, L.; Scotti, M.T.; de Aquino, T.M.; de Moura, R.O.; Mendonça, F.J.B.; et al. Synthesis, Structure-Activity Relationships (SAR) and in Silico Studies of Coumarin Derivatives with Antifungal Activity. *Int. J. Mol. Sci.* **2013**, *14*, 1293–1309. [[CrossRef](#)] [[PubMed](#)]
14. Li, D.; Wang, N.; Zhang, J.; Ma, S.; Zhao, Z.; Ellis, E.M. Hepatoprotective Effect of 7-Hydroxycoumarin against Methyl Glyoxal Toxicity via Activation of Nrf2. *Chem. Biol. Interact.* **2017**, *276*, 203–209. [[CrossRef](#)]
15. de Lima, F.O.; Nonato, F.R.; Couto, R.D.; Barbosa Filho, J.M.; Nunes, X.P.; Ribeiro dos Santos, R.; Soares, M.B.P.; Villarreal, C.F. Mechanisms Involved in the Antinociceptive Effects of 7-Hydroxycoumarin. *J. Nat. Prod.* **2011**, *74*, 596–602. [[CrossRef](#)]
16. Baccard, N.; Mechiche, H.; Nazeyrollas, P.; Manot, L.; Lamiable, D.; Devillier, P.; Millart, H. Effects of 7-Hydroxycoumarin (Umbelliferone) on Isolated Perfused and Ischemic-Reperfused Rat Heart. *Arzneimittelforschung* **2000**, *50*, 890–896. [[CrossRef](#)]
17. Jagadeesh, G.S.; Nagoor Meeran, M.F.; Selvaraj, P. Protective Effects of 7-Hydroxycoumarin on Dyslipidemia and Cardiac Hypertrophy in Isoproterenol-Induced Myocardial Infarction in Rats. *J. Biochem. Mol. Toxicol.* **2016**, *30*, 120–127. [[CrossRef](#)]
18. Alves, Q.L.; dos Anjos Moraes, R.; Froes, T.Q.; Castilho, M.S.; de Araújo, R.S.A.; Barbosa-Filho, J.M.; Meira, C.S.; Pereira Soares, M.B.; Silva, D.F. Inhibition of Intracellular  $\text{Ca}^{2+}$  Mobilization and Potassium Channels Activation Are Involved in the Vasorelaxation Induced by 7-Hydroxycoumarin. *Eur. J. Pharmacol.* **2020**, *887*, 173525. [[CrossRef](#)]
19. Mayet, J.; Hughes, A. Cardiac and Vascular Pathophysiology in Hypertension. *Heart* **2003**, *89*, 1104–1109. [[CrossRef](#)] [[PubMed](#)]
20. Vanhoutte, P.M.; Shimokawa, H.; Feletou, M.; Tang, E.H.C. Endothelial Dysfunction and Vascular Disease—A 30th Anniversary Update. *Acta Physiol.* **2017**, *219*, 22–96. [[CrossRef](#)]
21. Zhu, Y.; Qu, J.; He, L.; Zhang, F.; Zhou, Z.; Yang, S.; Zhou, Y. Calcium in Vascular Smooth Muscle Cell Elasticity and Adhesion: Novel Insights Into the Mechanism of Action. *Front. Physiol.* **2019**, *10*, 852. [[CrossRef](#)] [[PubMed](#)]
22. Santos, S.S.; Jesus, R.L.C.; Simões, L.O.; Vasconcelos, W.P.; Medeiros, I.A.; Veras, R.C.; Casais-E-Silva, L.L.; Silva, D.F. NO Production and Potassium Channels Activation Induced by Crotalus Durissus Cascavella Underlie Mesenteric Artery Relaxation. *Toxicol* **2017**, *133*, 10–17. [[CrossRef](#)] [[PubMed](#)]
23. Moraes, R.D.A.; Alves, Q.L.; Camargo, S.B.; de Azevedo Medeiros, C.F.; de Melo Jesus, A.; da Hora, V.R.S.; Stiz, D.S.; Corrêa, R.; Cechinel-Filho, V.; Silva, D.F. Itaconimides Derivatives Induce Relaxation in Mesenteric Artery and Negative Inotropism by Inhibition of  $\text{Ca}^{2+}$  Influx. *Pharmacol. Rep.* **2020**, *72*, 890–902. [[CrossRef](#)]
24. Cruz, L.F.; de Figueiredo, G.F.; Pedro, L.P.; Amorin, Y.M.; Andrade, J.T.; Passos, T.F.; Rodrigues, F.F.; Souza, I.L.A.; Gonçalves, T.P.R.; Dos Santos Lima, L.A.R.; et al. Umbelliferone (7-Hydroxycoumarin): A Non-Toxic Antidiarrheal and Antiulcerogenic Coumarin. *Biomed. Pharmacother.* **2020**, *129*, 110432. [[CrossRef](#)]
25. Nelson, M.T.; Quayle, J.M. Physiological Roles and Properties of Potassium Channels in Arterial Smooth Muscle. *Am. J. Physiol.* **1995**, *268*, C799–C822. [[CrossRef](#)]
26. Mondéjar-Parreño, G.; Cogolludo, A.; Perez-Vizcaino, F. Potassium ( $\text{K}^+$ ) Channels in the Pulmonary Vasculature: Implications in Pulmonary Hypertension Physiological, Pathophysiological and Pharmacological Regulation. *Pharmacol. Ther.* **2021**, *225*, 107835. [[CrossRef](#)]
27. van der Horst, J.; Manville, R.W.; Hayes, K.; Thomsen, M.B.; Abbott, G.W.; Jepps, T.A. Acetaminophen (Paracetamol) Metabolites Induce Vasodilation and Hypotension by Activating Kv7 Potassium Channels Directly and Indirectly. *Arterioscler. Thromb. Vasc. Biol.* **2020**, *40*, 1207–1219. [[CrossRef](#)] [[PubMed](#)]
28. Cox, R.H. Changes in the Expression and Function of Arterial Potassium Channels during Hypertension. *Vascul. Pharmacol.* **2002**, *38*, 13–23. [[CrossRef](#)]
29. Bratz, I.N.; Swafford, A.N.; Kanagy, N.L.; Dick, G.M. Reduced Functional Expression of  $\text{K}^+$  Channels in Vascular Smooth Muscle Cells from Rats Made Hypertensive with N[omega]-Nitro-L-Arginine. *Am. J. Physiol. Heart Circ. Physiol.* **2005**, *289*, H1284–H1290. [[CrossRef](#)]
30. Cox, R.H.; Lozinskaya, I.; Dietz, N.J. Differences in  $\text{K}^+$  Current Components in Mesenteric Artery Myocytes from WKY and SHR. *Am. J. Hypertens.* **2001**, *14*, 897–907. [[CrossRef](#)]

31. Nieves-Cintrón, M.; Syed, A.U.; Nystoriak, M.A.; Navedo, M.F. Regulation of Voltage-Gated Potassium Channels in Vascular Smooth Muscle during Hypertension and Metabolic Disorders. *Microcirculation* **2018**, *25*, 42–49. [[CrossRef](#)]
32. Tanaka, Y.; Igarashi, T.; Kaneko, H.; Yamaki, F.; Mochizuki, Y.; Aida, M.; Taniguchi, H.; Tanaka, H.; Shigenobu, K. NO-Mediated MaxiK(Ca) Channel Activation Produces Relaxation of Guinea Pig Aorta Independently of Voltage-Dependent L-Type Ca<sup>2+</sup> Channels. *Gen. Pharmacol.* **2000**, *34*, 159–165. [[CrossRef](#)]
33. Wellman, G.C. Ion Channels and Calcium Signaling in Cerebral Arteries Following Subarachnoid Hemorrhage. *Neurol. Res.* **2006**, *28*, 690–702. [[CrossRef](#)] [[PubMed](#)]
34. Meisher, K.D.; Cipkus, L.A.; Taylor, C.J. Mechanism of Action of Minoxidil Sulfate-Induced Vasodilation: A Role for Increased K<sup>+</sup> Permeability. *J. Pharmacol. Exp. Ther.* **1988**, *245*, 751–760.
35. Nielsen-Kudsk, J.E.; Boesgaard, S.; Aldershvile, J. K<sup>+</sup> Channel Opening: A New Drug Principle in Cardiovascular Medicine. *Heart* **1996**, *76*, 109–116. [[CrossRef](#)]
36. Zhang, Y.; Gao, Y.-J.; Zuo, J.; Lee, R.M.K.W.; Janssen, L.J. Alteration of Arterial Smooth Muscle Potassium Channel Composition and BKCa Current Modulation in Hypertension. *Eur. J. Pharmacol.* **2005**, *514*, 111–119. [[CrossRef](#)]
37. Tajada, S.; Ciudad, P.; Moreno-Domínguez, A.; Pérez-García, M.T.; López-López, J.R. High Blood Pressure Associates with the Remodelling of Inward Rectifier K<sup>+</sup> Channels in Mice Mesenteric Vascular Smooth Muscle Cells. *J. Physiol.* **2012**, *590*, 6075–6091. [[CrossRef](#)]
38. Hill, M.A.; Yang, Y.; Ella, S.R.; Davis, M.J.; Braun, A.P. Large Conductance, Ca<sup>2+</sup>-Activated K<sup>+</sup> Channels (BKCa) and Arteriolar Myogenic Signaling. *FEBS Lett.* **2010**, *584*, 2033–2042. [[CrossRef](#)]
39. El-Hajj, H.; Chandrasekhar, B.; Kadavil, E.A.; Oriowo, M.A. Interaction of BKCa Channel Modulators with Adrenergic Agonists in the Rat Aorta Is Influenced by Receptor Reserve. *Vascul. Pharmacol.* **2004**, *41*, 119–124. [[CrossRef](#)]
40. Goto, K.; Rummery, N.M.; Grayson, T.H.; Hill, C.E. Attenuation of Conducted Vasodilatation in Rat Mesenteric Arteries during Hypertension: Role of Inwardly Rectifying Potassium Channels. *J. Physiol.* **2004**, *561*, 215–231. [[CrossRef](#)]
41. Park, C.G. Hypertension and Vascular Aging. *Korean Circ. J.* **2006**, *36*, 477. [[CrossRef](#)]
42. Tajada, S.; Ciudad, P.; Colinas, O.; Santana, L.F.; López-López, J.R.; Pérez-García, M.T. Down-Regulation of CaV1.2 Channels during Hypertension: How Fewer CaV1.2 Channels Allow More Ca<sup>2+</sup> into Hypertensive Arterial Smooth Muscle. *J. Physiol.* **2013**, *591*, 6175–6191. [[CrossRef](#)]
43. Cox, R.H.; Rusch, N.J. New Expression Profiles of Voltage-Gated Ion Channels in Arteries Exposed to High Blood Pressure. *Microcirculation* **2002**, *9*, 243–257. [[CrossRef](#)]
44. Yang, Y.; Li, P.-Y.; Cheng, J.; Mao, L.; Wen, J.; Tan, X.-Q.; Liu, Z.-F.; Zeng, X.-R. Function of BKCa Channels Is Reduced in Human Vascular Smooth Muscle Cells from Han Chinese Patients with Hypertension. *Hypertension* **2013**, *61*, 519–525. [[CrossRef](#)] [[PubMed](#)]
45. Zhang, Y.; Liao, J.; Zhang, L.; Li, S.; Wu, Y.; Shi, L. BKCa Channel Activity and Vascular Contractility Alterations with Hypertension and Aging via B1 Subunit Promoter Methylation in Mesenteric Arteries. *Hypertens. Res.* **2018**, *41*, 96–103. [[CrossRef](#)]
46. Bühler, F.R.; Bolli, P.; Erne, P.; Kiowski, W.; Müller, F.B.; Hulthén, U.L.; Ji, B.H. Position of Calcium Antagonists in Antihypertensive Therapy. *J. Cardiovasc. Pharmacol.* **1985**, *7*, S21-7. [[CrossRef](#)] [[PubMed](#)]
47. Armstrong, C. Joint National Committee JNC8 Guidelines for the Management of Hypertension in Adults. *Am. Fam. Physician* **2014**, *90*, 503–504. [[PubMed](#)]
48. Iversen, B.M.; Arendshorst, W.J. Exaggerated Ca<sup>2+</sup> Signaling in Preglomerular Arteriolar Smooth Muscle Cells of Genetically Hypertensive Rats. *Am. J. Physiol.* **1999**, *276*, F260–F270. [[CrossRef](#)] [[PubMed](#)]
49. Tabet, F.; Savoia, C.; Schiffrin, E.L.; Touyz, R.M. Differential Calcium Regulation by Hydrogen Peroxide and Superoxide in Vascular Smooth Muscle Cells from Spontaneously Hypertensive Rats. *J. Cardiovasc. Pharmacol.* **2004**, *44*, 200–208. [[CrossRef](#)] [[PubMed](#)]
50. Pratt, P.F.; Bonnet, S.; Ludwig, L.M.; Bonnet, P.; Rusch, N.J. Upregulation of L-Type Ca<sup>2+</sup> Channels in Mesenteric and Skeletal Arteries of SHR. *Hypertension* **2002**, *40*, 214–219. [[CrossRef](#)]
51. Potje, S.R.; Isbatan, A.; Tostes, R.C.; Bendhack, L.M.; Dull, R.O.; Carvalho-de-Souza, J.L.; Chignalia, A.Z. Glypican 1 and Syndecan 1 Differently Regulate Noradrenergic Hypertension Development: Focus on IP3R and Calcium. *Pharmacol. Res.* **2021**, *172*, 105813. [[CrossRef](#)] [[PubMed](#)]
52. Xie, M.-J.; Zhang, L.-F.; Ma, J.; Cheng, H.-W. Functional Alterations in Cerebrovascular K<sup>+</sup> and Ca<sup>2+</sup> Channels Are Comparable between Simulated Microgravity Rat and SHR. *Am. J. Physiol. Heart Circ. Physiol.* **2005**, *289*, H1265–H1276. [[CrossRef](#)]
53. Ohya, Y.; Tsuchihashi, T.; Kagiya, S.; Abe, I.; Fujishima, M. Single L-Type Calcium Channels in Smooth Muscle Cells from Resistance Arteries of Spontaneously Hypertensive Rats. *Hypertension* **1998**, *31*, 1125–1129. [[CrossRef](#)]
54. Liao, J.; Zhang, Y.; Ye, F.; Zhang, L.; Chen, Y.; Zeng, F.; Shi, L. Epigenetic Regulation of L-Type Voltage-Gated Ca<sup>2+</sup> Channels in Mesenteric Arteries of Aging Hypertensive Rats. *Hypertens. Res.* **2017**, *40*, 441–449. [[CrossRef](#)] [[PubMed](#)]
55. Lozinskaya, I.M.; Cox, R.H. Effects of Age on Ca<sup>2+</sup> Currents in Small Mesenteric Artery Myocytes from Wistar-Kyoto and Spontaneously Hypertensive Rats. *Hypertension* **1997**, *29*, 1329–1336. [[CrossRef](#)]
56. Leung, F.P.; Yung, L.M.; Yao, X.; Laher, I.; Huang, Y. Store-Operated Calcium Entry in Vascular Smooth Muscle. *Br. J. Pharmacol.* **2008**, *153*, 846–857. [[CrossRef](#)]
57. Fellner, S.K.; Arendshorst, W.J. Store-Operated Ca<sup>2+</sup> Entry Is Exaggerated in Fresh Preglomerular Vascular Smooth Muscle Cells of SHR. *Kidney Int.* **2002**, *61*, 2132–2141. [[CrossRef](#)]



58. Leijten, P.A.; van Breemen, C. The Effects of Caffeine on the Noradrenaline-Sensitive Calcium Store in Rabbit Aorta. *J. Physiol.* **1984**, *357*, 327–339. [[CrossRef](#)] [[PubMed](#)]
59. Rossi, A.M.; Taylor, C.W. IP3 Receptors—Lessons from Analyses Ex Cellula. *J. Cell Sci.* **2018**, *132*, 222463. [[CrossRef](#)]
60. Ziolkowski, N.; Grover, A.K. Functional Linkage as a Direction for Studies in Oxidative Stress: Alpha-Adrenergic Receptors. *Can. J. Physiol. Pharmacol.* **2010**, *88*, 220–232. [[CrossRef](#)]
61. Konukoglu, D.; Uzun, H. Endothelial Dysfunction and Hypertension. *Adv. Exp. Med. Biol.* **2017**, *956*, 511–540. [[CrossRef](#)]
62. Waldron, G.J.; Ding, H.; Lovren, F.; Kubes, P.; Triggle, C.R. Acetylcholine-Induced Relaxation of Peripheral Arteries Isolated from Mice Lacking Endothelial Nitric Oxide Synthase. *Br. J. Pharmacol.* **1999**, *128*, 653–658. [[CrossRef](#)]
63. Comerma-Steffensen, S.; Risso, A.; Ascanio-Evanoff, E.; Zerpa, H. Endothelium-Dependent Relaxation Mechanisms Involve Nitric Oxide and Prostanoids in the Isolated Bovine Digital Vein. *J. Vet. Pharmacol. Ther.* **2019**, *42*, 361–367. [[CrossRef](#)]
64. Long, H.; Jiang, J.; Xia, J.; Jiang, R. Icariin Improves SHR Erectile Function via Inhibiting ENOS Uncoupling. *Andrologia* **2018**, *50*, e13084. [[CrossRef](#)]
65. Li, Q.; Youn, J.-Y.; Cai, H. Mechanisms and Consequences of Endothelial Nitric Oxide Synthase Dysfunction in Hypertension. *J. Hypertens.* **2015**, *33*, 1128–1136. [[CrossRef](#)]
66. Khan, S.I.; Andrews, K.L.; Jefferis, A.; Jennings, G.L.; Sampson, A.K.; Chin-Dusting, J.P.F. Vascular Dysfunction in the Stroke-Prone Spontaneously Hypertensive Rat Is Dependent on Constrictor Prostanoid Activity and Y Chromosome Lineage. *Clin. Sci.* **2018**, *132*, 131–143. [[CrossRef](#)]
67. Potje, S.R.; Grando, M.D.; Chignalia, A.Z.; Antoniali, C.; Bendhack, L.M. Reduced Caveolae Density in Arteries of SHR Contributes to Endothelial Dysfunction and ROS Production. *Sci. Rep.* **2019**, *9*, 6696. [[CrossRef](#)]
68. Vita, J.A. Endothelial Function. *Circulation* **2011**, *124*, e108–e114. [[CrossRef](#)]
69. Gielis, J.F.; Quirynen, L.; Briedé, J.J.; Roelant, E.; Cos, P.; Van Schil, P.E.Y. Pathogenetic Role of Endothelial Nitric Oxide Synthase Uncoupling during Lung Ischaemia-Reperfusion Injury. *Eur. J. Cardiothorac. Surg.* **2017**, *52*, 256–263. [[CrossRef](#)]
70. Touyz, R.M.; Schiffrin, E.L. Role of Calcium Influx and Intracellular Calcium Stores in Angiotensin II-Mediated Calcium Hyper-Responsiveness in Smooth Muscle from Spontaneously Hypertensive Rats. *J. Hypertens.* **1997**, *15*, 1431–1439. [[CrossRef](#)]
71. Romero, J.C.; Reckelhoff, J.F. State-of-the-Art Lecture. Role of Angiotensin and Oxidative Stress in Essential Hypertension. *Hypertension* **1999**, *34*, 943–949. [[CrossRef](#)]
72. Hernanz, R.; Martínez-Revelles, S.; Palacios, R.; Martín, A.; Cachofeiro, V.; Aguado, A.; García-Redondo, L.; Barrús, M.T.; de Batista, P.R.; Briones, A.M.; et al. Toll-like Receptor 4 Contributes to Vascular Remodelling and Endothelial Dysfunction in Angiotensin II-Induced Hypertension. *Br. J. Pharmacol.* **2015**, *172*, 3159–3176. [[CrossRef](#)]
73. Vanhoutte, P.M.; Feletou, M.; Taddei, S. Endothelium-Dependent Contractions in Hypertension. *Br. J. Pharmacol.* **2005**, *144*, 449–458. [[CrossRef](#)]
74. Qu, C.; Leung, S.W.S.; Vanhoutte, P.M.; Man, R.Y.K. Wy14643 Improves Vascular Function in the Aorta of the Spontaneously Hypertensive Rat Mainly by Activating Peroxisome Proliferator-Activated Receptors Alpha. *Eur. J. Pharmacol.* **2012**, *696*, 101–110. [[CrossRef](#)]
75. Silva, D.F.; Araújo, I.G.A.; Albuquerque, J.G.F.; Porto, D.L.; Dias, K.L.G.; Cavalcante, K.V.M.; Veras, R.C.; Nunes, X.P.; Barbosa-Filho, J.M.; Araújo, D.A.M.; et al. Rotundifolone-Induced Relaxation Is Mediated by BK(Ca) Channel Activation and Ca(v) Channel Inactivation. *Basic Clin. Pharmacol. Toxicol.* **2011**, *109*, 465–475. [[CrossRef](#)]
76. Langton, P.D.; Nelson, M.T.; Huang, Y.; Standen, N.B. Block of Calcium-Activated Potassium Channels in Mammalian Arterial Myocytes by Tetraethylammonium Ions. *Am. J. Physiol.* **1991**, *260*, H927–H934. [[CrossRef](#)]
77. Wang, S.-P.; Zang, W.-J.; Kong, S.-S.; Yu, X.-J.; Sun, L.; Zhao, X.-F.; Wang, S.-X.; Zheng, X.-H. Vasorelaxant Effect of Isopropyl 3-(3, 4-Dihydroxyphenyl)-2-Hydroxypropanoate, a Novel Metabolite from *Salvia Miltiorrhiza*, on Isolated Rat Mesenteric Artery. *Eur. J. Pharmacol.* **2008**, *579*, 283–288. [[CrossRef](#)]
78. Kawabata, A.; Kubo, S.; Nakaya, Y.; Ishiki, T.; Kuroda, R.; Sekiguchi, F.; Kawao, N.; Nishikawa, H. Distinct Roles for Protease-Activated Receptors 1 and 2 in Vasomotor Modulation in Rat Superior Mesenteric Artery. *Cardiovasc. Res.* **2004**, *61*, 683–692. [[CrossRef](#)]
79. Insuk, S.O.; Chae, M.R.; Choi, J.W.; Yang, D.K.; Sim, J.H.; Lee, S.W. Molecular Basis and Characteristics of KATP Channel in Human Corporal Smooth Muscle Cells. *Int. J. Impot. Res.* **2003**, *15*, 258–266. [[CrossRef](#)]
80. Sakata, K.; Karaki, H. Effects of a Novel Smooth Muscle Relaxant, KT-362, on Contraction and Cytosolic Ca<sup>2+</sup> Level in the Rat Aorta. *Br. J. Pharmacol.* **1991**, *102*, 174–178. [[CrossRef](#)]