

Knowledge, Self-Efficacy, and Correlates in Palliative and End-of-Life Care: Quantitative Insights from Final-Year Nursing and Medical Students in a Mixed-Methods Study

Germanus Natuhwera , Eve Namisango  and Peter Ellis

Abstract

Introduction: About a decade after the introduction of palliative care teaching for undergraduate nurses and medical students in Uganda, no research has examined students' knowledge and self-efficacy to provide palliative and end-of-life care and their correlates.

Aims: To: (1) estimate final-year undergraduate nursing and medical students' knowledge of and self-efficacy to provide palliative and end-of-life care, (2) identify correlates of knowledge and self-efficacy to provide palliative and end-of-life care.

Design: A multicentre cross-sectional quantitative study.

Setting/participants: Final-year undergraduate medical and nursing students in eight medical and nursing schools in Uganda. Instruments included biodata sheet, the Palliative Care Quiz for Nursing questionnaire and the Palliative Care Self-Efficacy scale. Statistical analyses were performed using STATA version 14.0.

Results: The mean age of the participants ($n=466$) was 24.45 ± 3.31 years. Participants' knowledge of palliative care scores was low in all domains 'Philosophy and principles of palliative care' 1.46 ± 0.93 (range: 0–4), 'Psychosocial and spiritual care' 0.61 ± 0.73 (range: 0–3) and 'Management of pain and other symptoms' 6.32 ± 1.75 (range: 0–13). Predictors of knowledge were Gender ($p=0.0242$), course of study ($p=0.0001$) and religion ($p=0.0338$). Participants had very low self-efficacy scores in the three domains of the Palliative Care Self-Efficacy scale.

Conclusion: Participants generally demonstrated limited knowledge and insufficient self-efficacy in providing palliative and end-of-life care. There is a need to integrate and strengthen practical, pedagogical and experiential teaching, review the palliative care curriculum. Future evaluative, longitudinal and interventional as well as qualitative studies are needed to gain deeper insights into this topic.

Keywords: end-of-life care, knowledge, medical, nursing, palliative care, self-efficacy, student, undergraduate

Received: 29 October 2024; revised manuscript accepted: 12 January 2025.

Palliative Care & Social Practice

2025, Vol. 19: 1–22

DOI: 10.1177/

26323524251316901

© The Author(s), 2025.
Article reuse guidelines:
sagepub.com/journals-
permissions

Correspondence to:
Germanus Natuhwera
Clinical and Administration
Departments, Little
Hospice Hoima, Hospice
Africa Uganda, P.O. Box
7757, Kampala, Uganda
Education Department,
Institute of Hospice and
Palliative Care in Africa,
P.O. Box 7757, Kampala,
Uganda
ngermans16@gmail.com
gnatuhwera@hospice-africa.org

Eve Namisango
Research and
Programmes Department,
African Palliative Care
Association, Kampala,
Uganda

Cicely Saunders Institute,
King's College London,
London, UK

Education Department,
Institute of Hospice and
Palliative Care in Africa,
Kampala, Uganda

Peter Ellis
Independent Nursing
Writer and Educator, and
Honorary Senior Research
Fellow, Canterbury Christ
Church University, United
Kingdom

Background

Over recent decades, the demand for palliative care (PC) has risen significantly in line with a rise in the global disease burden (both communicable and non-communicable) and an ageing population.¹⁻³ High-quality palliative and end-of-life care are now well recognized as an indispensable human right to dignity.³ The unmet need for palliative care services is still, however, huge, particularly in low-and-middle-income countries where at least 76% of adults require palliative care.⁴⁻⁶

Yet, globally, only 14%, of about 53 million adults and 4 million children, who need palliative care receive it.⁵ Furthermore, at least 25.7 million adults and 2.5 million children, including at least 80% of adults and 98% of children residing in low-and-middle-income countries, experience serious health-related suffering at the end of life.^{5,7} The need for palliative care at the end of life is projected to double by 2060.⁴

In Africa, disparities in access to palliative care are vast, with an estimated 5% of those in need able to access services.⁸ Cancer cases are rising exponentially and are projected to rise by 400% by the year 2050.⁹ It is estimated that in the World Health Organization Africa region, 2000 adults (≥ 18 years) per 100,000 of the population and 369 children (0–17 years) per 100,000 of the population require palliative care.⁵

In Uganda an estimated minimum of 400,000 adults and 86,000 children need palliative care every year.¹⁰ Furthermore, an estimated 84% and 80% of adult and children cancer patients respectively require palliative care (pain relief) as an integral component of their cancer care.⁷ But only 11% of Ugandans in need of palliative care can access services.^{5,11}

This phenomenon highlights an urgent need for a correspondingly significant increase in the availability of palliative care services.¹² Integrating education and training in palliative care and care for the dying (end-of-life care) within undergraduate health curricula is critical in preparing future nurses and medical doctors to provide high-quality palliative and end-of-life care.^{5,6,13,14}

Education has the potential to improve students' knowledge and promote positive attitudes towards palliative and end-of-life care.¹⁵⁻²⁰ Exposure to palliative and end-of-life training has been shown to improve nursing and medical students'

knowledge, skills, self-confidence, self-awareness and competence to provide holistic and person-centred care.²¹⁻²⁴ Numerous evaluation and interventional studies from high-income countries have corroborated that palliative care training orientates and improves students' knowledge and attitudes towards palliative care.²⁵⁻²⁹

A wealth of research evidence indicates nursing and medical students continue to graduate from training institutions possessing gross negative perceptions and being unprepared to provide culturally sensitive and person-centred palliative and end-of-life care.³⁰⁻³³ This unpreparedness usually manifests as inappropriate attitudes and insufficient or lacking knowledge, skills and confidence in essential domains such as principles of palliative care, pain and symptom management, provision of psychosocial and spiritual care, and communication with dying patients and their families.³⁴⁻⁴⁰ Very limited research has found good knowledge of palliative among undergraduates.⁴¹ Inadequate curricula, education and training have been cited as a major predictor for this common observation.^{42,43}

Around a decade since the introduction of palliative care teaching into the nursing, allied health and medical training curricula in Uganda, it remains unclear as to what extent, if at all, it is positively influencing nursing and medical students' knowledge and self-efficacy to provide palliative and end-of-life care or what characteristics of students are predictive of this. In the context of this study, self-efficacy is defined as students' ability to develop the appropriate attitude, skills and self-confidence necessary to apply learned knowledge to provide palliative and end-of-life care.

Furthermore, there is a dearth of documented research on the topic from Africa. In particular, no documented research on the topic was identified at the time of this study. A global systematic review conducted prior to this study revealed that nursing and medical students continue to graduate from training schools with low and inadequate knowledge of palliative and end-of-life care, ranging from 5.3% to 61%. Students' knowledge levels of palliative care are predominantly low with rates ranging from 5.3% to 61% across different regions of the World. Anecdotal observations in practice point to deficient levels of knowledge, negative perceptions and a lack of competence among students and qualified healthcare professionals.

Hence, this study sought to examine this under-researched topic. The evidence generated could reveal training needs and gaps in the scope and delivery of palliative care training, and suggest appropriate remedial action, including future research to optimize the impact of the course on students' training needs.

Aims of the study

The study aimed to

1. Estimate final-year undergraduate nursing and medical students' knowledge of palliative and end-of-life care.
2. Examine participants' self-efficacy to provide palliative and end-of-life care.
3. Examine the correlation between knowledge and self-efficacy, knowledge and socio-demographic and professional variables (independent variables).

Methods

Study design, population and context

This cross-sectional quantitative study was part of a larger multicentre parallel mixed-methods study of final-year undergraduate medical and nursing students in eight public and private tertiary education institutions in Uganda. The qualitative element of the study (preprint) is being considered for publication elsewhere.⁴⁴ The purpose of recruiting participants from different settings was to generate generalisability.

In Uganda, undergraduate medical students undergo training for 6 years including 1 year of clinical internship before they qualify to be awarded a Bachelor of Medicine and Bachelor of Surgery. Nursing students pursuing a Bachelor of Science in Nursing complete four years of training and an additional 1 year of internship in a clinical setting under supervision. Diploma-level participants were nursing and midwifery students in their final semester of a 2-year extension programme after initially attaining a 2.5-year certificate and practiced for at least 2 years. Regarding the palliative care course unit, based on the design of the curriculum, the course is allocated a whole semester and is taught in-; fourth year for undergraduate medical students, third year for undergraduate degree and diploma direct nursing students, second year for undergraduate diploma nursing students on extension and certificate level nursing students.

Sampling and sample size estimation

The sample size was estimated using Cochran's formula for unknown population⁴⁵ as $n = 384$. To increase the power of the study, an additional 38 participants, 10% was included to cater for 10% non-response rate, so total $n = 422$.


Data collection methods and instruments

Data collection took place between 21 November 2022 and 1 March 2023. Participants were accessed through their respective department within which the students are registered after securing institutional approval. A biographic data sheet including age, gender, discipline of study (nursing or medical), history of training in palliative care and or working with palliative care patients was used to collect participants' biographic data.

Students' knowledge of palliative care was assessed using the Palliative Care Quiz for Nursing questionnaire. According to Ross *et al.*,⁴⁶ the scale has a high internal content validity and acceptable reliability (test-retest = 0.56 and Kuder-Richardson 20 = 0.78). The questionnaire is self-administered and consists of 20 multiple-choice items (true/false/does not know/did not answer) to assess three main domains of palliative care: philosophy and principles of palliative care (4 items), control of pain and other symptoms (13 items) and psychosocial aspects of palliative care (3 items). These 20 items refer to knowledge applicable in clinical settings.

The questionnaire has been proven to be a valid and reliable tool (with adequate internal consistency in different languages) for assessing knowledge and identifying perceptions of palliative care among health professionals in both developed settings, for example, China⁴⁷ and developing countries, for example, Ethiopia.⁴⁸ Originally developed to assess knowledge in nursing, the questionnaire has further been proven to be an effective tool to assess knowledge of palliative care in other health cadres including active medical doctors and students.⁴⁹

Students' self-perceived efficacy was assessed using the Palliative Care Self-Efficacy scale, a validated 23-item self-assessment structured questionnaire that seeks to examine how clinicians/nurses manage various situations in a palliative care environment, for example, confidence in



MB ChB-Bachelor of Medicine & Bachelor of Surgery, BSN-Bachelor of Science in Nursing, Dip-N&M-Diploma in Nursing & Midwifery, ECN-Enrolled Certificate in Nursing

Figure 1. Participants' recruitment scheme.

BSN, Bachelor of Science in Nursing; Dip-N&M, Diploma in Nursing and Midwifery, ECN, Enrolled Certificate in Nursing; MB ChB, Bachelor of Medicine and Bachelor of Surgery.

communicating with the dying, managing symptoms, etc.⁵⁰

The scale has three distinct subscales covering palliative care domains: Communication Skills; Multidisciplinary Teams; and Pain and Symptom Management. Possible responses are rated on a Likert scale ranging from 1 to 4 with a score of 1 denoting 'need further basic instruction', a 2 denotes 'confident to perform with close supervision/coaching', a 3 suggests 'confident to perform with minimal consultation' and a 4 denotes 'confident to perform independently'. The entire scale has a strong validity, as demonstrated by a Cronbach's alpha coefficient of 0.92, and 0.87 and 0.91 for the subscales 'perceived capability to answer end-of-life care concerns of patients' and 'perceived capability to respond to patients' end-of-life symptoms', respectively.⁵⁰ Data were collected face-to-face by GN and research assistants who were trained and oriented by GN prior to commencing data collection.

Sampling and recruitment

Convenience sampling was employed to recruit participants who were readily accessible and willing to participate in the study. A total of 466 participants from eight teaching institutions volunteered to participate (Figure 1). Convenience sampling was preferred as the researcher could not get the sampling frame to fulfill the criteria for random sampling given the nature of the settings of the participants.

Eligible participants were: (i) nursing or medical students in the final year of their undergraduate training; (ii) Ugandans (to increase sample homogeneity and minimize possible skewed data variations due to cultural or contextual differences); (iii) able to grant their individual voluntary consent verbally and by signature; (iv) able to talk, read and write in English.

A prospective participant was excluded if they: (i) were non-finalist; (ii) non-Ugandan/international

students to minimize confounders related to home country factors such as prior experiential training or exposure to palliative care; (iii) declined to give their consent; (iv) were unable to hear, read and or write in English; (v) chose to opt out of the study.

Data handling and analysis

Completed questionnaires were screened manually for completeness and the data entered into a coded Microsoft Excel spreadsheet. The generated datasets were then exported into STATA version 14.0, StataCorp LP, College Station for analysis.

Participants' demographic characteristics and knowledge scores on the Palliative Care Quiz for Nursing scale were summarized using descriptive statistics, mainly frequencies, percentages, minimum and maximum (range), means and standard deviations. 'True' responses were scored as one, and 'False' or 'I don't know' responses scored as zero. For each questionnaire, the total score was computed by summing up the number of correctly answered items, with range of 0 (minimum score) to 20 (maximum score). The mean scores of each of the Palliative Care Quiz for Nursing items were computed by summing up the scores in the subscale and dividing the total score by the number of items. Hence, philosophy and principles had a theoretical score range (minimum–maximum) of 0–4, pain and symptom had a score range of 0–13, and psychosocial and spiritual care had a score range of 0–3.

Analysis of variance (ANOVA) and Bartlett's equal-variances tests were performed to examine relationships between students' palliative care knowledge scores on the Palliative Care Quiz for Nursing subscale and independent variables within and between groups. Weighted regression analysis was performed to further examine relationships between predictor variables (age, gender and course of study) and students' palliative care knowledge scores. Results were interpreted as odds ratios.

Descriptive statistics (proportions and percentages) were used to summarize students' responses on the Palliative Care Self-Efficacy scale. ANOVA and Bartlett's equal-variances tests were performed to examine the relationship between students' self-efficacy scores and predictor variables. The scores for each completed questionnaire

were summed up to get the total score. The lowest and highest possible scores for each questionnaire were 23 and 92, respectively.

A value of $p < 0.05$ was considered statistically significant for all the statistical tests. All study datasets are stored on GN's computer secured with a password.

Results

Four hundred sixty-six students completed the questionnaire compared with an estimated sample size of 422, translating to a response rate of 110%.

As seen in Table 1, of the 466 students that participated in the study, 253 (54.3%) were female and the majority were in the age range of 20–29 years 428 (91.9%). Their mean age was 24.45 ± 3.31 years and the age range was 20–44 years.

As can be seen in Table 2, a higher frequency of correct responses was observed in the '*management of pain and other symptoms*' subscale especially questions 2, 4, 8 and 18. However, the rest of the 10-scale items had below-average (50%) number of correct responses.

The least correctly answered scale items questions 3, 7 and 13 refer to pain management and the use of opioids.

In the '*philosophy and principles of palliative care*' subscale, a low number of correct and a higher number of incorrect/unsure responses was observed. Notably, the number of correct responses for all four scale question items was below average (50%).

The '*psychosocial and spiritual care*' subscale recorded the least number of correct responses. Of the three question items in the subscale, the highest number of correct responses were observed in Q11 (31.6%).

Participants' knowledge of palliative care was generally poor. Good knowledge scores were observed in only four subscales (Q2, Q4, Q8, Q18) in the '*management of pain and other symptoms*' category, with overall mean knowledge scores (6.32 ± 1.75). The rest of the subscales had knowledge scores below 50% (see Table 3).

Table 1. Socio-demographics of participants ($n = 466$).

Variable	Frequency (n)	Percentage (%)
Age years		
20–29	428	91.9
30–39	38	8.1
Mean \pm SD	24.45 \pm 3.31	
Gender		
Female	253	54.3
Male	213	45.7
Religion		
Catholic	160	34.3
Protestant	164	35.2
Pentecostal	57	12.2
Muslim	42	9.0
SDA	34	7.3
Other	9	1.9
Course of study		
Degree in medicine	225	48.2
Degree in nursing	49	10.5
Diploma in nursing	58	12.5
Certificate in nursing	134	28.8
Level of interaction with terminally ill patients		
Very frequently	57	12.2
Frequently	160	34.3
Occasionally	160	34.3
Rarely	79	17.0
Never	10	2.2
History or experience of working with palliative or terminally ill patients		
Yes	339	72.8
No	127	27.2
Where did you first learn about palliative care?		
Medical/nursing school	406	87.1
Workplace/hospital	28	6.0

(Continued)

Table 1. (Continued)

Variable	Frequency (n)	Percentage (%)
Family or friends	10	2.1
Media	11	2.4
Never	11	2.4
Have you had any training in palliative care in your institution?		
Yes	382	82.0
No/Never	84	18.0
When you hear of the term hospice, what do you first think of?		
Place for cancer patients	108	16.9
Place where all hope is lost	21	3.3
Place to go to when all hope is lost	63	9.9
Place for pain medication titration and control	99	15.5
Place for palliation and optimizing quality of life	326	51.1
Place for elderly patients	21	3.3
SDA, Seventh Day Adventis.		

In Table 4, in the ‘*philosophy and principles of palliative care*’ subscale, ANOVA show a statistically significant relationship for gender ($F=5.12$, $p=0.0242$) with males showing higher knowledge scores, and course of study ($F=7.68$, $p=0.0001$), with medical and diploma in nursing students having better knowledge than certificate and degree nursing colleagues. Bartlett’s test shows the mean knowledge score variances within and between groups are equal for age ($F=2.01$, $p=0.072$), gender ($F=5.12$, $p=0.217$), religion ($F=2.10$, $p=0.675$) and course of study ($p=0.301$).

In the ‘*management of pain and other symptoms*’ subscale, religion was statistically significant ($F=5$, $p=0.0338$). Muslims and Pentecostals demonstrated higher knowledge scores than Protestants and Catholics. Bartlett’s equal-variances test further shows that the mean variances within and between groups are significantly unequal ($p=0.040$).

In the ‘*psychosocial and spiritual care*’ subscale, the course of study shows a statistically significant positive association with knowledge of palliative care ($F=3.06$, $p=0.0279$). Notably, medical

students had higher scores and diploma in nursing students had the lowest scores.

In Table 5, in model one, *philosophy and principles of palliative care*, the odds of having knowledge of PC was 30% lower if the participant offered a certificate in the nursing course ($p=0.001$) OR 0.7008269 (95% CI: 0.5805999–0.8459498). Older students (30–39 years) were 32% ($p=0.050$) OR 1.3226 (95% CI: 0.9995–1.7499) more likely to have better knowledge of PC compared with younger students (20–29 years). Being male was 14% ($p=0.101$) OR 1.1386 (95% CI: 0.9759–1.3296) more likely to have better knowledge of PC compared with being female.

In model two, *management of pain and symptoms*, being older, male, and taking on courses other than degree in medicine was associated with reduced odds of having knowledge of palliative care.

In model three, ‘*psychosocial and spiritual care*’, the odds of having knowledge of palliative care was lower by 26% ($p=0.037$; 95% CI: 0.5531–0.9819) if the student was on a degree in the nursing course compared with if they were on a

Table 2. Frequency and percentage of responses of participants on the Palliative Care Quiz for Nursing scale ($n=466$).

Scale item	Subscale	Correct response (T)	Incorrect response (F/I)
		<i>n</i> (%)	<i>n</i> (%)
Philosophy and principles of palliative care			
Q1.	Palliative care is appropriate only in situations where there is evidence of a downward trajectory or deterioration. (F)	178 (38.2)	288 (61.8)
Q9.	The provision of palliative care requires emotional detachment. (F)	232 (49.8)	234 (50.2)
Q12.	The philosophy of palliative care is compatible with that of aggressive treatment. (T)	151 (32.4)	315 (67.6)
Q17.	The accumulation of losses makes burnout inevitable for those who work in palliative care. (F)	116 (24.9)	350 (75.1)
Management of pain and symptoms			
Q2.	Morphine is the standard used to compare the analgesic effect of other opioids. (T)	340 (73.0)	126 (27.0)
Q3.	The extent of the disease determines the method of pain treatment. (F)	55 (11.8)	411 (88.2)
Q4.	Adjuvant therapies are important in managing pain. (T)	411 (88.2)	55 (11.8)
Q6.	During the last days of life, drowsiness associated with electrolyte imbalance may decrease the need for sedation. (T)	225 (48.3)	241 (51.7)
Q7.	Drug addiction is a major problem when morphine is used on a long-term basis for the management of pain. (T)	90 (19.3)	376 (80.7)
Q8.	Individuals who are taking opioids should also follow a bowel regimen (laxative treatment). (T)	369 (79.2)	97 (20.8)
Q10.	During the terminal stages of an illness, drugs that can cause respiratory depression are appropriate for the treatment of severe dyspnoea. (T)	136 (29.2)	330 (70.8)
Q13.	The use of placebos is appropriate in the treatment of some types of pain. (F)	113 (24.2)	353 (75.8)
Q14.	High-dose codeine causes more nausea and vomiting than morphine. (T)	225 (48.3)	241 (51.7)
Q15.	Suffering and physical pain are identical. (F)	195 (41.9)	271 (58.1)
Q16.	Demerol (pethidine) is not an effective analgesic for the control of chronic pain. (T)	208 (44.6)	258(55.4)
Q18.	Manifestations of chronic pain are different from those of acute pain. (T)	397 (85.2)	69 (14.8)
Q20.	Pain thresholds are lowered by fatigue or anxiety. (T)	179 (38.4)	287 (61.6)
Psychosocial and spiritual care			
Q5.	It is crucial for family members to remain at the bedside until death occurs. (F)	62 (13.3)	404 (86.7)
Q11.	Men generally reconcile their grief more quickly than do women. (F)	147 (31.6)	319 (68.5)
Q19.	The loss of a distant relationship is easier to resolve than the loss of one that is close or intimate. (F)	74 (15.9)	392 (84.1)
Scoring: True (T) is a correct response (T)= 1 score, False (F) and 'I don't know' (I) are incorrect responses and each=0 score. PCQN, Palliative Care Quiz for Nursing.			

Table 3. Overall minimum and maximum, means, and mean knowledge scores of participants on the three conceptual categories of the PCQN scale.

PCQN conceptual categories	PCQN subscale	Mean (SD) knowledge scores	Minimum–Maximum total score (range)
Philosophy and principles of PC	Q1, Q9, Q12, Q17	1.46 ± 0.93	0–4
Management of pain and other symptoms	Q2–4, Q6–8, Q10, Q13–16, Q18, Q20	6.32 ± 1.75	0–13
Psychosocial and spiritual care	Q5, Q11, Q19	0.61 ± 0.73	0–3
Total score	Q1–20		0–20
PCQN, Palliative Care Quiz for Nursing.			

Table 4. Comparison of palliative care knowledge scores of the students according to the Palliative Care Quiz for Nursing with their socio-demographic characteristics ($n=466$).

PCQN subscale		ANOVA of mean scores between and within the group				Bartlett's equal-variances test between and within the groups	
Independent variable	<i>n</i>	% knowledge scores (mean ± SD)	Degrees of freedom	<i>F</i> -statistic	<i>p</i> Value	Chi-square (χ^2)	<i>p</i> Value
Philosophy and principles of palliative care: Q1, Q9, Q12, Q17							
Age			1	2.01	0.1569	3.2481	0.072
20–29	428	35.9 ± 22.8					
30–39	38	41.4 ± 28.0					
Gender			1	5.12	0.0242	1.5256	0.217
Female	253	34.1 ± 22.3					
Male	213	39.0 ± 24.2					
Religion			5	2.10	0.0646	3.1638	0.675
Catholic	160	33.8 ± 23.7					
Protestant	164	36.3 ± 22.2					
Pentecostal	57	40.4 ± 22.5					
Muslim	42	41.7 ± 26.3					
SDA	34	31.6 ± 23.2					
Other	9	50 ± 17.7					
Course of study			3	7.68	0.0001	3.6573	0.301
Medicine degree	225	40.0 ± 22.3					
Nursing degree	49	36.2 ± 25.5					
Nursing diploma	58	40.1 ± 25.2					

(Continued)

Table 4. (Continued)

PCQN subscale		ANOVA of mean scores between and within the group				Bartlett's equal-variances test between and within the groups	
Independent variable	<i>n</i>	% knowledge scores (mean ± SD)	Degrees of freedom	<i>F</i> -statistic	<i>p</i> Value	Chi-square (χ^2)	<i>p</i> Value
Nursing certificate	134	28.5 ± 21.5					
Management of pain and other symptoms: Q2–4, Q6–8, Q10, Q13–16, Q18, Q20							
Age			1	0.75	0.3854	0.0430	0.836
20–29	428	48.7 ± 13.4					
30–39	38	46.8 ± 13.8					
Gender			1	0.10	0.7580	0.2062	0.650
Female	253	48.4 ± 13.3					
Male	213	48.8 ± 13.8					
Religion			5	2.44	0.0338	11.6150	0.040
Catholic	160	47.0 ± 13.2					
Protestant	164	50.1 ± 13.4					
Pentecostal	57	49.5 ± 16.1					
Muslim	42	51.1 ± 11.9					
SDA	34	43.2 ± 9.5					
Other	9	51.3 ± 14.4					
Course of study			3	0.59	0.6241	7.2730	0.064
Medicine degree	225	49.3 ± 14.3					
Nursing degree	49	47.3 ± 14.6					
Nursing diploma	58	48.8 ± 12.9					
Nursing certificate	134	47.7 ± 11.7					
Psychosocial and spiritual care: Q5, Q11, Q19							
Age			1	0.06	0.8036	0.0186	0.892
20–29	428	20.3 ± 24.5					
30–39	38	19.0 ± 24.1					
Religion			5	1.14	0.3361	3.8937	0.565
Catholic	160	19.2 ± 24.7					
Protestant	164	21.8 ± 25.7					

(Continued)

Table 4. (Continued)

PCQN subscale		ANOVA of mean scores between and within the group				Bartlett's equal-variances test between and within the groups	
Independent variable	<i>n</i>	% knowledge scores (mean ± SD)	Degrees of freedom	<i>F</i> -statistic	<i>p</i> Value	Chi-square (χ^2)	<i>p</i> Value
Pentecostal	57	19.9 ± 22.6					
Muslim	42	14.3 ± 22.3					
SDA	34	26.5 ± 22.9					
Other	9	18.5 ± 17.6					
Course of study			3	3.06	0.0279	9.7220	0.021
Medicine degree	225	23.7 ± 26.6					
Nursing degree	49	17.0 ± 20.6					
Nursing diploma	58	15.5 ± 20.9					
Nursing certificate	134	17.7 ± 22.6					

ANOVA, analysis of variance; df, degrees of freedom; *F*, statistic; *p*, probability value; PCQN, Palliative Care Quiz for Nursing; SD, standard deviation of the sample; \bar{x} , mean knowledge score of the sample; χ^2 , Bartlett's Chi-square test statistic.

Table 5. Weighted logistic regression analysis to test correlations between students' knowledge of palliative care and independent/predictor variables.

Independent variables	Odds ratio	Std. error	<i>t</i> -Statistic	<i>p</i> Value	95% confidence interval
Philosophy and principles of palliative care					
Age group					
20–29*					
30–39	1.3226	0.1884	1.96	0.050	0.9995–1.7499
Gender					
Female*					
Male	1.1386	0.0898	1.65	0.101	0.9759–1.3296
Course of study					
Degree in medicine*					
Degree in nursing	0.8202	0.1090	–1.49	0.137	0.0632–1.0652
Diploma in nursing	0.9632	0.1153	–0.31	0.754	0.7611–1.2189
Certificate in nursing	0.7008	0.0671	–3.71	0.001	0.5806–0.8459
Management of pain and symptoms					
Age group					

(Continued)

Table 5. (Continued)

Independent variables	Odds ratio	Std. error	t-Statistic	p Value	95% confidence interval
20–29*					
30–39	0.8924	0.0904	-1.12	0.261	0.0731–1.0889
Gender					
Female*					
Male	0.9947	0.0569	-0.09	0.926	0.8888–1.1131
Course of study					
Degree in medicine*					
Degree in nursing	0.9177	0.0857	-0.92	0.358	0.7638–1.1026
Diploma in nursing	0.9894	0.0859	-0.12	0.902	0.8341–1.1736
Certificate in nursing	0.9409	0.0634	-0.91	0.366	0.8243–1.0740
Psychosocial and spiritual care					
Age group					
20–29*					
30–39	1.0262	0.1611	0.16	0.869	0.7531–1.3984
Gender					
Female*					
Male	0.9502	0.0858	-0.57	0.572	0.7952–1.1353
Course of study					
Degree in medicine*					
Degree in nursing	0.7369	0.1073	-2.10	0.037	0.5531–0.9819
Diploma in nursing	0.7735	0.1108	-1.79	0.074	0.5831–1.0259
Certificate in nursing	0.7924	0.0836	-2.21	0.028	0.6437–0.9755
Variables marked with an asterisk (*) represent the reference group.					

degree in medicine. The odds of having knowledge of palliative care was 21% lower ($p=0.028$) OR 0.7924 (95% CI: 0.6437–0.9755) if the student was doing a certificate in nursing course compared with if they were doing a degree in medicine course.

In Table 6, results indicate that participants had the lowest self-efficacy scores in all the eight items in the communication subscale with items regarding discussing death and dying, discussing impending death with the patient and with the

patient's family, bereavement, and answering the patient question 'how long have I got to live?' registering very low scores.

Somewhat better but low self-efficacy scores were observed in the patient management subscale. Notably, participants demonstrated low self-efficacy scores in their ability to prescribe appropriate and adequate pain control medication and provision of psychosocial care for the patient and their family. Similarly, results showed participants had low self-efficacy to provide palliative

Table 6. Participants' Palliative Care Self-Efficacy scale results.

Code no.	Patient/family interactions and clinical management	1	2	3	4
		n (%)	n (%)	n (%)	n (%)
CM: Communication					
CM1	Discussing the likely effects of cancer with the patient	92 (19.9)	136 (29.4)	127 (27.4)	108 (23.3)
CM2	Discussing the likely effects of cancer with the patient's family	47 (10.2)	147 (32.0)	147 (34.2)	108 (23.5)
CM3	Discussing the issues of death and dying	104 (22.9)	134 (29.5)	148 (32.6)	68 (15.0)
CM4	Discussing the patient's death (to occur) with the patient	133 (29.3)	154 (33.9)	114 (25.1)	53 (11.7)
CM5	Discussing the patient's death (to occur) with the patient's family	103 (22.4)	156 (33.9)	126 (27.4)	75 (16.3)
CM6	Discussing the patient's death with the family upon bereavement	90 (19.6)	142 (30.9)	147 (32.0)	80 (17.4)
CM7	Answering the patient's questions 'How long have I got to live?'	127 (27.6)	140 (30.4)	115 (25.0)	79 (17.1)
CM8	Answering the patient's questions 'Will there be much suffering or pain?'	79 (17.2)	126 (27.5)	145 (31.6)	109 (23.8)
PM: Patient management					
PM1	In my ability to assess the patient's needs	64 (13.9)	111 (24.2)	171 (37.3)	113 (24.6)
PM2	In my knowledge of the aetiology of common symptoms experienced by palliative care patients	64 (13.8)	112 (24.2)	166 (35.9)	121 (26.1)
PM3	In my ability to manage common symptoms experienced in palliative care patients	61 (13.3)	122 (26.5)	168 (36.5)	109 (23.7)
PM4	In my ability to prescribe appropriate and adequate pain control medication	66 (14.3)	129 (28.0)	180 (39.1)	86 (18.7)
PM5	In my knowledge of the therapeutic and side effects of analgesic agents	61 (13.2)	104 (22.5)	160 (34.6)	138 (29.8)
PM6	In my ability to provide psychological care for the palliative care patient and their family	95 (20.7)	142 (30.9)	132 (28.8)	90 (19.6)
PM7	In my ability to provide social care for the palliative care patient and their family	81 (17.6)	134 (29.1)	165 (35.9)	80 (17.4)
PM8	In my ability to provide spiritual care for the palliative care patient and their family	87 (19.1)	115 (25.3)	135 (29.7)	118 (25.9)
MT: Multidisciplinary team					
MT1	Working in a multidisciplinary palliative care/healthcare team	77 (16.9)	127 (27.9)	151 (33.2)	100 (22.0)

(Continued)

Table 6. (Continued)

Code no.	Patient/family interactions and clinical management	1	2	3	4
		n (%)	n (%)	n (%)	n (%)
MT2	Appropriately referring palliative care patients for physiotherapy	86 (18.8)	127 (27.8)	141 (30.6)	103 (22.5)
MT3	Appropriately referring palliative care patients for occupational therapy	79 (17.3)	131 (28.7)	153 (33.6)	93 (20.4)
MT4	Appropriately referring palliative care patients for complementary therapies	88 (19.3)	153 (33.5)	134 (29.3)	82 (18.0)
MT5	Appropriately referring palliative care patients to a lymphoedema service	146 (31.7)	140 (30.4)	118 (25.6)	57 (12.4)
MT6	Appropriately referring palliative care patients for psychiatric evaluation	81 (17.7)	141 (30.8)	139 (30.4)	97 (21.2)
MT7	Appropriately referring palliative care patients to a spiritual advisor	73 (15.8)	108 (23.4)	128 (27.8)	152 (33.0)

Scoring: Need further basic instruction = 1, Confident to perform with close supervision / coaching = 2, Confident to perform with minimal consultation = 3, Confident to perform independently = 4.

care in all the seven items in the multidisciplinary teamwork subscale. Participants scored the lowest scores in the 'referring palliative care patients for complementary therapies and to a lymphoedema service' subscale items.

In Table 7, ANOVA was performed to investigate the relationship between students' self-efficacy scores and selected predictor variables. The course of study ($p < 0.0001$), level of interaction with terminally ill patients ($p < 0.0001$), history of working with terminally ill patients ($p < 0.0001$), an environment where one first learned about palliative care ($p < 0.0001$) and history of training in palliative care ($p < 0.0001$) all showed statistically significant relationships with students' self-efficacy scores. Bartlett's equal-variances test showed statistically significant results in only two variables; course of study ($p < 0.036$) and history or experience of working with palliative or terminally ill patients ($p < 0.001$). Certificate-level nursing participants had better self-efficacy scores than their diploma and degree in nursing and medicine counterparts.

Discussion

The main aim of this study was to examine final-year undergraduate nursing and medical students' knowledge and perceptions of, and

orientation toward, palliative care. The study found participants had low knowledge of palliative care with correct response scores ranging between 11.0% and 49.8% in all but 3 of the 20 items of the Palliative Care Quiz for Nursing questionnaire.

Regarding participants' demographics, statistically significant predictors of knowledge of palliative care were found to be gender ($p = 0.0242$) and course of study ($p = 0.0001$) in the '*philosophy and principles of palliative care*' subscale. Males and degrees in medicine and diplomas in nursing students showed higher knowledge scores of palliative care than their certificate and degree in nursing colleagues. Notably, Escribano et al.⁵¹ found that females had better knowledge of and self-efficacy to provide end-of-life care than their male counterparts.

Previous studies conducted in high-income countries found that females had higher knowledge of palliative care than their male counterparts.⁵²⁻⁵⁴ In the management of pain and other symptoms subscale, religion ($p = 0.0338$) was statistically significant with Muslims and Pentecostals having higher knowledge scores than Protestants and Catholics. Similarly, in the Psychosocial and Spiritual Care subscale, the course of study was significant ($p = 0.0279$) with

Table 7. Analysis of variance results on the relationship between students' self-efficacy scores and predictor variables.

Predictor variable	Analysis of variance			Bartlett's equal-variances test	
	Mean (SD)	F-statistic	p Value	Chi-square	p Value
Course of study		8.47	$p < 0.0001$	8.56	$p < 0.036$
Medicine degree	58.6 (14.9)				
Nursing degree	49.6 (16.8)				
Nursing diploma	54.3 (12.9)				
Nursing certificate	60.5 (12.6)				
Level of interaction with terminally ill patients		6.45	$p < 0.0001$	0.86	$p > 0.930$
Very frequently	62.1 (13.8)				
Frequently	60.2 (13.8)				
Occasionally	56.8 (14.5)				
Rarely	52.2 (14.9)				
Never	50.4 (14.0)				
History or experience of working with palliative or terminally ill patients?		33.92	$p < 0.0001$	14.2	$p < 0.001$
Yes	60.0 (12.9)				
No	51.7 (14.6)				
Where did you first learn about palliative care?		5.85	$p < 0.0001$	2.59	$p > 0.629$
Medical/nursing school	58.6 (13.4)				
Workplace/hospital	54.0 (13.8)				
Family or friends	52.0 (16.3)				
Media	56.4 (10.0)				
Never	39.4 (12.9)				
Have you had any training in palliative care in your institution?		33.84	$p < 0.0001$	3.07	$p > 0.080$
Yes	59.5 (13.7)				
No/Never	49.6 (15.8)				

medical and diploma nursing students having higher knowledge scores. This latter observation could be linked to prior exposure to the clinical setting given that most of the diploma nursing students are already qualified certificate-level nurses. Clinical exposure and higher level of education have been reported as predictors of knowledge of palliative and end-of-life care

among nursing and medical students in previous studies.^{19,37,42,54–56}

The low and insufficient knowledge of medical students observed in this study is documented in research conducted elsewhere among undergraduate nursing and medical students in high-income countries.^{53,54,57,58} This study found no

association between age and students' knowledge of palliative care, contrary to a wide body of evidence from previous research.^{35,52,54,59,60}

All the students in this study demonstrated low to poor understanding of palliative care concepts with the lowest mean knowledge scores in the *philosophy and principles of palliative care* (1.46 ± 0.93) and psychosocial and spiritual care (0.61 ± 0.73) subscales, while generally, students had better knowledge scores in the management of pain and other symptoms subscale (6.32 ± 1.75). This could be due to the predominant disconnect between theory and practice. To note, in the qualitative findings of the study,⁴⁴ students blamed the training which they said is mainly theory (classroom) based with virtually no or very limited opportunities for them to experience learning in a real-life clinical setting. Similar findings of inadequate to poor knowledge of palliative care among nursing students are reported in previous studies in both middle-income and high-income and countries^{37,61–66} and medical students in high-income countries.^{35,41,67–71} This underlines a global need for concerted action strengthening palliative and end-of-life training.

Regarding perceptions and self-efficacy to provide palliative care, the results of the Palliative Care Self-Efficacy scale found student participants scored low in nearly all the scale items. This could be linked to, and stemming, from deficiencies in training in the classroom; notably training that is predominantly theoretical when numerous evaluation and intervention studies, all in high-income countries, have demonstrated that palliative care training integrating problem-based learning improved students' knowledge, attitudes and skills.^{26–29}

Low self-efficacy to provide palliative care to patients and families with life-limiting illnesses has been widely reported in studies with medical and nursing graduates.^{22,33,40,57,64,66,68,72,73} Numerous studies corroborate that this is due to inadequate training about palliative and end-of-life care which has been widely reported in nursing,^{37,53,56,74–76} medicine,^{77–79} and high-income countries.^{27,67,69,80–82} Intervention and evaluation studies have shown how education and training increase knowledge of, and confidence to provide palliative and end-of-life care.^{26,28,55,83,84} Surprisingly, certificate-level nursing students showed higher self-efficacy scores compared to diploma and degree nursing and degree medicine

students. Hypothetically, it may be that the former has more clinical exposure (hands-on) than the latter. In their study in Portugal, Martins et al.⁸⁵ found that self-efficacy was the variable that had the most significant impact on qualified nurses' and doctors' ability to provide end-of-life care.

Notably, all participants scored the lowest scores in all three elements of the Psychosocial and Spiritual Care subscale and Palliative Care Self-Efficacy scale, particularly handling death work, that is, discussions around death, dying and bereavement, further underlining the training gap. This finding is consistent with previous studies conducted with qualified health professionals.^{36,86} However, some studies conducted found that qualified health professionals were moderately prepared and qualified to provide end-of-life care.^{85,87,88} Previous research has demonstrated adequate education and training improve students' capacity to handle death work.^{54,66,89} In fact, adequate education and training were found to reduce emotional distress, death avoidance and anxiety related to death and dying in nursing students,^{57,90} and among medical students.^{27,29,91,92} Adesina et al.⁴² in their study reported a negative correlation between training and students' knowledge of, and confidence and competence to provide end-of-life care.

Strengths and limitations of the study

The study has some salient strengths. First, this is one of the first studies to examine knowledge of and self-efficacy to provide palliative and end-of-life care and their correlates among undergraduate nursing and medical students in Uganda. Second, the multicentre nature of the study, with participants from different courses makes the findings more generalizable. Third, validated measurements were used to collect data, this adds to the reliability, dependability and validity of the study findings. Fourth, although the study employed non-probability convenience sampling, a large sample size of 466 increased the statistical power of convenience data, and so minimization of type II error.

The study has a few limitations. First, the cross-sectional and multicentre design could potentially be associated with contextual differences such as variations in course content and scope which may to some extent, contribute to differences in students' knowledge and perceptions of and

orientation towards PC. Yet, this study may not be able to adequately explain such contextual and causal differences and relationships.

Second, this is a one-point inquiry, hence, it is not possible to give clear accounts for some salient observations and causal trends. For example, the influence of religion and other important tenets of culture on students' knowledge, perceptions and orientation towards PC, and also the extent to which the curriculum contributes to the study findings. Hence, limited ability to control confounders.

Third, the study was conducted in only eight teaching institutions, hence, the findings may not be wholly generalizable or a true representative and reflection of the reality in other non-participating institutions.

Fourth, the authors acknowledge the inadequacy of question items to assess students' knowledge of 'spiritual care' aspect in the 'psychosocial and spiritual care domain of the PCQN scale', though the aspect was further adequately explored in the qualitative part.

Conclusions and recommendations

The study findings highlight participants generally had low and inadequate knowledge of and low self-efficacy to provide palliative care, underlining deficiencies in training and perhaps curricula. There is an urgent need to integrate and strengthen practical, pedagogical and experiential teaching to foster skills development but also to allow for translation of knowledge into practice. In particular, students should be placed, or rotated through clinical settings including hospices, cancer centres and wards that look after patients with palliative care need for them to experience learning in a real-life setting.

The study found participants had serious low self-efficacy or self-confidence, most notably in handling death work or end-of-life care. Teaching and training should urgently address existing gaps in death work, including difficult conversations around death and dying. Importantly, there is a need to review the palliative care curriculum to establish its adequacy in meeting the training needs of students.

Evaluative, more robust qualitative study designs, for example, grounded theory and ethnography

and longitudinal studies are needed to further examine how culture and the palliative care course and mode of delivery orientate and impact students' knowledge and perceptions of palliative and end-of-life care.

Declarations

Ethics approval and consent to participate

The study was approved by Hospice Africa Uganda Research Ethics Committee (HAUREC), protocol number HAU-2022-03-A. Ethical approval was granted on 20th September 2022. Administrative clearance was obtained from Makerere University School of Medicine Research Ethics Committee (SOMREC) on 28th October 2022. Written permission to access and enroll participants in the study was obtained from the res administration of each of the participating study sites. Voluntary written informed consent was obtained from each of the study participants.

Consent for publication

Not applicable.

Author contributions

Germanus Natuhwera: Conceptualization; Data curation; Formal analysis; Investigation; Methodology; Project administration; Validation; Visualization; Writing – original draft; Writing – review & editing.

Eve Namisango: Data curation; Formal analysis; Supervision; Validation; Visualization; Writing – review & editing.

Peter Ellis: Conceptualization; Methodology; Supervision; Validation; Writing – review & editing.

Acknowledgements

The authors thank the following individuals who supported data collection for this research. Mr. Kintu Herbert, Mulago School of Nursing and Midwifery, Kampala, Uganda. Dr. Rogers Abaho, Kampala International University and Medical Officer, Bushenyi District Local Government, Bushenyi, Uganda. Dr. Victor Niwamanyire, Makerere University College of Health Sciences, Kampala, Uganda. Dr. Kibet Emmanuel, Mbarara University of Science and Technology, Mbarara, Uganda. Dr. Robert Alinda, Uganda Christian University and Mulago National Referral Hospital, Kampala, Uganda. Ms. Mercy

Akoragye, Student Midwife, Mayanja Training Institute, Mbarara, Uganda.

Funding

The authors received no financial support for the research, authorship, and/or publication of this article.

Competing interests

The authors declare that there is no conflict of interest.

Availability of data and materials

Requests for data collected and analysed in this study should be directed to the corresponding author at: gnatuhwera@hospice-africa.org or ngermans16@gmail.com. Satisfactory reason of the need for the data must be supplied before data can be released.

ORCID iDs

Germanus Natuhwera  <https://orcid.org/0000-0003-4474-4063>

Eve Namisango  <https://orcid.org/0000-0001-5032-4714>

Supplemental material

Supplemental material for this article is available online.

References

1. Etkind SN, Bone AE, Gomes B, et al. How many people will need palliative care in 2040? Past trends, future projections and implications for services. *BMC Med* 2017; 15: 102.
2. Pereira SM, Hernandez-Marrero P, Pasman HR, et al. Nursing education on palliative care across Europe: results and recommendations from the EAPC taskforce on preparation for practice in palliative care nursing across the EU based on an online-survey and country reports. *Palliat Med* 2021; 35: 130–141.
3. World Health Organization. Palliative care, <https://www.who.int/news-room/fact-sheets/detail/palliative-care> (2020, accessed 13 August 2024).
4. Sleeman KE, de Brito M, Etkind S, et al. The escalating global burden of serious health-related suffering: projections to 2060 by world regions, age groups, and health conditions. *Lancet Glob Health* 2019; 7: e883–e892.
5. Worldwide Hospice Palliative Care Alliance and World Health Organization. *Global atlas of palliative care*. 2nd ed., <http://www.thewhpc.org/resources/global-atlas-of-palliative-care> (2020, last accessed 19 December 2024).
6. Abu-Odah H, Molassiotis A and Liu J. Lessons learned from countries that have introduced palliative care services into their national health system: a narrative review. *J Palliat Care* 2021; 37: 55–66.
7. Knaut F, Farmer P, Krauker E, et al. Alleviating the access abyss in palliative care and pain relief-an imperative of universal health coverage: the Lancet Commission Report. *Lancet* 2018; 39(10128): 1391–1454.
8. Downing J, Grant L, Leng M, et al. Understanding models of palliative care delivery in sub-Saharan Africa: learning from programs in Kenya and Malawi. *J Pain Symptom Manage* 2015; 50(3): 362–370.
9. Harding R and Higginson IJ. Palliative care in sub-Saharan Africa. *Lancet* 2005; 365, 1971–1977.
10. Namukwaya E, Mwaka AD, Namisango E, et al. Current state of palliative care in Uganda. In: Silbermann M (ed.) *Palliative care for chronic cancer patients in the community*. Cham: Springer, 2021, pp. 267–278. https://doi.org/10.1007/978-3-030-54526-0_24.
11. Palliative Care Association of Uganda. *Essential palliative care package for universal health coverage in Uganda: do not leave children and adults with life limiting illnesses in UHC*. Kampala: PCAU, 2021.
12. Logie DE and Harding R. An evaluation of a morphine public health programme for cancer and AIDS pain relief in sub-Saharan Africa. *BMC Public Health* 2005; 5, 82.
13. Ferrell B, Malloy P, Mazanec P, et al. CARES: AACN's new competencies and recommendations for educating undergraduate nursing students to improve palliative care. *J Prof Nurs* 2016; 32: 327–333.
14. Brooks LA, Manias E and Nicholson P. Communication and decision-making about end-of-life care in the intensive care unit. *Am J Crit Care* 2017; 26(4): 336–341.
15. Chover-Sierra E, Martínez-Sabater A and Lapeña-Moñux YR. An instrument to measure nurses' knowledge in palliative care: validation of the Spanish version of Palliative Care Quiz for Nurses. *PLoS One* 2017; 12: e0177000.
16. Pastrana T, Wüller J, Weyers S, et al. Insights from a community-based palliative care course: a qualitative study. *BMC Palliat Care* 2021; 20(1): 106.
17. Schallenburger M, Schwartz J, Batzler YN, et al. Handling the desire to die- evaluation of an

- elective course for medical students. *BMC Med Educ* 2024; 24(1): 279.
18. Fristedt S, Grynne A, Melin-Johansson C, et al. Registered nurses and undergraduate nursing students' attitudes to performing end-of-life care. *Nurse Educ Today* 2021; 98: 104772.
 19. Cao W, Li C, Zhang Q, et al. Perceptions on the current content and pedagogical approaches used in end-of-life care education among undergraduate nursing students: a qualitative, descriptive study. *BMC Med Educ* 2022; 22(1): 553.
 20. Haroen H, Mirwanti R and Sari CWM. Knowledge and attitude toward end-of-life care of nursing students after completing the multi-methods teaching and learning palliative care nursing course. *Sustainability* 2023; 15: 4382.
 21. Schulz-Quach C, Wenzel-Meyburg U and Fetz K. Can elearning be used to teach palliative care? – medical students' acceptance, knowledge, and self-estimation of competence in palliative care after elearning. *BMC Med Educ* 2018; 18(1): 1–7.
 22. Pieters J, Dolmans DHJM, Verstegen DML, et al. Palliative care education in the undergraduate medical curricula: students' views on the importance of, their confidence in, and knowledge of palliative care. *BMC Palliat Care* 2019; 18(1): 72.
 23. Amroud MS, Raeissi P, Hashemi SM, et al. Investigating the challenges and barriers of palliative care delivery in Iran and the World: a systematic review study. *J Educ Health Promot* 2021; 10: 246.
 24. Wang W, Wu C, Bai D, et al. A meta-analysis of nursing students' knowledge and attitudes about end-of-life care. *Nurse Educ Today* 2022; 119: 105570.
 25. Centeno C, Ballesteros M, Carrasco JM, et al. Does palliative care education matter to medical students? The experience of attending an undergraduate course in palliative care. *BMJ Support Palliat Care* 2016; 6: 128–134.
 26. Moody K, McHugh M, Baker R, et al. Providing pediatric palliative care education using problem-based learning. *J Palliat Med* 2018; 21(1): 22–27.
 27. Wallace CL, Cohen HL and Jenkins DA. Transforming students' attitudes and anxieties toward death and loss: the role of prior death experiences. *OMEGA* 2019; 79(1): 52–71.
 28. Kudubes AA and Murat Bektas M. The effect of web-based pediatric palliative care education on the palliative care knowledge level and practices of nursing students. *Perspect Psychiatr Care* 2020; 56: 533–540.
 29. Moehl K, Wright RM, Shega J, et al. How to teach medical students about pain and dementia: e-learning, experiential learning, or both? *Pain Med* 2020; 21(10): 2117–2122.
 30. Aslakson RA, Wyskiel R, Thornton I, et al. Nurse-perceived barriers to effective communication regarding prognosis and optimal end-of-life care for surgical ICU patients: a qualitative exploration. *J Palliat Med* 2012; 15(8): 910–915.
 31. Strang S, Bergh I, Ek K, et al., Swedish nursing students' reasoning about emotionally demanding issues in caring for dying patients. *Int J Palliat Nurs* 2014; 20(4): 194–200.
 32. O'Brien MR, Kinloch K, Groves KE, et al. Meeting patients' spiritual needs during end-of-life care: a qualitative study of nurses' and healthcare professionals' perceptions of spiritual care training. *J Clin Nurs* 2018; 28: 182–189.
 33. Hökkä M, Lehto JT, Kyngäs H, et al. Finnish nursing students' perceptions of the development needs in palliative care education and factors influencing learning in undergraduate nursing studies – a qualitative study. *BMC Palliat Care* 2022; 21(1): 40.
 34. Leung SY and Wong EL. Assessing medical students' confidence towards provision of palliative care: a cross-sectional study. *Int J Environ Res Public Health* 2021; 18(15): 8071.
 35. Abuhammad S, Muflih S, Alazzam S, et al. Knowledge of pediatric palliative care among medical students in Jordan: a cross-sectional study. *Ann Med Surg (Lond)* 2021; 64: 102246.
 36. Alshammari F, Sim J, Lapkin S, et al. Registered nurses' knowledge, attitudes and beliefs about end-of-life care in non-specialist palliative care settings: a mixed studies review. *Nurse Educ Pract* 2022; 59: 103294.
 37. Alwawi AA, Abu-Odah H and Bayuo J. Palliative care knowledge and attitudes towards end-of-life care among undergraduate nursing students at Al-Quds University: implications for Palestinian education. *Int J Environ Res Public Health* 2022; 19(15): 9563.
 38. Wang Y and Yuan H. Nursing undergraduates' experiences of a simulation-centred educational program in hospice care in Macao: qualitative research. *Int J Nurs Sci* 2022; 9(4): 504–511.
 39. Yoong SQ, Wang W, Seah ACW, et al. Nursing students' experiences with patient death and palliative and end-of-life care: a systematic review and meta-synthesis. *Nurse Educ Pract* 2023; 69: 103625.

40. Camara C, Rosengarten L and Callum J. Experiences of nursing students providing end of life care for children and young people: a focus group study. *Nurse Educ Today* 2024; 137: 106147.
41. Argyra E, Sifaka I, Moutzouri A, et al. How does an undergraduate pain course influence future physicians' awareness of chronic pain concepts? A comparative study. *Pain Med* 2015; 16: 301–311.
42. Adesina O, DeBellis A and Zannettino L. Third-year Australian nursing students' attitudes, experiences, knowledge, and education concerning end-of-life care. *Int J Palliat Nurs* 2014; 20(8): 395–401.
43. McIlpatrick S, Connolly M, Collins R, et al. Evaluating a dignity care intervention for palliative care in the community setting: community nurses' perspectives. *J Clin Nurs* 2017; 26(23–24): 4300–4312.
44. Natuhwera G, Ellis P and Namisango E. Final-year undergraduate nursing and medical students' knowledge and perception of, orientation and perceived self-efficacy to provide palliative and end-of-life care. Qualitative findings from a parallel mixed methods study. *BMC Med Educ* [Preprint]2024: 1–40.
45. Cochran WG. *Sampling techniques*. 2nd ed. New York, NY: John Wiley & Sons Inc., 1977.
46. Ross MM, McDonald B and McGuinness J. The palliative care quiz for nursing (PCQN): the development of an instrument to measure nurses' knowledge of palliative care. *J Adv Nurs* 1996; 23(1): 126–137.
47. Hao Y, Zhan L and Huang M. Nurses' knowledge and attitudes towards palliative care and death: a learning intervention. *BMC Palliat Care* 2021; 20: 50.
48. Zeru T, Berihu H, Gerensea H, et al. Assessment of knowledge and attitude towards palliative care and associated factors among nurses working in selected Tigray hospitals, northern Ethiopia: a cross-sectional study. *Pan Afr Med J* 2020; 35: 121.
49. Nair M, Kumar D, Mahajan R, et al. Knowledge, attitudes, and practices regarding palliative care: a mixed-methods study from Bihar, India. *J Palliat Care* 2021; 36(1): 9–11.
50. Mason S and Ellershaw J. Assessing undergraduate palliative care education: validity and reliability of two scales examining perceived efficacy and outcome expectancies in palliative care. *Med Educ* 2004; 38(10): 1103–1110.
51. Escribano S, Cabañero-Martínez MJ, Fernández-Alcántara M, et al. Efficacy of a standardised patient simulation programme for chronicity and end-of-life care training in undergraduate nursing students. *Int J Environ Res Public Health* 2021; 18(21): 11673.
52. Usta E, Aygin D and Sağlam E. Knowledge and opinions of nursing students on palliative care: a university example. *J Hum Sci* 2016; 13(3): 4405.
53. Dimoula M, Kotronoulas G, Katsaragakis S, et al. Undergraduate nursing students' knowledge about palliative care and attitudes towards end-of-life care: a three-cohort, cross-sectional survey. *Nurse Educ Today* 2019; 74: 7–14.
54. Jiang Q, Lu Y, Ying Y, et al. Attitudes and knowledge of undergraduate nursing students about palliative care: an analysis of influencing factors. *Nurse Educ Today* 2019; 80(1): 15–21.
55. Morrison LJ, Thompson BM and Gill AC. A required third-year medical student palliative care curriculum impacts knowledge and attitudes. *J Palliat Med* 2012; 15(7): 784–789.
56. Cao M, Li WX, Wan M, et al. Nursing students knowledge of palliative care at a medical university in China: a cross-sectional study. *Open Access Lib J* 2023; 10: 1–15.
57. Zhou Y, Li Q and Zhang W. Undergraduate nursing students' knowledge, attitudes and self-efficacy regarding palliative care in China: a descriptive correlational study. *Nurs Open* 2020; 8: 343–353.
58. Cleary AS. Graduating nurses' knowledge of palliative and end-of-life care. *Int J Palliat Nurs* 2020; 26(1): 5–12.
59. Al Qadire M. Knowledge of palliative care: an online survey. *Nurse Educ Today* 2014; 34: 714–718.
60. Karkada S and Malathi Nayak BS. Awareness of palliative care among diploma nursing students. *Indian J Palliat Care* 2011; 17(1): 20–23.
61. Brajtman S, Forthergill-Bourbonnais F, Casey A, et al. Providing direction for change: assessing Canadian nursing students learning needs. *Int J Palliat Nurs* 2007; 13(5): 213–221.
62. Al-Khawaldeh OA, Al-Hussami M and Darawad M. Knowledge and attitudes regarding pain management among Jordanian nursing students. *Nurse Educ Today* 2013; 33(4): 339–345.
63. Bassah N, Seymour J and Cox K. A modified systematic review of research evidence about education for pre-registration nurses in palliative care. *BMC Palliat Care* 2014; 13: 56.

64. Harazneh L, Ayed A, Fashafsheh, et al. Knowledge of palliative care among bachelors nursing students. *J Health Med Nurs* 2015; 18: 25–32.
65. Polat Ü and Karataş T. Nursing students' knowledge of palliative care at a university in Turkey. *Res Theory Nurs Pract* 2018; 32(3): 276–295.
66. Dobrowolska B, Mazurb E, Pilewska-Kozak A, et al. Predicted difficulties, educational needs, and interest in working in end-of-life care among nursing and medical students. *Nurse Educ Today* 2019; 83: 104194.
67. Weber M, Schmiedel S, Nauck F, et al. Knowledge and attitude of final-year medical students in Germany towards palliative care: an inter-institutional questionnaire-based study. *BMC Palliat Care* 2011; 10: 19.
68. Eyigor S. Fifth-year medical students' knowledge of palliative care and their views on the subject. *J Palliat Med* 2013; 16(8): 941–946.
69. Korzeniewska-Eksterowicz A, PrzysBo A, Kwziarska B, et al. The impact of pediatric palliative care education on medical students' knowledge and attitudes. *ScientificWorldJournal* 2013; 2013: 498082.
70. Ferri P, Di Lorenzo R, Stifani S, et al. Nursing student attitudes toward dying patient care: a European multicenter cross-sectional study. *Acta Biomed* 2021; 92(S2): e2021018.
71. Gerlach C, Mai SS, Schmidtman I, et al. Palliative care in undergraduate medical education - consolidation of the learning contents of palliative care in the final academic year. *GMS J Med Educ* 2021; 38(6): Doc103.
72. Abu-El-Noor NI and Abu-El-Noor MK. Attitude of Palestinian nursing students toward caring for dying patients: a call for change in health education policy. *J Holist Nurs* 2016; 34(2): 193–199.
73. Sukcharoen P, Polruk J, Lukthitikul S, et al. Undergraduate nursing students' experiences of palliative care in the intensive care unit. *BMC Nurs* 2023; 22(1): 251.
74. Callahan MF, Breakwell S and Suhayda R. Knowledge of palliative and end-of-life care by student registered nurse anesthetists. *AANA J* 2011; 79(Suppl. 4): S15–S20.
75. Ayed A, Sayej S, Harazneh L, et al. The nurses' knowledge and attitudes towards the palliative care. *J Educ Pract* 2015; 6(4): 22–28.
76. Chari S, Gupta M, Choudhary M, et al. Knowledge and attitude of nursing students towards palliative care: role of focused training. *Int J Biomed Res* 2016; 7(9): 629–632.
77. Head BA, Schapmire TJ, Earnshaw L, et al. Improving medical graduates' training in palliative care: advancing education and practice. *Adv Med Educ Pract*. 2016; 7: 99–113.
78. Richman PS, Saft HL, Messina CR, et al. Palliative and end-of-life educational practices in US pulmonary and critical care training programs. *J Crit Care* 2016; 31: 172–177.
79. Çıtlı R, Okan İ, Önder Y, et al. Evaluation of the awareness of medical students on palliative care. *Bezimalen Sci* 2018; 6: 100–107.
80. Li L, Hong FF and Liu WJ. The investigation on the cognition in palliative care of nursing undergraduates. *Chin Gen Pract Nurs* 2015; 13(28): 2776–2778.
81. Aboshaiqah AE. Predictors of palliative care knowledge among nursing students in Saudi Arabia: a cross-sectional study. *J Nurs Res* 2020; 28(1): e60.
82. Al Khalailah M and Al Qadire M. Pain management in Jordan: nursing students' knowledge and attitude. *Br J Nurs* 2013; 22(21): 34–40.
83. Bailey C and Hewison A. The impact of a 'Critical Moments' workshop on undergraduate nursing students' attitudes to caring for patients at the end of life: an evaluation. *J Clin Nurs* 2014; 23: 3555–3563.
84. Valsangkar S, Trupti N, Bodhare TN, et al. Evaluation of knowledge among interns in a medical college regarding palliative care in people living with HIV/AIDS and the impact of a structured intervention. *Indian J Palliat Care* 2011; 17(1): 6–10.
85. de Lurdes da Costa Martins M, Rodrigues AP, Marques CDCP, et al. Do spirituality and emotional intelligence improve the perception of the ability to provide care at the end of life? The role of knowledge and self-efficacy. *Palliat Support Care* 2024; 22(5): 1109–1117.
86. Shi H, Shan B, Zheng J, et al. Knowledge and attitudes toward end-of-life care among community health care providers and its influencing factors in China: a cross-sectional study. *Medicine (Baltimore)* 2019; 98(45): e17683.
87. Rodenbach R, Kavalieratos D, Tamber A, et al. Coaching palliative care conversations: evaluating the impact on resident preparedness and goals of-care conversations. *J Palliat Med* 2020; 23(2): 220–225.

88. Gelfman LP, Morrison RS, Moreno J, et al. Palliative care as essential to a hospital system's pandemic preparedness planning: how to get ready for the next wave. *J Palliat Med* 2021; 24(5): 656–658.
89. Pawłowski L, Wyszadko A, Janiszewska J, et al. The impact of palliative medicine education on medical students' perception of palliative care. *Psychoonkologia* 2015; 1: 19–26 [In Polish].
90. Hagelin CL, Melin-Johansson C, Henoch I, et al. Factors influencing attitude toward care of dying patients in first-year nursing students. *Int J Palliat Nurs* 2016; 22: 28–36.
91. Wong WY. The concept of death and the growth of death awareness among university students in Hong Kong: a study of the efficacy of death education programmes in Hong Kong universities. *OMEGA* 2017; 74(3): 304–328.
92. Xu F, Huang K, Wang Y, et al. A questionnaire study on the attitude towards death of the nursing interns in eight teaching hospitals in Jiangsu, China. *Biomed Res Int* 2019; 2019: 3107692.

Visit Sage journals online
[journals.sagepub.com/
home/pcr](https://journals.sagepub.com/home/pcr)

 Sage journals