

**Conclusion.** The current state of room movement with CDI patients is suboptimal, resulting in increased infection risk for subsequent patients. A multi-faceted intervention to address this problem is essential to achieve an optimal goal of 1–2 rooms per hospital stay for a CDI patient. Quality improvement projects include (1) notification of patient movement to nursing leadership, (2) alerts to nursing supervisors of potential CDI patients before room transfer, and (3) earlier diagnosis of community-onset *C. difficile*.

**Disclosures.** All authors: No reported disclosures.

### 1156. Observation of Stethoscope Sanitation Practices in an Emergency Department Setting

Rajiv Vasudevan, BS<sup>1</sup>; Sean Mojaver, BS<sup>1</sup>; Kay-Won Chang, MD<sup>2</sup>; Alan Maisel, MD<sup>3</sup>; Romteen Sedighi, BS<sup>1</sup> and Punam Chowdhury, MD<sup>3</sup>; <sup>1</sup>Cardiology, University of California, San Diego, San Diego, California, <sup>2</sup>Internal Medicine, University of California, San Diego, San Diego, California, <sup>3</sup>Internal Medicine, VA San Diego Healthcare System, La Jolla, California

**Session:** 135. Healthcare Epidemiology: Environmental and Occupational Health  
**Friday, October 5, 2018: 12:30 PM**

**Background.** Nosocomial infections pose significant risk to patients and hygienic standards may be easily overlooked, especially in fast-paced emergency department settings. Studies have shown that stethoscopes can harbor pathogens, which can be transferred to patients when proper sanitary measures are not taken. Survey-based studies have mostly assessed stethoscope hygiene, but they may not accurately represent cleaning practice. This study aimed to accurately assess cleaning practice through observation of stethoscope cleaning and hand hygiene among medical providers in an emergency department setting.

**Methods.** Four hundred twenty-six provider–patient encounters were observed in the emergency department of the VA San Diego Healthcare System. The frequency and methods of stethoscope and hand hygiene practices were anonymously observed and recorded. Stethoscope hygiene was recorded during and after each encounter if cleaning took place for at least 15 seconds. Hand hygiene data were also gathered before and after each encounter. Data analysis was performed to determine the frequency of these practices.

**Results.** Of 426 encounters, 115 involved the use of a personal stethoscope. In 15 of 115 encounters (13.0%), the provider placed a glove over the stethoscope before patient contact. Following patient interaction, 13 of 115 encounters (11.3%) involved stethoscope hygiene with an alcohol swab. Stethoscope hygiene with water and hand towel before patient interaction was observed in 5 of 115 encounters (4.3%). Hand sanitizer use or hand washing was observed in 213 of 426 encounters (50.0%) before patient interaction. Gloves were also used prior to patient interaction in 206 of 426 (48.4%) encounters, some overlapping with those who used hand sanitizer or washed their hands. Hand sanitizer or hand washing was used in 332 of 426 encounters (77.9%) after patient interaction.

**Conclusion.** Only 11.3% of encounters involving the use of a personal stethoscope had stethoscope sanitation with an alcohol swab. In 4.3% post-encounters, there was an attempt to clean stethoscopes with water and towel. Rates of appropriate hand hygiene were also lower than expected. Stethoscope and hand hygiene standards need to be raised to reduce infection risk and further studies need to be conducted to set guidelines for adequate stethoscope sanitation.

**Disclosures.** All authors: No reported disclosures.

### 1157. Optimal Isolation Periods Based on Time Interval to Elimination of Scabies After Topical Agent in the Real Clinical Practice

Jebyung Park, MD<sup>1</sup>; Se Yoon Park, MD<sup>1</sup>; Jaijun Han, MD<sup>1</sup>; So Young Lee, BSN<sup>2</sup>; Gil Eun Kim, BSN<sup>2</sup>; Yeon Su Jeong, BSN<sup>2</sup>; Jin Hwa Kim, MSN<sup>2</sup>; Eunyoung Lee, MD<sup>1</sup>; Eun Jung Lee, MD<sup>1</sup>; Shi Nae Yu, MD<sup>3</sup>; Tark Kim, MD<sup>4</sup>; Min Hyok Jeon, MD<sup>3</sup>; Eun Ju Choo, MD<sup>4</sup> and Tae Hyong Kim, MD, PhD<sup>1</sup>; <sup>1</sup>Division of Infectious Diseases, Department of Internal Medicine, Soonchunhyang University Seoul Hospital, Seoul, Korea, Republic of (South), <sup>2</sup>Infection Control Team, Soonchunhyang University Seoul Hospital, Seoul, Korea, Republic of (South), <sup>3</sup>Division of Infectious Diseases, Soonchunhyang University Cheonan Hospital, Cheonan, Korea, Republic of (South), <sup>4</sup>Division of Infectious Diseases, Soonchunhyang University Bucheon Hospital, Bucheon, Korea, Republic of (South)

**Session:** 135. Healthcare Epidemiology: Environmental and Occupational Health  
**Friday, October 5, 2018: 12:30 PM**


**Background.** Scabies reemerged globally as the ninth of most highly prevalent skin diseases from 1990 to 2010. Existing topical agents, like permethrin and lindane are highly effective showing 95% or more therapeutic response. Patient isolation is only recommended for one day after treating with topical agents. However, there is little evidence-based recommendation on optimal isolation period after treatment in healthcare settings to prevent secondary infection.

**Methods.** All patients who were diagnosed from 2008 to 2017 with scabies at a referral university hospital in Seoul, Republic of Korea were analyzed. We investigated the time interval between symptom onset and diagnosis of scabies. The period from the application of topical agents and clinical resolution was also analyzed.

**Results.** A total of 23 patients were diagnosed with scabies. There was no crusted scabies such as Norwegian scabies. Seventy percent of these patients were referred from long-term care facilities. Median number of treatment was three times per patient (Interquartile range [IQR], 2–3). Time interval between the onset of symptoms and the diagnosis of scabies was median 4 days (IQR, 2–14). Median isolation period was

13 days (IQR, 8–15). Dermatologist confirmed successful treatment of 16 patients, the median time until confirmation was 15 days (IQR, 8–17). Three patients discontinued isolation before dermatologist's confirmation. After re-evaluation by a dermatologist, one of the three was re-isolated due to persistent scabies infestation.

**Conclusion.** Scabicides are highly efficacious, but repeated microscopic examination for confirming elimination is strongly recommended. Optimal isolation period should be individualized based on the repeated dermatologic examination.



**Figure 1.** A timeline of cured scabies patients confirmed by a dermatologist.

**Disclosures.** All authors: No reported disclosures.

### 1158. Needlestick Injuries and Other Potential Exposures to Bloodborne Pathogens Among Police Officers in a City Police Department, 2011–2016

Marie De Perio, MD<sup>1</sup>; Kerton Victory, PhD<sup>2</sup> and Matthew Groenewold, PhD<sup>1</sup>; <sup>1</sup>National Institute for Occupational Safety and Health, Centers for Disease Control and Prevention, Cincinnati, Ohio, <sup>2</sup>National Institute for Occupational Safety and Health, Centers for Disease Control and Prevention, Atlanta, Georgia

**Session:** 135. Healthcare Epidemiology: Environmental and Occupational Health  
**Friday, October 5, 2018: 12:30 PM**

**Background.** The approximately 850,000 police officers nationwide are at risk of bloodborne diseases through needlestick injuries and other exposure incidents because of the nature of their work. In response to a request for a health hazard evaluation, we determined the incidence and circumstances of needlestick injuries and other potential exposures to bloodborne pathogens among police officers at a city department from 2011 to 2016.

**Methods.** We analyzed data extracted from the city's centralized human resource database on all needlestick injuries and other potential exposure incidents from January 1, 2011 to December 31, 2016 and characterized their circumstances. We calculated the annual incidence of needlestick injuries using two methods. We ran a Poisson regression model to determine the trend in the annual incidence over time using SAS 9.4.

**Results.** We identified 13 needlestick injuries and 37 additional potential exposure incidents. Needlestick injuries most commonly occurred during pat-down searches of a suspect and during search of a suspect's property or vehicle. Nine of 11 source persons with documented test results after a needlestick injury were found to have hepatitis C infection. The annual incidence of needlestick injuries ranged from 0 to 5.07 per 1,000 police officers and from 0 to 2.45 per 10,000 reactive calls for service and did not appear to have a significant trend. Most exposure incidents consisted of spitting incidents, human bites, and other contact with blood. Of 22 source persons with blood drawn, four had hepatitis C infection, two have HIV infection, and one had both. None of the incidents reportedly led to transmission of hepatitis B, hepatitis C, or HIV.

**Conclusion.** Police officers in this department are at risk for needlestick injuries and other exposures to bloodborne pathogens. We recommended improvements to engineering, administrative, and personal protective equipment controls, including training on safe searching techniques and sharps evidence collection and provision of nitrile gloves.

**Disclosures.** All authors: No reported disclosures.

### 1159. Influenza Symptoms in Vaccinated Healthcare Workers in an H3N2-Dominant Season

Allison Bartlett, MD, MS<sup>1</sup>; Jessica P. Ridgway, MD, MS<sup>2</sup>; Rachel Marrs, DNP, RN, CIC<sup>3</sup>; Amanda M. Henson, MS, MLS(ASCP), CIC<sup>3</sup>; Vera Chu, MS, MLS(ASCP), CIC<sup>3</sup>; Aurea Enriquez, M(ASCP), CIC<sup>3</sup>; Cynthia Murillo, M(ASCP), CIC<sup>3</sup>; Demetria Runjo, MPH<sup>3</sup>; Alexandra Seguin, BSN, RN<sup>3</sup>; Kathleen G. Beavis, MD<sup>4</sup>; Caroline Guenette, MS<sup>5</sup> and Emily Landon, MD<sup>6</sup>; <sup>1</sup>Department of Pediatrics, Section Infectious Disease, University of Chicago/Comer Children Hospital, Chicago, Illinois, <sup>2</sup>Section of Infectious Diseases and Global Health, University of Chicago Medicine, Chicago, Illinois, <sup>3</sup>Infection Control Program, The University of Chicago Medicine, Chicago, Illinois, <sup>4</sup>Department of Pathology, University of Chicago, Chicago, Illinois, <sup>5</sup>Department of Occupational Medicine, The University of Chicago Medicine,