



Genome Sequences of Two Choline-Utilizing Methanogenic Archaea, *Methanococcoides* spp., Isolated from Marine Sediments

 Gordon Webster,^a  Alex J. Mullins,^a Andrew J. Watkins,^b  Edward Cunningham-Oakes,^a  Andrew J. Weightman,^a
 Eshwar Mahenthiralingam,^a  Henrik Sass^b

^aMicrobiomes, Microbes and Informatics Group, Organisms and Environment Division, School of Biosciences, Cardiff University, Cardiff, Wales, United Kingdom

^bSchool of Earth and Ocean Sciences, Cardiff University, Cardiff, Wales, United Kingdom

ABSTRACT The genomes of two *Methanococcoides* spp. that were isolated from marine sediments and are capable of carrying out methanogenesis from choline and other methylotrophic substrates were sequenced. The average nucleotide identity and *in silico* DNA-DNA hybridization analyses demonstrate that they represent species different from those previously described.

The genus *Methanococcoides* comprises four described and characterized species, *M. methylutens* (1), *M. burtonii* (2), *M. alaskense* (3), and *M. vulcani* (4), as well as several other strains isolated from marine and mangrove sediments (5–7). To date, all *Methanococcoides* species are obligate methylotrophs able to utilize a range of methylated compounds for methanogenesis and belong to the diverse methanogen family *Methanosarcinaceae* (8).

Two *Methanococcoides* strains (with >98% 16S rRNA gene similarity to *M. methylutens*) (5) were isolated from sediments of Aarhus Bay, Baltic Sea (AM1), and Napoli Mud Volcano, eastern Mediterranean Sea (NM1), in artificial seawater (ASW) supplemented with methylamine, using deep-agar shake tubes and a dilution-to-extinction series with antibiotics to inhibit bacterial growth at 25°C (5).

For genome sequencing, each strain was grown in 2 × 10 ml ASW with 10 mM trimethylamine in tubes fitted with rubber stoppers and caps. Cells collected by centrifugation were washed with ASW and transferred to Lysing Matrix E tubes, and DNA was extracted using the FastDNA Spin kit (MP Biomedicals) (9). Sequencing was performed on an Illumina NextSeq 500 platform using a Nextera XT DNA library preparation kit. For each genome, 0.5 to 1.0 million 2 × 150-bp paired-end reads were generated. Illumina adaptors were trimmed with Trim Galore version 0.4.2, quality was assessed using FastQC version 0.10.1, and contigs were assembled *de novo* using SPAdes version 3.9.1. The genome assemblies had 30× (AM1) and 64× (NM1) coverage. The genome metrics for the two assemblies are as follows: for AM1, 2.48 Mbp, 42.66% G+C content, an N_{50} value of 433,016 bp, and 46 contigs; and for NM1, 2.34 Mbp, 43.18% G+C content, an N_{50} value of 241,508 bp, and 45 contigs. Both genome sizes are close to those reported for other *Methanococcoides* species (Table 1). A total of 2,445 (AM1) and 2,292 (NM1) coding DNA sequences (CDS) were identified using the Prokka version 1.12-beta genome annotation tool (8). The AM1 and NM1 genomes contained 6 and 7 rRNAs, 44 and 46 tRNAs, and 3 and 2 predicted CRISPR regions and several cytochromes, respectively. The entire operon encoding methyl coenzyme M reductase (Mcr) and genes for methanogenesis (*fmd*, *ftr*, *mch*, *mtd*, *mer*, *mtrABCDEFGH*, and *hdrABCDE*) were present. The genomes contained evidence of monomethylamine, dimethylamine, trimethylamine, and methanol metabolism, with genes encoding methanol-corrinoid protein comethyltransferase (Mta), monomethylamine methyltrans-

Citation Webster G, Mullins AJ, Watkins AJ, Cunningham-Oakes E, Weightman AJ, Mahenthiralingam E, Sass H. 2019. Genome sequences of two choline-utilizing methanogenic archaea, *Methanococcoides* spp., isolated from marine sediments. *Microbiol Resour Announc* 8:e00342-19. <https://doi.org/10.1128/MRA.00342-19>.

Editor Frank J. Stewart, Georgia Institute of Technology

Copyright © 2019 Webster et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Gordon Webster, websterg@cardiff.ac.uk.

Received 28 March 2019

Accepted 4 April 2019

Published 2 May 2019

TABLE 1 Pairwise ANI and *in silico* DDH between the novel strains and other *Methanococcoides* species and their respective genome sizes

Strain genome (description and source or accession no.)	Genome size (Mbp)	Pairwise ANI (%) with genome of ^a :		Pairwise DDH (%) with genome of ^b :	
		AM1	NM1	AM1	NM1
AM1	2.48		90.4		39.0
NM1	2.34	90.4		39.0	
<i>M. methylutens</i> DSM 2657 ^T (JRHO00000000)	2.51	90.1	89.7	37.7	37.1
<i>M. burtonii</i> DSM 6242 ^T (CP000300)	2.58	84.5	85.1	21.0	21.6
<i>M. vulcani</i> SLH33 ^T (FOHQ00000000)	2.31	90.6	95.8	39.8	64.7
<i>M. methylutens</i> MM1 (CP009518)	2.39	85.6	85.5	26.2	26.7
<i>M. methylutens</i> DSM 2657 ^T (this study)	2.50	90.2	89.8	37.6	37.1
<i>M. vulcani</i> SLH33 ^T (this study)	2.32	90.6	95.9	39.7	64.8
<i>Methanococcoides</i> sp. strain EBM-47 (anaerobic digester metagenome; MPPA00000000)	2.18	83.7	83.3	15.2	15.8

^a Average nucleotide identity (ANI) values of <95% indicate different species.

^b *In silico* DNA-DNA hybridization (DDH) values of <70% indicate different species.

ferase (Mtm), dimethylamine methyltransferase (Mtb), and trimethylamine methyltransferase (Mtt), along with the corresponding corrinoid protein genes. Both strains had genes encoding methylsulfide methyltransferase-associated sensors predicted to be involved in two-component regulation systems (10).

Methanococcoides strains were subject to average nucleotide identity (ANI) analysis using PyANI (<https://github.com/widowquinn/pyani>) and *in silico* DNA-DNA hybridization (DDH) with the Genome-to-Genome Distance Calculator (GGDC) 2.1 (11). The AM1 genome possessed ANI values below 95% compared to previously recognized species, while the NM1 genome identity was just above 95% compared to *M. vulcani* (Table 1). An ANI value of <95% has been proposed for species delineation (12). The ANI comparisons indicated that AM1 represents a novel species, while NM1 is closely related to *M. vulcani*. However, analysis by *in silico* DDH produced values for both strains below the 70% species threshold compared with genomes from described *Methanococcoides* species, including NM1 with *M. vulcani* (Table 1). The ANI and DDH values together suggested that AM1 and NM1 represent two phylogenetically different species of *Methanococcoides* from those described previously and warrant further characterization.

Data availability. The genome sequences and Illumina raw sequence reads have been deposited at the European Nucleotide Archive (ENA) under the ENA project/study number PRJEB31721. The accession numbers for the genomes are CAAGSW010000000 for AM1 and CAAGTW010000000 for NM1. The genomes of *M. methylutens* DSM 2657 and *M. vulcani* SLH33 were also resequenced and submitted under accession numbers CAAGSM010000000 and CAAGSJ010000000, respectively.

ACKNOWLEDGMENTS

This research was funded by the Natural Environment Research Council (NERC) grant NE/F018983/1, awarded to H. Sass and A. J. Weightman. A. J. Mullins and E. Cunningham-Oakes acknowledge funding from the Biotechnology and Biological Sciences Research Council (BBSRC) South West doctoral training partnership (grant BB/M009122/1). G. Webster and E. Mahenthalingam acknowledge support from the Wellcome Trust Institutional Strategic Support Fund. Sequencing was carried out at the Cardiff School of Biosciences Genomics Research Hub. All analysis was done using the Cloud Infrastructure for Microbial Bioinformatics (CLIMB), funded by the Medical Research Council (MRC) grant MR/L015080/1.

REFERENCES

- Sowers KR, Ferry JG. 1983. Isolation and characterization of a methylotrophic marine methanogen, *Methanococcoides methylutens* gen. nov., sp. nov. *Appl Environ Microbiol* 45:684–690.
- Franzmann PD, Springer N, Ludwig W, Conway De Macario E, Rohde M. 1992. A methanogenic archaeon from Ace Lake, Antarctica: *Methanococcoides burtonii* sp. nov. *Syst Appl Microbiol* 15:573–581. [https://doi.org/10.1016/S0723-2020\(11\)80117-7](https://doi.org/10.1016/S0723-2020(11)80117-7).
- Singh N, Kendall MM, Liu Y, Boone DR. 2005. Isolation and characteriza-

- tion of methylotrophic methanogens from anoxic marine sediments in Skan Bay, Alaska: description of *Methanococcoides alaskense* sp. nov., and emended description of *Methanosarcina baltica*. *Int J Syst Evol Microbiol* 55:2531–2538. <https://doi.org/10.1099/ijs.0.63886-0>.
4. L'Haridon S, Chalopin M, Colombo D, Toffin L. 2014. *Methanococcoides vulcani* sp. nov., a marine methylotrophic methanogen that uses betaine, choline and *N,N*-dimethylethanolamine for methanogenesis, isolated from a mud volcano, and emended description of the genus *Methanococcoides*. *Int J Syst Evol Microbiol* 64:1978–1983. <https://doi.org/10.1099/ijs.0.058289-0>.
 5. Watkins AJ, Roussel EG, Webster G, Parkes RJ, Sass H. 2012. Choline and *N,N*-dimethylethanolamine as direct substrates for methanogens. *Appl Environ Microbiol* 78:8298–8303. <https://doi.org/10.1128/AEM.01941-12>.
 6. Watkins AJ, Roussel EG, Parkes RJ, Sass H. 2014. Glycine betaine as a direct substrate for methanogens (*Methanococcoides* spp.). *Appl Environ Microbiol* 80:289–293. <https://doi.org/10.1128/AEM.03076-13>.
 7. Lyimo TJ, Pol A, Jetten MSM, Op den Camp HJM. 2009. Diversity of methanogenic archaea in a mangrove sediment and isolation of a new *Methanococcoides* strain. *FEMS Microbiol Lett* 291:247–253. <https://doi.org/10.1111/j.1574-6968.2008.01464.x>.
 8. Sowers KR, Johnson JL, Ferry JG. 1984. Phylogenetic relationships among the methylotrophic methane-producing bacteria and emendation of the family *Methanosarcinaceae*. *Int J Syst Evol Microbiol* 34:444–450. <https://doi.org/10.1099/00207713-34-4-444>.
 9. Webster G, Newberry CJ, Fry JC, Weightman AJ. 2003. Assessment of bacterial community structure in the deep sub-seafloor biosphere by 16S rDNA-based techniques: a cautionary tale. *J Microbiol Methods* 55:155–164. [https://doi.org/10.1016/S0167-7012\(03\)00140-4](https://doi.org/10.1016/S0167-7012(03)00140-4).
 10. Molitor B, Stassen M, Modi A, El-Mashtoly SF, Laurich C, Lubitz W, Dawson JH, Rother M, Frankenberg-Dinkel N. 2013. A heme-based redox sensor in the methanogenic archaeon *Methanosarcina acetivorans*. *J Biol Chem* 288:18458–18472. <https://doi.org/10.1074/jbc.M113.476267>.
 11. Meier-Kolthoff JP, Auch AF, Klenk H-P, Göker M. 2013. Genome sequence-based species delimitation with confidence intervals and improved distance functions. *BMC Bioinformatics* 14:60. <https://doi.org/10.1186/1471-2105-14-60>.
 12. Goris J, Konstantinidis KT, Klappenbach JA, Coenye T, Vandamme P, Tiedje JM. 2007. DNA-DNA hybridization values and their relationship to whole-genome sequence similarities. *Int J Syst Evol Microbiol* 57:81–91. <https://doi.org/10.1099/ijs.0.64483-0>.