



RESEARCH

# IFSO Bariatric Endoscopy Committee Evidence-Based Review and Position Statement on Endoscopic Sleeve Gastroplasty for Obesity Management

Barham K. Abu Dayyeh<sup>1,10</sup> · Christine Stier<sup>2</sup> · Aayed Alqahtani<sup>3</sup> · Reem Sharaiha<sup>4</sup> · Mohit Bandhari<sup>5</sup> · Silvana Perretta<sup>6</sup> · Singh Pichamol Jirapinyo<sup>7</sup> · Gerhard Prager<sup>8</sup> · Ricardo V. Cohen<sup>9</sup>

Received: 8 September 2024 / Revised: 11 September 2024 / Accepted: 16 September 2024 / Published online: 1 November 2024  
© The Author(s) 2024

## Abstract

**Background** Obesity is a significant global health issue. Metabolic and bariatric surgery (MBS) is the gold standard in the treatment of obesity due to its proven effectiveness and safety in the short and long term. However, MBS is not suitable for all patients. Some individuals are at high surgical risk or refuse surgical treatment, while others do not meet the criteria for MBS despite having obesity-related comorbidities. This gap has driven the development of endoscopic solutions like endoscopic sleeve gastroplasty (ESG), which offers a less invasive alternative that preserves organ function and reduces risks. A recent IFSO International Delphi consensus study highlighted that multidisciplinary experts agree on the utility of ESG for managing obesity in patients with class I and II obesity and for those with class III obesity who do not wish to pursue or qualify for MBS. This IFSO Bariatric Endoscopy Committee position statement aims to augment these consensus statements by providing a comprehensive systematic review of the evidence and delivering an evidence-based position on the value of ESG within the spectrum of obesity management.

**Methods** A comprehensive systematic review followed the Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) and Cochrane guidelines.

**Results** *Systematic Review:* The systematic review included 44 articles encompassing 15,714 patients receiving ESG. The studies varied from large case series to cohort studies and a randomized controlled trial (RCT). The mean baseline BMI was 37.56 kg/m<sup>2</sup>. The review focused on weight loss outcomes and safety data.

*Meta-analysis:*

Time point	Mean %EWL	Mean %TBWL
6 months	<b>48.04</b>	<b>15.66</b>
12 months	<b>53.09</b>	<b>17.56</b>
18 months	<b>57.98</b>	<b>16.25</b>
24 months	<b>46.57</b>	<b>15.2</b>
36 months	<b>53.18</b>	<b>14.07</b>
60 months	<b>45.3</b>	<b>15.9</b>

These results demonstrate significant weight loss following ESG.

**Safety:** The pooled serious adverse event (SAE) rate was 1.25%. This low rate of SAEs indicates that ESG is a relatively safe procedure.

**Quality of Evidence:** The quality of evidence from the included observational studies was assessed as very low, primarily due to the inherent limitations associated with observational study designs, such as potential biases and lack of randomization. In contrast, the quality of evidence from the single randomized controlled trial was rated as MODERATE, reflecting a more robust study design that provides a higher level of evidence despite some limitations.

Extended author information available on the last page of the article

**Conclusions** The IFSO Bariatric Endoscopy Committee, after conducting a comprehensive systematic review and meta-analysis, endorses endoscopic sleeve gastroplasty (ESG) as an effective and valuable treatment for obesity. ESG is particularly beneficial for patients with class I and II obesity, as well as for those with class III obesity who are not suitable candidates for metabolic bariatric surgery. ESG provides significant weight loss outcomes and demonstrates a favorable safety profile with a low rate of serious adverse events. Despite the limitations of the included observational studies, the randomized controlled trial included in the analysis reinforces the efficacy and safety of ESG and provides an evidence-based foundation for the position statement. Thus, the IFSO position statement supports and provides an evidence base for the role of ESG within the broader spectrum of obesity management.

**Keywords** Obesity · Endoscopic Sleeve Gastroplasty · Meta-analysis

## Introduction

Obesity rates are galloping, though regional, cultural, and socioeconomic factors contribute to disparities in distribution, prevalence, and incidence across the globe [1]. Still, the World Health Organization (WHO) estimates that 1.9 billion people are overweight, with 650 million having obesity as of 2016 [2]. In the United States, around 40% of the population currently live with obesity [3, 4], and prediction models estimate that this number will increase to 51% by 2030 [5]. After unsuccessful non-invasive therapies, metabolic and bariatric surgery (MBS) is the gold-standard treatment to address moderate to severe obesity. Most recently, it has also been proposed for mild obesity if it is associated with refractory metabolic diseases [6]. MBS is effective and safe in the short and long term, promoting sustained weight loss and reliable reduction in all-cause mortality rates [7].

Data show that MBS procedures have increased over decades [8, 9]. However, the rate of obesity growth is outpacing the growth in surgical interventions [10]. In addition, several patients refuse surgical treatment, others are at high surgical risk, and some suffer from overweight or mild obesity but are still not eligible for MBS. Nevertheless, obesity-related complications increase in states of overweight and mild obesity [11]. Altogether, a gap between the needs of patients with obesity and what we can offer in terms of medical and surgical interventions exists.


This unmet need has driven the development of endoscopic solutions to address obesity, particularly when MBS is not feasible or indicated. Endoscopic bariatric therapies offer several advantages, including organ preservation, an improved risk profile, reduced healthcare utilization, and decreased burden of compliance on the patient. These benefits potentially enable the scalability of procedural offerings to effectively combat excess adiposity. Endoscopic sleeve gastroplasty (ESG) is one such solution that has gained global adoption from patients and providers in the past few years. In its current clinically adopted and regulatory approved form (Fig. 1) [12], ESG employs the Apollo Overstitch™ platform (Boston Scientific, Marlborough,

MA, USA)—a full-thickness endoscopic suturing device to create apposition of the anterior against the posterior wall of the stomach, passing through the greater curvature [13, 14]. The Overstitch™ platform is currently the only US FDA-approved endoscopic suturing device for an obesity indication. Suturing starts at the transition between the gastric body and antrum, moving proximally toward the fundus, which is typically partially reduced with the preservation of a small pouch to allow fundal accommodation. Thus, it tubularizes the gastric body, altering satiety and satiation [15]. Although different stitching patterns have been proposed and discussed [16–18], the above-mentioned anatomic principles are consistent across centers and providers; thus, the procedure is clinically mature, homogeneous, and reproducible [19].

The evidence concerning the efficacy and safety of the ESG has been mounting. More than 200 international medical articles have been published on this topic, with study designs varying from large cases series [20] to cohort studies [21] and, more recently, also includes an open-label, multicenter randomized trial with 24 months follow-up [22]. The procedure is currently employed clinically in all continents, and more than 40,000 clinical procedures have been performed to date. A recent IFSO International Delphi consensus conference highlighted that multidisciplinary experts agree on the utility of ESG for managing obesity in patients with class I and II obesity and for those with class III obesity who do not wish to pursue or qualify for MBS in the context of a comprehensive multidisciplinary obesity program [23]. This IFSO Bariatric Endoscopy Committee position statement aims to augment these consensus statements by providing a comprehensive systematic review of the evidence and delivering an evidence-based position on the value of ESG within the spectrum of obesity care.

## Methods

This position statement is derived after a comprehensive systematic review to retrieve all available data on the outcomes of ESG. All the Preferred Reporting Items for Systematic



**Fig. 1** Schematic representation of the endoscopic sleeve gastropasty procedure

Reviews and Meta-analyses (PRISMA) [24] and Cochrane Handbook for Systematic Reviews of Interventions [25] guidelines were rigorously followed for this position statement's systematic review and meta-analysis portion. Two independent researchers (VOB and NJ) conducted all literature searches and a third independent reviewer adjudicated discrepancies. After defining the eligibility criteria, final inclusion was determined by consensus with two additional researchers (RK and BAD). One researcher collected data from the included studies using a standardized shared spreadsheet, and another independently validated the data extraction. Methodologists' names and affiliations are in the "Acknowledgements" section.

The risk of bias in the included studies was assessed using the Joanna Briggs Institute Critical Appraisal checklist for case series [26], the New-Castle Ottawa scale for cohort studies [27], and both JADAD score [28] and a modified Cochrane

Collaboration Risk of Bias tool (available from <https://www.ncbi.nlm.nih.gov/books/NBK132494/bin/appf-fm1.pdf>).


We used the Review Manager (Version 5.4, the Cochrane Collaboration, 2020) for pooling comparative data and the Comprehensive Meta-analysis software (Version 4, Biostat, Englewood, NJ, USA, 2022) to pool non-comparative data. Means and standard deviations (SDs) were estimated from medians and ranges based on previously validated mathematical formulas [29]. The estimation of standard deviation based on interquartile ranges, or 95% confidence interval (CI), followed the instructions in the Cochrane Handbook (Chapter 06, Section 6–5–2) [25]. If the article did not provide any measure for dispersion or sample size, we attempted to obtain them by emailing the authors. If unsuccessful, we proceeded with data input based on the SD of articles with similar sample sizes and time points (per Cochrane Handbook's guidance).

Continuous variables were expressed preferably as means and standard deviation, while categorical ones were expressed as rates or frequencies. A  $p$ -value  $< 0.05$  was considered statistically significant for a 95%CI. As a measure of effect, we employed main difference (MD) with fixed-effect mode analysis to compare data. Then, we assessed for heterogeneity among studies with the Higgins test ( $I^2$ ).  $I^2$  higher than 50% indicated high heterogeneity, and sensitivity analyses utilizing forest plots were undertaken to assess for outliers. If no true outliers were identified, the heterogeneity was considered true, and we switched from fixed to random-effect mode analysis to mitigate its impact on the summary estimate.

Using the results from the critical appraisal/risk of bias assessment and the meta-analysis, we evaluated the quality of the current evidence using the Grading of Recommendations, Assessment, Development, and Evaluations (GRADE) approach [30]. This standardized methodology analyzes data per outcome and uses several aspects of the studies (study design, risk of bias, imprecision, inconsistency, indirectness, publication bias, magnitude of effect, dose–response gradient, impact of residual confounding on the summary estimate) to classify the quality of the pooled evidence into 4 different categories: VERY LOW, LOW, MODERATE, and HIGH. This assessment demonstrates our certainty on how close the actual effect is to the effect estimated in our meta-analysis. All the data was input into the GRADEpro GDT online software (GRADEpro Guideline Development Tool, McMaster University, and Evidence Prime, 2022) for analysis and generation of the overall quality of evidence.

Finally, considering all the information gathered from the systematic literature review and meta-analysis, balancing the benefits and harms of the therapy, clinicians' values and preferences, resource utilization, and cost-effectiveness, the committee determined the final position statement and level of support.

**Fig. 2** PRISMA flowchart for the literature screening and inclusion/exclusion process for the overall outcomes of ESG (non-comparative analysis)



Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(7): e1000097. doi:10.1371/journal.pmed1000097

## Results

### Safety and Efficacy of ESG

#### Outcomes of ESG

**Systematic Review** Two independent researchers (VOB and NJ) ran separate literature searches assessing eligible studies. We searched MEDLINE (PubMed), EMBASE, and gray literature from January 1, 2013 (the year ESG was described), to October 1, 2022. The final strategy was as follows:

- *MEDLINE (PubMed): (total weight loss) OR (total body weight loss) OR (excess weight loss) OR (absolute weight loss) OR (excess body weight loss) OR (responders rate) OR (adverse event) OR (BMI reduc-*

*tion) OR (BMI decrease) OR (complication) AND (endoscopy) OR (endoscopic) OR (transoral\*)OR (per-oral\*)OR (incisionless) AND (sleeve) OR ( overstitch) OR (gastroplasty) OR (gastric plication) OR (gastric imbrication) AND (overweight) OR (obesity) AND ("2013/01/01"[Date—Publication]: "3000"[Date—Publication])*

- *EMBASE: endoscopic AND sleeve AND gastroplasty OR (apollo AND overstitch) AND [embase]/lim NOT ([embase]/lim AND [medline]/lim) AND ('article'/it OR 'article in press'/it OR 'conference review'/it OR 'note'/it OR 'review'/it)*

The eligibility criteria included:

- Articles published online from 1 Jan. 2013 until 1 Oct. 2022 (last search update);

**Table 1** Baseline data of the 44 articles included in the meta-analysis of the ESG outcomes

Author (year)	Study design	Single vs. multicenter	Retrospective vs. prospective	Population (total sample)	Intervention (n)	Comparison (n)	Observations	Inclusion criteria (BMI)	Age	Sex	Mean baseline BMI (kg/m <sup>2</sup> )	Type II Diabetes
Abu Dayyeh, B. K., et al. [31]	Case series	Single center	Prospective	25	ESG	-	Study of gastric physiology	BMI > 30 and < 40	47.6 (10)	21 women	35.5 (2.6)	1
Abu Dayyeh, B. K., et al. [22]	Randomized clinical trial	9 centers	Prospective	209	ESG (85)	Lifestyle intervention (124)	Open-label FDA-regulated trial	BMI > 30 and < 40	47.3 (9.3) ESG and 45.7 (10) lifestyle intervention	68 (88%) ESG and 91 (84%) lifestyle intervention	35.5 (2.6) ESG and 35.7 (2.6) lifestyle intervention	18 ESG and 36 lifestyle intervention
Alqahtani, A., et al. [20]	Case series	Single center	Prospective	1000	ESG	-	-	Ineligible for or refuse bariatric surgery	34.4 (9.5)	897 (89.7%) women	33.3 (4.5)	17
Alqahtani, A., et al. [32]	Case series	Single center	Prospective	109	ESG	-	Patients aged < 21 years old	BMI > 120% of the 95th percentile	17.6 (2.2)	99 (91.7%) women	33 (4.7)	NR
Alqahtani, A. R., et al. [33]	Cohort (propensity score-matched)	Single center	Retrospective	6036	ESG (3018)	LSG (3018)	-	BMI > 27.5	33.8 (9.6) ESG and 33.9 (9.7) LSG	2686 (89%) women for both groups	32.5 (3.1) ESG and 32.9 (3.5) LSG	112 ESG
Asokkumar, R., et al. [34]	Case series	Single center	Prospective	35	ESG	-	-	BMI > 27.5	43.6 (11.3)	20 (52.7%) women	34 (4.9)	8 (23%)
Badurdeen, D., et al. [35]	Cohort (propensity score-matched)	3 centers	Prospective	52	ESG (26)	ESG + Liraglutide (26)	-	BMI > 27	41.15 (10.64) ESG and 40.65 (8.69) ESG + Liraglutide	16 women ESG and 17 women ESG + Liraglutide	35.56 (1.68) ESG and 35.83 (2.33) ESG + Liraglutide	16 (8 and 8)
Barrichello, S., et al. [36]	Case series	7 centers	Retrospective	193	ESG	-	-	BMI > 25	42.3 (9.6)	148 women	34.11 (2.97)	NR
Bhandari, M., et al. [37]	Case series	Single center	Retrospective	53	ESG	-	-	BMI > = 28	40.54 (13.79)	43 (81.1%)	34.78 (5.20)	10
Callahan, Z. M., et al. [38]	Case series	Single center	Retrospective	10	ESG	-	Study on GI tract suturing including 10 ESG cases	NR	50.2 (12.2)	10 women	NR	NR
Carr, P., et al. [39]	Cohort	Single center	Prospective	61	ESG (16)	LSG (45)	-	BMI > 26 with comorbidity or BMI > 30	41.4 (10.4) vs 40.4 (9.0)	31 women (57.4%) ESG and 59 (71.1%) LSG	35.5 (5.2) vs 40.7 (5.6)	0 ESG and 2 LSG
Cheskin, L. J., et al. [40]	Cohort (propensity score-matched)	Single center	Retrospective	386	ESG (105)	Lifestyle intervention (281)	-	NR	47.58 (11.97) ESG and 48.17 (12.18) Lifestyle intervention	75 (71.42%) women ESG and 189 (67.2%) women Lifestyle intervention	40.5 (7.89) ESG and 39.85 (7.62) lifestyle intervention	NR
Espinete-Coll, E., et al. [41]	Case series	Single center	Prospective	38	ESG	-	Focus on persistence of sutures	BMI > 27	47 (5.5)*	30 women	37.6 (3)*	NR

**Table 1** (continued)

Author (year)	Study design	Single vs. multicenter	Retrospective vs. prospective	Population (total sample)	Intervention (n)	Comparison (n)	Observations	Inclusion criteria (BMI)	Age	Sex	Mean baseline BMI (kg/m <sup>2</sup> )	Type II Diabetes
Espinete-Coll, E., et al. [42]	Cohort	2 centers	Retrospective	88	ESG (standard stitching pattern = "TbP")	ESG with different stitching patterns ("Lp" and "TbP")	Comparison of 3 different stitching patterns	BMI > 27	Overall 46.1 (12.3)	61 women	Overall 39.40 (4.69)	Overall 11 (12.5%)
Farha, J., et al. [43]	Cohort	2 centers	Retrospective	247	ESG (98)	ESG with fundal suturing (149)	-	BMI > 30	44.9 (9.4) vs 47.2 (11.5)	84 (85.7%) and 107 (71.8%)	38.3 (5.6) vs 39.4 (7.3)	Overall 16
Fayad, L., et al. [44]	Cohort	Single center	Retrospective	137	ESG (54)	LSG (83)	-	NR	48 (12) ESG and 47.75 (6.16) LSG*	31 women (57.4%) ESG and 59 (71.1%) LSG	43.07 (8.85) ESG and 44.12 (5.78) LSG*	3.7% ESG and 20.48% LSG
Fayad, L., et al. [45]	Cohort	Single center	Retrospective	105	ESG (58)	IGB (47)	-	BMI > 27 for IGB and BMI > 30 for ESG	48.2 (11.8) ESG and 47.7 (12.4) IGB	34 women (58.6%) ESG and 46 (97.9%) IGB	41.5 (8.2) ESG and 34.5 (6.7) IGB	3 (5.2) ESG and 4 (8.5) IGB
Fiorillo, C., et al. [46]	Cohort	Single center	Retrospective	46	ESG (23)	LSG (23)	Focus on quality of life after 6 months	NR	41 (2) ESG and 37 (4.5) LSG*	16 women ESG and 17 LSG	39.5 (2) ESG and 41 (1.27) LSG*	2 ESG and 3 LSG
Ghoz, H., et al. [47]	Case series	Single center	Retrospective	20	ESG	-	Focus on nutritional deficiencies	NR	46.2 (14.1)	17 women	36.4 (4.1)	2
Glaysheer, M. A., et al. [48]	Cohort	Single center	Prospective	32	ESG without longitudinal compression (9)	ESG with longitudinal compression (23)	-	BMI > 30	45 (12) ESG and 43 (10) ESG with compression	5 women ESG and 18 ESG with compression	36.62 (3.72) ESG and 36.42 (3.27) ESG with compression*	NR
Graus Morales, J., et al. [49]	Case series	Single center	Prospective	148	ESG	-	ESG with modified stitching pattern	BMI > 30	41.53 (10)	121 women	35.11 (5.5)	NR
Gudur, A. R., et al. [50]	Cohort	Multicenter (database)	Retrospective	36,323	ESG (6,053)	LSG (30,270)	MBSAQIP database cohort study	-	47.47 (11.44) and 44.87 (11.94)	5116 and 24926 women	40.54 (8.65) and 42.8 (6.17)	1040 and 5775
Hajifathalian, K., et al. [51]	Case series	Single center	Prospective	118	ESG	-	Focus on NAFLD scores	BMI > 30 and NAFLD	46 (13)	80 (68%) women	40 (7)	35
Hill, C., et al. [52]	Case series	Single center	Prospective	21	ESG	-	Focus on learning curve	NR	47.7 (11.2)	13 women	41.8 (8.5)	NR
Jagtap, N., et al. [53]	Case series	Single center	Prospective	26	ESG	-	Focus on NAFLD scores	BMI > 27.5 and NAFLD	41.5 (9.58)	16 women	36.55 (5.07)	13
James, T. W., et al. [54]	Case series	Single center	Retrospective	100	ESG	-	Non-academic setting	NR	45 (9)	86 women	38.41 (5.44)	4

**Table 1** (continued)

Author (year)	Study design	Single vs. multicenter	Retrospective vs. prospective	Population (total sample)	Intervention (n)	Comparison (n)	Observations	Inclusion criteria (BMI)	Age	Sex	Mean baseline BMI (kg/m <sup>2</sup> )	Type II Diabetes
Kumar, N., et al. [55]	Case series	Multicenter (NS)	Prospective	122	ESG (23 in phase 1, 22 in phase 2, 77 in phase 3)	-	Focus on technical refinement over time	BMI > 30	Phase 1 37.7 (1.9); Phase 2 39.2 (1.6); Phase 3 41.3 (1.1)	Phase 1 19 women; phase 2 20 women; phase 3 59 women	Phase 1 34.2 (1.1); Phase 2 34.3 (1.1); Phase 3 36.1 (0.6)	NR
Li, R., et al. [56]	Case series	Single center	Prospective	24	ESG	-	Focus on high-risk cases	BMI > 50, severe comorbidities, or impenetrable abdomen	55.6 (9.2)	6 women	49.9 (14.4)	15
Lopez-Nava, G., et al. [57]	Case series	Single center	Retrospective	435	ESG	-		BMI > 30	48.5 (10.2)	314 women	38.9 (5.3)	NR
Lopez-Nava, G., et al. [58]	Cohort	2 centers	Prospective	24	ESG (12)	LSG (12)	Focus on enterohormonal changes	BMI > 30	49.3 (2.4) ESG and 50.5 (1.9) LSG	9 women ESG and 9 women LSG	38.3 (1.8) ESG and 39.2 (1.5) LSG	0 ESG and 3 preDM LSG
Lopez-Nava, G., et al. [59]	Case series	3 centers	Prospective	248	ESG	-	-	NR	44.5 (10)	181 women	37.8 (5.6)	NR
Manos, T., et al. [60]	Case series	Single center	Retrospective	191	ESG	-	Single-channel endoscope device (Overstitch SX)	BMI > 30	36.9 (no SD)	173 women	33.7 (4.18)*	NR
Matteo, M. V., et al. [61]	Case series	Single center	Prospective	18	ESG	-	Patients > 65 years old	BMI > 30	67 (4.5)	10 women	41.2 (5.9)	4
Maydeo, A., et al. [16]	Case series	Single center	Prospective	58	ESG	-	Different stitching pattern ("accordion")	BMI > 28	42.1 (8.7)	55 women	37.88 (5.76)	17
Mehta, A., et al. [62]	Case series	Single center	Prospective	50	ESG	-	Focus on quality of life and mental health	NR	49.5 (14)	37 women	38.5 (5.8)	9
Neto, M. G., et al. [63]	Case series	4 centers	Prospective	233	ESG	-	-	BMI > 30 and < 40	41.1 (10.5)	170 women	34.7 (2.6)	12
Neto, M. G., et al. [64]	Case series	Multicenter (NS)	Retrospective	1828	ESG	-	Clinical consensus gathering 47 Brazilian endoscopists	NR	NR	NR	NR	NR
Novikov, A. A., et al. [65]	Cohort	Single center	Retrospective	278	ESG (91)	LSG (120) and LAGB (67)	-	BMI > 30	43.86 (11.26) ESG, 40.71 (11.95) LSG, and 41.94 (13.31) LAGB	62 women ESG, 94 women LSG, and 54 women LAGB	38.61 (6.98) ESG, 47.22 (7.84) LG, and 44.98 (6.45) LAGB	20 ESG, 31 LSG, and 15 LAGB
Pizzicannella, M., et al. [66]	Case series	Single center	Prospective	133	ESG	-	Focus on durability of sutures and their correlation with weight loss	NR	NR	NR	43.2 (8.6)	NR

**Table 1** (continued)

Author (year)	Study design	Single vs. multicenter	Retrospective vs. prospective	Population (total sample)	Intervention (n)	Comparison (n)	Observations	Inclusion criteria (BMI)	Age	Sex	Mean baseline BMI (kg/m <sup>2</sup> )	Type II Diabetes
Rapakka, B., et al. [67]	Cohort	2 centers	Prospective	41	ESG (23)	IGB (18)	-	NR	47.69 (5.06) ESG and 41.06 (8.81) IGB	20 women ESG and 18 IGB	41.21 (5.38) ESG and 34.5 (4.46)	NR
Sarkar, A., et al. [68]	Case series	6 centers	Retrospective	91	ESG	-	Focus on new bariatric endoscopy programs	BMI > 30	39.7 (11.6)	56 women	38.7 (4.4)*	46
Sartoretto, A., et al. [69]	Case series	3 centers	Retrospective	112	ESG	-	-	BMI > 27	45.1 (11.7)	77 women	37.9 (6.7)	14
Saumoy, M., et al. [70]	Case series	Single center	Prospective	128	ESG	-	Focus on learning curve	BMI > 30	43.62 (11.37)	86 women	38.92 (6.95)	NR
Sharaiha, R. Z., et al. [71]	Case series	Single center	Retrospective	216	ESG	-	Long-term follow-up	BMI > 30 or > 27 with comorbidities	46 (13)	146 women	39 (6)	67

\*Calculated field

- ESG performed with the Apollo Overstitch device (no restriction as to stitching pattern);
- No language restriction;
- Full-text articles only;
- Study designs case series with sample  $\geq 10$ , cohort studies, case–control studies, and randomized trials. For the non-comparative meta-analysis, we extracted results from the ESG cohort from comparative studies;
- To avoid overestimating the real sample, only the most recent or the most representative (larger sample) study was considered for each center if repeated data was suspected;
- Studies describing outcomes at predetermined time points: 6, 12, 18, 24, 36, > 36 months;
- Studies reporting efficacy and/or safety data.

The initial search retrieved 3015 records. After screening titles and abstracts, 100 articles were selected for full-text assessment. Finally, 44 articles were included in the qualitative and quantitative analyses. Figure 2 shows the screening and inclusion/exclusion flowchart.

## Descriptive Analysis

### Baseline and Demographic Data

Among the 44 articles, we identified 29 case series, 14 cohort studies, and 1 randomized controlled trial (RCT). Among the cohort studies, 7 compared ESG to LSG, 1 compared ESG to lifestyle intervention alone, 1 compared ESG versus ESG plus anti-obesity medication (liraglutide), 2 compared ESG and intragastric balloons, and 3 compared ESG cohorts with different stitching patterns. Table 1 summarizes the overall and baseline data of the qualitative analysis of the included studies.

Among the 44 articles, the total sample included 49,848 patients (15,714 ESG and 34,134 controls, including LSG, laparoscopic adjustable gastric banding, and IGB). At baseline, the mean age and BMI were 44.24 (SE 1.405, 95%CI 41.48–46–99, 41 articles  $n = 13,562$ ) and 37.56 (SE 0.45 95%CI 36.66–38.46, 42 articles,  $n = 13,876$ ), respectively. Most patients were female (11,449 females, 83.2% and 2304 males, 16.8%, 42 articles,  $n = 13,753$ ).

**Risk of Bias/Critical Appraisal Assessment** All included studies were assessed for their risk of bias using specific tools based on the study design. Case series were evaluated using the Joanna Briggs Institute Critical Appraisal Checklist. Ten items are scored based on the perceived risk, and the scoring is positive. The scale ranges from 0 to 10, with 0 being the highest risk of bias and 10 being the lowest. The included case series (29 articles) had a mean score of  $7.5 \pm 1.8$ . Reporting of outcomes, and follow-up, and statistical analyses were the two topics with the worst positive scoring (16/29, 55.2%).

**Table 2** Critical appraisal and risk of bias assessment for the included case series

CASE SERIES		JBI Critical appraisal checklist domains											
Author, year	Study design	Were there clear criteria for inclusion in the case series	Was the condition measured in a standard, reliable way for all participants included in the case series?	Were valid methods used for identification of the condition for all participants included in the case series?	Did the case series include consecutive participants?	Did the case series have complete inclusion of participants?	Was there clear reporting of the demographics of the participants in the study?	Was there transparent reporting of clinical information of the participants?	Were the outcomes or follow-up results of cases clearly reported?	Was there transparent reporting of the demographic information of the pre-senting site(s)/ clinic(s)?	Was statistical analysis appropriate?	Obs.	Total of "yes" (max=10)
Articles (n) = 29	Abu Dayyeh, B. K., et al. [31]	Case series	yes	yes	yes	yes	yes	yes	no	yes	no		8
	Alqahtani, A., et al. [20]	Case series	yes	yes	yes	yes	yes	yes	no	yes	no		8
	Alqahtani, A., et al. [32]	Case series	yes	yes	yes	yes	yes	yes	yes	yes	no	Pediatric population	9
	Asokumar, R., et al. [34]	Case series	yes	yes	yes	yes	yes	yes	no	yes	no		8
	Barrichello, S., et al. [36]	Case series	yes	yes	yes	yes	yes	yes	no	yes	no		8
	Bhandari, M., et al. [37]	Case series	yes	yes	yes	yes	yes	yes	no	yes	no		8
	Callahan, Z. M., et al. [38]	Case series	yes	yes	yes	yes	yes	yes	no	yes	no		8
	Espinet-Coll, E., et al. [41]	Case series	yes	yes	yes	yes	yes	yes	no	yes	no		7
	Ghoz, H., et al. [47]	Case series	yes	yes	yes	yes	yes	yes	no	no	no	no	6

## CASE SERIES

	Case series	yes	yes	yes	yes	yes	yes	yes	no	yes	no	8
Graus Morales, J., et al. [17, 49]												
Hajifathal- ian, K., et al. [51]	Case series	yes	yes	yes	yes	yes	yes	yes	no	yes	no	8
Hill, C., et al. [52]	Case series	yes	yes	yes	yes	yes	yes	yes	no	yes	no	Only for adverse events, no effi- cacy data
Jagtap, N., et al. [53]	Case series	yes	yes	yes	yes	yes	yes	yes	no	yes	yes	9
James, T. W., et al. [54]	Case series	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	10
Kumar, N., et al. [55]	Case series	yes	unclear	unclear	no	unclear	no	no	yes	no	yes	3
Li, R., et al. [56]	Case series	yes	yes	yes	no	unclear	yes	yes	yes	yes	yes	8
Lopez- Nava, G., et al. [57]	Case series	yes	yes	yes	unclear	no	yes	no	yes	yes	yes	7
Lopez- Nava, G., et al. [59]	Case series	yes	yes	yes	yes	unclear	yes	unclear	yes	yes	yes	8
Manos, T., et al. [60]	Case series	yes	yes	yes	unclear	unclear	yes	no	yes	no	yes	6
Matteo, M. V., et al. [61]	Case series	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	10
Maydeo, A., et al. [16]	Case series	yes	yes	unclear	unclear	unclear	yes	yes	unclear	yes	yes	7
Mehta, A., et al. [62]	Case series	unclear	unclear	unclear	unclear	unclear	yes	yes	yes	yes	yes	6

 Springer

CASE SERIES	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100	101	102	103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120	121	122	123	124	125	126	127	128	129	130	131	132	133	134	135	136	137	138	139	140	141	142	143	144	145	146	147	148	149	150	151	152	153	154	155	156	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	295	296	297	298	299	300	301	302	303	304	305	306	307	308	309	310	311	312	313	314	315	316	317	318	319	320	321	322	323	324	325	326	327	328	329	330	331	332	333	334	335	336	337	338	339	340	341	342	343	344	345	346	347	348	349	350	351	352	353	354	355	356	357	358	359	360	361	362	363	364	365	366	367	368	369	370	371	372	373	374	375	376	377	378	379	380	381	382	383	384	385	386	387	388	389	390	391	392	393	394	395	396	397	398	399	400	401	402	403	404	405	406	407	408	409	410	411	412	413	414	415	416	417	418	419	420	421	422	423	424	425	426	427	428	429	430	431	432	433	434	435	436	437	438	439	440	441	442	443	444	445	446	447	448	449	450	451	452	453	454	455	456	457	458	459	460	461	462	463	464	465	466	467	468	469	470	471	472	473	474	475	476	477	478	479	480	481	482	483	484	485	486	487	488	489	490	491	492	493	494	495	496	497	498	499	500	501	502	503	504	505	506	507	508	509	510	511	512	513	514	515	516	517	518	519	520	521	522	523	524	525	526	527	528	529	530	531	532	533	534	535	536	537	538	539	540	541	542	543	544	545	546	547	548	549	550	551	552	553	554	555	556	557	558	559	560	561	562	563	564	565	566	567	568	569	570	571	572	573	574	575	576	577	578	579	580	581	582	583	584	585	586	587	588	589	590	591	592	593	594	595	596	597	598	599	600	601	602	603	604	605	606	607	608	609	610	611	612	613	614	615	616	617	618	619	620	621	622	623	624	625	626	627	628	629	630	631	632	633	634	635	636	637	638	639	640	641	642	643	644	645	646	647	648	649	650	651	652	653	654	655	656	657	658	659	660	661	662	663	664	665	666	667	668	669	670	671	672	673	674	675	676	677	678	679	680	681	682	683	684	685	686	687	688	689	690	691	692	693	694	695	696	697	698	699	700	701	702	703	704	705	706	707	708	709	710	711	712	713	714	715	716	717	718	719	720	721	722	723	724	725	726	727	728	729	730	731	732	733	734	735	736	737	738	739	740	741	742	743	744	745	746	747	748	749	750	751	752	753	754	755	756	757	758	759	760	761	762	763	764	765	766	767	768	769	770	771	772	773	774	775	776	777	778	779	780	781	782	783	784	785	786	787	788	789	790	791	792	793	794	795	796	797	798	799	800	801	802	803	804	805	806	807	808	809	810	811	812	813	814	815	816	817	818	819	820	821	822	823	824	825	826	827	828	829	830	831	832	833	834	835	836	837	838	839	840	841	842	843	844	845	846	847	848	849	850	851	852	853	854	855	856	857	858	859	860	861	862	863	864	865	866	867	868	869	870	871	872	873	874	875	876	877	878	879	880	881	882	883	884	885	886	887	888	889	890	891	892	893	894	895	896	897	898	899	900	901	902	903	904	905	906	907	908	909	910	911	912	913	914	915	916	917	918	919	920	921	922	923	924	925	926	927	928	929	930	931	932	933	934	935	936	937	938	939	940	941	942	943	944	945	946	947	948	949	950	951	952	953	954	955	956	957	958	959	960	961	962	963	964	965	966	967	968	969	970	971	972	973	974	975	976	977	978	979	980	981	982	983	984	985	986	987	988	989	990	991	992	993	994	995	996	997	998	999	1000
Neto, M. G., et al. [63]	Case series	yes	yes	yes	yes	unclear	unclear	unclear	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																								

**Table 3** Critical appraisal and risk of bias assessment for the included cohort studies

COHORT STUDIES										
Author, year	Study design	New Castle-Ottawa Scale for cohort studies domains								
		Representativeness of the exposed cohort (1)	Selection of the non-exposed cohort (1)	Ascertainment of exposure (1)	Demonstration that outcome of interest was not present at start of study (1)	Comparability of cohorts based on the design or analysis (2)	Assessment of outcome (1)	Was follow-up long enough for outcomes to occur (1)	Adequacy of follow-up of cohorts (1)	Total score (max = 9)
Alqahtani, A. R., et al. [33]	Cohort	1	0	0	1	1	1	0	1	5
Badurdeen, D. et al. [35]	Cohort	1	0	0	1	0	1	0	1	4
Carr, P., et al. [39]	Cohort	1	0	0	1	1	1	0	1	5
Cheskin, L. J., et al. [40]	Cohort	1	0	1	1	1	1	0	1	6
Espinet-Coll, E., et al. [41]	Cohort	1	0	1	1	1	1	0	1	6
Farha, J., et al. [43]	Cohort	1	1	0	1	1	1	1	1	7
Fayad, L., et al. [44]	Cohort	1	1	0	1	1	1	0	1	6
Fayad, L., et al. [45]	Cohort	1	0	0	1	1	1	0	1	5
Fiorillo, C., et al. [46]	Cohort	1	0	1	1	1	1	0	1	6
Glaysner, M. A., et al. [48]	Cohort	1	1	1	1	1	1	1	1	9
Gudur, A. R., et al. [50]	Cohort	1	1	1	1	1	1	1	1	9
Lopez-Nava, G., et al. [58]	Cohort	1	1	1	1	1	1	1	1	9
Novikov, A. A., et al. [65]	Cohort	1	1	1	1	0	1	1	0	6
Rapaka, B., et al. [67]	Cohort	1	1	1	1	0	1	1	0	6
<b>Summary</b>		<b>1 (0)</b>	<b>0.42 (0.51)</b>	<b>0.5 (0.51)</b>	<b>1 (0)</b>	<b>0.78 (0.42)</b>	<b>1 (0)</b>	<b>0.35 (0.49)</b>	<b>0.85 (0.36)</b>	<b>6.07 (1.43)</b>

**Table 4** Critical appraisal and risk of bias assessment for the included randomized clinical trial

Author, year	Study design	Cochrane risk of bias tool				JADAD score					Withdrawals	Total
		Selection bias:		Reporting bias:	Other Bias -	Performance bias	Detection bias	Attrition bias	Randomization	Blinding		
		Random Sequence generation	Allocation Concealment									
Abu Dayyeh, B. K., et al. [22]	Randomized clinical trial	Low	Low	Low	Low	High	High	High	2	0	1	3

For cohort studies, we employed the New-Castle Ottawa scale that assesses 8 topics for bias. The scale ranges from 0 to 9, with 0 being the highest risk of bias and 9 being the lowest. The 14 included cohort articles scored an average of  $6.07 \pm 1.43$ . The selection of a non-exposed cohort and the duration of follow-up were the two topics with the worst scoring, thus most subject to bias ( $0.42 \pm 0.51$  and  $0.35 \pm 0.49$ ).

For the single RCT [22], the JADAD score was 3, which is the maximum score for open-label trials. As to the modified Cochrane risk of bias tool, the trial was at low risk for selection and reporting bias. However, we detected a high risk of other biases: performance, detection, and attrition. The GRADE assessment of the quality of evidence later weighted the impact of those biases. Tables 2, 3, and 4 summarize the assessment of biases for case series, cohorts, and RCT, respectively.

**Meta-analysis** Forty-two articles reported %Excess Weight Loss (%EWL) and/or %Total Body Weight Loss (%TBWL) at least in one time point of interest (6, 12, 18, 24, 36, > 36 months). Two articles [50, 64] only reported safety outcomes. Continuous variables (%EWL and %TBWL) were pooled using the CMA software, and the results are presented ahead of time according to time points. Categorical variables were pooled using absolute numbers to calculate pooled rates. Four articles reported the responder rate as  $\geq 5\%$  TWL at 12 months, and 9 reported it as  $\geq 10\%$  TBWL. The pooled rates were 422/478 (88.3%) and 632/768 (82.3%). Forty articles reported the SAEs rate (according to the FDA definition from <https://www.fda.gov/safety/reporting-serious-problems-fda/what-serious-adverse-event>). Among 15,398 ESG procedures, 194 events fulfilled the criteria for SAE for a pooled rate of 1.25%. Table 5 shows all outcomes of the included studies according to follow-up time points, and Fig. 2 graphically depicts weight loss outcomes over time.

### Summary of Weight Loss Outcomes After ESG (Table 6 and Fig. 3)

### Quality of Evidence Assessment

All pooled outcomes were assessed for the quality of evidence according to the GRADE methodology. Since this analysis included only non-comparative data, all endpoints were rated as VERY LOW quality of evidence. Table 7 depicts the GRADE assessment.

**Table 5** Outcomes of the studies included in the non-comparative meta-analysis

Author (year)	Population (total sample)	Intervention (n)	Comparison (n)	%EWL	n	6 months	n	12 months	n	18 months	n	24 months	n	≥36 months	%TBWL
Abu Dayyeh, B. K., et al. [31]	25	ESG	—	—	25	53 (17)	10	54 (40)	8	45 (41)	—	—	—	—	—
Abu Dayyeh, B. K., et al. [22]	209	ESG (85)	Lifestyle intervention (124)	—	—	—	77	49.2 (32)	—	—	50	41 (32)	—	—	—
Alqahtani, A., et al. [20]	1000	ESG	—	—	369	64.3 (56.2)	216	67.5 (52.3)	54	64.7 (55.4)	—	—	—	—	—
Alqahtani, A., et al. [32]	109	ESG	—	—	82	80.1 (63.3)	43	87.1 (59.5)	24	70.9 (55.5)	17	63.8 (52.3)	—	—	—
Alqahtani, A. R., et al. [33]	6036	ESG (3018)	LSG (3018)	—	2490	67.0 (28.6)	2243	77.1 (24.6)	—	—	1911	75.2 (47.9)	854	59.7 (57.1)	—
Asokkumar, R., et al. [34]	35	ESG	—	—	—	—	—	—	—	—	—	—	—	—	—
Badurdeen, D., et al. [35]	52	ESG (26)	ESG + Liraglutide (26)	—	26	69.94 (6.3)	—	—	—	—	—	—	—	—	—
Barrichello, S., et al. [36]	193	ESG	—	—	181	56.15 (22.93)	121	59.41 (25.69)	—	—	—	—	—	—	—
Bhandari, M., et al. [37]	53	ESG	—	—	—	—	—	—	—	—	—	—	—	—	—
Callahan, Z. M., et al. [38]	10	ESG	—	—	—	—	3	17.6 (47.3)	—	—	4	12.7 (16.9)	—	—	—
Carr, P., et al. [39]	61	ESG (16)	LSG (45)	—	13	51 (11)	9	57 (32)	—	—	—	—	—	—	—
Cheskin, L. J., et al. [40]	386	ESG (105)	Lifestyle intervention (281)	—	—	—	—	—	—	—	—	—	—	—	—

**Table 5** (continued)

Author (year)	Population (total sample)	Intervention (n)	Comparison (n)	%EWL	n	6 months	n	12 months	n	18 months	n	24 months	n	≥36 months	%TBWL
Espinete–Coll, E., et al. [41]	38	ESG	–	–	–	–	–	48.3 (18.5)*	–	–	–	–	–	–	–
Espinete–Coll, E., et al. [18]	88	ESG (standard stitching pattern = "TBp")	ESG with different stitching patterns ("Lp" and "TMp")	–	–	–	88	46.41 (20.6)	–	–	–	–	–	–	–
Farha, J., et al. [43]	247	ESG (98)	ESG with fundal suturing (149)	82	54.7 (19.2)	57	65.3 (21.1)	–	–	–	–	–	–	–	–
Fayad, L., et al. [44]	137	ESG (54)	LSG (83)	99	37.7 (17.3)	66	40.6 (23.5)	–	–	–	–	–	–	–	–
Fayad, L., et al. [45]	105	ESG (58)	IGB (47)	–	–	–	–	–	–	–	–	–	–	–	–
Fiorillo, C., et al. [46]	46	ESG (23)	LSG (23)	23	39.05 (20.7)*	–	–	–	–	–	–	–	–	–	–
Choz, H., et al. [47]	20	ESG	–	–	–	–	–	–	–	–	–	–	–	–	–
Glaysheer, M. A., et al. [48]	32	ESG without longitudinal compression (9)	ESG with longitudinal compression (23)	5	42.4 (18.1)*	–	–	–	–	–	–	–	–	–	–
				7	65.6 (23.9)*	–	–	–	–	–	–	–	–	–	–
Graus Morales, J., et al. [17, 49]	148	ESG	–	148	64.93 (51)	148	75.4 (85)	72	79.25 (43)	–	–	–	–	–	–
Gudur, A. R., et al. [50]	36,323	ESG (6,053)	LSG (30,270)	–	–	–	–	–	–	–	–	–	–	–	–
Hajifathalian, K., et al. [51]	118	ESG	–	114	45.3 (29.39)*	100	47.8 (32.65)*	–	–	–	78	45.5 (33.11)*	–	–	–
Hill, C., et al. [52]	21	ESG	–	–	–	–	–	–	–	–	–	–	–	–	–

**Table 5** (continued)

Author (year)	Population (total sample)	Intervention (n)	Comparison (n)	%EWL	n	6 months	n	12 months	n	18 months	n	24 months	n	≥36 months	%TBWL
Jagtap, N., et al. [53]	26	ESG	—	—	26	32.67 (19.51)	26	51.33 (17.33)	—	—	—	—	—	—	—
James, T. W., et al. [54]	100	ESG	—	—	34	48.9 (19.9)	12	66.1 (21.5)	—	—	—	—	—	—	—
Kumar, N., et al. [55]	122	ESG (23 in phase 1, 22 in phase 2, 77 in phase 3)	—	—	—	—	—	—	—	—	—	—	—	—	—
Li, R., et al. [56]	24	ESG	—	—	12	25 (9.1)	7	29.1 (17.9)	—	—	—	—	—	—	—
Lopez-Nava, G., et al. [57]	435	ESG	—	—	—	—	—	—	—	—	—	—	—	—	—
Lopez-Nava, G., et al. [58]	24	ESG (12)	LSG (12)	—	—	—	—	—	—	—	—	—	—	—	—
Lopez-Nava, G., et al. [59]	248	ESG	—	—	—	—	—	—	—	—	—	—	—	—	—
Manos, T., et al. [60]	191	ESG	—	—	84	41.6 (20**)	69	34.7 (22**)	—	—	—	—	—	—	—
Matteo, M. V., et al. [61]	18	ESG	—	—	18	39.25 (5.5)*	12	38.25 (10.97)*	10	40.25 (13.26)*	10	41 (8.08)*	—	—	—
Maydeo, A., et al. [16]	58	ESG	—	—	52	42.8 (13.1)	—	—	—	—	—	—	—	—	—
Mehia, A., et al. [62]	50	ESG	—	—	47	46.9 (22.4)	39	50.5 (24.9)	47	47.7 (26.5)	—	—	—	—	—
Neto, M. G., et al. [63]	233	ESG	—	—	178	47.1 (18)	123	54.8 (17.4)	—	—	—	—	—	—	—
Neto, M. G., et al. [64]	1828	ESG	—	—	—	—	—	—	—	—	—	—	—	—	—
Novikov, A. A., et al. [65]	278	ESG (91)	LSG (120) and LAGB (67)	—	—	—	—	—	—	—	—	—	—	—	—
Pizzicannella, M., et al. [66]	133	ESG	—	—	87	34.5 (19.8)	41	34.3 (21.9)	—	—	—	—	—	—	—
Rapaka, B., et al. [67]	41	ESG (23)	IGB (18)	—	23**	16.17 (5.69)	—	—	—	—	—	—	—	—	—

Table 5 (continued)

Author (year)	Popula- tion (total sample)	Intervention (n)	Comparison (n)	%EWL	n	6 months	n	12 months	n	18 months	n	24 months	n	≥36 months	%TBWL
Sarkar, A., et al. [68]	91	ESG	—	—	52	35.6 (20**)	—	—	—	—	—	—	—	—	—
Sartoretto, A., et al. [69]	112	ESG	—	—	52	50.3 (22.4)	—	—	—	—	—	—	—	—	—
Saumoy, M., et al. [70]	128	ESG	—	—	—	—	—	—	—	—	—	—	—	—	—
Sharaiha, R. Z., et al. [71]	216	ESG	—	—	—	—	142	47.9 (33.11)*	—	—	—	—	36 months = 68 60 months = 56	45.1 (42.67)* 45.3 (47.32)*	—
<b>Total</b>	<b>49848</b>	<b>—</b>	<b>—</b>	<b>—</b>	<b>4329</b>	<b>48.04 (SE 3.59)</b>	<b>3652</b>	<b>53.09 (SE 4.15)</b>	<b>215</b>	<b>57.98 (SE 7.38)</b>	<b>2070</b>	<b>46.57 (SE 9.85)</b>	<b>36 months = 922</b>	<b>36 months = 53.18 (SE 7.25)</b>	<b>—</b>
Author (year)	n	6 months	n	12 months	n	18 months	n	24 months	n	≥36 months	12-month Responder rate (≥5%TBWL)	12-month Responder rate (≥10%TBWL)	SAEs	Obs.	
Abu Dayyeh, B. K., et al. [31]	—	—	—	—	—	—	—	—	—	—	—	—	3/25	—	
Abu Dayyeh, B. K., et al. [22]	—	—	77	13.6 (8.0)	—	—	—	50	11.4 (8.4)	—	70/77	48/77	6/150	—	
Alqahtani, A., et al. [20]	369	13.7 (6.8)	216	15 (7.7)	54	14.8 (8.5)	—	—	—	—	193/216	—	24/1000	—	
Alqahtani, A., et al. [32]	82	14.4 (6.5)	43	16.2 (8.3)	24	15.4 (9.2)	17	13.7(8)	—	—	—	—	1/109	—	
Alqahtani, A. R., et al. [33]	2490	15.1 (6.1)	2243	19.2 (7.7)	—	—	1911	16.2 (9.7)	854	14 (12.1)	—	—	14/3018	—	
Asokkumar, R., et al. [34]	10	16.2 (4.9)	—	—	—	—	—	—	—	—	—	—	0/35	—	
Badurdeen, D. et al. [35]	26	20.51 (1.68)	—	—	—	—	—	—	—	—	—	—	1/52	—	

Table 5 (continued)

Author (year)	n	6 months	n	12 months	n	18 months	n	24 months	n	≥36 months	12-month Responder rate (≥5%TBWL)	12-month Responder rate (≥10%TBWL)	SAEs	Obs.
Barrichello, S., et al. [36]	181	14.25 (5.26)	121	15.06 (5.22)	—	—	—	—	—	—	—	—	2/193	
Bhandari, M., et al. [37]	42	14.25 (6.17)	42	19.94 (4.89)	—	—	—	—	—	—	—	—	0	37/42 (88%) had >15% TBWL at 12 months
Callahan, Z. M., et al. [38]	—	—	—	—	—	—	—	—	—	—	—	—	2/10	
Carr, P., et al. [39]	13	15 (6)	9	18 (11)	—	—	—	—	—	—	—	—	0	ESG 78% >25% EWL and 55.6% >20%TBWL at 12months
Cheskin, L. J., et al. [40]	63	17.7 (6.4)	43	20.6 (8.3)	—	—	—	—	—	—	41/43	39/43	5/105	28/105 pre-sented >20% TBWL
Espinete-Coll, E., et al. [41]	—	—	—	17.1 (3.1)*	—	—	—	—	—	—	—	36/38	1/38	
Espinete-Coll, E., et al. [18]	—	—	88	17.36 (6.09)	—	—	—	—	—	—	—	84/88	0/88	
Farha, J., et al. [43]	82 99	17.3 (4.5) 16.2 (7.0)	57 66	21.3 (6.2) 17.5 (10.2)	—	—	—	—	—	—	—	—	6/247	
Fayad, L., et al. [44]	35	17.1 (6.5)	—	—	—	—	—	—	—	—	—	—	3/54	39/54 had >15% TBWL 72.2% at 6 months
Fayad, L., et al. [45]	25	19.5 (5.7)	21	21.3 (6.6)	—	—	—	—	—	—	—	—	3/58	
Fiorillo, C., et al. [46]	23	13.87 (6.55)*	—	—	—	—	—	—	—	—	—	—	0/23	

Table 5 (continued)

Author (year)	n	6 months	n	12 months	n	18 months	n	24 months	n	≥36 months	12-month Responder rate (≥5%TBWL)	12-month Responder rate (≥10%TBWL)	SAEs	Obs.
Ghoz, H., et al. [47]	13	13.7 (9.3)	10	16.2 (10.4)	—	—	—	—	—	—	—	—	NR	
Glaysner, M. A., et al. [48]	57	12.4 (3.1)* 20.5 (5)*	—	—	—	—	—	—	—	—	—	—	0/32	
Graus Morales, J., et al. [17, 49]	148	15.45 (5.9)	148	17.53 (7.57)	72	18.66 (7.3)	—	—	—	—	—	—	2/148	
Gudur, A. R., et al. [50]	—	—	—	—	—	—	—	—	—	—	—	—	86/6053	
Hajifathalian, K., et al. [51]	114	14.6 (7.07)*	100	15.6 (8.92)*	—	—	78	15.5 (10.13)*	—	—	—	75/100	0/118	60 (74%) >10% TBWL at 2 years; improvement in ALT, AST, HSI, NAFLD fib score, HbA1c, HOMA-IR and Leptin
Hill, C., et al. [52]	—	—	—	—	—	—	—	—	—	—	—	—	1/31	Learning curve plateau at 7 cases
Jagtap, N., et al. [53]	26	11.33 (4.99)	26	18.07 (3.35)	—	—	—	—	—	—	—	—	0/26	23/26 (88.4%) >15%TBWL at 12 months; improvement in ALT, HSI, NAFLD fib score, FIB-4, and APRI at 12 months
James, T. W., et al. [54]	34	16.41 (5.4)	12	23.1 (7.5)	—	—	—	—	—	—	—	12/12	2/100	

Table 5 (continued)

Author (year)	n	6 months	n	12 months	n	18 months	n	24 months	n	≥36 months	12-month Responder rate (≥5%TBWL)	12-month Responder rate (≥10%TBWL)	SAEs	Obs.
Kumar, N., et al. (2018).	Phase 2 NO SAM- PLE 12	17.3 (1.7) 16 (0.8)	Phase 2 Phase 3 Phase 3 (20) (44)	17.3 (2.6) 17.6 (2.1)	–	–	–	–	–	–	–	–	0/122	
Li, R., et al. [56]	12	11.3 (4.7)	7	12.2 (8.9)	–	–	–	–	–	–	–	–	1/24	
Lopez–Nava, G., et al. [57]	Class I = 99 Class II = 77 Class III = 84 146	14.9 (6.5) 16.8 (6.3) 16.8 (6.3)	Class I = 50 Class II = 77 Class III = 84 146	17.1 (6.7) 19 (8) 22.2 (9.3)	–	–	–	–	–	–	–	42/50 67/77 78/84	6/435	
Lopez–Nava, G., et al. [59]	12	13.3 (7)	–	–	–	–	–	–	–	–	–	–	0/12	
Lopez–Nava, G., et al. [59]	215	15.17 (7.66)*	–	–	57	18.6 (11.15)*	–	–	–	–	–	48/57	5/248	
Manos, T., et al. [61]	84	22.4 (8**)	69	18.7 (8**)	–	–	–	–	–	–	–	–	2/191	
Matteo, M. V., et al. [61]	18	14.97 (4.56)*	12	15.27 (5.25)*	10	16 (7.38)*	10	15.55 (6.93)*	–	–	–	–	0/18	
Maydeo, A., et al. [16]	52	17.1 (4.3)	–	–	–	–	–	–	–	–	–	–	0/58	
Mehta, A., et al. [62]	47	14.8 (5.3)	39	15.5 (7.2)	47	15.4 (9.8)	–	–	–	–	–	–	0/50	Significant improvement in depression scores

Table 5 (continued)

Author (year)	n	6 months	n	12 months	n	18 months	n	24 months	n	≥36 months	12-month Responder rate (≥5%TBWL)	12-month Responder rate (≥10%TBWL)	SAEs	Obs.
Neto, M. G., et al. [63]	178	16.9 (6.2)	123	19.7 (5.7)	—	—	—	—	—	—	—	—	0/233	
Neto, M. G., et al. [64]	—	—	—	—	—	—	—	—	—	—	—	—	15/1828	
Novikov, A. A., et al. [65]	61	14.37 (7)	28	17.57 (8.1)	—	—	—	—	—	—	—	—	1/91	
Pizzicannella, M., et al. [66]	87	13.2 (7.4)	41	13.1 (8.1)	—	—	—	—	—	—	—	—	—	Experience improved the proportion of intact ESG at 12 months
Rapaka, B., et al. [67]	—	—	—	—	—	—	—	—	—	—	—	—	—	Delay in T50 correlated with %TBWL at 3 months
Sarkar, A., et al. [68]	52	17.4 (6.5**)	—	—	—	—	—	—	—	—	—	—	—	
Sartoretto, A., et al. [69]	52	14.9 (6.1)	—	—	—	—	—	—	—	—	—	—	3/112	
Saumoy, M., et al. [70]	74	13.43 (7.4)	60	15.8 (9.5)	—	—	—	—	—	—	—	—	2/128	Learning curve: efficiency at 29 and mastery at 55 cases
Sharaitha, R. Z., et al. [71]	—	—	142	15.6 (9.11)*	—	—	—	—	36 months = 68	14.9 months (11.77)*	118/142	103/142	3/216	
									60 months = 56	15.9 months (16.79)*				
<b>Total</b>	<b>5227</b>	<b>15.66 (SE 0.35)</b>	<b>4118</b>	<b>17.56 (SE 0.39)</b>	<b>207</b>	<b>16.25 (SE 0.95)</b>	<b>2123</b>	<b>15.2 (SE 0.93)</b>	<b>36 months = 922</b>	<b>36 months = 14.07 (SE 0.39)</b>	<b>422/478 (88.3%)</b>	<b>632/768 (82.3%)</b>	<b>194/15,398 (1.25%)</b>	

\* Calculated fields

\*\* Inputted data

## Direct Comparative ESG Studies

### Systematic Review

Two independent researchers (VOB and NJ) ran separate literature searches assessing eligible studies. We searched MEDLINE (PubMed), EMBASE, and gray literature from January 1, 2013 (the year ESG was described), to October 1, 2022. The step-by-step construction of the search strategy is provided in SUPPL 3. The final strategy was as follows:

- MEDLINE (PubMed): *(excess weight) OR (overweight) OR (obesity) AND (endoscopy) OR (endoscopic) OR (transoral\*) OR (peroral\*) OR (incisionless) AND (sleeve) OR (overstitch) OR (gastroplasty) OR (gastric plication) OR (gastric imbrication) AND (lifestyle) OR (diet) OR (exercise) OR (counseling) OR (sham) OR (placebo)*
- EMBASE: *endoscopic AND sleeve AND gastroplasty OR (apollo AND overstitch) AND [embase]/lim NOT ([embase]/lim AND [medline]/lim) AND ('article'/it OR 'article in press'/it OR 'conference review'/it OR 'note'/it OR 'review'/it)*

The eligibility criteria included:

- Articles published online from 01/JAN/2013 until 01/OCT/2022 (last search update);
- ESG performed with the Apollo Overstitch device (no restriction as to stitching pattern);
- No language restriction;
- Full-text articles only;
- Comparative study designs: cohort studies, case–control studies, and randomized trials;
- Studies reporting efficacy and/or safety data.

The initial search retrieved 537 records. After screening titles and abstracts, 13 articles were selected for full-text assessment. Finally, only 2 articles were included in the qualitative and quantitative analyses. Figure 4 shows the screening and inclusion/exclusion flowchart.

**Descriptive Analysis** Cheskin et al. [40] and Abu Dayyeh et al. [22] were eligible studies for directly comparing ESG and lifestyle intervention. The first was a case-matched (1 ESG: 2–3 controls) cohort study comparing ESG plus low-intensity diet and lifestyle therapy (LIDLT) versus high-intensity diet and lifestyle therapy (HIDLT). This study included patients with obesity class 1 or higher. For both groups, patients paid out-of-pocket for the treatment (total cost ESG: US\$ 16,000; total cost

HIDLT: US\$ 3200). A total of 386 patients (105 ESG, 281 control) were included in the study, with similar baseline characteristics. The final follow-up visit was at 12 months.

Abu Dayyeh et al. [22] was a multicenter, US FDA-regulated, open-label, randomized trial comparing ESG plus lifestyle interventions to lifestyle intervention alone (MERIT Trial). Only patients with obesity classes I and II (BMI 30–40 kg/m<sup>2</sup>) were included and allocated to ESG or control group in a 1–1.5 ratio. After 52 weeks, compliant control patients crossed over to ESG. Two hundred and nine patients (85 ESG, 124 control) were enrolled and had similar baseline characteristics. The primary endpoints were %EWL and %TBWL at 12 months, but there was an extended follow-up at 24 months for the intervention group and a 12-month follow-up for the control group crossing over to intervention.

The baseline data and the critical appraisal/risk of bias assessment for the studies are summarized in Tables 1, 3, and 4.

**Meta-analysis** The two studies differ in design (cohort vs. RCT) and population (non-specified obesity vs. class I and II). According to the Cochrane Handbook [25], data from different study designs should not be pooled when few eligible studies exist. Therefore, we analyzed data from Cheskin et al. 2020 and Abu Dayyeh et al. 2022 separately. Since we could not pool data from different studies, heterogeneity, and sensitivity analyses do not apply.

#### A) Outcomes from MERIT Trial [22]

- At 12 months, the mean difference in weight loss outcomes compared to moderate-intensity lifestyle control was.

MD (%EWL): 46.00 [38.05–53.95, 95%CI] – Fig. 5

MD (%TBWL): 13.10 [11.08–15.12, 95%CI] – Fig. 6

SAE rate was 2% without mortality or need for intensive care or surgical intervention

The quality of evidence Abu Dayyeh et al. generated was MODERATE according to the GRADE methodology. Overall, data coming from a single study (imprecision) and the absence of double blinding were the two factors downgrading the quality of evidence. Table 8 summarizes the GRADE assessment.

#### B) Outcomes from Cheskin et al. (ADD REF here)


- At 12 months, the mean difference in weight loss outcomes compared to high-intensity lifestyle control [40].

**Table 6** Summary of weight loss outcomes after ESG

Time	Mean %EWL	Mean %TBWL
6 months	48.04% (SE 3.59, 95%CI 40.98–55.09, 24 articles, 4329 patients)	15.66% (SE 0.35, 95%CI 14.95–16.36, 33 articles, 5227 patients)
12 months	53.09% (SE 4.15, 95%CI 44.95–61.23, 21 articles, 3652 patients)	17.56% (SE 0.39, 95%CI 16.8–18.32, 27 articles, 4118 patients)
18 months	57.98% (SE 7.38, 95%CI 43.5–72.46, 6 articles, 215 patients)	16.25% (SE 0.95, 95%CI 14.38–18.13, 5 articles, 207 patients)
24 months	46.57% (SE 9.85, 95%CI 27.26–65.88, 6 articles, 2070 patients)	15.2% (SE 0.93, 95%CI 13.36–17.04, 6 articles, 2123 patients)
36 months	53.18% (SE 7.25, 2 articles, 922 patients)	14.07% (SE 0.39, 2 articles, 922 patients)
60 months	45.3 (SD 47.32, 1 article, 56 patients)	15.9 (SD 16.79, 1 article, 56 patients)

\*SE standard errors, CI confidence interval, EWL excess weight loss, TBWL total body weight loss

**Fig. 3** Figure displaying the mean percentage of excess weight loss (%EWL) and total body weight loss (%TBWL) over different time points following endoscopic sleeve gastroplasty (ESG) with standard error bars. The sample sizes and the number of articles at each time point are incorporated above the bars for clarity and additional context



MD (%TBWL): 6.3 [3.12–9.48, 95%CI] – Figure 7  
Adverse events rate in the ESG group was 4.8%, with no mortality, need for intensive care, or surgical intervention.

The quality of evidence Cheskin et al. generated was VERY LOW according to the GRADE methodology. Overall, data from a single study (imprecision) and a non-randomized study design (selection bias) led to the final quality of evidence. Table 9 summarizes the GRADE assessment.

## IFSO Bariatric Endoscopy Position Statement and Future Direction

Based on a comprehensive systematic review and meta-analysis, the IFSO Bariatric Endoscopy Committee endorses endoscopic sleeve gastroplasty (ESG) as an effective and

valuable intervention for managing obesity. ESG is particularly beneficial for patients with class I and II obesity, as well as for those with class III obesity who are not suitable candidates for traditional MBS. This minimally invasive procedure not only achieves significant weight loss outcomes in the short and mid-terms but also maintains a favorable safety profile, as evidenced by a low incidence of serious adverse events.


The systematic review encompassed numerous observational studies, which, despite being categorized as very low quality, consistently reported positive and similar outcomes from different cohorts and practice settings globally, indicating reproducibility, generalizability, and maturity of ESG. Additionally, including a single randomized controlled trial in the meta-analysis provided moderate quality evidence, further substantiating the efficacy and safety of ESG. This dual-source evidence base enhances the

**Table 7** GRADE assessment of the quality of evidence for the non-comparative meta-analysis


Certainty assessment							№ of patients		Effect		Certainty	Importance
№ of studies	Study design	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	ESG	no treatment	Relative (95% CI)	Absolute (95% CI)		
6-month %EWL												
24	observational studies	very serious	not serious	not serious	not serious	strong association	4329	-	-	-	⊕○○○○ Very low	IMPORTANT
6-month %TBWL												
33	observational studies	very serious	not serious	not serious	not serious	strong association	5227	-	-	-	⊕○○○○ Very low	IMPORTANT
12-month %EWL												
Certainty assessment							№ of patients		Effect		Certainty	Importance
№ of studies	Study design	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	ESG	no treatment	Relative (95% CI)	Absolute (95% CI)		
21	observational studies	very serious	not serious	not serious	not serious	strong association	3652	-	-	-	⊕○○○○ Very low	CRITICAL
12-month %TBWL												
27	observational studies	very serious	not serious	not serious	not serious	strong association	4118	-	-	-	⊕○○○○ Very low	CRITICAL
18-month %EWL												
6	observational studies	very serious <sup>a</sup>	not serious	not serious	not serious	strong association	215	-	-	-	⊕○○○○ Very low	IMPORTANT
18-month %TBWL												
5	observational studies	very serious <sup>a</sup>	not serious	not serious	not serious	strong association	207	-	-	-	⊕○○○○ Very low	IMPORTANT
24-month %EWL												
6	observational studies	very serious <sup>a</sup>	not serious	not serious	not serious	strong association	2070	-	-	-	⊕○○○○ Very low	CRITICAL
24-month %TBWL												
6	observational studies	very serious <sup>a</sup>	not serious	not serious	not serious	strong association	2123	-	-	-	⊕○○○○ Very low	CRITICAL
36-month %EWL												
2	observational studies	very serious <sup>a</sup>	not serious	not serious	not serious	strong association	922	-	-	-	⊕○○○○ Very low	IMPORTANT
36-month %TBWL												
2	observational studies	very serious <sup>a</sup>	not serious	not serious	not serious	strong association	922	-	-	-	⊕○○○○ Very low	IMPORTANT
60-month %EWL												
1	observational studies	very serious <sup>a</sup>	not serious	not serious	not serious	strong association	56	-	-	-	⊕○○○○ Very low	IMPORTANT
60-month %TBWL												
1	observational studies	very serious <sup>a</sup>	not serious	not serious	not serious	strong association	56	-	-	-	⊕○○○○ Very low	IMPORTANT
12-month 5%TBWL responder rate												
4	observational studies	very serious <sup>a</sup>	not serious	not serious	not serious	strong association	422/478 (88.3%)	-	-	-	⊕○○○○ Very low	NOT IMPORTANT
12-month 10%TBWL responder rate												
9	observational studies	very serious <sup>a</sup>	not serious	not serious	not serious	strong association	632/768 (82.3%)	-	-	-	⊕○○○○ Very low	CRITICAL
SAE rate												
40	observational studies	very serious <sup>a</sup>	not serious	not serious	not serious	none	194/15398 (1.3%)	-	-	-	⊕○○○○ Very low	CRITICAL

(a) Data from case series

**Fig. 4** PRISMA flowchart for the literature screening and inclusion/exclusion process for only comparing ESG and lifestyle intervention



Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PLoS Med* 6(7): e1000097. doi:10.1371/journal.pmed1000097




**Fig. 5** Forest plot for %EWL at 12 months in comparing ESG vs. lifestyle intervention for patients with class I and II obesity

robustness of the findings, lending more significant support to the recently published IFSO International Delphi Consensus statement on the position of ESG in the spectrum of obesity care [23].







It is important to emphasize the crucial role of integrating and complementing any obesity intervention, such as ESG, with a comprehensive and longitudinal healthy living program. This program should include a healthy diet, physical

activity, adequate sleep, and mindfulness to maintain the weight loss benefits and maximize the overall impact on health by the intervention. By incorporating ESG into a comprehensive program, healthcare providers can offer a broader spectrum of options for obesity management tailored to the needs and circumstances of individual patients. This integrated approach enhances the effectiveness of ESG but also ensures long-term health benefits for patients.




**Fig. 6** Forest plot for %TBWL at 12 months in comparing ESG vs. lifestyle intervention for patients with class I and II obesity

**Table 8** GRADE assessment of the quality of evidence for comparing ESG vs. lifestyle intervention for patients with mild and moderate obesity





Certainty assessment							№ of patients		Effect		Certainty	Importance
№ of studies	Study design	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	ESG plus lifestyle modification	Lifestyle modification alone	Relative (95% CI)	Absolute (95% CI)		
%EWL (follow-up: mean 12 months)												
1	randomized trials	serious <sup>a</sup>	not serious	not serious	serious <sup>b</sup>	strong association	77	110	-	MD <b>46 % higher</b> (38.05 higher to 53.95 higher)	 Moderate	CRITICAL
%TBWL (follow-up: mean 12 months)												
1	randomized trials	serious <sup>a</sup>	not serious	not serious	serious <sup>c</sup>	strong association	77	110	-	MD <b>13.1 % higher</b> (11.08 higher to 15.12 higher)	 Moderate	CRITICAL
25%EWL Responders Rate (follow-up: mean 12 months)												
1	randomized trials	serious <sup>a</sup>	not serious	not serious	serious <sup>d</sup>	strong association	59/77 (76.6%)	13/110 (11.8%)	<b>OR 24.46</b> (11.17 to 53.53)	<b>65 more per 100</b> (from 48 more to 76 more)	 Moderate	IMPORTANT
5%TBWL Responders Rate (follow-up: mean 12 months)												
1	randomized trials	serious <sup>a</sup>	not serious	not serious	serious <sup>d</sup>	strong association	70/77 (90.9%)	18/110 (16.4%)	<b>OR 51.11</b> (20.23 to 129.12)	<b>75 more per 100</b> (from 63 more to 80 more)	 Moderate	IMPORTANT
10%TBWL Responders Rate (follow-up: mean 12 months)												
1	randomized trials	serious <sup>a</sup>	not serious	not serious	serious <sup>d</sup>	strong association	48/77 (62.3%)	6/110 (5.5%)	<b>OR 28.69</b> (11.17 to 73.68)	<b>57 more per 100</b> (from 34 more to 76 more)	 Moderate	IMPORTANT
SAEs rate (follow-up: mean 12 months)												
1	randomized trials	serious <sup>a</sup>	not serious	not serious	serious <sup>d</sup>	strong association	9/77 (11.7%)	0/110 (0.0%)	<b>OR 30.65</b> (1.76 to 535.05)	<b>0 fewer per 100</b> (from 0 fewer to 0 fewer)	 Moderate	CRITICAL

CI confidence interval, MD mean difference, OR odds ratio. (a) Open-label trial (detection bias) and loss to follow-up rates (20% intervention, 29% control group). Attrition bias present. (b) Single RCT with 77 patients in the intervention arm and 110 in the control arm. Large SDs: mean  $49.2 \pm 32$  versus  $3.2 \pm 18.6$ . (c) Single RCT with 77 patients in the intervention arm and 110 in the control arm. Large SD: mean  $13.6 \pm 8$  versus  $0.8 \pm 5$ . (d) Single RCT with 77 patients in the intervention arm and 110 in the control arm

**Fig. 7** Forest plot for %TBWL at 12 months in the comparison of ESG vs. high-intensity lifestyle intervention for patients with obesity (all classes)



**Table 9** GRADE assessment of the quality of evidence for comparing ESG vs. lifestyle intervention for patients with mild and moderate obesity. Question: ESG plus lifestyle modification compared to Lifestyle modification alone for obesity

Certainty assessment							№ of patients		Effect		Certainty	Importance
№ of studies	Study design	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	ESG plus lifestyle modification	Lifestyle modification alone	Relative (95% CI)	Absolute (95% CI)		
%TBWL (follow-up: mean 12 months)												
1	observational studies	very serious <sup>a</sup>	not serious	not serious	serious <sup>b</sup>	all plausible residual confounding would reduce the demonstrated effect	43	100	-	MD 6.3% higher (3.12 higher to 9.48 higher)	 Very low	CRITICAL
>5% TBWL Responders rate (follow-up: mean 12 months)												
1	observational studies	very serious <sup>a</sup>	not serious	not serious	serious <sup>b</sup>	all plausible residual confounding would reduce the demonstrated effect	94/105 (89.5%)	219/281 (77.9%)	OR 2.42 (1.22 to 4.80)	116 more per 1,000 (from 32 more to 165 more)	 Very low	IMPORTANT
>10% TBWL Responders rate (follow-up: mean 12 months)												
1	observational studies	very serious <sup>a</sup>	not serious	not serious	serious <sup>b</sup>	all plausible residual confounding would reduce the demonstrated effect	71/105 (67.6%)	143/281 (50.9%)	OR 2.02 (1.26 to 3.23)	168 more per 1,000 (from 57 more to 261 more)	 Very low	IMPORTANT
SAEs rate (follow-up: mean 12 months)												
1	observational studies	very serious <sup>a</sup>	not serious	not serious	serious <sup>b</sup>	all plausible residual confounding would reduce the demonstrated effect	5/105 (4.8%)	0/281 (0.0%)	OR 30.81 (1.69 to 562.21)	0 fewer per 1,000 (from 0 fewer to 0 fewer)	 Very low	CRITICAL

CI confidence interval, MD mean difference, OR odds ratio. (a) Selection bias (matched controls). Channeling bias. Confounding variables (socio-economic status), high loss to follow-up rates. (b) Single study, small total sample size/small number of events, large SDs

## Future Direction

In this report, we investigated endoscopic sleeve gastropasty utilizing the Apollo Overstitch™ platform (Boston Scientific, Marlborough, MA, USA) based on the maturity of the technique, regulatory approvals ([https://www.accessdata.fda.gov/cdrh\\_docs/pdf21/DEN210045.pdf](https://www.accessdata.fda.gov/cdrh_docs/pdf21/DEN210045.pdf), <https://www.nice.org.uk/guidance/ipg783>) and

cost-effectiveness [72–74]. Other endoscopic gastric remodeling techniques, including the Primary Obesity Surgery Endoluminal 2.0 (USGI Medical, San Clemente, CA), Endomina™ Gastric Plication (Endo Tools, Gosselies, Belgium), and the Endozip™ automated suturing device (Caesarea, Israel), are at different stages of clinical trials and evidence generation and are demonstrating similar safety and efficacy profiles. The committee will

update its position statement to reflect and incorporate the evolving evidence base as the clinical evidence continues to mature for these procedures.

Advancements in obesity management medication now offer effective options for selected patients. The value proposition and comparative effectiveness of ESG compared to, or in addition to, obesity pharmacotherapies is an active area of investigation. Observational studies have demonstrated the benefits of combining or sequencing ESG with obesity pharmacotherapies, particularly in enhancing the durability of the response [75].

However, given the limited follow-up in the existing literature—typically extending to five years or less—additional data is required to better understand the different archetypes of response to ESG over the long term. This data will also help in defining optimal personalized approaches to maximize the durability of the response and improve long-term health outcomes.

**Acknowledgements** Acknowledgements: The IFSO Bariatric and Metabolic Endoscopy Committee would like to extend their sincere appreciation to the following experts for their invaluable contributions to the systematic review, evidence extraction, evidence grading, meta-analyses, and the drafting of tables and figures: Dr. Vitor O. Brunaldi (VOB): Methodologist and researcher at the University of São Paulo, specializing in endoscopic techniques. Dr. Nitin Jagtap (NJ): Expert in systematic reviews and meta-analyses at the Asian Institute of Gastroenterology. Dr. Rakesh Kalapala (RK): Senior Consultant at the Asian Institute of Gastroenterology, India. His content expertise ensured the clinical relevance and accuracy of this work. Dr. Barham Abu Dayyeh (BAD): Professor of Medicine and Chair of the IFSO Bariatric and Metabolic Endoscopy Committee during the preparation of this document. IFSO Scientific Committee: For critical review and multiple rounds of revisions of this document prior to approval. IFSO Executive Board: For critical review and multiple rounds of revisions of this document prior to approval.

**Author Contribution** B.K.A.: Conception of the study, critical review of the data, drafting of the manuscript, tables, and figures. Chair, IFSO Bariatric Endoscopy Committee at the time of study initiation. C.S.: Conception of the study, critical review of the data and manuscript. Chair, IFSO Bariatric Endoscopy Committee at the time of study conclusion. A.A.: Conception of the study, review of the manuscript. Member, IFSO Bariatric Endoscopy Committee. R.S.: Conception of the study, review of the manuscript. Member, IFSO Bariatric Endoscopy Committee. M.B.: Conception of the study, review of the manuscript. Member, IFSO Bariatric Endoscopy Committee. S.P.: Conception of the study, review of the manuscript. Member, IFSO Bariatric Endoscopy Committee. S.P.J.: Conception of the study, review of the manuscript. Member, IFSO Bariatric Endoscopy Committee. G.P.: Critical review of the data and manuscript. IFSO President during the conduct of the study. R.C.: Critical review of the data and manuscript. IFSO President at the conclusion of the study.

**Funding** Open access funding provided by SCEL, Statewide California Electronic Library Consortium.

**Data Availability** No datasets were generated or analysed during the current study.

## Declarations

**Competing Interests** Barham K. Abu Dayyeh: Received research support and educational consulting fees from Boston Scientific; previously Apollo Endosurgery. Additionally, received research support from USGI Medical. Christine Stier: Consulting and educational engagements with Boston Scientific. Aayed Alqahtani: Consulting and educational engagements with Boston Scientific. Reem Sharaiha: Consulting and educational engagements with Boston Scientific. Mohit Bandhari: Consulting and educational engagements with Boston Scientific. Silvana Perretta: Consulting and educational engagements with Boston Scientific. Sigh Pichamol Jirapinyo: Consulting and educational engagements with Boston Scientific. Gerhard Prager: Declares no conflicts of interest related to this study. Ricardo Cohen: Declares no conflicts of interest related to this study.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

1. Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128.9 million children, adolescents, and adults. *Lancet*. 2017;390:2627–42.
2. World Health Organization. Overweight and obesity - global observatory data. <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>.
3. Hales CM, Carroll MD, Fryar CD, et al. Prevalence of obesity among adults and youth: United States, 2015–2016. *NCHS Data Brief*. 2017;288:1–8.
4. Hales CM, Fryar CD, Carroll MD, et al. Trends in obesity and severe obesity prevalence in US youth and adults by sex and age, 2007–2008 to 2015–2016. *JAMA*. 2018;319:1723–5.
5. Finkelstein EA, Khavjou OA, Thompson H, et al. Obesity and severe obesity forecasts through 2030. *Am J Prev Med*. 2012;42:563–70.
6. Eisenberg D, Shikora SA, Aarts E, et al. American Society for Metabolic and Bariatric Surgery (ASMBS) and International Federation for the Surgery of Obesity and Metabolic Disorders (IFSO): indications for metabolic and bariatric surgery. *Surg Obes Relat Dis*. 2022;18:1345–56.
7. Carlsson LMS, Sjöholm K, Jacobson P, et al. Life expectancy after bariatric surgery in the Swedish obese subjects study. *N Engl J Med*. 2020;383:1535–43.
8. Angrisani L, Santonicola A, Iovino P, et al. Bariatric Surgery Survey 2018: similarities and disparities among the 5 IFSO chapters. *Obes Surg*. 2021;31:1937–48.
9. Angrisani L, Santonicola A, Iovino P, et al. IFSO Worldwide Survey 2016: primary, endoluminal, and revisional procedures. *Obes Surg*. 2018;28:3783–94.

10. Ogden CL, Carroll MD, Kit BK, et al. Prevalence of childhood and adult obesity in the United States, 2011–2012. *JAMA*. 2014;311:806–14.
11. Whitlock G, Lewington S, Sherliker P, et al. Body-mass index and cause-specific mortality in 900 000 adults: collaborative analyses of 57 prospective studies. *Lancet* (London, England). 2009;373:1083–96.
12. Abu Dayyeh BK, Rajan E, Gostout CJ. Endoscopic sleeve gastrectomy: a potential endoscopic alternative to surgical sleeve gastrectomy for treatment of obesity. *Gastrointest Endosc*. 2013;78:530–5.
13. Galvao-Neto MDP, Grecco E, Souza TFd, et al. Endoscopic sleeve gastrectomy - minimally invasive therapy for primary obesity treatment. *Arquivos brasileiros de cirurgia digestiva : ABCD = Brazilian Archiv of Digestive Surgery*. 2016;29Suppl 1:95–97.
14. Lopez-Nava G, Galvão MP, Bautista-Castaño I, et al. Endoscopic sleeve gastrectomy: how i do it? *Obes Surg*. 2015;25:1534–8.
15. Vargas EJ, Rizk M, Gomez-Villa J, et al. Effect of endoscopic sleeve gastrectomy on gastric emptying, motility and hormones: a comparative prospective study. *Gut* 2022. <https://doi.org/10.1136/gutjnl-2022-327816>
16. Maydeo A, Patil G, Dalal A, et al. An Indian experience of endoscopic treatment of obesity by using a novel technique of endoscopic sleeve gastrectomy (accordion procedure). *J Assoc Physicians India*. 2020;68:14–7.
17. Graus Morales J, Crespo Perez L, Marques A, et al. Modified endoscopic gastrectomy for the treatment of obesity. *Surg Endosc*. 2018;32:3936–42.
18. Espinet-Coll E, Nebreda-Durán J, Galvao-Neto M, et al. Suture pattern does not influence outcomes of endoscopic sleeve gastrectomy in obese patients. *Endoscopy international open*. 2020;8:E1349–58.
19. López-Nava Breviere G, Bautista-Castaño I, Fernández-Corbelle JP, et al. Endoscopic sleeve gastrectomy (the Apollo method): a new approach to obesity management. *Rev Esp Enferm Dig*. 2016;108:201–6.
20. Alqahtani A, Al-Darwish A, Mahmoud AE, et al. Short-term outcomes of endoscopic sleeve gastrectomy in 1000 consecutive patients. *Gastrointest Endosc*. 2019;89:1132–8.
21. Cheskin LJ, Hill C, Adam A, et al. Endoscopic sleeve gastrectomy versus high-intensity diet and lifestyle therapy: a case-matched study. *Gastrointest Endosc*. 2020;91:342–349.e1.
22. Abu Dayyeh BK, Bazerbach F, Vargas EJ, et al. Endoscopic sleeve gastrectomy for treatment of class 1 and 2 obesity (MERIT): a prospective, multicentre, randomised trial. *Lancet*. 2022;400:441–51.
23. Salminen P, Kow L, Aminian A, et al. IFSO consensus on definitions and clinical practice guidelines for obesity management-an international delphi study. *Obes Surg*. 2024;34:30–42.
24. Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ* (Clinical research ed). 2009;339:b2535–b2535.
25. Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, Welch VA. *Cochrane Handbook for Systematic Reviews of Interventions* (Version 6.3, updated February 2022). *Cochrane*. 2022. Available from: <https://training.cochrane.org/handbook>.
26. The Joanna Briggs Institute. *Joanna Briggs Institute Reviewers' Manual*: 2016 ed. Australia: the joanna briggs institute. 2016. Available from: <https://jbi.global/research/handbook>.
27. Wells GA, Shea B, O'Connell D, et al. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses [Internet]. 2021. Available from: [http://www.ohri.ca/programs/clinical\\_epidemiology/oxford.asp](http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp).
28. Jadad AR, Moore RA, Carroll D, et al. Assessing the quality of reports of randomized clinical trials: is blinding necessary? *Control Clin Trials*. 1996;17:1–12.
29. Hozo SP, Djulbegovic B, Hozo I. Estimating the mean and variance from the median, range, and the size of a sample. *BMC Med Res Methodol*. 2005;5:13–13.
30. Schünemann H, Brożek J, Guyatt G, Oxman A. *GRADE handbook for grading quality of evidence and strength of recommendations*. The GRADE Working Group. 2013. Available from: <https://gdt.gradepro.org/app/handbook/handbook.html>.
31. Abu Dayyeh BK, Acosta A, Camilleri M, et al. Endoscopic sleeve gastrectomy alters gastric physiology and induces loss of body weight in obese individuals. *Clin Gastroenterol Hepatol*. 2017;15:37–43.e1.
32. Alqahtani A, Elahmedi M, Alqahtani YA, et al. Endoscopic sleeve gastrectomy in 109 consecutive children and adolescents with obesity: two-year outcomes of a new modality. *Am J Gastroenterol*. 2019;114:1857–62.
33. Alqahtani AR, Elahmedi M, Aldarwish A, et al. Endoscopic gastrectomy versus laparoscopic sleeve gastrectomy: a noninferiority propensity score-matched comparative study. *Gastrointest Endosc*. 2022;96:44–50.
34. Asokkumar R, Lim CH, Tan AS, et al. Safety and early efficacy of endoscopic sleeve gastrectomy (ESG) for obesity in a multi-ethnic Asian population in Singapore. *JGH Open*. 2021;5:1351–6.
35. Badurdeen D, Hoff AC, Hedjoudje A, et al. Endoscopic sleeve gastrectomy plus liraglutide versus endoscopic sleeve gastrectomy alone for weight loss. *Gastrointest Endosc*. 2021;93:1316–1324.e1.
36. Barrichello S, Hourneaux de Moura DT, Hourneaux de Moura EG, et al. Endoscopic sleeve gastrectomy in the management of overweight and obesity: an international multicenter study. *Gastrointest Endosc* 2019;90:770–780.
37. Bhandari M, Jain S, Mathur W, et al. Endoscopic sleeve gastrectomy is an effective and safe minimally invasive approach for treatment of obesity: first Indian experience. *Digestive endoscopy : official journal of the Japan Gastroenterological Endoscopy Society* 2019. <https://doi.org/10.1111/den.13508>
38. Callahan ZM, Su B, Kuchta K, et al. Endoscopic suturing results in high technical and clinical success rates for a variety of gastrointestinal pathologies. *J Gastrointest Surg*. 2020;24:278–87.
39. Carr P, Keighley T, Petocz P, et al. Efficacy and safety of endoscopic sleeve gastrectomy and laparoscopic sleeve gastrectomy with 12+ months of adjuvant multidisciplinary support. *BMC Prim Care*. 2022;23:26.
40. Cheskin LJ, Hill C, Adam A, et al. Endoscopic sleeve gastrectomy versus high-intensity diet and lifestyle therapy: a case-matched study. *Gastrointest Endosc*. 2020;91:342–349.e1.
41. Espinet-Coll E, Díaz-Galán P, Nebreda-Durán J, et al. Persistence of sutures and gastric reduction after endoscopic sleeve gastrectomy: radiological and endoscopic assessment. *Obes Surg*. 2022;32:1969–79.
42. Espinet-Coll E, Nebreda-Durán J, Galvao-Neto M, et al. Suture pattern does not influence outcomes of endoscopic sleeve gastrectomy in obese patients. *Endosc Int Open*. 2020;8:E1349–e1358.
43. Farha J, McGowan C, Hedjoudje A, et al. Endoscopic sleeve gastrectomy: suturing the gastric fundus does not confer benefit. *Endoscopy*. 2021;53:727–31.
44. Fayad L, Adam A, Schweitzer M, et al. Endoscopic sleeve gastrectomy versus laparoscopic sleeve gastrectomy: a case-matched study. *Gastrointest Endosc*. 2019;89:782–8.
45. Fayad L, Cheskin LJ, Adam A, et al. Endoscopic sleeve gastrectomy versus intragastric balloon insertion: efficacy, durability, and safety. *Endoscopy* 2019. <https://doi.org/10.1055/a-0852-3441>

46. Fiorillo C, Quero G, Vix M, et al. 6-month gastrointestinal quality of life (QoL) results after endoscopic sleeve gastroplasty and laparoscopic sleeve gastrectomy: a propensity score analysis. *Obes Surg.* 2020;30:1944–51.
47. Ghaz H, Bryant M, Fritz H, et al. Endoscopic sleeve gastroplasty and postprocedural nutritional deficiencies: results from a single center exploratory study. *Eur J Gastroenterol Hepatol.* 2021;33:e1039–41.
48. Glaysher MA, Moekotte AL, Kelly J. Endoscopic sleeve gastroplasty: a modified technique with greater curvature compression sutures. *Endosc Int Open.* 2019;7:E1303–e1309.
49. Graus Morales J, Crespo Pérez L, Marques A, et al. Modified endoscopic gastroplasty for the treatment of obesity. *Surg Endosc.* 2018;32:3936–42.
50. Gudur AR, Geng C, Kshatri S, et al. Comparison of endoscopic sleeve gastroplasty versus surgical sleeve gastrectomy: a metabolic and bariatric surgery accreditation and quality improvement program database analysis. *Gastrointest Endosc* 2022. <https://doi.org/10.1016/j.gie.2022.07.017>
51. Hajifathalian K, Mehta A, Ang B, et al. Improvement in insulin resistance and estimated hepatic steatosis and fibrosis after endoscopic sleeve gastroplasty. *Gastrointest Endosc.* 2021;93:1110–8.
52. Hill C, El Zein M, Agnihotri A, et al. Endoscopic sleeve gastroplasty: the learning curve. *Endosc Int Open.* 2017;5:E900–e904.
53. Jagtap N, Kalapala R, Katakwar A, et al. Endoscopic sleeve gastroplasty - minimally invasive treatment for non-alcoholic fatty liver disease and obesity. *Indian J Gastroenterol.* 2021;40:572–9.
54. James TW, Reddy S, Vulpis T, et al. Endoscopic sleeve gastroplasty is feasible, safe, and effective in a non-academic setting: short-term outcomes from a community gastroenterology practice. *Obes Surg.* 2020;30:1404–9.
55. Kumar N, Abu Dayyeh BK, Lopez-Nava Breviere G, et al. Endoscopic sutured gastroplasty: procedure evolution from first-in-man cases through current technique. *Surg Endosc.* 2018;32:2159–64.
56. Li R, Veltzke-Schlieker W, Adler A, et al. Endoscopic sleeve gastroplasty (ESG) for high-risk patients, high body mass index (> 50 kg/m<sup>2</sup>) patients, and contraindication to abdominal surgery. *Obes Surg.* 2021;31:3400–9.
57. Lopez-Nava G, Laster J, Negi A, et al. Endoscopic sleeve gastroplasty (ESG) for morbid obesity: how effective is it? *Surg Endosc.* 2022;36:352–60.
58. Lopez-Nava G, Negi A, Bautista-Castaño I, et al. Gut and metabolic hormones changes after endoscopic sleeve gastroplasty (ESG) vs. laparoscopic sleeve gastrectomy (LSG). *Obes Surg.* 2020;30:2642–51.
59. Lopez-Nava G, Sharaiha RZ, Vargas EJ, et al. Endoscopic sleeve gastroplasty for obesity: a multicenter study of 248 patients with 24 months follow-up. *Obes Surg.* 2017;27:2649–55.
60. Manos T, Costil V, Karsenty L, et al. Safety of endoscopic sleeve gastroplasty with a single-channel endoscope. *Obes Surg.* 2022;32:3074–8.
61. Matteo MV, Bove V, Pontecorvi V, et al. Outcomes of endoscopic sleeve gastroplasty in the elder population. *Obes Surg* 2022. <https://doi.org/10.1007/s11695-022-06232-4>
62. Mehta A, Hajifathalian K, Shah SL, et al. Quality of life, mental health, and weight loss outcomes following endoscopic sleeve gastroplasty. *J Gastrointestinal Surg : Off J Soc Surg Alimentary Tract.* 2022;26:469–71.
63. Neto MG, Moon RC, de Quadros LG, et al. Safety and short-term effectiveness of endoscopic sleeve gastroplasty using over-stitch: preliminary report from a multicenter study. *Surg Endosc.* 2020;34:4388–94.
64. Neto MG, Silva LB, de Quadros LG, et al. Brazilian consensus on endoscopic sleeve gastroplasty. *Obes Surg.* 2021;31:70–8.
65. Novikov AA, Afaneh C, Saumoy M, et al. Endoscopic sleeve gastroplasty, laparoscopic sleeve gastrectomy, and laparoscopic band for weight loss: how do they compare? *J Gastrointest Surg.* 2018;22:267–73.
66. Pizzicannella M, Lapergola A, Fiorillo C, et al. Does endoscopic sleeve gastroplasty stand the test of time? Objective assessment of endoscopic ESG appearance and its relation to weight loss in a large group of consecutive patients. *Surg Endosc.* 2020;34:3696–705.
67. Rapaka B, Maselli DB, Lopez-Nava G, et al. Effects on physiologic measures of appetite from intragastric balloon and endoscopic sleeve gastroplasty: results of a prospective study. *Chin Med J (Engl).* 2022;135:1234–41.
68. Sarkar A, Tawadros A, Andalib I, et al. Safety and efficacy of endoscopic sleeve gastroplasty for obesity management in new bariatric endoscopy programs: a multicenter international study. *Ther Adv Gastrointest Endosc.* 2022;15:26317745221093884.
69. Sartoretto A, Sui Z, Hill C, et al. Endoscopic sleeve gastroplasty (ESG) is a reproducible and effective endoscopic bariatric therapy suitable for widespread clinical adoption: a large, international multicenter study. *Obes Surg.* 2018;28:1812–21.
70. Saumoy M, Schneider Y, Zhou XK, et al. A single-operator learning curve analysis for the endoscopic sleeve gastroplasty. *Gastrointest Endosc.* 2018;87:442–7.
71. Sharaiha RZ, Hajifathalian K, Kumar R, et al. Five-year outcomes of endoscopic sleeve gastroplasty for the treatment of obesity. *Clin Gastroenterol Hepatol.* 2021;19:1051–1057.e2.
72. Kelly J, Menon V, O'Neill F, et al. UK cost-effectiveness analysis of endoscopic sleeve gastroplasty versus lifestyle modification alone for adults with class II obesity. *Int J Obes (Lond).* 2023;47:1161–70.
73. Saumoy M, Gandhi D, Buller S, et al. Cost-effectiveness of endoscopic, surgical and pharmacological obesity therapies: a micro-simulation and threshold analyses. *Gut.* 2023;72:2250–9.
74. Haseeb M, Chhatwal J, Xiao J, et al. Semaglutide vs endoscopic sleeve gastroplasty for weight loss. *JAMA Netw Open.* 2024;7:e246221.
75. Gala K, Ghosn W, Brunaldi V, et al. Outcomes of concomitant antiobesity medication use with endoscopic sleeve gastroplasty in clinical US settings. *Obes Pillars.* 2024;11:100112.

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

## Authors and Affiliations

**Barham K. Abu Dayyeh**<sup>1,10</sup> · **Christine Stier**<sup>2</sup> · **Aayed Alqahtani**<sup>3</sup> · **Reem Sharaiha**<sup>4</sup> · **Mohit Bandhari**<sup>5</sup> · **Silvana Perretta**<sup>6</sup> · **Sigh Pichamol Jirapinyo**<sup>7</sup> · **Gerhard Prager**<sup>8</sup> · **Ricardo V. Cohen**<sup>9</sup>

✉ Barham K. Abu Dayyeh  
abudayyeh.barham@mayo.edu

Christine Stier  
christinestier@gmail.com

Aayed Alqahtani  
qahtani@yahoo.com

Reem Sharaiha  
rzs9001@med.cornell.edu

Mohit Bandhari  
drmohitbandhari@gmail.com

Silvana Perretta  
silvana.perretta@ircad.fr

Sigh Pichamol Jirapinyo  
pjirapinyo@bwh.harvard.edu

Gerhard Prager  
gerhard.prager@meduniwien.ac.at

Ricardo V. Cohen  
ricardo.cohen@prestador.haoc.com.br

<sup>1</sup> Mayo Clinic, Rochester, USA

<sup>2</sup> University Medical Centre Mannheim, Mannheim, Germany

<sup>3</sup> King Abdulaziz University, Jeddah, Saudi Arabia

<sup>4</sup> Cornell University, Ithaca, USA

<sup>5</sup> Sri Aurobindo Medical College and PG Institute, Indore, India

<sup>6</sup> IRCAD, Strasbourg, France

<sup>7</sup> Brigham and Women's Hospital, Boston, USA

<sup>8</sup> Medical University of Vienna, Vienna, Austria

<sup>9</sup> The Center for Obesity and Diabetes, Oswaldo Cruz German Hospital, San Paolo, Brazil

<sup>10</sup> Gastroenterology and Advanced Endoscopy, Cedars-Sinai Health System, Los Angeles, USA