

## Right hepatectomy for combined primary neuroendocrine and hepatocellular carcinoma. A case report<sup>☆</sup>

Ahmed Aboelenen, Amira Kamal El-Hawary, Nirmene Megahed, Khaled refaat Zalata, Eman M. El-Salk, Marwa Abdel Fattah, Mohamed El Sorogy\*, Ahmed Shehta

Gastroenterology Surgical Center, Mansoura University, Egypt



### ARTICLE INFO

#### Article history:

Received 10 October 2013

Received in revised form 24 October 2013

Accepted 30 October 2013

Available online 11 November 2013

#### Keywords:

Neuroendocrine tumor

Hepatocellular carcinoma

Right hepatectomy

### ABSTRACT

**INTRODUCTION:** Cases of primary neuroendocrine tumors in the liver combined with hepatocellular carcinoma are scarce. Such cases could present either as combined-type tumor or collision type.

**PRESENTATION OF CASE:** A 51-year-old man presented with a mass in the right hemiliver. Serum level of alpha-fetoprotein was slightly elevated (2.3 ng/ml), with normal CA19-9 and CA125. The patient underwent right hepatectomy. The resected specimen showed a well-defined and heterogeneous gray-white to brown friable tumor, 20 cm in diameter. Microscopically, the tumor consisted predominantly of monotonous small- to medium-sized neoplastic cells arranged in trabeculae separated by sinusoidal spaces. Immunohistochemically, the tumor cells were strongly positive for synaptophysin and focally positive for chromogranin-A. Interestingly, the tumor cells showed patchy positive coarse granular staining of HerPar-1 involving about 1% of the tumor cells. Glypican-3 staining was negative. These immunohistochemical findings supported the diagnosis of combined high grade neuroendocrine carcinoma and hepatocellular carcinoma.

**DISCUSSION:** Cases of primary neuroendocrine tumors in the liver combined with hepatocellular carcinoma are scarce. The uniqueness of this case lies in the fact that the neuroendocrine carcinoma component comprised more than 99% of the tumor area, and the minor hepatocellular carcinoma component was detected only by the immunohistochemical staining for HepPar-1.

**CONCLUSION:** To the best of our knowledge, this is the first case of combined neuroendocrine carcinoma and hepatocellular carcinoma in Egypt.

© 2013 The Authors. Published by Elsevier Ltd on behalf of Surgical Associates Ltd. All rights reserved.

## 1. Introduction

Primary hepatic neuroendocrine carcinoma (NEC) is rare and its origin is not clearly understood. An admixture of hepatocellular carcinoma (HCC) and neuroendocrine carcinoma is particularly rare.<sup>1</sup> There are 2 types of primary mixed NEC and HCC in the liver. The combined-type tumor in which both NEC and HCC components intermingle with each other and cannot be clearly separated in the transitional area within a single tumor nodule,<sup>2</sup> and the collision-type in which the tumor have two histologically distinct tumors simultaneously involving the same organ with no transition between them.<sup>3</sup> We herein describe a case of combined primary high grade NEC and HCC of the liver in which the NEC component

comprised more than 99% of the tumor area, and only small nests of HCC were detected only by immunohistochemical staining for HepPar-1.


## 2. Case report

A 51-year-old man presented to our outpatient clinic with a 9-month history of dull aching abdominal pain and a medical history of hepatitis C. Periodic hepatic ultrasound follow-up showed a mass in the right hemiliver. A CT scan of the abdomen revealed a 20 cm × 15 cm mass occupying most of the hemiliver with cystic degeneration and calcification and multiple dilated collaterals related to the mass (Fig. 1). The mass was compressing the right portal vein, yet no invasion detected (Fig. 2). CT volumetry estimated the residual left hemiliver volume including the middle hepatic vein to be 1300 cc. Serum level of alpha-fetoprotein (AFP) was slightly elevated (2.3 ng/ml). The patient had normal liver enzymes, total serum bilirubin of 1.2 mg/dl, serum albumin of 3.3 mg/dl and INR of 1.1. CA19-9 and CA125 were normal. CT of chest and extrahepatic abdomen showed no other lesions.

\* This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike License, which permits non-commercial use, distribution, and reproduction in any medium, provided the original author and source are credited.

Corresponding author. Tel.: +20 1003448984; fax: +20 502243220.

E-mail addresses: [sorogy@gmail.com](mailto:sorogy@gmail.com), [msorogy@mans.edu.eg](mailto:msorogy@mans.edu.eg) (M.E. Sorogy).




**Fig. 1.** Arterial phase abdominal CT scan showing large right hemiliver mass with areas of cystic degeneration. The residual left hemiliver including the middle hepatic vein being shadowed.

The patient underwent laparotomy through a J shaped incision (Fig. 3). On exploration the mass was found to be capsulated, 20 cm × 15 cm occupying most of the right hemiliver with multiple dilated collaterals extending from the mass to the diaphragm and the hepatic flexure of the colon. No ascites was found. No peritoneal nodules or significant lymph nodes. Right hepatectomy was done. Blood loss was 2500 cc and the patient received 4 units of blood intraoperatively due to presence of collaterals and large size of the mass making mobilization and resection of the right lobe technically challenging. The patient had an eventless post-operative course and was discharged on the 6th post-operative day.

The resected specimen measured 20 cm × 15 cm and weighed 4500 g. Cut section revealed a well-defined heterogeneous gray-white to brown friable mass 7.5 cm in diameter (Fig. 4). The surrounding liver was non-cirrhotic.

Microscopically, the tumor consisted predominantly of monotonous small- to medium-sized neoplastic cells arranged in trabeculae separated by sinusoidal spaces. Tumor cells had round to oval punctuate nuclei within conspicuous nucleoli and scanty cytoplasm (Fig. 5A). The tumor was accompanied by wide areas of



**Fig. 3.** Operative photography of the mass showing dilated veins extending from the mass to the hepatic flexure of the colon.

geographic necrosis with high mitotic rate that ranged between 2 and 20/10 HPF. The whole morphological picture was reminiscent of high grade NEC. The surrounding liver tissue revealed chronic hepatitis without cirrhotic changes.


Immunohistochemically, the tumor cells were strongly positive for synaptophysin (Fig. 5D) and focally positive for chromogranin-A. Interestingly, the tumor cells showed patchy positive coarse granular staining for HerPar-1 in less than 1% of tumor cells (Fig. 5B and C) while Glycican-3staining was negative. The immunohistochemical findings supported the diagnosis of combined high grade NEC and HCC. Follow up period was short but no recurrence detected up to 6 months after surgery.

### 3. Discussion


Cases of primary neuroendocrine tumors in the liver combined with hepatocellular carcinoma are scarce.<sup>2,4</sup> It can present either as combined-type or collision type.<sup>5</sup> We herein describe a case of combined primary high grade NEC and HCC of the liver in which the NEC component comprised more than 99% of the tumor area, and a small nest of HCC was detected only by immunohistochemical staining for HepPar-1. In the case reported by Yang et al.,<sup>5</sup> the tumor consisted predominantly of small- to medium-sized



**Fig. 2.** CT portography and hepatic venography showing the relation of the mass to the portal and hepatic veins.



**Fig. 4.** Cut section of the specimen.



**Fig. 5.** (A) The tumor was formed predominantly of small to medium-sized monotonous cells arranged in trabecular pattern separated by highly vascular stroma. There was no histological evidence of HCC (100 $\times$ ). (B) Less than 1% of the tumor cells showed granular cytoplasmic staining for Hep-Par-1. The cells showed no histological evidence of HCC differentiation by H&E (100 $\times$ ). (C) Higher magnification of Hep-Par-1 immunostaining. (D) The same cells showed cytoplasmic positive staining for synaptophysin.

neoplastic cells arranged in solid sheets separated by dense fibrovascular septa which were reminiscent of NEC. In addition, a small island of polygonal tumor cells with abundant eosinophilic granular cytoplasm and round nuclei arranged in trabecular structures consistent with moderately differentiated HCC were detected in H&E sections. However, in our case, we could not find the large polygonal cells with morphological features of HCC in H&E sections. Those cells were highlighted only after immuno-histochemistry for HepPar-1 which showed patchy cytoplasmic granular staining in less than 1% of tumor cells. Interestingly, those cells were also positive for synaptophysin.

Yamaguchi et al.<sup>2</sup> reported that some tumor cells of the HCC in the combined tumor may be immunoreactive for neuroendocrine markers. This supports the postulations of Yang et al.,<sup>5</sup> Ishida et al.<sup>6</sup> and Tazi et al.<sup>3</sup> that combined NEC-HCC originates from poorly differentiated HCC that underwent extensive neuroendocrine differentiation. Other studies postulated that these combined tumors arise from stem cells with divergent differentiation.<sup>7,8</sup>

In our case, the surrounding liver tissue was non-cirrhotic which goes in concordance with most studies that showed that about 20–40% of primary NEC-HCC developed in non-cirrhotic livers.<sup>9,10</sup>

On the contrary, pure HCCs are well known to be a direct complication of liver cirrhosis.

Follow up of our patient for 6 months revealed no evidence of recurrence. Despite the fact that patients with pure high grade neuroendocrine carcinomas usually die within few months after diagnosis,<sup>6</sup> reports of some combined NEC-HCC cases showed longer survival without recurrences for up to 28 months.<sup>5</sup> However, the number of reported NEC-HCC is still rare and further studies are still needed to substantiate prognosis and treatment of such cases.

To the best of our knowledge, this is the first case of combined NEC and HCC in Egypt. The uniqueness of this case lies in the fact that the NEC component comprised more than 99% of the tumor area, and the minor HCC component was detected only by immunohistochemical staining for HepPar-1.

### **Conflict of interest statement**

None.

### **Funding**

None.

### **Ethical approval**

A written informed consent was obtained from the patient for publication of the case report. A copy of the written consent is available for review by the Editor-in-Chief of this journal on request.

### **Author contributions**

- (1) The acquisition of data, or analysis and interpretation of data (Amira Kamal El-Hawary, Nirmeen Megahed, Eman M El-Salk, Marwa Abdel Fattah).

- (2) Drafting the article or revising it critically for important intellectual content (Amira Kamal El-Hawary, Mohamed El Sorogy, Ahmed Shehta).
- (3) Final approval of the version to be submitted (Ahmed Aboelenen, Khaled refaat Zalata).

### **Acknowledgement**

The author thanks Mr. Ebrahim Attia for his technical assistance.

### **References**

1. Nakanishi C, Sato K, Ito Y, Abe T, Akada T, Muto R, et al. Combined hepatocellular carcinoma and neuroendocrine carcinoma with sarcomatous change of the liver after transarterial chemoembolization. *Hepatol Res* 2012;42(11):1141–5.
2. Yamaguchi R, Nakashima O, Ogata T, Hanada K, Kumabe T, Kojiro M. Hepatocellular carcinoma with an unusual neuroendocrine component. *Pathol Int* 2004;54:861–5.
3. Tazi E, Essadi I, M'rabi H. Hepatocellular carcinoma and high grade neuroendocrine carcinoma: a case report and review of the literature. *World J Oncol* 2011;2(1):37–40.
4. Barsky SH, Linnola I, Triche TJ, Costa J. Hepatocellular carcinoma with carcinoid features. *Hum Pathol* 1984;15(9):892–4.
5. Yang C, Wen MC, Jan YJ, Wang J, Wu CC. Combined primary neuroendocrine carcinoma and hepatocellular carcinoma of the liver. *J Chin Med Assoc* 2009;72(August (8)):430–3.
6. Ishida M, Seki K, Tatsuzawa A, Katayama K, Hirose K, Azuma T, et al. Primary hepatic neuroendocrine carcinoma coexisting with hepatocellular carcinoma in hepatitis C liver cirrhosis: report of a case. *Surg Today* 2003;33:214–8.
7. Haratake J, Hashimoto H. An immunohistochemical analysis of 13 cases with combined hepatocellular and cholangiocellular carcinoma. *Liver* 1995;15(1):9–15.
8. Theise ND, Yao JL, Harada K, Hytiroglou P, Portmann B, Thung SN, et al. Hepatic 'stem cell' malignancies in adults: four cases. *Histopathology* 2003;43(3):263–71.
9. Monto A, Wright TL. The epidemiology and prevention of hepatocellular carcinoma. *Semin Oncol* 2001;28(5):441–9.
10. Nzeako UC, Goodman ZD, Ishak KG. Hepatocellular carcinoma in cirrhotic and noncirrhotic livers. A clinico-histopathologic study of 804 North American patients. *Am J Clin Pathol* 1996;105(1):65–75.

### **Open Access**

This article is published Open Access at [sciedirect.com](http://sciedirect.com). It is distributed under the [IJSCR Supplemental terms and conditions](#), which permits unrestricted non commercial use, distribution, and reproduction in any medium, provided the original authors and source are credited.