

Non-genomic effects of nuclear receptors: insights from the anucleate platelet

Amanda J. Unsworth[†], Gagan D. Flora[†], and Jonathan M. Gibbins*

School of Biological Sciences, Institute of Cardiovascular and Metabolic Research, Harborne Building, Whiteknights, Reading RG6 6AS, Berkshire, UK

Received 9 November 2017; revised 3 January 2018; editorial decision 17 January 2018; accepted 13 February 2018; online publish-ahead-of-print 14 February 2018

Abstract

Nuclear receptors (NRs) have the ability to elicit two different kinds of responses, *genomic* and *non-genomic*. Although genomic responses control gene expression by influencing the rate of transcription, non-genomic effects occur rapidly and independently of transcriptional regulation. Due to their anucleate nature and mechanistically well-characterized and rapid responses, platelets provide a model system for the study of any non-genomic effects of the NRs. Several NRs have been found to be present in human platelets, and multiple NR agonists have been shown to elicit anti-platelet effects by a variety of mechanisms. The non-genomic functions of NRs vary, including the regulation of kinase and phosphatase activity, ion channel function, intracellular calcium levels, and production of second messengers. Recently, the characterization of mechanisms and identification of novel binding partners of NRs have further strengthened the prospects of developing their ligands into potential therapeutics that offer cardio-protective properties in addition to their other defined genomic effects.

Keywords

Nuclear receptors • Non-genomic • Platelets • Thrombosis

1. Introduction

Nuclear receptors (NRs) represent the family of mammalian proteins associated with the transcriptional regulation in human tissues and include the androgen receptor (AR), oestrogen receptor (ER), glucocorticoid receptor (GR), farnesoid X receptor (FXR), liver X receptor (LXR), peroxisome proliferator-activated receptors (PPARs), retinoic acid receptor (RAR), retinoid X receptor (RXR), and the vitamin D receptor (VDR). Upon activation by their lipophilic ligands, NRs regulate several fundamental biological processes, such as cell proliferation, differentiation, metabolism, and homeostasis (Table 1).^{1,2} Any deviation from their normal function can lead to the pathological manifestations, such as cancer, diabetes, arthritis, and obesity.³

NRs have the ability to function in both *genomic* and *non-genomic* ways. Whilst historically associated with the regulation of transcription and control of gene expression (genomic), more recently non-genomic roles for the NRs have been identified that occur independently of transcriptional regulation. Unlike the genomic functions, which can occur over minutes or hours, these events occur in the time frame of seconds to a few minutes, which is considered too rapid to be attributed to the biosynthesis of mRNA or protein, and is often unaffected by the inhibitors of transcription or translation.⁴ The non-genomic functions of NRs vary^{5–7} and whilst it is thought these functions are initiated by physical

interactions of NRs with cofactors and binding partners that initiate rapid signalling events, the exact mechanisms are not well understood. One possible explanation is that the cellular localization of NRs influences the availability of cofactors and substrates, which leads to varying combinations of binding partner interactions. For instance, localization of NRs to the cytosol, plasma membrane, or other intracellular organelles such as mitochondria increases the likelihood of initiation of non-genomic effects, whilst genomic functions may be restricted to when NRs are localized in the nucleus.^{8–11} The formation of different multi-protein signalling complexes with different localization and distribution patterns across different cell types could offer a high degree of cell and tissue selective action for the NRs but as of yet these are poorly defined.^{10,11} Given the underlying differences between the genomic and non-genomic activities, non-genomic effects are more easily observed in cell types that lack a functional nucleus such as erythrocytes and platelets.

Emerging evidence indicates that platelets are also involved in roles beyond those described in haemostasis and thrombosis. For instance, granule secretion following platelet activation results in the release of an array of chemokines, cytokines, growth factors, anti-inflammatory factors, and several other biologically active molecules into the vicinity of injured tissues that contribute towards the progression of numerous diseases including inflammatory conditions (e.g. atherosclerosis and rheumatoid arthritis),^{12,13} type 2 diabetes,¹⁴ and cancer cell metastasis.¹⁵

* Corresponding author. Tel: +44 (0)118 3787082; fax: +44 (0) 118 9310180, E-mail: j.m.gibbins@reading.ac.uk

[†] The first two authors contributed equally to the study.

© The Author(s) 2018. Published by Oxford University Press on behalf of the European Society of Cardiology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

Table 1 NRs and their biological functions

Nuclear receptor	Ligands	Biological function
GR	Natural: glucocorticoid Synthetic: RU38486, A348441	Lipolysis Glucose metabolism
ER	Natural: oestrogen, including oestrone (E1), oestradiol (E2) and oestriol (E3)	Development of the female reproductive system and secondary sexual characteristics
AR	Natural: dihydrotestosterone, testosterone Synthetic: mibolerone	Development of the male reproductive system and secondary sexual characteristics
LXR	Natural: oxysterols Synthetic: T0901317, GW3965	Lipid and carbohydrate metabolism
FXR	Natural: bile acids Synthetic: GW4064, farnesol, CDCA	Bile acid homeostasis
PPAR α	Natural: polyunsaturated fatty acids Synthetic: fibrates (gemfibrozil, fenofibrate, clofibrate)	Fatty acid oxidation and lipid metabolism
PPAR β	Natural: unsaturated/saturated fatty acids, eicosanoids, prostacyclin Synthetic: GW501516	Cholesterol metabolism
PPAR γ	Natural: 15-deoxy-12, 14 prostaglandin J2 (15d-PGJ2) Synthetic: thiazolidinedione (ciglitazone, pioglitazone, rosiglitazone)	Lipid and glucose metabolism
RAR	Natural: atRA	Cell growth, differentiation and organogenesis
RXR	Natural: 9-cis-retinoic acid, docosahexaenoic acid Synthetic: methoprene acid, rexinoids (LG100268)	Cellular proliferation and differentiation, glucose, fatty acid and cholesterol metabolism
VDR	Natural: calcitriol Synthetic: maxacalcitol, calcipotriol	Calcium homeostasis, cell proliferation and differentiation

Thus, platelets are highly active cells with diverse functions despite lacking genomic DNA. Although devoid of a nucleus, platelets still contain different forms of RNA (mRNA, rRNA, tRNA, and miRNA) and components of the transcription and translation machinery that are derived from megakaryocytes during thrombopoiesis.¹⁶ There is a growing consensus that these RNAs are not subjected to a random transfer by megakaryocytes but are specifically sorted and are competent for translation within platelets.^{17,18} Moreover, there is evidence to suggest that platelet-derived microparticles may deliver platelet mRNAs into other nucleated cells, such as monocytes and endothelial cells, where they then undergo translation.¹⁹ Components of this transcription machinery found to be present inside platelets include the intracellular NRs. Due to their nucleated nature and mechanistically well characterized and rapid responses, such as aggregation and adhesion, platelets provide an excellent model system to study the acute non-genomic effects of the NRs.^{20,21}

2. NRs are acute regulators of platelet function

On the basis of the mechanisms of action in the nucleated cells, NRs are classified into two classes: type I or the steroid hormone receptors and type II or the non-steroid receptors. Platelets are known to express

both the classes of these receptors. This includes the AR,^{22,23} ER,^{22,24–26} GR,^{27,28} FXR,^{29,30} LXR,^{30,31} PPARs,^{32–40} RAR,⁴¹ RXR,^{42,43} and VDR.^{44,45}

Both natural and synthetic ligands for these NRs have been shown to alter platelet function through a variety of mechanisms as described below and summarized in *Table 2*.

3. Type I NRs

Mechanisms by which type I NRs (GR, ER, and AR) regulate platelet functions are poorly understood. This might be attributed to the variations in the plasma levels of the steroid hormones targeting these NRs (especially in females and under certain pathological conditions).^{49–51} This might lead to an inaccurate assessment of the role type I NRs play in modulating platelet functions in acute vs. chronic studies and might account for the existing contradictory published data. The effects of ligands of type I NRs on platelet function currently published are described below.

3.1 Glucocorticoid receptor

GRs are activated by glucocorticoid and anti-inflammatory hormones that regulate inflammation and glucose homeostasis.⁵² Prednisolone, a synthetic glucocorticoid derived from cortisol, has been shown to

Table 2 A summary of NRs identified in platelets and their modes of action

Nuclear receptor	Ligands	Effect on platelet function	Mechanisms of action
GR ^{27,28}	<ul style="list-style-type: none"> • Prednisolone 	<ul style="list-style-type: none"> • Negative regulation of platelet secondary mediator regulated effects (ADP and TXA₂) <ul style="list-style-type: none"> • <i>In vitro</i> • Human platelets 	Mechanism is unknown
ER ^{25,26}	<ul style="list-style-type: none"> • Oestrogen—oestrone (E1), oestradiol (E2) and oestriol (E3) 	<ul style="list-style-type: none"> • Reduction in platelet responsiveness however, conflicting results exist <ul style="list-style-type: none"> • <i>In vitro, ex vivo, in vivo</i> • Human, mouse platelets 	Mechanism is unknown
AR ^{46–48}	<ul style="list-style-type: none"> • Testosterone • Dihydrotestosterone 	<ul style="list-style-type: none"> • Potentiation of platelet aggregation <ul style="list-style-type: none"> • <i>In vitro</i> and <i>ex vivo</i> • Human and rat platelets 	Mechanism is unknown
LXR ³¹	<ul style="list-style-type: none"> • GW3965 • T0901317 • 24(S)-OH-cholesterol • 27-OH-cholesterol 	<ul style="list-style-type: none"> • Inhibition of platelet function and thrombosis <ul style="list-style-type: none"> • <i>In vitro</i> and <i>in vivo</i> • Human and mouse platelets • Conversion of platelets to the procoagulant state <ul style="list-style-type: none"> • <i>In vitro</i> • Human platelets 	<ul style="list-style-type: none"> • Reduced phosphorylation of early GPVI signalling components—Syk, LAT and PLCγ2 • Increase LXR-Syk and LXR-PLCγ2 interaction • Formation of coated platelets, including PS exposure, mitochondrial membrane depolarization (see Figure 1)
FXR ²⁹	<ul style="list-style-type: none"> • GW4064 • Chenodeoxycholic acid • 6α-ethyl-chenodeoxycholic acid 	<ul style="list-style-type: none"> • Inhibition of platelet function, thrombosis and haemostasis <ul style="list-style-type: none"> • <i>In vitro</i> and <i>in vivo</i> • Human, mouse platelets • Conversion of platelets to the procoagulant state <ul style="list-style-type: none"> • <i>In vitro</i> • Human platelets 	<ul style="list-style-type: none"> • Cyclophilin D-dependent formation of coated platelets and closure of surface integrins • Associated with PS exposure and mitochondrial membrane depolarization • Augmented cGMP levels which promote PKG activity and phosphorylation of VASP S239 (see Figure 1)
PPAR α ³³	<ul style="list-style-type: none"> • Fenofibrate • Statins 	<ul style="list-style-type: none"> • Inhibition of platelet function <ul style="list-style-type: none"> • <i>In vitro</i> • Human, mouse platelets 	<ul style="list-style-type: none"> • Increase in cAMP levels • PPARα–PKCα interaction and attenuation of PKCα (see Figure 2)
PPAR β/δ ³⁵	<ul style="list-style-type: none"> • GW0742 • L-165041 	<ul style="list-style-type: none"> • Inhibition of platelet function <ul style="list-style-type: none"> • <i>In vitro</i> • Human, mouse platelets 	<ul style="list-style-type: none"> • Increase in cAMP levels • PPARα–PKCα interaction and attenuation of PKCα (see Figure 2)
PPAR γ ^{38,39}	<ul style="list-style-type: none"> • 15d-PGJ2 • Thiazolidinediones (rosiglitazone, ciglitazone, pioglitazone) 	<ul style="list-style-type: none"> • Inhibition of platelet function, thrombosis and haemostasis <ul style="list-style-type: none"> • <i>In vitro</i> and <i>in vivo</i> • Human, mouse platelets 	<ul style="list-style-type: none"> • Inhibition in phosphorylation of Syk and LAT to reduce GPVI signalling • Reduced PPARγ–Syk and PPARγ–LAT interaction upon PPARγ ligand treatment • Negative regulation of integrin αIIbβ3 outside-in via up-regulation of PKA activity and inhibition β3 phosphorylation (see Figure 2)
RAR ⁴¹	<ul style="list-style-type: none"> • atRA 	<ul style="list-style-type: none"> • Inhibition of cytoskeletal rearrangements and platelet spreading <ul style="list-style-type: none"> • <i>In vitro</i> • Human platelets 	Disruption of RAR α –Arp2/3 interactions. (see Figure 3)

Continued

Table 2 Continued

Nuclear receptor	Ligands	Effect on platelet function	Mechanisms of action
RXR ^{42,43}	<ul style="list-style-type: none"> 9-<i>cis</i>-retenoic acid Methoprene acid Docosahexaenoic acid 	<ul style="list-style-type: none"> Inhibition of platelet function, thrombosis and haemostasis <ul style="list-style-type: none"> <i>In vitro</i> and <i>in vivo</i> Human, mouse platelets 	<ul style="list-style-type: none"> RXR–Gq interaction and negative regulation of Rac activation to inhibit GPCR-mediated platelet activation Up-regulation of PKA activity and phosphorylation of VASP S157 in cAMP- and NFκβ-dependent manner (see Figure 3)
VDR ⁴⁵	<ul style="list-style-type: none"> Vitamin D and its metabolites 	<ul style="list-style-type: none"> Low vitamin D plasma levels cause high mean platelet volume, a marker of platelet hyperactivity <ul style="list-style-type: none"> <i>In vivo</i> Human platelets 	Mechanism is unknown

attenuate platelet function.^{27,28} Prednisolone-treated platelets displayed reduced aggregation and thromboxane B₂ (TxB₂) release in response to stimulation by either ADP or the TxA₂ mimetic U46619, which was reversed following treatment with a GR antagonist mifepristone.²⁷ This inhibition was not found to be associated with up-regulation of cyclic nucleotides—cyclic adenosine monophosphate (cAMP) or cyclic guanosine monophosphate (cGMP), key inhibitory mediators of platelet activity.²⁸ Platelet adhesion and thrombus formation under flow on collagen *in vitro* were also found to be diminished following prednisolone treatment which is likely an outcome of reduced platelet responses to ADP and TxA₂, secondary mediators of platelet activation that support platelet adhesion and thrombus growth.²⁸ Prednisolone has also shown to modulate platelet–monocyte interactions following stimulation by ADP, which is attributed to an attenuation of platelet activity and not to inhibition of monocytes.²⁸ It should be noted, however, that alternative GR ligands—dexamethasone, fludrocortisone, and triamcinolone have not been shown to elicit anti-platelet effects under the experimental conditions used in these studies.^{27,28} This difference in activation is thought to be due to the formation of a heterodimeric complex between GR and the mineralocorticoid receptor that is susceptible to the differential activation by specific receptor ligands. The mechanism underlying the negative regulation of secondary mediator signalling by GR and its ligand prednisolone is yet to be fully explored although evidence suggests that this might be mediated through the regulation of signalling events downstream of the P2Y₁₂ receptor.²⁸

3.2 Oestrogen receptor

Oestradiol-17β (E2) and ERs are not only well known for their role in reproductive and sexual development but also known to directly influence cardiovascular health.⁵³ Human platelets have been shown to express ERβ but not ERα.²² Studies investigating the effects of several forms of oestrogen, including oestrone (E1), oestradiol (E2), and oestril (E3) on platelet function have yielded conflicting results. In one study, acute treatment of platelets *ex vivo* with either E1 or E3 was found to increase aggregation to adrenaline or ADP.²⁴ In contrast, chronic treatment with oestrogen, in an alternative study, investigating oestrogen replacement therapy (3 months), found a significant decrease in adrenaline-induced platelet aggregation and ATP release in patients receiving the therapy compared to the control groups.²⁵ In further support of this, chronic treatment with high levels of oestradiol in mice was

found to cause a marked decrease in platelet responsiveness both *ex vivo* and *in vivo*, with both an increase in bleeding time and resistance to thromboembolism being observed.²⁶ However, it is important to note these effects on platelet reactivity are due to modulation of expression of platelet proteins (such as β1 tubulin) during haematopoiesis that then alter platelet production and activation,²⁶ rather than a direct non-genomic effect on platelet function.

3.3 Androgen receptor

The AR, activated by either testosterone or dihydrotestosterone, has been identified in platelets,²² but little is known regarding its potential role in the regulation of platelet function. Some studies have specified that the aggregation response of platelets isolated from male rats was stronger in comparison to female rats owing to higher levels of androgenic steroids⁴⁶ as platelet aggregation was found to be reduced following castration in male rats and the reversal of these effects following treatment with testosterone.⁴⁷ Pilo *et al.* also reported that acute treatment of rat or human PRP with testosterone potentiates platelet aggregation induced by ADP, adrenaline, collagen, arachidonic acid, and calcium ionophore indicating its rapid non-genomic responses.⁴⁸ Two independent studies also confirmed that testosterone causes a significant increase in TXA₂ receptor density on the platelet surface, thereby, indirectly increasing platelet responsiveness.^{54,55} However, inhibition of platelet aggregation has also been observed following treatment with testosterone, although this was found to be attributed to endothelial NO synthesis and therefore not necessarily a direct effect of testosterone on the platelet AR and platelet activity.²³

4. Type II NRs

4.1 Liver X receptor

LXR receptors are implicated in the regulation of fatty acid, cholesterol, and glucose homeostasis. Endogenous ligands for the LXR receptors include oxysterols such as 22(R)-hydroxycholesterol, 24(S)-hydroxycholesterol, 27-hydroxycholesterol, and several synthetic ligands including GW3965 and T0901317 have also been developed.^{56,57} Like some of the other NRs, ligands for LXR have been described to have anti-inflammatory and atheroprotective properties, and LXRβ has been shown to be expressed in platelets.³¹ Treatment of platelets with the

synthetic agonist GW3965 results in inhibition of platelet activation, with attenuation of aggregation, calcium mobilization, secretion and integrin activation observed following stimulation by collagen, collagen-related peptide (CRP-XL; a GPVI collagen receptor-specific agonist), and thrombin. In analysis of thrombosis, GW3965-treated mice were also found to form smaller, less stable thrombi following laser injury of the cremaster arterioles. LXR has also been shown to interact with several components of the GPVI signalling pathway following treatment with GW3965, including Syk and PLC γ 2, and treatment with LXR ligands is associated with decreased phosphorylation and signalling.³¹ In support of this, another study reported the ability of endogenous LXR ligand 22(R)-OH-cholesterol [but not its stereoisomer 22(S)-OH-cholesterol] to inhibit collagen-induced platelet aggregation and shape change.⁵⁸

During thrombus formation, two distinct populations of platelets appear, coaggregated platelets, which support thrombus growth, and loosely attached procoagulant platelets that expose phosphatidylserine

and support coagulation. Conversion to the procoagulant state is also thought to be associated with platelet hyper-reactivity, a trait often observed in patients with an increased risk of thrombosis including those with pathological conditions, such as hyperlipidaemia, obesity, and high plasma cholesterol levels. Treatment of platelets with LXR ligands, GW3965 and T0901317, and natural ligands, 27-OH-cholesterol and 24-(S)-hydroxyl-cholesterol, has also been shown to cause platelet inhibition to several agonists through the conversion of platelets to procoagulant coated platelets.³⁹ LXR ligand-stimulated coated platelets not only expose phosphatidylserine at the membrane surface but also retain high levels of fibrinogen (which is converted to fibrin) and other alpha granule components at the platelet membrane (Figure 1). Conversion to the coated platelet state is thought to support coagulation but renders the platelet, through closure of integrin α IIb β 3, unresponsive to platelet agonists, which was also observed in platelets following treatment with LXR agonists. The mechanism by which this occurs in LXR agonist (GW3965)-




Figure 1 LXR and FXR ligands negatively regulate platelet function through inhibition of platelet signalling and formation of procoagulant coated platelets. Treatment of platelets with LXR ligands results in reduced tyrosine phosphorylation of key GPVI signalling molecules Syk, LAT and PLC γ 2. An increase in the level of LXR–Syk and LXR–PLC γ 2 interaction is also observed. Exposure to LXR ligands also renders platelets into a procoagulant state characterized by the exposure of phosphatidylserine and α -granule contents on the platelet surface, which is coupled with depolarization of the mitochondrial membrane potential, reduced calcium mobilization and down-regulation in the affinity of integrin α IIb β 3, ultimately resulting in the inhibition of platelet aggregation. Similarly, incubation of platelets with FXR ligands can lead to platelet swelling and conversion to procoagulant coated platelets which is dependent on cyclophilin D activity. Additionally, FXR ligands are able to increase cGMP levels that promotes the activity of PKG and phosphorylation of VASP S239 and thereby suppresses platelet activation.

treated platelets appears to be via deregulation of intracellular calcium signalling, depolarization of the mitochondrial membrane potential independently of cyclophilin D, and generation of reactive oxygen species (ROS).³⁹ It is therefore possible that the platelet dysfunction observed in patients with high cholesterol, hyperlipidaemia, metabolic syndrome, and obesity could be attributed to altered LXR signalling in platelets.

4.2 Farnesoid X receptor

The bile acid receptor, FXR, which is recognized to regulate bile acid and cholesterol homeostasis has been identified in both human and mouse platelets. Treatment of platelets with synthetic FXR ligand GW4064 was found to cause a decrease in sample turbidity,^{29,39} which was later confirmed to be due to platelet swelling and conversion of platelets to a pro-coagulant state, forming coated-platelets.³⁹ Synthetic and natural FXR ligand-dependent formation of coated platelets, prior to platelet agonist stimulation, results in phosphatidylserine exposure, retention of fibrinogen, fibrin and alpha granule proteins at the platelet surface, cyclophilin D-dependent depolarization of the mitochondrial membrane, sustained calcium signalling, generation of reactive oxygen species, and closure of integrins at the platelet surface.³⁹ This closure of platelet integrins is believed to underlie the observed reduction in platelet aggregation to platelet agonists. Although the initial kinetics of thrombus formation was increased in mouse *in vivo* models of thrombosis, consistent with a pro-coagulant state, thrombus stability was significantly decreased following treatment with the FXR ligand GW4064, in agreement with reduced integrin function, which is essential for stable thrombus formation.²⁹ Treatment with FXR ligands was also found to be associated with an increase in intracellular levels of cGMP in platelets, indicative of deregulation of intracellular signalling (Figure 1). Platelets from FXR-deficient mice

were found to be unresponsive to the actions of FXR agonists, confirming the selective non-genomic actions of these ligands to the FXR.²⁹

4.3 Peroxisome proliferator-activated receptors

PPARs represent three NR isoforms, PPAR α , PPAR β , and PPAR γ , which are involved in cell development, differentiation, cholesterol and fatty acid metabolism, and glucose homeostasis. All three isoforms of PPARs, upon binding to their ligands, are capable of heterodimerizing with the RXR⁴³ and all have been identified to have acute, non-genomic, negative-regulatory effects in human platelets.

4.3.1 PPAR α

The treatment of platelets with ligands of PPAR α such as fenofibrate or statins (simvastatin) has been shown to inhibit ADP-stimulated platelet activation by increasing intracellular levels of cAMP via a PPAR α -dependent mechanism. In support of this, the observed inhibition can be reversed following treatment with PPAR α antagonist GW6471.³³ This dependence on PPAR α is further reinforced by experiments that show fenofibrate-induced inhibition of platelet activation and increased bleeding time in mice does not occur in mice deficient in PPAR α . Fenofibrate-induced inhibition of platelet activity was found to be mediated through up-regulation of cAMP levels via inhibition of PKC α , a key mediator of platelet signalling, through interaction between PPAR α and PKC α . This interaction is believed to sequester PKC α away from its substrates and thereby attenuates platelet functions (Figure 2).³³ These findings identify PPAR α as a key mediator of statin and fenofibrate-mediated anti-platelet activity.




Figure 2 Non-genomic regulation of platelets by PPAR ligands. PPAR α ligands, fenofibrate or simvastatin and PPAR β/δ ligands, GW0742 and L-165041 cause a reduction in intracellular calcium mobilization and platelet activation. This inhibition was found to be mediated by augmented levels of cAMP and attenuation of PKC α through its interaction with PPAR α or PPAR β/δ , which limits its availability to facilitate signalling downstream of PKC α . Treatment with PPAR γ ligands inhibits platelet activation to collagen through an inhibition in phosphorylation of Syk and LAT that mediate signalling initiated by the collagen receptor GPVI. Negative regulation of integrin $\alpha_{IIb}\beta_3$ outside-in signalling was observed as an outcome of up-regulation of PKA activity and inhibition in phosphorylation of β_3 and subsequent downstream signalling molecules—Syk, PLC γ_2 , PKC substrates, FAK and PI3K substrates.

4.3.2 PPAR β/δ

Studies using synthetic ligands for PPAR β/δ , GW0742, and L-165041 have identified negative-regulation of platelet activity arbitrated through PPAR β/δ . Incubation with PPAR β/δ ligands showed inhibition of platelet aggregation and mobilization of intracellular calcium following stimulation by several platelet agonists.³⁴ PPAR β/δ can also be activated by the prostaglandin PGI₂, and therefore some of the inhibitory effects of PGI₂ on platelet activity could also be mediated through PPAR β/δ in addition to the prostaglandin IP receptor but this has yet to be tested.³⁴ Similar to PPAR α , treatment of platelets with synthetic ligands of PPAR β/δ have been shown to cause an increase in intracellular cAMP levels and PKC α has been identified as a potential binding partner of the receptor indicating a plausible mechanism by which PPAR β/δ regulates platelet reactivity (Figure 2)³⁵ PPAR β/δ ligands have been shown to decrease plaque formation and attenuate the progression of atherosclerosis.⁵⁹ As platelets play a key role in the initiation and progression of atherosclerosis, anti-platelet effects of PPAR β/δ ligands may partly explain such observed reduction in the development of atherosclerosis.

4.3.3 PPAR γ

PPAR γ is the most widely studied of the PPAR family in platelets. This is mainly because of its direct involvement with numerous cardiovascular diseases, such as diabetes mellitus, atherosclerosis, and thrombosis.^{60–62} Synthetic ligands of PPAR γ , the thiazolidinediones (pioglitazone, rosiglitazone, lobeglitazone, etc.), are currently in use for the treatment of type 2 diabetes and have been observed clinically to have cardioprotective properties. The anti-platelet activity of PPAR γ ligands may provide a mechanistic basis that in part underlies these observations. For example, a clinical study conducted on patients suffering from coronary heart disease and taking rosiglitazone reported its long-term antiplatelet effects with down-regulation of P-selectin exposure and granule secretion.⁶³ Exposure of platelets *ex vivo* to the endogenous (15d-PGJ₂) and synthetic (rosiglitazone and ciglitazone) ligands of PPAR γ has been shown to inhibit platelet activation to a variety of platelet agonists, including the G-protein-coupled receptor agonists—thrombin and ADP,³² GPVI agonists—collagen and CRP-XL,³⁸ and the adhesion receptor integrin $\alpha_{IIb}\beta_3$ agonist fibrinogen.³⁹ PPAR γ ligands, 15d-PGJ₂ or rosiglitazone, inhibit platelet responses including granule secretion and TxB₂ synthesis in response to thrombin or ADP.³² These ligands have also been shown to reduce GPVI agonist-stimulated platelet aggregation, granule secretion, and mobilization of intracellular calcium, via inhibition of early GPVI signalling events such as phosphorylation of Syk and LAT.³⁸ PPAR γ was also found to interact with Syk and LAT upon stimulation with collagen in the absence of PPAR γ ligands; however, this interaction is disrupted on treatment with PPAR γ ligands. These ligands have also been shown to inhibit integrin $\alpha_{IIb}\beta_3$ outside-in signalling through the up-regulation of PKA activity. PPAR γ ligand-dependent inhibition of β_3 phosphorylation and other downstream signalling molecules of the integrin $\alpha_{IIb}\beta_3$ signalling pathway including Syk, PLC γ_2 , PKC, FAK, and PI3K indicates several different mechanisms by which PPAR γ ligands can negatively regulate platelet function.³⁹ This negative regulation of platelet activity has also been found to result in an inhibition of thrombus formation *in vivo* in animal models following treatment with another synthetic PPAR γ ligand, pioglitazone.³⁷

PPAR γ is also implicated in a mechanism by which statins mediate acute anti-platelet effects.^{33,36} Treatment of human whole blood with simvastatin has been shown to cause a reduction in platelet aggregation to ADP. This inhibition of platelet function was attributed to an increase

in intracellular cAMP levels which is associated with PPAR γ activity and its association with and inhibition of PKC α (Figure 2).³³ In addition, treatment of platelets with simvastatin was also found to inhibit collagen-induced platelet aggregation, granule secretion, integrin activation, and Ca²⁺ mobilization in a PPAR γ -dependent manner. This was found to involve PPAR γ -dependent mediation of mitogen-activated protein kinase (MAPKs, i.e. p38 MAPK, ERK) signalling by increasing association of MAPKs with the receptor resulting in an increase of cAMP formation that is associated with an increase in VASP Ser157 phosphorylation and inhibition of Akt phosphorylation.³⁶

4.4 Retinoic acid receptor

RARs play a critical role in numerous biological processes, including development, reproduction, immunity, organogenesis, and homeostasis.⁶⁴ Three forms of RAR exist—RAR α , RAR β , and RAR γ . Of these, RAR α is ubiquitously distributed and has been reported to be robustly expressed in human platelets, whilst the other two isoforms have tissue-specific distribution.⁶⁵ RARs are activated by retinoids which are metabolites of vitamin A and several synthetic ligands also exist.⁶⁴ In platelets, RAR α has been observed to directly interact with actin-related protein-2/3 complex (Arp2/3) subunit 5 (Arp2/3s5) which is required for the regulation of platelet cytoskeletal processes. Treatment of platelets with the endogenous RAR α ligand all-*trans*-retinoic acid (atRA) disrupts the RAR α –Arp2/3 interactions resulting in an inhibition of both cytoskeletal rearrangements and platelet spreading (Figure 3).⁴¹ Recent developments have reported that RAR α is capable of regulating protein synthesis (including microtubule-associated protein-1 light chain 3 beta 2) in human platelets by binding to a subset of mRNAs and blocking translation. Schwertz *et al.*⁶⁶ found that platelets treated with RAR α ligand atRA for several hours displayed significantly altered levels of protein synthesis compared to controls.

4.5 Retinoid X receptor

RXR is regarded as one of the most important receptors in this superfamily. Most likely due to its ability to interact with almost a quarter of the known human NRs (PPAR's, LXR, FXR, PXR, etc.) and form heterodimers,⁶⁷ although, the presence of RXR homodimers has also been reported.⁶⁸ RXR is well characterized and is involved in the regulation of some of the most vital and fundamental biological processes including cell proliferation, differentiation and death, haematopoiesis, metabolism (glucose, fatty acid, and cholesterol), and pattern formation during embryogenesis.⁶⁹ Human platelets have been shown to express RXR α and RXR β (but the presence or absence of RXR γ has not been established) and are known to form heterodimers with PPAR α , PPAR γ , and LXR in platelets.⁴³ Treatment of platelets with the endogenous ligand of RXR, 9-*cis*-retinoic acid or the synthetic ligand, methoprene acid, results in inhibition of platelet function stimulated by Gq-coupled GPCRs—ADP, U46619⁴² or thrombin and also, GPVI-mediated platelet activation via stimulation by collagen or CRP-XL.⁴³ Regulation of GPCR-mediated platelet activation by RXR has been associated with its binding to G α_q in a ligand-dependent manner that inhibits Gq-induced Rac activation and intracellular Ca²⁺ mobilization.⁴² Exposure to RXR ligands has also been shown to reduce integrin $\alpha_{IIb}\beta_3$ outside-in signalling and cytoskeletal rearrangements. The negative regulation of several platelet activation pathways and processes results in robust inhibition of thrombosis and haemostasis *in vivo*.⁴³ As seen with several other NRs, treatment with RXR ligands has also been shown to up-regulate PKA activity and VASP S157 phosphorylation via a process that is dependent on cAMP and also involves NF κ B (Figure 3). This suggests that RXR ligands inhibit platelet function using several different




Figure 3 Inhibition of platelet function by RAR and RXR ligands is mediated through Arp2/3- and Gq-induced Rac activation and up-regulation of PKA activity, respectively. RAR ligand atRA disrupts the RAR α –Arp2/3 interaction resulting in an inhibition of cytoskeletal rearrangements and platelet spreading. RXR ligands, 9cRA and methoprene acid inhibit platelet activation to a range of platelet agonists that include GPCR agonists (ADP, U46619 or thrombin) and GPVI agonists (collagen or CRP-XL). Interaction of RXR with Gq and subsequent negative regulation of Rac activation is one of the probable explanations for the reduction in GPCR-mediated platelet activation. These ligands have also been shown to up-regulate PKA activity in a cAMP- and NF κ B-dependent manner providing a more generalized mechanism of inhibition.

inhibitory mechanisms which is likely to reflect its ability to form heterodimers with several different NRs in platelets.⁴³

4.6 Vitamin D receptor

The VDR is another ligand-activated transcription factor that mediates the actions of vitamin D and its metabolites. VDR is also known to form a heterodimer with the RXR and regulate calcium homeostasis, cell growth and differentiation, detoxification of xenobiotics, and modulation of adaptive and innate immunity.⁷⁰ Although anticoagulant effects of vitamin D have been reported and VDR signalling has been characterized in monocytes and vascular cells, the role for the VDR in platelet function remains unknown. Human platelets have been found to express the VDR. Biochemical fractionation studies along with immuno-electron microscopy analysis identified the VDR to be localized in the soluble and mitochondrial compartment.⁴⁴ Although little is known about the role for vitamin D and the VDR in platelet function, a patient study identified a strong association between low vitamin D plasma levels and a high mean platelet volume, a marker of platelet hyperactivity.⁴⁵

5. Future perspectives

5.1 Could NRs offer anti-platelet therapeutic targets?

The role of platelets in controlling haemostasis and initiating thrombosis is well known. This makes them important therapeutic targets for the treatment of cardiovascular diseases, particularly atherothrombosis.⁷¹ Despite significant advances in the development of antithrombotic therapeutics, they are associated with increased bleeding risk and their efficacy

is often compromised in patients suffering from several conditions, such as hypertension and diabetes.^{11,72,73} Therefore, more refined and effective therapeutics that ensure a balance between the treatment of thrombosis and related complications is needed.

A key step forward would be using our current knowledge of the molecular mechanisms governing platelet functions as the basis for the development of more effective and safer anti-platelet therapies. Both natural and synthetic ligands of NRs have been shown to exhibit non-genomic effects to alter platelet function through a variety of mechanisms, several of which appear to be shared by different NR family members.^{21,74} For instance, RXR, FXR, PPAR α , PPAR β , and PPAR γ receptors have been shown to be involved in the regulation of platelet inhibitory signalling pathways by either increasing cAMP or cGMP levels or directly modulating PKA/PKG activity. LXR and PPAR γ can negatively regulate signalling downstream of the collagen receptor via interactions with different components of the GPVI signalling cascade. Given the potential of heterodimeric receptor interactions between RXR and other NRs, the idea of cross-talk becomes even more pronounced.⁴³

5.2 Important considerations

Development of NRs as anti-platelet therapeutic targets requires a few important considerations. First, the majority of the studies conducted so far focus on understanding the acute effects of NR ligands on platelet function; therefore, prior to further development it would be important to study the implications of chronic exposure of platelets to NR ligands. As NRs can regulate the expression of multiple genes, in various cell types it is therefore highly likely that chronic exposure to NR ligands could lead to systemic effects that might indirectly affect platelet function.⁷⁵

Table 3 Commercially available nuclear receptor drugs

Nuclear receptor	Disease	Drug generic name (marketed drug)
GR	Metabolic and immunological Disorders	Dexamethasone (Dexasone), Prednisolone (Orapred) ^{94,95}
ER	Breast cancer, obesity	Tamoxifen (Nolvadex), Raloxifene (Evista) ^{96,97}
PPAR α	Dyslipidaemia, atherosclerosis	Fenofibrate (Tricor) ⁹⁸
PPAR γ	Diabetes, obesity	Pioglitazone (Actos), Rosiglitazone (Avandia) ^{99,100}
RAR	Leukaemia, acne	13- <i>cis</i> -retinoic acid (Isotretinoin) ¹⁰¹
RXR	Leukaemia, Kaposi sarcoma, eczema	9- <i>cis</i> -retinoic acid (Alitretinoin), Bexarotene (Targretin) ^{102–104}
VDR	Osteoporosis, calcium homeostasis	Calcitriol (Calcijex), Paricalcitol (Zemplar) ^{105,106}

Secondly, although, there exists a clear distinction between genomic and non-genomic effects, the existence of mRNA in platelets and their limited ability to perform translation raises the possibility^{17,76} that there are interactions between NRs and mRNA in platelets as in nucleated cells. While the differences in the timescales taken to elicit these genomic-like effect (hours) in comparison to the non-genomic effects (minutes) still enable the differentiation between the two regulatory mechanisms. Future studies should consider including inhibitors of translation which will help further differentiate between genomic and truly non-genomic actions of these receptors, or indeed determine whether NR ligands are capable of regulating protein translation in platelets and characterizing whether any changes in protein levels have functional effects. Schwertz *et al.*⁶⁶ recently described such a mechanism demonstrating RAR α -dependent translational control in human platelets, which resulted in the synthesis of several transcripts. Whether other NRs (such as RXR and PPARs), identified in platelets, can also replicate such a mechanism is still unknown. Moreover, evaluating whether genomic and non-genomic regulation can facilitate cross-talk between the different NRs in platelets requires further investigation. Development of NRs as anti-platelet therapies would require careful balancing of their genomic vs. non-genomic effects not only in platelets but also systemically.

Finally, it is important to note that NRs share a significant level of structural similarity with each other, making them potentially promiscuous in nature.^{77–81} Studies examining the genomic roles for the NRs have shown, for example that 15d-PGJ₂ is an endogenous ligand for PPAR γ but it can also act as an antagonist for FXR⁸² and phytanic acid has the ability to activate both PPAR α and RXR.⁸³ Guggulsterone is regarded as an ER agonist but an antagonist to FXR, GR, and AR,⁸⁴ although it does not appear to function as a non-genomic FXR antagonist in platelets (L.A.M, A.J.U, J.M.G, unpublished observations). Similarly, LG100754 is a highly specific RXR: PPAR γ agonist while it acts as a strong antagonist of RXR homodimers.⁸⁵ This makes the selective targeting of the NRs even more challenging and as such identification of ligands that function in a receptor- and gene-specific manner is important. Future work will be required to establish how and when NR heterodimers regulate platelet activity, and for each NR to establish its role in normal physiological processes.

6. Conclusions

Platelets are known to act as direct contributors towards the progression of CVDs such as atherosclerosis,⁸⁶ and their activity becomes considerably enhanced in cases of hyperlipidaemia,⁸⁷ obesity,⁸⁸ diabetes

mellitus,⁸⁹ or hypertension.⁹⁰ Many NRs have been found to be expressed in human platelets, including AR, ER, GR, FXR, LXR, PPARs, RAR, RXR, and VDR, and agonists for several of these receptors have been shown to elicit anti-platelet effects by a variety of mechanisms. NRs including PPARs, LXR, and FXR ligands have all been reported to have anti-atherosclerotic effects,^{91,92} coupling this with their anti-platelet effects; there exists the possibility of a potentially new paradigm of treatment that can target a range of pathophysiological conditions whilst also offering platelet-targeted anti-thrombotic activity. Of FDA-approved drugs, 13% function by targeting NRs for the treatment of numerous pathological conditions⁹³ (Table 3), and as a result, the effects on platelets might be a likely consequence associated with the administration of these drugs. It is important for the future development and use of NR agonists that their acute and long-term effects on platelet function are fully understood.

Authors' contributions

A.J.U, G.D.F, and J.M.G wrote the review. A.J.U and G.D.F contributed equally.

Conflict of interest: none declared.

Funding

This work was supported by the British Heart Foundation (RG/15/2/31224) and a Felix Scholarship.

References

- Bain DL, Heneghan AF, Connaghan-Jones KD, Miura MT. Nuclear receptor structure: implications for function. *Annu Rev Physiol* 2007;**69**:201–220.
- Kiss M, Czimmerer Z, Nagy L. The role of lipid-activated nuclear receptors in shaping macrophage and dendritic cell function: from physiology to pathology. *J Allergy Clin Immunol* 2013;**132**:264–286.
- Khan S, Lingrel JB. Thematic minireview series on nuclear receptors in biology and diseases. *J Biol Chem* 2010;**285**:38741–38742.
- Falkenstein E, Norman AWW, Wehling M. Mannheim classification of nongenomically initiated (rapid) steroid action(s). *J Clin Endocrinol Metab* 2000;**85**:2072–2075.
- Hammes SR, Levin ER. Extranuclear steroid receptors: nature and actions. *Endocr Rev* 2007;**28**:726–741.
- Losel RM, Falkenstein E, Feuring M, Schultz A, Tillmann HC, Rossol-Haseroth K, Wehling M. Nongenomic steroid action: controversies, questions, and answers. *Physiol Rev* 2003;**83**:965–1016.
- Nadal A, Diaz M, Valverde MA. The estrogen trinity: membrane, cytosolic, and nuclear effects. *News Physiol Sci* 2001;**16**:251–255.
- Boonyaratankornkit V, Edwards DP. Receptor mechanisms mediating non-genomic actions of sex steroids. *Semin Reprod Med* 2007;**25**:139–153.

9. Ordóñez-Morán P, Muñoz A. Nuclear receptors: genomic and non-genomic effects converge. *Cell Cycle (Georgetown, Tex)* 2009;**8**:1675–1680.
10. McKenna NJ, O'Malley BW. Combinatorial control of gene expression by nuclear receptors and coregulators. *Cell* 2002;**108**:465–474.
11. Nathan AS, Sen S, Yeh RW. The risk of bleeding with the use of antiplatelet agents for the treatment of cardiovascular disease. *Expert Opin Drug Saf* 2017;**16**:561–572.
12. Boilard E, Nigrovic PA, Larabee K, Watts GFM, Coblyn JS, Weinblatt ME, Massarotti EM, Remold-O'Donnell E, Fardale RW, Ware J, Lee DM. Platelets amplify inflammation in arthritis via collagen-dependent microparticle production. *Science* 2010;**327**:580–583.
13. Morrell CN, Aggrey AA, Chapman LM, Modjeski KL. Emerging roles for platelets as immune and inflammatory cells. *Blood* 2014;**123**:2759–2767.
14. Santilli F, Simeone P, Liani R, Davi G. Platelets and diabetes mellitus. *Prostaglandins Other Lipid Mediat* 2015;**120**:28–39.
15. Borsig L. The role of platelet activation in tumor metastasis. *Expert Rev Anticancer Ther* 2008;**8**:1247–1255.
16. Schubert S, Weyrich AS, Rowley JW. A tour through the transcriptional landscape of platelets. *Blood* 2014;**124**:493–502.
17. Rowley JW, Schwertz H, Weyrich AS. Platelet mRNA: the meaning behind the message. *Curr Opin Hematol* 2012;**19**:385.
18. Cecchetti L, Tolley ND, Michetti N, Bury L, Weyrich AS, Gresle P. Megakaryocytes differentially sort mRNAs for matrix metalloproteinases and their inhibitors into platelets: a mechanism for regulating synthetic events. *Blood* 2011;**118**:1903–1911.
19. Risitano A, Beaulieu LM, Vitseva O, Freedman JE. Platelets and platelet-like particles mediate intercellular RNA transfer. *Blood* 2012;**119**:6288–6295.
20. Bishop-Bailey D. The platelet as a model system for the acute actions of nuclear receptors. *Steroids* 2010;**75**:570–575.
21. Jones CI, Barrett NE, Moraes LA, Gibbins JM, Jackson DE. Endogenous inhibitory mechanisms and the regulation of platelet function. *Methods Mol Biol* 2012;**788**:341–366.
22. Khetawat G, Faraday N, Nealen ML, Vijayan KV, Bolton E, Noga SJ, Bray PF. Human megakaryocytes and platelets contain the estrogen receptor beta and androgen receptor (AR): testosterone regulates AR expression. *Blood* 2000;**95**:2289–2296.
23. Campelo AE, Cutini PH, Massheimer VL. Testosterone modulates platelet aggregation and endothelial cell growth through nitric oxide pathway. *J Endocrinol* 2012;**213**:77–87.
24. Akaraseenont P, Tripatara P, Chotewuttakorn S, Palo T, Thaworn A. The effects of estrone, estradiol and estriol on platelet aggregation induced by adrenaline and adenosine diphosphate. *Platelets* 2006;**17**:441–447.
25. Bar J, Tepper R, Fuchs J, Pardo Y, Goldberger S, Ovadia J. The effect of estrogen replacement therapy on platelet aggregation and adenosine triphosphate release in postmenopausal women. *Obstet Gynecol* 1993;**81**:261–264.
26. Valera MC, Gratacap MP, Gourdy P, Lenfant F, Cabou C, Toutain CE, Marcellin M, Saint Laurent N, Sie P, Sixou M, Arnal JF, Payrastre B. Chronic estradiol treatment reduces platelet responses and protects mice from thromboembolism through the hematopoietic estrogen receptor alpha. *Blood* 2012;**120**:1703–1712.
27. Moraes LA, Paul-Clark MJ, Rickman A, Flower RJ, Goulding NJ, Perretti M. Ligand-specific glucocorticoid receptor activation in human platelets. *Blood* 2005;**106**:4167–4175.
28. Liverani E, Banerjee S, Roberts W, Naseem KM, Perretti M. Prednisolone exerts exquisite inhibitory properties on platelet functions. *Biochem Pharmacol* 2012;**83**:1364–1373.
29. Moraes LA, Unsworth AJ, Vajyapuri S, Ali MS, Sasikumar P, Sage T, Flora GD, Bye AP, Kriek N, Dorchie E, Molendi-Coste O, Dombrowicz D, Staels B, Bishop-Bailey D, Gibbins JM. Farnesoid X receptor and its ligands inhibit the function of platelets. *Arterioscler Thromb Vasc Biol* 2016;**36**:2324–2333.
30. Unsworth AJ, Bye AP, Tannetta DS, Desbrouwer MJR, Kriek N, Sage T, Allan HE, Crescente M, Yaqoob P, Warner TD, Jones CI, Gibbins JM. Farnesoid X receptor and liver X receptor ligands initiate formation of coated platelets. *Arterioscler Thromb Vasc Biol* 2017;**37**:1482–1493.
31. Spyridon M, Moraes LA, Jones CI, Sage T, Sasikumar P, Bucci G, Gibbins JM. LXR as a novel antithrombotic target. *Blood* 2011;**117**:5751–5761.
32. Akbiyik F, Ray DM, Gettings KF, Blumberg N, Francis CW, Phipps RP. Human bone marrow megakaryocytes and platelets express PPARgamma, and PPARgamma agonists blunt platelet release of CD40 ligand and thromboxanes. *Blood* 2004;**104**:1361–1368.
33. Ali FY, Armstrong PC, Dhanji AR, Tucker AT, Paul-Clark MJ, Mitchell JA, Warner TD. Antiplatelet actions of statins and fibrates are mediated by PPARs. *Arterioscler Thromb Vasc Biol* 2009;**29**:706–711.
34. Ali FY, Davidson SJ, Moraes LA, Traves SL, Paul-Clark M, Bishop-Bailey D, Warner TD, Mitchell JA. Role of nuclear receptor signaling in platelets: antithrombotic effects of PPARbeta. *FASEB J* 2006;**20**:326–328.
35. Ali FY, Hall MG, Desvergne B, Warner TD, Mitchell JA. PPARbeta/delta agonists modulate platelet function via a mechanism involving PPAR receptors and specific association/repression of PKCalpha—brief report. *Arterioscler Thromb Vasc Biol* 2009;**29**:1871–1873.
36. Du H, Hu H, Zheng H, Hao J, Yang J, Cui W. Effects of peroxisome proliferator-activated receptor gamma in simvastatin antiplatelet activity: influences on cAMP and mitogen-activated protein kinases. *Thromb Res* 2014;**134**:111–120.
37. Li D, Chen K, Sinha N, Zhang X, Wang Y, Sinha AK, Romeo F, Mehta JL. The effects of PPAR-gamma ligand pioglitazone on platelet aggregation and arterial thrombus formation. *Cardiovasc Res* 2005;**65**:907–912.
38. Moraes LA, Spyridon M, Kaiser WJ, Jones CI, Sage T, Atherton RE, Gibbins JM. Non-genomic effects of PPARgamma ligands: inhibition of GPVI-stimulated platelet activation. *J Thromb Haemost* 2010;**8**:577–587.
39. Unsworth A, Kriek N, Bye A, Naran K, Sage T, Flora G, Gibbins JM. PPARgamma agonists negatively regulate alphaIIb beta3 integrin outside-in signaling and platelet function through up-regulation of protein kinase A activity. *J Thromb Haemost* 2017;**15**:356–369.
40. Unsworth AJ, Kriek N, Bye AP, Naran K, Sage T, Flora GD, Gibbins JM. PPARgamma agonists negatively regulate alphaIIb beta3 integrin outside-in signalling and platelet function through upregulation of protein kinase A activity. *J Thromb Haemost* 2017;**15**:356–369.
41. Rondina MT, Freitag M, Pluthero FG, Kahr WH, Rowley JW, Kraiss LW, Franks Z, Zimmerman GA, Weyrich AS, Schwertz H. Non-genomic activities of retinoic acid receptor alpha control actin cytoskeletal events in human platelets. *J Thromb Haemost* 2016;**14**:1082–1094.
42. Moraes LA, Swales KE, Wray JA, Damazo A, Gibbins JM, Warner TD, Bishop-Bailey D. Nongenomic signaling of the retinoid X receptor through binding and inhibiting Gq in human platelets. *Blood* 2007;**109**:3741–3744.
43. Unsworth AJ, Flora GD, Sasikumar P, Bye AP, Sage T, Kriek N, Crescente M, Gibbins JM. RXR ligands negatively regulate thrombosis and hemostasis. *Arterioscler Thromb Vasc Biol* 2017;**37**:812–822.
44. Silvagno F, De Vivo E, Attanasio A, Gallo V, Mazzucco G, Pescarmona G. Mitochondrial localization of vitamin D receptor in human platelets and differentiated megakaryocytes. *PLoS One* 2010;**5**:e8670.
45. Cumhuri Cure M, Cure E, Yuce S, Yazici T, Karakoyun I, Efe H. Mean platelet volume and vitamin D level. *Ann Lab Med* 2014;**34**:98–103.
46. Johnson M, Ramey E, Ramwell PV. Sex and age differences in human platelet aggregation. *Nature* 1975;**253**:355–357.
47. Johnson M, Ramey E, Ramwell P. Androgen-mediated sensitivity in platelet aggregation. *Am J Physiol Heart Circ Physiol* 1977;**232**:H381–H385.
48. Pilo R, Aharony D, Raz A. Testosterone potentiation of ionophore and ADP induced platelet aggregation: relationship to arachidonic acid metabolism. *Thromb Haemost* 1981;**46**:538–542.
49. Frye CA, Rhodes ME. The role and mechanisms of steroid hormones in approach-avoidance behavior. In Elliot AJ. (ed.). *Handbook of Approach and Avoidance Motivation*. Abingdon: Routledge; 2008. p109–126.
50. Richard A, Rohrmann S, Zhang L, Eichholzer M, Basaria S, Selvin E, Dobs AS, Kanarek N, Menke A, Nelson WG, Platz EA. Racial variation in sex steroid hormone concentration in black and white men: a meta-analysis. *Andrology* 2014;**2**:428–435.
51. Güncü GN, Tözüm TF, Çağlayan F. Effects of endogenous sex hormones on the periodontium—review of literature. *Aust Dent J* 2005;**50**:138–145.
52. Bledsoe RK, Montana VG, Stanley TB, Delves CJ, Apolito CJ, McKee DD, Consler TG, Parks DJ, Stewart EL, Willson TM, Lambert MH, Moore JT, Pearce KH, Xu HE. Crystal structure of the glucocorticoid receptor ligand binding domain reveals a novel mode of receptor dimerization and coactivator recognition. *Cell* 2002;**110**:93–105.
53. Murphy E. Estrogen signaling and cardiovascular disease. *Circ Res* 2011;**109**:687–696.
54. Ajayi AA, Mathur R, Halushka PV. Testosterone increases human platelet thromboxane A2 receptor density and aggregation responses. *Circulation* 1995;**91**:2742–2747.
55. Matsuda K, Ruff A, Morinelli TA, Mathur RS, Halushka PV. Testosterone increases thromboxane A2 receptor density and responsiveness in rat aortas and platelets. *Am J Physiol Heart Circ Physiol* 1994;**267**:H887–H893.
56. Gabbi C, Warner M, Gustafsson J-Å. Action mechanisms of Liver X Receptors. *Biochem Biophys Res Commun* 2014;**446**:647–650.
57. Wójcicka G, Jamroz-Wiśniewska A, Horoszewicz K, Beltowski J. Liver X receptors (LXRs). Part I: structure, function, regulation of activity, and role in lipid metabolism. Receptory wątrobowe X (LXR). Część I: budowa, funkcja, regulacja aktywności i znaczenie w metabolizmie lipidów. *Journal Postepy Hig Med Dosw* 2015;**61**:736–759.
58. Schaffer S, Tandon R, Zipse H, Siess W, Schmidt A, Jamasbi J, Karshovska E, Steglich W, Lorenz R. Stereoselective platelet inhibition by the natural LXR agonist 22 (R)-OH-cholesterol and its fluorescence labelling with preserved bioactivity and chiral handling in macrophages. *Biochem Pharmacol* 2013;**86**:279–285.
59. Lee CH, Chawla A, Urbiztondo N, Liao D, Boisvert WA, Evans RM, Curtiss LK. Transcriptional repression of atherogenic inflammation: modulation by PPARdelta. *Science* 2003;**302**:453–457.
60. Beckman JA, Creager MA, Libby P. Diabetes and atherosclerosis: epidemiology, pathophysiology, and management. *JAMA* 2002;**287**:2570–2581.
61. Chinetti G, Fruchart J-C, Staels B. Peroxisome proliferator-activated receptors (PPARs): nuclear receptors at the crossroads between lipid metabolism and inflammation. *Inflamm Res* 2000;**49**:497–505.
62. Moraes LA, Piqueras L, Bishop-Bailey D. Peroxisome proliferator-activated receptors and inflammation. *Pharmacol Ther* 2006;**110**:371–385.
63. Sidhu JS, Cowan D, Tooze JA, Kaski J-C. Peroxisome proliferator-activated receptor-gamma agonist rosiglitazone reduces circulating platelet activity in patients without diabetes mellitus who have coronary artery disease. *Am Heart J* 2004;**147**:1032–1037.

64. Duong V, Rochette-Egly C. The molecular physiology of nuclear retinoic acid receptors. From health to disease. *Biochim Biophys Acta* 2011;**1812**:1023–1031.
65. Dolle P. Developmental expression of retinoic acid receptors (RARs). *Nucl Recept Signal* 2009;**7**:e006.
66. Schwertz H, Rowley JW, Zimmerman GA, Weyrich AS, Rondina MT. Retinoic acid receptor- α regulates synthetic events in human platelets. *J Thromb Haemost* 2017;**15**:2408–2418.
67. Evans RM, Mangelsdorf DJ. Nuclear receptors, RXR, and the big bang. *Cell* 2014;**157**:255–266.
68. Sato Y, Ramalanjaona N, Huet T, Potier N, Osz J, Antony P, Peluso-Iltis C, Poussin-Courmontagne P, Ennifar E, Mély Y, Dejaegere A, Moras D, Rochel N. The “Phantom Effect” of the Retinoid LG100754: structural and functional insights. *PLoS One* 2010;**5**:e15119.
69. Ahuja H, Szanto A, Nagy L, Davies P. The retinoid X receptor and its ligands: versatile regulators of metabolic function, cell differentiation and cell death. *J Biol Regul Homeost Agents* 2003;**17**:29–45.
70. Kato S. The function of vitamin D receptor in vitamin D action. *J Biochem* 2000;**127**:717–722.
71. Badimon L, Padró T, Vilahur G. Atherosclerosis, platelets and thrombosis in acute ischaemic heart disease. *Eur Heart J Acute Cardiovasc Care* 2012;**1**:60–74.
72. Hankey GJ, Eikelboom JW. Aspirin resistance. *Lancet* 2006;**367**:606–617.
73. Ajan R, Grant PJ. The role of antiplatelets in hypertension and diabetes mellitus. *J Clin Hypertens* 2011;**13**:305–313.
74. Bye AP, Unsworth AJ, Gibbins JM. Platelet signaling: a complex interplay between inhibitory and activatory networks. *J Thromb Haemost* 2016;**14**:918–930.
75. Huang P, Chandra V, Rastinejad F. Structural overview of the nuclear receptor superfamily: insights into physiology and therapeutics. *Annu Rev Physiol* 2010;**72**:247–272.
76. Zimmerman GA, Weyrich AS. Signal-dependent protein synthesis by activated platelets new pathways to altered phenotype and function. *Arterioscler Thromb Vasc Biol* 2008;**28**:s17–s24.
77. Ng HW, Perkins R, Tong W, Hong H. Versatility or promiscuity: the estrogen receptors, control of ligand selectivity and an update on subtype selective ligands. *Int J Environ Res Public Health* 2014;**11**:8709–8742.
78. Kwon SY, Kim IS, Bae JE, Kang JW, Cho YJ, Cho NS, Lee SW. Pathogen inactivation efficacy of Mirasol PRT System and Intercept Blood System for non-leucoreduced platelet-rich plasma-derived platelets suspended in plasma. *Vox Sang* 2014;**107**:254–260.
79. Noy N. Ligand specificity of nuclear hormone receptors: sifting through promiscuity. *Biochemistry* 2007;**46**:13461–13467.
80. Sepe V, Festa C, Renga B, Carino A, Cipriani S, Finamore C, Masullo D, del Gaudio F, Monti MC, Fiorucci S, Zampella A. Insights on FXR selective modulation. Speculation on bile acid chemical space in the discovery of potent and selective agonists. *Sci Rep* 2016;**19**:19008.
81. Krasowski MD, Ni A, Hagey LR, Ekins S. Evolution of promiscuous nuclear hormone receptors: IXR, FXR, VDR, PXR, and CAR. *Mol Cell Endocrinol* 2011;**334**:39–48.
82. Xu X, Lu Y, Chen L, Chen J, Luo X, Shen X. Identification of 15d-PGJ2 as an antagonist of farnesoid X receptor: molecular modeling with biological evaluation. *Steroids* 2013;**78**:813–822.
83. Hellgren LI. Phytanic acid—an overlooked bioactive fatty acid in dairy fat? *Ann N Y Acad Sci* 2010;**1190**:42–49.
84. Burris TP, Montrose C, Houck KA, Osborne HE, Bocchinfuso WP, Yaden BC, Cheng CC, Zink RW, Barr RJ, Hepler CD, Krishnan V, Bullock HA, Burris LL, Galvin RJ, Bramlett K, Staybrook KR. The hypolipidemic natural product guggulsterone is a promiscuous steroid receptor ligand. *Mol Pharmacol* 2005;**67**:948–954.
85. Cesario RM, Klausung K, Razzaghi H, Crombie D, Rungta D, Heyman RA, Lala DS. The retinoid LG100754 is a novel RXR: pPAR γ agonist and decreases glucose levels in vivo. *Mol Endocrinol* 2001;**15**:1360–1369.
86. Schulz C, Massberg S. Platelets in atherosclerosis and thrombosis. *Handb Exp Pharmacol* 2012;**210**:111–133.
87. Wang N, Tall AR. Cholesterol in platelet biogenesis and activation. *Blood* 2016;**127**:1949–1953.
88. Trayhurn P, Wood IS. Signalling role of adipose tissue: adipokines and inflammation in obesity. *Biochem Soc Trans* 2005;**33**:1078–1081.
89. Schneider DJ. Factors contributing to increased platelet reactivity in people with diabetes. *Diabetes Care* 2009;**32**:525–527.
90. El Haouari M, Rosado JA. Platelet function in hypertension. *Blood Cells Mol Dis* 2009;**42**:38–43.
91. Calkin A, Tontonoz P. LXR signaling pathways and atherosclerosis. *Arterioscler Thromb Vasc Biol* 2010;**30**:1513.
92. Mencarelli A, Fiorucci S. FXR, an emerging therapeutic target for the treatment of atherosclerosis. *J Cell Mol Med* 2010;**14**:79–92.
93. Overington JP, Al-Lazikani B, Hopkins AL. How many drug targets are there? *Nat Rev Drug Discov* 2006;**5**:993–996.
94. Sundahl N, Bridelance J, Libert C, De Bosscher K, Beck IM. Selective glucocorticoid receptor modulation: new directions with non-steroidal scaffolds. *Pharmacol Ther* 2015;**152**:28–41.
95. Kadmiel M, Cidlowski JA. Glucocorticoid receptor signaling in health and disease. *Trends Pharmacol Sci* 2013;**34**:518–530.
96. Muchmore DB. Raloxifene: a selective estrogen receptor modulator (SERM) with multiple target system effects. *Oncologist* 2000;**5**:388–392.
97. Maximov PY, Lee TM, Craig Jordan V. The discovery and development of selective estrogen receptor modulators (SERMs) for clinical practice. *Curr Clin Pharmacol* 2013;**8**:135–155.
98. Filippatos T, Milionis HJ. Treatment of hyperlipidaemia with fenofibrate and related fibrates. *Expert Opin Investig Drugs* 2008;**17**:1599–1614.
99. Kersten S, Desvergne B, Wahli W. Roles of PPARs in health and disease. *Nature* 2000;**405**:421–424.
100. Ahmadian M, Suh JM, Hah N, Liddle C, Atkins AR, Downes M, Evans RM. PPAR [gamma] signaling and metabolism: the good, the bad and the future. *Nat Med* 2013;**9**:557–566.
101. Layton A. The use of isotretinoin in acne. *Dermatoendocrinol* 2009;**1**:162–169.
102. Walmsley S, Northfelt DW, Melosky B, Conant M, Friedman-Kien AE, Wagner B; Group PGNAS. Treatment of AIDS-related cutaneous Kaposi’s sarcoma with topical alitretinoin (9-cis-retinoic acid) gel. *J Acquir Immune Defic Syndr* 1999;**22**:235–246.
103. Ghasri P, Scheinfeld N. Update on the use of alitretinoin in treating chronic hand eczema. *Clin Cosmet Investig Dermatol* 2010;**3**:59–65.
104. Njar VCO. Retinoids in clinical use. In Ottow E, Weinmann H. (eds). *Nuclear Receptors as Drug Targets*. Weinheim, Germany: Wiley-VCH Verlag GmbH & Co. KGaA; 2008. p389–407.
105. Wu-Wong JR. Potential for vitamin D receptor agonists in the treatment of cardiovascular disease. *Br J Pharmacol* 2009;**158**:395–412.
106. Makishima M, Yamada S. Targeting the vitamin D receptor: advances in drug discovery. *Expert Opin Ther Pat* 2005;**15**:1133–1145.