

Cesarean Section Rate in Singleton Primiparae and Related Factors in Beijing, China

Geng Song¹, Yu-Mei Wei¹, Wei-Wei Zhu², Hui-Xia Yang¹

¹Department of Obstetrics and Gynecology, Peking University First Hospital, Beijing 10034, China

²Exchange and Cooperation Division, National Institute of Hospital Administration, Beijing 100191, China

Abstract

Background: The cesarean section rate (CSR) has been a main concern worldwide. The present study aimed to investigate the CSR in Beijing, China, and to analyze the related factors of CS delivery.

Methods: An observational study was conducted in 15 medical centers in Beijing using a systemic cluster sampling method. In total, 15,194 pregnancies were enrolled in the study between June 20, 2013 and November 30, 2013. Independent *t*-tests and Pearson's Chi-square test were used to examine differences between two groups, and related factors of the CSR were examined by multivariable logistic regression.

Results: The CSR was 41.9% (4471/10,671) in singleton primiparae. Women who were more than 35 years old had a 7.4-fold increased risk of CS delivery compared with women <25 years old (odds ratio [OR] = 7.388, 95% confidence interval [CI] = 5.561–9.816, *P* < 0.001). Prepregnancy obese women had a 2-fold increased risk of CS delivery compared with prepregnancy normal weight women (OR = 2.058, 95% CI = 1.640–2.584, *P* < 0.001). The excessive weight gain group had a 1.4-fold increased risk of CS delivery compared with the adequate weight gain group (OR = 1.422, 95% CI = 1.289–1.568, *P* < 0.001). Gestational diabetes mellitus (GDM) women and DM women had an increased risk of CS delivery (1.2- and 1.7-fold, respectively) compared with normal blood glucose women. Women who were born in rural areas had a lower risk of CS delivery than did those who were born in urban areas (OR = 0.696, 95% CI = 0.625–0.775, *P* < 0.001). The risk of CS delivery gradually increased with a decreasing education level. Neonates weighing 3000–3499 g had the lowest CSR (36.2%). Neonates weighing <2500 g had a 2-fold increased risk of CS delivery compared with neonates weighing 3000–3499 g (OR = 2.020, 95% CI = 1.537–2.656, *P* < 0.001). Neonates weighing ≥4500 g had an 8.3-fold increased risk of CS delivery compared with neonates weighing 3000–3499 g (OR = 8.313, 95% CI = 4.436–15.579, *P* < 0.001).

Conclusions: Maternal age, prepregnancy body mass index, gestational weight gain, blood glucose levels, residence, education level, and singleton fetal birth weight are all factors that might significantly affect the CSR.

Key words: Blood Glucose Levels; Cesarean Section Rate; Gestational Weight Gain; Prepregnancy Body Mass Index; Related Factors

INTRODUCTION

Cesarean section rate (CSR) has been a main concern worldwide in the recent decades. In 1985, the World Health Organization (WHO) suggested a CSR approximately 15% as being appropriate.^[1] CSs can reduce maternal/neonatal mortality, but overuse may be associated with an increased risk of severe maternal outcomes, such as increased risk of death, Intensive Care Unit admission, blood transfusion, and hysterectomy.^[2]

According to a large cross-sectional study conducted in 24 countries around the world between 2004 and 2008 by the World Health Organization Global Survey on Maternal and Perinatal Health,^[2] Chinese health

facilities had the highest CSR of 46.2%, while the rate was 25.7% in Asia. A recent study showed that, between 2008 and 2014, the overall annual rate of cesarean deliveries increased in China, reaching 34.9%,^[3] but there was major geographic variation in the rates and trends over time, with the rates declining in some of the largest urban areas.

Address for correspondence: Dr. Hui-Xia Yang,

Department of Obstetrics and Gynecology, Peking University First Hospital, No. 1 Xianmen Street, Xicheng District, Beijing 10034, China
E-Mail: yanghuixia@bjmu.edu.cn

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

© 2017 Chinese Medical Journal | Produced by Wolters Kluwer - Medknow

Received: 04-07-2017 **Edited by:** Peng Lyu

How to cite this article: Song G, Wei YM, Zhu WW, Yang HX. Cesarean Section Rate in Singleton Primiparae and Related Factors in Beijing, China. *Chin Med J* 2017;130:2395-401.

Access this article online

Quick Response Code:



Website:
www.cmj.org

DOI:
10.4103/0366-6999.216415

Chinese health-care providers have recognized this problem and made great efforts in the recent years, the rate of CS has shown a downward trend in some hospitals.^[4]

The purpose of this study was to investigate the CSR in Beijing, analyze the related factors, and attempt to identify ways to lower the CSR.

METHODS

Ethical approval

The study was conducted in accordance with the *Declaration of Helsinki* and was approved by the Ethics Committee of Peking University First Hospital (No. 2013[578]). Informed written consent was obtained from all patients prior to their enrollment in this study.

Study population

The sampling frame for this study consisted of all public hospitals that offer delivery services in Beijing. Random seed and sampling intervals were chosen and sorted by the number of deliveries in 2012; 15 hospitals in Beijing were chosen as clusters by a systemic cluster sampling method (listed in the acknowledgments). There were 2 specialist hospitals and 13 general hospitals as well as 8 tertiary hospitals and 7 secondary hospitals, which well represented the hospitals in Beijing. The eligibility criteria included all women delivering after 28 weeks of pregnancy between June 20 and November 30, 2013. The exclusion criteria were as follows: missing data on major items, such as 75 g oral glucose tolerance test (OGTT) results, birth weight, gestational age, and delivery mode.

The study developed a questionnaire to collect and record the data, including the high risk factors of CS delivery, such as maternal age, prepregnancy body mass index (BMI), gestational weight gain (GWG), blood glucose levels, residence, average monthly income, education level, and fetal birth weight.

Definitions

Gestational diabetes mellitus

GDM is diagnosed when any one value met or exceeded 5.1 mmol/L at 0 h, 10.0 mmol/L at 1 h, or 8.5 mmol/L at 2 h in the OGTT at 24–28 weeks. The gestational diabetes mellitus (GDM) diagnostic criteria followed the new guideline established from 2011 in China.^[5]

Prepregnancy body mass index categories

According to the standard of the Working Group on Obesity in China,^[6] the prepregnancy BMI categories were classified as follows: underweight, $\text{BMI} < 18.5 \text{ kg/m}^2$; normal weight, $18.5 \leq \text{BMI} \leq 23.9 \text{ kg/m}^2$; overweight, $24 \leq \text{BMI} \leq 27.9 \text{ kg/m}^2$; and obese, $\text{BMI} \geq 28 \text{ kg/m}^2$.

Guidelines for excessive gestational weight gain during pregnancy

According to the Institute of Medicine (IOM),^[7] underweight women ($\text{BMI} < 18.5 \text{ kg/m}^2$) should gain 12.5–18 kg; normal weight women ($18.5 \leq \text{BMI} \leq 24.9 \text{ kg/m}^2$) should gain

11.5–16 kg; overweight women ($25 \leq \text{BMI} \leq 29.9 \text{ kg/m}^2$) should gain 7–11.5 kg; and obese women ($\text{BMI} \geq 30 \text{ kg/m}^2$) should gain 5–9 kg. According to each BMI category, GWG less than the lower limit is defined as inadequate weight gain, more than the upper limit is defined as excessive weight gain, and within the limits is defined as adequate weight gain.

Statistical analysis

SPSS 16.0 (SPSS, Inc., Chicago, IL, USA) was used for the statistical analysis. Continuous variables were expressed as the mean \pm standard deviation and were tested by independent *t*-tests. Categorical variables were presented as numbers and percentages; Pearson's Chi-square test was applied to examine differences between the groups. Related factors of the CSR were examined by multivariable logistic regression. $P < 0.05$ was considered statistically significant.

RESULTS

Maternal characteristics

In this study, we enrolled 15,194 women in the initial sample, of which 14,941 women had singleton pregnancies and 253 women had multiple pregnancies; in addition, 10,824 were nulliparous and 4370 were multiparous. Compared with the singleton pregnancy group, the multiple birth group had a significantly higher risk of CSR (42.5% vs. 85.8%, $P < 0.001$). For the multiparous women, one of the main causes of CS was a history of CS, might have biased the conclusion. Therefore, we excluded the multiple birth group and the multiparous group and performed the following calculations for the 10,671 singleton primiparae.

In the 10,671 singleton primiparae, the mean age of the women was 27.6 ± 4.0 years, and the mean gestational age was 39.1 ± 1.6 weeks. Of the 4471 women who delivered their infants by CS, the CSR was 41.9%.

For the 10,671 singleton primiparae, the demographic data of continuous variables comparing the CS delivery and vaginal delivery groups are shown in Table 1, including maternal age, gestational weeks, prepregnancy BMI, GWG during pregnancy, and fetal birth weight. We found that the CS delivery group had a higher maternal age, prepregnancy BMI, GWG, and fetal birth weight, and that the vaginal delivery group had a higher gestational age.

Risk factors for cesarean section delivery

Data of the 10,671 singleton primiparae were used to analyze the related factors of CS delivery. We used CS delivery as the dependent variable and eight categorical variables as independent variables to perform the single-factor analysis. The eight categorical variables included were as follows: maternal age, prepregnancy BMI, GWG, blood glucose levels, residence, average monthly income, education level, and fetal birth weight [Figure 1 and Table 2]. Except for one variable (average monthly income) with no significant difference, all the other variables presented with $P < 0.05$. In total, 9453 cases without missing information were included

Table 1: Comparison of maternal age, gestational weeks, and other demographic data of the samples in CS delivery and vaginal delivery groups

Variables	CS delivery (n = 4471)		Vaginal delivery (n = 6200)		t	P
	n	Values	n	Values		
Maternal age (years)	4471	28.4 ± 4.3	6200	27.0 ± 3.7	-17.170	<0.001
Gestational weeks	4471	38.9 ± 1.7	6200	39.2 ± 1.5	7.028	<0.001
Prepregnancy BMI (kg/m ²)	4431	22.1 ± 3.5	6162	21.0 ± 2.9	-17.308	<0.001
GWG (kg)	4209	16.5 ± 5.6	5848	15.7 ± 5.0	-7.476	<0.001
Fetal birth weight (g)	4471	3401.5 ± 537.8	6200	3313.3 ± 413.6	-9.183	<0.001

Data were shown as mean ± SD. CS: Cesarean section; BMI: Body mass index; GWG: Gestational weight gain; SD: Standard deviation.




Figure 1: Cesarean section rate and related factors in single-factor analysis. * $P < 0.001$; † $P < 0.05$. In the fetal birth weight (g) chart, except for group 2500–2999 g and group 3000–3499 g, all the other two groups had $P < 0.05$. BMI: Body mass index; GWG: Gestational weight gain; GDM: Gestational diabetes mellitus; DM: Diabetes mellitus.

in the logistic regression analysis. Using the enter method in the logistic regression, all seven variables were statistically significant. Hence, we considered the related factor of CS delivery to be maternal age, prepregnancy BMI, GWG, blood glucose level, residence, education level, and singleton fetal birth weight.

As shown in Table 3, we found that the CSR increased with age. Women who were more than 35 years old had a 7.4-fold increased risk of CS delivery compared with women <25 years old (odds ratio [OR] = 7.388, 95% confidence interval [CI] = 5.561–9.816, $P < 0.001$). As the prepregnancy BMI increased, the CSR increased accordingly; prepregnancy obese women had a 2-fold increased risk of CS delivery compared with prepregnancy normal weight women (OR = 2.058, 95% CI = 1.640–2.584, $P < 0.001$). When we divided the women by GWG group during pregnancy, the inadequate weight gain group had a similar CSR to the adequate weight gain group (OR = 0.926, 95%

CI = 0.804–1.065, $P = 0.281$), but the excessive weight gain group had a 1.4-fold increased risk of CS delivery compared with the adequate weight gain group (OR = 1.422, 95% CI = 1.289–1.568, $P < 0.001$). Compared with the normal blood glucose women, the GDM women and DM women had an increased risk of CS delivery (1.2- and 1.7-fold, respectively). Women who were born in rural areas had a lower risk of CS delivery than those who were born in urban areas (OR = 0.696, 95% CI = 0.625–0.775, $P < 0.001$). The risk of CS delivery gradually increased with the decrease in education level; women graduating from junior school or lower had a higher risk of CS delivery than did women graduating from graduate school and above (OR = 1.403, 95% CI = 1.380–1.729, $P < 0.001$). The relationship between fetal birth weight and CS delivery appeared to be like a “U shape”: neonates weighing 3000–3499 g had the lowest CSR, the farther away from this weight group (more or less), the higher the CSR was. Neonates weighing <2500 g had a

Table 2: Single-factor analysis of the related factors of CSR

Variables	Total number (missing)	CS delivery, n (%)	Vaginal delivery, n (%)	CSR (%)	Statistics	P
Maternal age					-14.688*	<0.001
<25 years	10,671 (0)	759 (17.0)	1528 (24.6)	33.2		
25–29 years		2052 (45.9)	3055 (49.3)	40.2		
30–34 years		1282 (28.7)	1493 (24.1)	46.2		
≥35 years		378 (8.5)	124 (2.0)	75.3		
Prepregnancy BMI (kg/m ²)					-14.969*	<0.001
<18.5	10,593 (78)	514 (11.6)	1074 (17.4)	32.4		
18.5–23.9		2839 (64.1)	4256 (69.1)	40.0		
24–27.9		799 (18.0)	666 (10.8)	54.5		
≥28		279 (6.3)	166 (2.7)	62.7		
GWG group					-11.094*	<0.001
Inadequate	10,039 (632)	530 (12.6)	978 (16.8)	35.1		
Adequate		1245 (29.6)	2158 (37.0)	36.6		
Excessive		2426 (57.7)	2702 (46.3)	47.3		
Blood glucose levels					64.377†	<0.001
Normal	10,671 (0)	3405 (76.2)	5092 (82.1)	40.1		
GDM		987 (22.1)	1056 (17.0)	48.3		
DM		79 (1.8)	52 (0.8)	60.3		
Education					-2.561*	0.010
Graduate and above	10,012 (659)	470 (11.1)	690 (11.9)	40.5		
College		2703 (64.0)	3470 (59.9)	43.8		
High school		617 (14.6)	890 (15.4)	40.9		
Junior school or lower		433 (10.3)	739 (12.8)	36.9		
Residence					95.935†	<0.001
Urban	10,620 (51)	3310 (74.3)	4032 (65.4)	45.1		
Rural		1145 (25.7)	2133 (34.6)	34.9		
Average monthly income (RMB, Yuan)					-1.106*	0.269
<1000	10,140 (531)	2918 (68.3)	4,65 (69.3)	41.8		
1000–2999		1268 (29.7)	1686 (28.7)	42.9		
3000–4999		82 (1.9)	107 (1.8)	43.4		
≥5000		6 (0.1)	8 (0.1)	42.9		
Fetal birth weight (g)					-10.969*	<0.001
<2500	10,671 (0)	204 (4.6)	168 (2.7)	54.8		
2500–2999		582 (13.0)	976 (15.7)	37.4		
3000–3499		1681 (37.6)	2969 (47.9)	36.2		
3500–3999		1447 (32.4)	1831 (29.5)	44.1		
4000–4499		490 (11.0)	241 (3.9)	67.0		
≥4500		67 (1.5)	15 (0.2)	81.7		

*Statistics shown as Z; †Statistics shown as χ^2 . CS: Cesarean section; CSR: Cesarean section rate; GDM: Gestational diabetes mellitus; DM: Diabetes mellitus; BMI: Body mass index; GWG: Gestational weight gain.

2-fold increased risk of CS delivery compared with neonates weighing 3000–3499 g ($OR = 2.020$, 95% $CI = 1.537$ – 2.656 , $P < 0.001$), and neonates weighing ≥ 4500 g had an 8.3-fold increased risk of CS delivery compared with neonates weighing 3000–3499 g ($OR = 8.313$, 95% $CI = 4.436$ – 15.579 , $P < 0.001$), as shown in Figure 1.

DISCUSSION

The rate of CS in this study was 41.9%, which was lower than that in other studies in urban areas, namely, 65.6% in a study in Tianjin^[8] from 2009 to 2011 and 64.1% in 2008 in a national cross-sectional survey.^[9] However, 41.9% was still a very high value. In the present study, we found some factors influencing the CSR. Maternal age, prepregnancy BMI,

GWG, blood glucose levels, residence, education level, and singleton fetal birth weight were all factors that significantly affected the CSR.

The CSR increased with maternal age, and the same trend has been found in Korea^[10] and the USA^[11] [Figure 2], as the total CSR was 36% in Korea in 2012 and 32.7% in the USA in 2013. Older maternal age was one of the reasons for the increased CSR because it increased the development of maternal diseases, which increased ante- and intra-partum complications. In a study in England,^[12] there was an 80% increase in the number of women over the age of 40 years between 2006 and 2011, and the overall CSR increased from 34.6% in 2006 to 53.7% in 2011, comprising an increase in both elective and emergency cesarean sections. Subsequent to the

Table 3: Multivariate analysis of the related factors of CSR

Variables	β	SE	P	OR	95% CI
Maternal age			<0.001		
<25 years				1.000	
25–29 years	0.232	0.064	<0.001	1.261	1.113–1.428
30–34 years	0.507	0.073	<0.001	1.660	1.440–1.913
≥ 35 years	2.000	0.145	<0.001	7.388	5.561–9.816
Prepregnancy BMI (kg/m ²)			<0.001		
<18.5	-0.156	0.065	0.017	0.856	0.753–0.972
18.5–23.9				1.000	
24–27.9	0.449	0.065	<0.001	1.567	1.380–1.780
≥ 28	0.722	0.116	<0.001	2.058	1.640–2.584
GWG group			<0.001		
Inadequate	-0.077	0.072	0.281	0.926	0.804–1.065
Adequate				1.000	
Excessive	0.352	0.050	<0.001	1.422	1.289–1.568
Blood glucose levels			<0.001		
Normal				1.000	
GDM	0.187	0.057	<0.001	1.206	1.079–1.348
DM	0.541	0.209	0.010	1.717	1.140–2.587
Residence			<0.001		
Urban				1.000	
Rural	-0.362	0.055	<0.001	0.696	0.625–0.775
Education			0.005		
Graduate and above				1.000	
College	0.219	0.071	0.002	1.244	1.082–1.431
High school	0.290	0.093	0.002	1.336	1.112–1.605
Junior school or lower	0.338	0.107	0.002	1.403	1.138–1.729
Fetal birth weight (g)			<0.001		
<2500	0.703	0.140	<0.001	2.020	1.537–2.656
2500–2999	0.094	0.067	0.163	1.098	0.963–1.253
3000–3499				1.000	
3500–3999	0.287	0.051	<0.001	1.332	1.205–1.472
4000–4499	1.201	0.091	<0.001	3.322	2.777–3.974
≥ 4500	2.118	0.320	<0.001	8.313	4.436–15.579
Constant	-1.246	0.099	<0.001	0.288	

BMI: Body mass index; GWG: Gestational weight gain; GDM: Gestational diabetes mellitus; DM: Diabetes mellitus. CS: Cesarean section; CSR: Cesarean section rate; OR: Odd ratio; CI: Confidence interval.




Figure 2: Relationship between cesarean section rate and maternal age in China, Korea, and the USA. * $P < 0.05$, China 2013 group versus Korea 2012 group; † $P < 0.05$, USA 2013 group versus China 2013 group; ‡ $P < 0.05$, Korea 2012 group versus USA 2013 group.

implementation of the two-child policy in 2015 in China, we are facing the same problem now and must pay more attention to the complications of older maternal age, including the CSR.

Overweight people are at increased risk for chronic conditions, such as cardiovascular disease, hypertension, and diabetes. Overweight or obesity before/during pregnancy might have an influence on both mother and child. Previous investigations have found a positive association between maternal prepregnancy BMI and cesarean delivery.^[13-15] A study showed that a nulliparous woman with a BMI >30 kg/m² is six times more likely to have a cesarean delivery than is a nulliparous woman whose BMI was <20 kg/m²;^[16] the data showed that women whose BMI is ≥ 28 kg/m² have a 2.1-fold increased risk of CS delivery compared with women whose BMI is between 18.5 and 24 kg/m². Data of 292,568 cases from 1993 to 2005 in China reveal that maternal overweight and a high GWG or a GWG above the IOM recommendation is associated with adverse outcomes, such as hypertensive disorders complicating the pregnancy, cesarean delivery, macrosomia, and large-for-gestational-age infants.^[17] We also found that maternal obesity was closely related to the CSR; the excessive weight gain group had a much higher CSR than did the adequate weight gain group and the inadequate

weight gain group. Health-care providers should inform women that maintaining a proper weight before pregnancy and achieving adequate weight gain during pregnancy will help to reduce the CSR.

Compared with normal blood glucose women, GDM women and DM women had a much higher CSR (40.1%, 48.3%, and 60.3%, respectively; all $P < 0.001$). It is well established that GDM is associated with a number of adverse outcomes;^[18] CSR is just one of these. The incidence of GDM has risen from 2–6% to 15–20%;^[19] how to maintain a proper blood glucose level during pregnancy has become more important,^[20] as it will concurrently lower the CSR.

Some studies have shown that CSs without medical indications may have contributed to the high CSR,^[21-23] and the introduction of the one-child policy in 1979 may be one of the reasons. Parents who expect to have only one child might prefer a CS to vaginal delivery because it is free from pain and anxiety. However, since 2015, in which the two-child policy was inducted, more parents are inclined to have two children, and thus, they prefer a vaginal delivery for their first child. In the recent years, the CSR in primiparae might be lower. However, there are still a considerable number of women with a CS history who want to have their second child, and most of them will again choose a CS. Therefore, reducing the primary CSR is the most important way of decreasing the CSR.^[24]

Undoubtedly, there were some limitations in this study. In the multiparous women, one of the main reasons for a CS delivery was a previous CS delivery history, but this factor was not properly incorporated at the beginning of the study; consequently, we did not obtain accurate data regarding the previous CS delivery history. Hence, we only performed the analysis in singleton primiparae. This shortcoming will be rectified in future studies.

In conclusion, maternal age, prepregnancy BMI, GWG, blood glucose levels, residence, education level, and singleton fetal birth weight are all factors that might significantly affect the CSR. Maintaining a fetal birth weight from 3000–3499 g yielded the lowest CSR.

Acknowledgments

We would like to thank the following 15 hospitals that participated in the study: Tongzhou Maternal and Child Health Hospital of Beijing, Peking University First Hospital, Peking University Third Hospital, Aviation General Hospital, Chinese People's Liberation Army General Hospital of the Beijing Military Region, Navy General Hospital, Beijing Hospital of Chinese Traditional and Western Medicine, Pinggu District Maternal and Child Health Hospital of Beijing, Beijing Daxing District Hongxing Hospital, Beijing Chuiyangliu Hospital, Miyunxian Hospital, Peking University Shougang Hospital, Changping District Hospital of Chinese Traditional Medicine, Beijing Jingmei Group General Hospital, and Beijing No. 6 Hospital.

Financial support and sponsorship

The study was supported by grants from the World Diabetes Foundation (No. WDF 10-517 and No. WDF 14-908).

Conflicts of interest

There are no conflicts of interest.

REFERENCES

1. World Health Organization. Appropriate technology for birth. *Lancet* 1985;2:436-7. doi: 10.1016/S0140-6736(85)92750-3.
2. Lumbiganon P, Laopaiboon M, Gülmezoglu AM, Souza JP, Taneepanichskul S, Ruyan P, *et al*. Method of delivery and pregnancy outcomes in Asia: The WHO global survey on maternal and perinatal health 2007-08. *Lancet* 2010;375:490-9. doi: 10.1016/S0140-6736(09)61870-5.
3. Li HT, Luo S, Trasande L, Hellerstein S, Kang C, Li JX, *et al*. Geographic variations and temporal trends in cesarean delivery rates in China, 2008-2014. *JAMA* 2017;317:69-76. doi: 10.1001/jama.2016.18663.
4. Liu Y, Wang X, Zou L, Ruan Y, Zhang W. An analysis of variations of indications and maternal-fetal prognosis for caesarean section in a tertiary hospital of Beijing: A population-based retrospective cohort study. *Medicine (Baltimore)* 2017;96:e5509. doi: 10.1097/MD.0000000000005509.
5. Yang HX. Diagnostic criteria for gestational diabetes mellitus (WS 331-2011). *Chin Med J* 2012;125:1212-3. doi: 10.3760/cma.j.isn.0366-6999.2012.07.004.
6. Chen C, Lu FC; Department of Disease Control Ministry of Health, PR China. The guidelines for prevention and control of overweight and obesity in Chinese adults. *Biomed Environ Sci* 2004;17 Suppl: 1-36.
7. Institute of Medicine. *Weight Gain during Pregnancy: Reexamining the Guidelines*. Washington, DC: National Academies Press; 2009.
8. Li N, Liu E, Guo J, Pan L, Li B, Wang P, *et al*. Maternal prepregnancy body mass index and gestational weight gain on pregnancy outcomes. *PLoS One* 2013;8:e82310. doi: 10.1371/journal.pone.0082310.
9. Feng XL, Xu L, Guo Y, Ronsmans C. Factors influencing rising caesarean section rates in China between 1988 and 2008. *Bull World Health Organ* 2012;90:30-9, 39A. doi: 10.2471/BLT.11.090399.
10. Chung SH, Seol HJ, Choi YS, Oh SY, Kim A, Bae CW. Changes in the cesarean section rate in Korea (1982-2012) and a review of the associated factors. *J Korean Med Sci* 2014;29:1341-52. doi: 10.3346/jkms.2014.29.10.1341.
11. Martin JA, Hamilton BE, Osterman MJ, Curtin SC, Matthews TJ. Births: Final data for 2013. *Natl Vital Stat Rep* 2015;64:1-65.
12. Moore EK, Irvine LM. The impact of maternal age over forty years on the caesarean section rate: Six year experience at a busy district general hospital. *J Obstet Gynaecol* 2014;34:238-40. doi: 10.3109/01443615.2013.838546.
13. Vahratian A, Siega-Riz AM, Savitz DA, Zhang J. Maternal pre-pregnancy overweight and obesity and the risk of cesarean delivery in nulliparous women. *Ann Epidemiol* 2005;15:467-74. doi: 10.1016/j.annepidem.2005.02.005.
14. Deruelle P, Servan-Schreiber E, Riviere O, Garabedian C, Vendittelli F. Does a body mass index greater than 25kg/m2 increase maternal and neonatal morbidity? A French historical cohort study. *J Gynecol Obstet Hum Reprod* 2017. pii: S2468-784730152-6. doi: 10.1016/j.jogoh.2017.06.007.
15. Xiong C, Zhou A, Cao Z, Zhang Y, Qiu L, Yao C, *et al*. Association of pre-pregnancy body mass index, gestational weight gain with cesarean section in term deliveries of China. *Sci Rep* 2016;6:37168. doi: 10.1038/srep37168.
16. Young TK, Woodmansee B. Factors that are associated with cesarean delivery in a large private practice: The importance of prepregnancy body mass index and weight gain. *Am J Obstet Gynecol* 2002;187:312-8. doi: 10.1067/mob.2002.126200.
17. Liu Y, Dai W, Dai X, Li Z. Prepregnancy body mass index and gestational weight gain with the outcome of pregnancy: A 13-year study of 292,568 cases in China. *Arch Gynecol Obstet* 2012;286:905-11. doi: 10.1007/s00404-012-2403-6.
18. Wendland EM, Torloni MR, Falavigna M, Trujillo J, Dode MA, Campos MA, *et al*. Gestational diabetes and pregnancy outcomes – A

systematic review of the World Health Organization (WHO) and the International Association of Diabetes in Pregnancy Study Groups (IADPSG) diagnostic criteria. *BMC Pregnancy Childbirth* 2012;12:23. doi: 10.1186/1471-2393-12-23.

19. Zhu WW, Yang HX, Wang C, Su RN, Feng H, Kapur A. High prevalence of gestational diabetes mellitus in Beijing: Effect of maternal birth weight and other risk factors. *Chin Med J* 2017;130:1019-25. doi: 10.4103/0366-6999.204930.
20. Song G, Wang C, Yang HX. Diabetes management beyond pregnancy. *Chin Med J* 2017;130:1009-11. doi: 10.4103/0366-6999.204938.
21. Zhang J, Liu Y, Meikle S, Zheng J, Sun W, Li Z. Cesarean delivery on maternal request in Southeast China. *Obstet Gynecol* 2008;111:1077-82. doi: 10.1097/AOG.0b013e31816e349e.
22. Liu X, Landon MB, Cheng W, Chen Y. Cesarean delivery on maternal request in China: What are the risks and benefits? *Am J Obstet Gynecol* 2015;212:817.e1-9. doi: 10.1016/j.ajog.2015.01.043.
23. Stützer PP, Berlit S, Lis S, Schmahl C, Sütterlin M, Tuschy B. Elective caesarean section on maternal request in Germany: Factors affecting decision making concerning mode of delivery. *Arch Gynecol Obstet* 2017;295:1151-6. doi: 10.1007/s00404-017-4349-1.
24. American College of Obstetricians and Gynecologists (College); Society for Maternal-Fetal Medicine, Caughey AB, Cahill AG, Guise JM, Rouse DJ. Safe prevention of the primary cesarean delivery. *Am J Obstet Gynecol* 2014;210:179-93. doi: 10.1016/j.ajog.2014.01.026.