

Harmful alcohol drinking among HIV-positive people in Nepal: an overlooked threat to anti-retroviral therapy adherence and health-related quality of life

Khem Narayan Pokhrel^a, Kalpana Gaulee Pokhrel^b, Sanjeev Raj Neupane^b and Vidya Dev Sharma^c

^aDepartment of HIV, Nutrition, and Health, Health Research and Social Development Forum, Kathmandu, Nepal; ^bDepartment of HIV and Nutrition, Integrated Development Foundation, Kathmandu, Nepal; ^cDepartment of Psychiatry and Mental Health, Institute of Medicine, Tribhuvan University, Kathmandu, Nepal

ABSTRACT

Background: People living with Human Immunodeficiency Virus (HIV) often suffer from alcohol-use disorders resulting in their poor health and treatment outcomes. Little is known about the association of harmful alcohol drinking with their adherence to anti-retroviral therapy (ART) and health-related quality of life (QOL) in low-resource settings.

Objective: This study aimed to investigate associations between harmful alcohol drinking, adherence to ART and health-related QOL in HIV-positive people, stratified by gender, in Nepal.

Methods: We conducted a cross-sectional study of 682 HIV-positive people on ART to measure their self-reported harmful alcohol drinking and non-adherence to ART in the previous month of data collection. We also measured health-related QOL using a WHOQOL-HIV BREF scale. The association between harmful alcohol drinking and non-adherence to ART was examined using multiple logistic regressions. Additionally, multiple linear regressions examined association between harmful alcohol drinking and QOL.

Results: Harmful alcohol drinking was associated with non-adherence to ART among men (AOR: 2.48, 95% CI: 1.50, 4.11, $p < 0.001$) and women (AOR: 2.52, 95% CI: 1.32, 4.80, $p = 0.005$). Men were more likely to have lower score for the psychological ($\beta = -0.55$, $p = 0.021$) and level of independence ($\beta = -0.68$, $p = 0.018$) domains when they had harmful alcohol drinking. Moreover, women were more likely to have lower scores for the physical ($\beta = -1.01$, $p = 0.015$), social relations ($\beta = -0.82$, $p = 0.033$), environmental ($\beta = -0.88$, $p = 0.011$), and spiritual ($\beta = -1.30$, $p = 0.005$) domains of QOL when they had harmful alcohol drinking.

Conclusions: Harmful alcohol drinking had a negative association with ART adherence and QOL in both HIV-positive men and women in Nepal. Screening for alcohol-use disorders and community-based counseling services should be provided while delivering ART services to improve treatment adherence and QOL.

ARTICLE HISTORY

Received 11 November 2017
Accepted 12 February 2018

RESPONSIBLE EDITOR

Jennifer Stewart Williams,
Umeå University, Sweden

KEYWORDS


HIV; harmful alcohol drinking; anti-retroviral therapy; quality of life; Nepal

Background

Access to anti-retroviral therapy (ART) improved the life expectancies of people living with Human Immunodeficiency Virus (HIV) globally [1,2]. However, their medication adherence remains salient. Many suffer from harmful alcohol drinking disorders, this being one of the common behavioral disorders [3]. In HIV-positive people, alcohol consumption increases the risk of HIV disease progression, leading to poor health outcomes [4]. Harmful alcohol drinking eventually results in negative consequences for medication adherence [5] and poor health-related quality of life (QOL) [6].

Harmful alcohol drinking among HIV-positive people has been associated with non-adherence to ART in both high-income countries (HICs) and in low- and middle-income countries (LMICs) [7–9]. In a study in Peru, harmful alcohol drinking had inverse

effects on ART adherence among men who have sex with men (MSM) [10]. In this study, 31.9% of the HIV-positive people practiced drinking alcohol at a harmful level, as measured by the alcohol-use disorders identification test (AUDIT). In the United States, HIV-positive people experienced poor social and emotional support from family and friends when they continued consuming alcohol. Such poor support has been shown to have a negative impact on adherence to ART [11]. In addition, HIV-positive people who use alcohol frequently can experience more conflict with their partners, which can impact negatively on spousal support and lead to poor compliance with ART [12]. Harmful alcohol drinking can also reduce coping skills and increase the likelihood of opportunistic infections [13]. Moreover, HIV-positive people who used illicit substances also had alcohol-use disorders resulting in non-adherence to ART [14].

CONTACT Khem Narayan Pokhrel  pratikjee@gmail.com  Department of HIV, Nutrition, and Health, Health Research and Social Development Forum, PO Box 24133, Kathmandu, Nepal

© 2018 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

People living with HIV who use alcohol tend to have a poor health-related QOL. Research in the United States shows that alcohol-use disorders can impact negatively on physical functioning and on the emotional well-being of HIV-positive people, which can then decrease mental health-related QOL [15]. In that study, HIV-positive people who suffered from alcohol-use disorders were less likely to have perceived social support, which resulted in poor QOL in the psychological domain. Moreover, another study has shown that HIV-positive people who usually did not seek social support, experienced poor psychological QOL [16]. Alcohol use also increases the risk of substance use such as heroin, which may have deleterious effects on health-related QOL in both physical and mental health dimensions [5,17].

Gender difference also has implications in relation to the association between alcohol use and health outcomes. In the United States, men having sex with men (MSM) had a higher prevalence of HIV because of unprotected sexual practices resulting from harmful alcohol drinking behavior [18]. Women in the United States also tended to receive poor social support, which had negative effects on adherence to ART when they suffered from alcohol-use disorders [19].

HIV-positive people who use alcohol, poorly engage in HIV services and this affects their continuum of care. A systematic review found that alcohol users were more likely to have missed at least one or more cascades of continuum of care such as attending treatment services, ART compliance, routine investigation of viral loads, and CD4 count [20]. Additionally, alcohol users also engage in substance use as a coping mechanism for reducing stress related to HIV, which further deteriorates their health [21].

HIV-positive people had non-adherence to ART when they consumed alcohol in Nepal [22,23]. Alcohol consumption, however, was measured using a single-item scale in Nepal to examine the association of alcohol use and non-adherence using AIDS Clinical Trial Group (ACTG) questionnaires. Little is known about the harmful effects of drinking alcohol and its association with ART adherence in Nepal. A measure that is more comprehensive would be helpful in covering harmful alcohol drinking that comprises dose, frequency, and hazardous alcohol use. Alcohol use is common in both rural and urban areas of the country because homemade alcohol and brewed alcohol are easily available [24]. Additionally, no evidence is available about the role of harmful alcohol drinking on the QOL of HIV-positive people stratified by gender in South Asia. This study was conducted on HIV-positive people in Nepal. The aims were, first, to investigate associations between harmful alcohol drinking and adherence to ART

among HIV-positive people stratified by gender, and, second, to examine the association between harmful alcohol drinking and health-related QOL in HIV-positive people stratified by gender.

Methods

Study setting

This study was conducted in Nepal, a low-income country situated in South Asia, which harbors a population of 28 million. About 40,000 HIV-positive people are living in the country. Among them, approximately 12,000 were on ART in 2015 [25].

Study design and participants

The design was cross-sectional. The study districts were selected by convenience sampling method and participants were conveniently recruited from the various non-governmental organizations (NGOs) involved in providing HIV services. The participants who took ART for at least one year at the time of data collection and those who were diagnosed as HIV-positive within five years were included in the study sample.

Sampling procedures

We calculated the sample size using G*Power statistical software. First, we took the reference prevalence of non-adherence to ART as 21% among HIV-positive people who received community-based psychosocial support [26]. Second, a previous study found a 10% difference in non-adherence to ART among HIV-positive people who received psychosocial support compared to those who did not receive the support [27]. We assumed the same differences of 10% non-adherence to ART between participants who would enroll in a community home-based care program and those who would not enroll in the program. This resulted in an estimated prevalence of non-adherence to ART among the control group as 31%. With p-value for significant of less than 0.05 (two-tailed) and the power of 0.80, the calculated sample size was 638 (intervention: 319; control: 319). In total, we recruited 720 participants, of them, 682 were considered in the analysis.

Measures

Harmful alcohol drinking

Participants were asked about their regular drinking behavior. The extent of harmful drinking was measured using the Alcohol Use Disorders Identification Test (AUDIT), which is a 10-item scale designed to measure alcohol use covering amount, frequency, and

hazardous drinking [10,28]. The available alcoholic drinks such as locally brewed alcohol; bottled wine, beer, whisky, vodka, etc. were standardized as one unit for each 10 gm of alcohol content. For example, if an individual consumed 200 ml of locally brewed alcohol with concentration of 25% it implies that he/she consumed five units of alcohol. The scale has already been validated and applied in Nepal [29], and the Cronbach's alpha was 0.78 in this study. A cumulative total score ranges from 0 to 40. Scores of eight or more were categorized as being harmful alcohol drinking.

ART non-adherence

The questionnaires developed by the AIDS Clinical Trial Group (ACTG) were applied to assess the missed pills. The data collection considered participants who missed at least one dose in the past month as being non-adherent to ART [30,31].

Health-related QOL

Participants' health-related QOL was measured using the WHO Quality of Life-HIV BREF (WHOQOL-HIV BREF) scale [32]. The scale measures physical, psychological, level of independence, social relations, environmental, and spiritual/religious domains of life. Each item score ranges from 1 to 5 on a Likert scale. A score of '1' indicates a low and negative direction and '5' indicates a high and positive direction. Six domains of QOL were calculated using various facets of the scale. The scores were arranged reversely, which had a negative direction (pain and discomfort, negative feelings, dependence on medication, death and dying). The total scores for each domain range from 4 to 20. Higher scores indicate a better QOL. The values of Cronbach's alpha for the domains were: physical 0.78; psychological 0.84; level of independence 0.82; social relations 0.86; environmental 0.88; and spiritual/religion 0.83. The scale was developed by World Health Organization (WHO) and it has been widely used in various countries including Nepal [32–34].

Socio-demographics

Participants were interviewed and asked for self-reported information regarding age, gender, marital status education, employment status, and physical symptoms. Marital status was categorized as married and unmarried/widow/single. Education level was grouped in two categories: illiterate and primary level education; or more. For employment status, we categorized the participants as employed/self-employed and unemployed. Further participants

who reported opportunistic infections were categorized as having physical symptoms and we considered those who did not report any symptoms related to opportunistic infection as having no physical symptoms.

HIV-clinical staging

The records about clinical staging were retrieved from the ART centers of the selected districts. The first and second clinical stages were categorized as early stage, and the third and fourth stages were categorized as advanced [35].

Procedure for data collection

A set of questionnaires was first translated from English to Nepali and then back translated to English by two independent translators, to ensure consistency in meaning and interpretation. Finally, some modifications were made before finalizing the questionnaires for pre-testing. Pre-testing of questionnaires was done among 48 participants in Kathmandu. Necessary modifications were made in the questionnaire after the pre-testing. We approached 720 HIV-positive people on ART and 682 participants (94.7%) responded to the questions. The pre-tested participants were not included in the study. The baseline data collection was conducted in February and March 2015.

Data analysis

Data were analyzed descriptively using chi-squared and Fisher's exact test for categorical variables and a t-test for age. Multiple logistic regressions examined the association between harmful alcohol drinking and non-adherence to ART. Additionally, multiple linear regressions examined the association between harmful alcohol drinking and the quality of life (QOL) domains. In each model, important covariates such as age, education, marital status, occupation, physical symptoms, and HIV-clinical staging were adjusted, as studies have indicated that these factors can influence ART adherence [30,31]. All data were analyzed using STATA version 12.1.

Ethical considerations

The study was approved by the Research Ethics Committee of the Graduate School of Medicine, the University of Tokyo and the Nepal Health Research Council. Prior to data collection, permission was obtained from the selected agencies providing community home-based care and support services for HIV-positive people in Nepal. Participants were ensured about the confidentiality of information and their voluntary participation. Moreover, written

informed consent was obtained from the participants on a designated form. The participants were also asked for their thumbprint if they were not able to write their name. During the interview, if participants were found to have a severe alcohol-use disorder, they were referred to non-governmental organizations and nearby hospitals.

Results

Socio-demographic characteristics and harmful alcohol drinking of participants

Of 682 participants, 354 were men and 328 were women. About 25.1% of the men and 29.3% of the women were non-adherent to ART. ART adherent and non-adherent participants were not significantly differences in relation to their age, education, marital status, employment, HIV-clinical staging, or physical symptoms. Among them, 35.0% of the men and 15.5% of the women had harmful alcohol drinking. Compared with adherent men, more non-adherent men had harmful alcohol drinking (51.1% vs. 29.6%, $p < 0.001$). A significantly higher proportion of non-adherent women had harmful alcohol drinking compared to those who were ART adherent (22.7% vs. 12.5%, $p = 0.013$) (see Table 1).

Multiple logistic regression: association of harmful alcohol drinking with non-adherence to ART stratified by gender

Men were more likely to be non-adherent to ART when they had harmful alcohol drinking (AOR: 2.48,

95% CI: 1.50, 4.11, $p < 0.001$). Also, harmful alcohol drinking was positively associated with non-adherence to ART among women (AOR: 2.52, 95% CI: 1.32, 4.80, $p = 0.001$) (see Table 2).

Harmful alcohol drinking by QOL domains

Compared to those who did not have harmful alcohol drinking, men had lower scores for psychological (mean: 12.3 vs. 12.9, $p = 0.013$), level of independence (mean: 12.0 vs. 12.8, $p = 0.008$), environmental (mean: 11.6 vs. 12.2, $p = 0.010$), and spiritual (mean: 11.1 vs. 12.1, $p = 0.001$) domains when they had harmful alcohol drinking.

Similarly, women who had harmful alcohol drinking had lower scores of physical (mean: 12.3 vs. 13.1, $p = 0.041$) and spiritual (mean: 10.6 vs. 11.7, $p < 0.001$) domains compared to those who did not have harmful alcohol drinking (see Table 3).

Multiple linear regression: association of harmful alcohol drinking and QOL domains stratified by gender

After controlling age, marital status, education level, employment status, physical symptoms, and HIV-clinical staging, harmful alcohol drinking was negatively associated with QOL domains. Among men, harmful alcohol drinking was associated with the psychological ($\beta = -0.55$, $p = 0.021$), level of independence ($\beta = -0.68$, $p = 0.018$), environmental ($\beta = -0.55$, $p = 0.022$), and spiritual ($\beta = -0.98$, $p = 0.004$) domains. Additionally, women were more likely to have lower scores for physical

Table 1. Sociodemographic characteristics and harmful alcohol drinking among ART adherent and non-adherent participants.

| Variables | Total n (%) | Men (n = 354) | | P-value | Women (n = 328) | | P-value |
|---|-------------|---------------------------------|------------------------------------|---------|---------------------------------|------------------------------------|---------|
| | | ART adherent (n = 265) n (%) | ART non-adherent (n = 89) n (%) | | ART adherent (n = 232) n (%) | ART non-adherent (n = 96) n (%) | |
| Age (mean, SD)^a | 36.3 (8.2) | 36.9 (8.8) | 36.7 (7.8) | 0.845 | 35.5 (8.1) | 36.1 (8.0) | 0.893 |
| Education level^b | | | | | | | |
| Illiterate | 289 (42.4) | 91 (34.3) | 36 (40.4) | 0.265 | 111 (47.8) | 51 (53.1) | 0.384 |
| ≥Primary level | 393 (57.6) | 174 (65.7) | 53 (59.6) | | 121 (52.2) | 45 (46.9) | |
| Marital status^b | | | | | | | |
| Married | 554 (81.2) | 225 (84.9) | 64 (71.9) | 0.016 | 189 (81.4) | 76 (79.2) | 0.828 |
| Unmarried/widow/single | 128 (18.8) | 40 (15.1) | 25 (28.1) | | 43 (18.6) | 20 (20.8) | |
| Employment status^b | | | | | | | |
| Employed/self-employed | 448 (65.7) | 147 (55.5) | 63 (70.8) | 0.060 | 162 (69.8) | 76 (79.2) | 0.085 |
| Unemployed | 234 (34.3) | 118 (44.5) | 26 (29.2) | | 70 (30.2) | 20 (20.8) | |
| HIV-clinical staging^b | | | | | | | |
| Early stage | 550 (80.6) | 219 (86.6) | 67 (77.0) | 0.039 | 192 (84.6) | 72 (75.8) | 0.062 |
| Advanced stage | 132 (19.4) | 34 (13.4) | 20 (23.0) | | 35 (15.4) | 23 (24.2) | |
| Physical symptoms^b | | | | | | | |
| Yes | 146 (21.4) | 44 (16.6) | 23 (25.8) | 0.050 | 52 (22.4) | 27 (28.1) | 0.271 |
| No | 536 (78.6) | 221 (83.4) | 66 (74.2) | | 180 (76.6) | 69 (71.9) | |
| Harmful alcohol drinking^b | | | | | | | |
| Yes | 175 (25.7) | 78 (29.6) | 46 (51.1) | <0.001 | 29 (12.5) | 22 (22.7) | 0.013 |
| No | 507 (74.3) | 186 (70.4) | 44 (48.9) | | 202 (87.5) | 75 (77.3) | |

^aIndependent sample t-test.

^bChi-squared test.

Table 2. Logistic regression analysis: the association between harmful alcohol drinking and non-adherence to ART stratified by gender.

| Characteristics | ART non-adherence | | | |
|--|---------------------------|---------|---------------------------|---------|
| | Men | | Women | |
| | AOR ^a (95% CI) | P-value | AOR ^a (95% CI) | P-value |
| Harmful alcohol drinking | 2.48 (1.50, 4.11) | <0.001 | 2.52 (1.32, 4.80) | 0.005 |
| Marital status (married) | 1.00 (0.96, 1.03) | 0.964 | 0.99 (0.96, 1.02) | 0.696 |
| Age | 0.95 (0.56, 1.62) | 0.862 | 0.82 (0.49, 1.37) | 0.842 |
| Education (\leq primary) | 0.98 (0.57, 1.69) | 0.962 | 0.76 (0.46, 1.26) | 0.298 |
| Employment status (employed/self-employed) | 0.53 (0.27, 1.04) | 0.067 | 0.83 (0.44, 1.57) | 0.580 |
| HIV-clinical staging (advanced) | 0.56 (0.32, 0.99) | 0.047 | 0.69 (0.38, 1.26) | 0.239 |
| Physical symptoms (present) | 1.23 (0.41, 3.66) | 0.707 | 0.79 (0.29, 2.13) | 0.653 |

^aAdjusted odds Ratio.

Table 3. Descriptive analysis: harmful alcohol drinking by QOL domains stratified by gender.

| Variable | Physical | | | | Psychological | | | | Level of independence | | | |
|------------------|-----------------|---------|------------|---------|---------------|---------|------------|---------|-----------------------|---------|------------|---------|
| | Men | | Women | | Men | | Women | | Men | | Women | |
| | Mean (SD) | P-value | Mean (SD) | P-value | Mean (SD) | P-value | Mean (SD) | P-value | Mean (SD) | P-value | Mean (SD) | P-value |
| HAD ^a | | | | | | | | | | | | |
| No | 13.1 (2.7) | 0.091 | 13.1 (2.7) | 0.041 | 12.9 (2.1) | 0.013 | 12.5 (2.2) | 0.577 | 12.8 (2.6) | 0.008 | 12.4 (2.6) | 0.274 |
| Yes | 12.6 (2.8) | | 12.3 (2.6) | | 12.3 (2.2) | | 12.3 (2.2) | | 12.0 (2.5) | | 12.0 (2.6) | |
| Variable | Social relation | | | | Environmental | | | | Spiritual | | | |
| | Men | | Women | | Men | | Women | | Men | | Women | |
| | Mean (SD) | P-value | Mean (SD) | P-value | Mean (SD) | P-value | Mean (SD) | P-value | Mean (SD) | P-value | Mean (SD) | P-value |
| HAD ^a | | | | | | | | | | | | |
| No | 12.5 (2.5) | 0.235 | 12.1 (2.8) | 0.136 | 11.6 (2.1) | 0.010 | 11.3 (2.3) | 0.062 | 11.1 (3.0) | 0.001 | 10.6 (2.6) | <0.001 |
| Yes | 12.8 (2.4) | | 12.6 (2.4) | | 12.2 (2.1) | | 11.9 (2.2) | | 12.1 (2.9) | | 11.7 (3.0) | |

HAD = Harmful alcohol drinking.

^aIndependent sample t-test.

($\beta = -1.01$, $p = 0.015$), social relations ($\beta = -0.82$, $p = 0.033$), environmental ($\beta = -0.88$, $p = 0.011$), and spiritual ($\beta = -1.30$, $p = 0.005$) domains when they had harmful alcohol drinking (see Tables 4 and 5).

Discussion

In this study, HIV-positive men and women in Nepal were more likely to be non-adherent to ART when they had harmful alcohol drinking. Additionally, harmful alcohol drinking was negatively associated

Table 4. Multiple regression analysis: the association of harmful alcohol drinking with the QOL domains (physical, psychological, and level of independence) stratified by gender.

| Variables | Physical | | | | Psychological | | | | Level of Independence | | | |
|--|----------|------|---------|------|---------------|------|---------|------|-----------------------|------|---------|------|
| | Men | | Women | | Men | | Women | | Men | | Women | |
| | β | SE | β | SE | β | SE | β | SE | β | SE | β | SE |
| Harmful alcohol drinking | -0.41 | 0.31 | -1.01* | 0.41 | -0.55* | 0.24 | -0.39 | 0.34 | -0.68* | 0.29 | -0.63 | 0.40 |
| Marital status (married) | 0.12 | 0.42 | -0.13 | 0.38 | 0.01 | 0.32 | -0.09 | 0.31 | 0.31 | 0.39 | -0.29 | 0.37 |
| Age | 0.02 | 0.02 | 0.04* | 0.02 | 0.02 | 0.02 | 0.04* | 0.02 | 0.00 | 0.02 | 0.05 | 0.02 |
| Education (\geq primary) | 0.89** | 0.31 | 0.20 | 0.30 | 0.54 | 0.24 | 0.25 | 0.25 | 0.17 | 0.29 | 0.11 | 0.29 |
| Employment status (Employed/self-employed) | -0.19 | 0.31 | 0.24 | 0.34 | 0.50* | 0.24 | 0.12 | 0.28 | 0.10 | 0.29 | 0.24 | 0.33 |
| Clinical stage (advanced) | -0.45 | 0.72 | -1.53* | 0.62 | -0.35 | 0.55 | -1.20* | 0.51 | -0.17 | 0.68 | -0.64 | 0.60 |
| Physical symptoms (present) | -0.54 | 0.68 | 0.34 | 0.55 | -0.26 | 0.52 | 0.23 | 0.45 | -0.70 | 0.64 | -0.23 | 0.53 |

*p-value<0.05, **p-value<0.01.

Table 5. Multiple regression analysis: the association of harmful alcohol drinking with QOL domains (physical, psychological, and level of independence) stratified by gender.

| Variables | Social relation | | | | Environmental | | | | Spiritual | | | |
|--|-----------------|------|---------|------|---------------|------|---------|------|-----------|------|---------|------|
| | Men | | Women | | Men | | Women | | Men | | Women | |
| | β | SE | β | SE | β | SE | B | SE | β | SE | β | SE |
| Harmful alcohol drinking | -0.28 | 0.28 | -0.82* | 0.38 | -0.55* | 0.24 | -0.88* | 0.34 | -0.98** | 0.33 | -1.30** | 0.46 |
| Marital status (married) | 0.11 | 0.37 | 0.03 | 0.36 | 0.22 | 0.32 | -0.04 | 0.32 | 0.07 | 0.45 | -0.09 | 0.43 |
| Age | 0.02 | 0.02 | 0.05 | 0.02 | -0.01 | 0.02 | 0.03 | 0.02 | 0.01 | 0.02 | -0.01 | 0.02 |
| Education (\geq primary) | 0.20 | 0.28 | 0.05 | 0.28 | 0.18 | 0.24 | -0.14 | 0.25 | 0.57 | 0.34 | 0.09 | 0.34 |
| Employment status (Employed/self-employed) | 0.30 | 0.28 | 0.41 | 0.31 | 0.18 | 0.24 | 0.26 | 0.28 | 0.39 | 0.34 | -0.04 | 0.38 |
| Clinical stage (advanced) | -0.68 | 0.65 | -1.18* | 0.57 | -0.42 | 0.56 | -1.42** | 0.52 | -0.89* | 0.78 | -1.28 | 0.69 |
| Physical symptoms (present) | -0.04 | 0.61 | 0.38 | 0.51 | -0.23 | 0.52 | 0.31 | 0.46 | 0.37 | 0.74 | 0.74 | 0.62 |

*p-value <0.05, **p-value <0.001.

with the psychological, level of independence, environmental, and spiritual domains of QOL among men. Women were more likely to have poor QOL for physical, social relations, environmental, and spiritual domains when they had harmful alcohol drinking.

HIV-positive men and women who had harmful alcohol drinking were more likely to be non-adherent to ART. With regard to gender differences, both men and women had non-adherence. The strength of the association in women was slightly higher compared to men. In the Nepali context, alcohol drinking behavior might have impaired women to access to HIV services compared to men. It is possible that both men and women might not have sought family support and might have experienced poor coping skills because they suffered from other mental health and substance-use disorders such as use of heroin and cocaine.

Higher frequency of alcohol consumption induces a stronger maladaptive coping strategy among HIV-positive people because the alcohol may be associated with other behavioral disorders that lead to non-adherence to ART [13]. Alcohol users may also be less likely to seek support from their family, friends, and caregivers, which may lead to non-adherence to ART [36]. Moreover, they may also simultaneously engage in any substance use in order to cope with the treatment side-effects which may also lead to poor treatment compliance [37]. Our results are consistent with the studies conducted in African and Latin American countries, which reported the negative implication of alcohol use on ART adherence [38]. This study provides evidence about the positive association between harmful alcohol drinking and non-adherence to ART in both women and men.

Harmful alcohol drinking had negative effects on the QOL domains of both HIV-positive men and women or simply HIV-positive people. They might have endured poor self-efficacy and not sought appropriate care for opportunistic infections and physical symptoms [39]. Alcohol further increases the risk of the HIV disease progression such as leading to a decrease in CD4 count and an increase in viral load [40]. Eventually, alcohol use might have deteriorated the physical condition of HIV-positive people leading to their poor physical QOL.

In this study, we observed gender differences in the QOL among men and women. Women were more likely to have a poor QOL in respect to social relations and spiritual domains. The lower level of support in women might have had a negative role in those who sought support from their spouse and family [41]. Men were more likely to have a lower QOL in respect to the psychological, level of independence, environment and spiritual domains. Compared to women, Nepalese HIV-positive men might have had problems with multiple substance use and poor engagement in HIV services as reported by the studies in South Africa, where men were

more likely to engage in substance use that has negative effect on their use of continuum of care services [42].

This study has some limitations. Social desirability bias may have affected our results as participants self-reported their status regarding missing pills and alcohol use. To minimize this as much as possible the interviews were conducted in private places. Furthermore, we counted consumed pills out of the total prescribed for the previous month. The study districts were selected conveniently to reach the participants from NGOs, and therefore selection bias may limit generalizability of the findings. However, the results can be applied in the settings where participants have similar characteristics. The results might have also been biased by other substance-use disorders and mental health disorders. These factors impact negatively on engagement in HIV services. To minimize this as much as possible we recruited participants from similar settings.

Despite these limitations, this study is novel being the first to examine these issues separately in HIV-positive men and women in Nepal. This study also adds new evidence about the negative association between harmful alcohol drinking, ART adherence, and QOL in both men and women.

Conclusion

Harmful alcohol drinking was negatively associated with ART adherence and QOL of HIV-positive people in Nepal. When designing services for ART counseling, alcohol-use disorders should be integrated along with psychosocial counseling. Health services providers should pay particular attention to address alcohol-use disorders while providing screening and treatment for HIV-related consequences. Frontline health workers should be provided with training material that are community-based and tailor made so that they can screen for and treat these disorders in their work settings. Additionally, support mechanisms should be established at family and community levels to address alcohol-use disorders, improve ART adherence and enhance the QOL for HIV-positive people. More studies are warranted to investigate causal relationships between harmful alcohol drinking and ART adherence taking into account factors such as depression, stress, and substance-use disorders.

Acknowledgments

All authors gratefully acknowledge Bobby Singh, Niru Lama, and Milan Khadka for their kind assistance in the arrangement of fieldwork. We also honor the support from the various NGOs and mutual support groups for giving us permission to conduct this work. Our thanks go to Mr. Gregory King for his kind assistance in editing the English of this manuscript. Finally, we would like to show our sincere gratitude to the HIV-positive people who participated in this study.

Author contributions

KNP was involved in the study design, data analysis, and manuscript preparation. VDS provided the technical guidance and supervised the whole research work in the field. KGP and SRN were involved in data collection, data entry, data processing, and manuscript preparation. All authors were involved in the finalization of this manuscript.

Disclosure statement

No potential conflict of interest was reported by the authors.

Ethics and consent

The study received approval from Research Ethics Committee, Graduate School of Medicine, the University of Tokyo, Japan and Nepal Health Research Council, Nepal. Participants provided their written informed consent, and their confidentiality and voluntary participation was ensured.

Funding information

Authors received no funding support for this study.

Paper context

Nepal has been experiencing concentrated epidemics of HIV among most at-risk populations for the past two decades. More than 12,000 out of an estimated 40,000 HIV-positive people are taking anti-retroviral therapy (ART). Among them, harmful alcohol drinking is high in HIV-positive people. Evidence of association between harmful alcohol drinking and ART adherence is limited in Nepal and South Asia. Additionally, the relationship between harmful alcohol drinking and non-adherence by gender is also not well understood. This study investigates association between harmful alcohol drinking (using the AUDIT measure) non-adherence to ART and health-related quality of life in HIV-positive people in Nepal.

References

- [1] Anti-retroviral Therapy Collaboration. Life expectancy of individuals on combination antiretroviral therapy in high-income countries: a collaborative analysis of 14 cohort studies. *Lancet*. 2008;372:293–299.
- [2] Bor J, Herbst AJ, Newell M-L, et al. Increases in adult life expectancy in rural South Africa: valuing the scale-up of HIV treatment. *Science*. 2013;339:961–965.
- [3] Galvan FH, Bing EG, Fleishman JA, et al. The prevalence of alcohol consumption and heavy drinking among people with HIV in the United States: results from the HIV cost and services utilization study. *J Stud Alcohol*. 2002;63:179–186.
- [4] Ghebremichael M, Paintsil E, Ickovics JR, et al. Longitudinal association of alcohol use with HIV disease progression and psychological health of women with HIV. *AIDS Care*. 2009;21:834–841.
- [5] Chander G, Lau B, Moore RD. Hazardous alcohol use: a risk factor for non-adherence and lack of suppression in HIV infection. *J Acquir Immune Defic Syndr*. 2006;43:411–417.
- [6] Liu C, Ostrow D, Detels R, et al. Impacts of HIV infection and HAART use on quality of life. *Qual Life Res*. 2006;15:941–949.
- [7] Braithwaite RS, McGinnis KA, Conigliaro J, et al. A temporal and dose-response association between alcohol consumption and medication adherence among veterans in care. *Alcohol Clin Exp Res*. 2005;29:1190–1197.
- [8] Nakimuli-Mpungu E, Bass JK, Alexandre P, et al. Depression, alcohol use and adherence to antiretroviral therapy in sub-Saharan Africa: a systematic review. *AIDS Behav*. 2012;16:2101–2118.
- [9] Jaquet A, Ekouevi DK, Bashi J, et al. Alcohol use and non-adherence to antiretroviral therapy in HIV-infected patients in West Africa. *Addiction*. 2010;105:1416–1421.
- [10] Ferro EG, Weikum D, Vagenas P, et al. Alcohol use disorders negatively influence antiretroviral medication adherence among men who have sex with men in Peru. *AIDS Care*. 2015;27:93–104.
- [11] Samet JH, Horton NJ, Meli S, et al. A randomized controlled trial to enhance antiretroviral therapy adherence in patients with a history of alcohol problems. *Antivir Ther*. 2005;10:83–93.
- [12] Malow R, Dévieux JG, Stein JA, et al. Depression, substance abuse and other contextual predictors of adherence to antiretroviral therapy (ART) among Haitians. *AIDS Behav*. 2013;17:1221–1230.
- [13] Pence BW, Thielman NM, Whetten K, et al. Coping strategies and patterns of alcohol and drug use among HIV-infected patients in the United States Southeast. *AIDS Patient Care STDS*. 2008;22:869–877.
- [14] Mellins CA, Havens JF, McDonnell C, et al. Adherence to antiretroviral medications and medical care in HIV-infected adults diagnosed with mental and substance abuse disorders. *AIDS Care*. 2009;21:168–177.
- [15] Jia H, Uphold CR, Wu S, et al. Predictors of changes in health-related quality of life among men with HIV infection in the HAART era. *AIDS Patient Care STDS*. 2005;19:395–405.
- [16] Moak ZB, Agrawal A. The association between perceived interpersonal social support and physical and mental health: results from the National Epidemiological Survey on Alcohol and Related Conditions. *J Public Health (Oxf)*. 2010;32:191–201.
- [17] Grant BF, Stinson FS, Dawson DA, et al. Prevalence and co-occurrence of substance use disorders and independent mood and anxiety disorders: results from the National Epidemiologic Survey on Alcohol and Related Conditions. *Arch Gen Psychiatry*. 2004;61:807–816.
- [18] Millett GA, Peterson JL, Wolitski RJ, et al. Greater risk for HIV infection of black men who have sex with men: a critical literature review. *Am J Public Health*. 2006;96:1007–1019.
- [19] Berg KM, Demas PA, Howard AA, et al. Gender differences in factors associated with adherence to antiretroviral therapy. *J Gen Intern Med*. 2004;19:1111–1117.
- [20] Vagenas P, Azar MM, Copenhaver MM, et al. The impact of alcohol use and related disorders on the HIV continuum of care: a systematic review: alcohol and the HIV continuum of care. *Curr HIV/AIDS Rep*. 2015;12:421–436.

- [21] Mayston R, Kinyanda E, Chishinga N, et al. Mental disorder and the outcome of HIV/AIDS in low-income and middle-income countries: a systematic review. *Aids*. 2012;26:S117–135.
- [22] Bam K, Rajbhandari RM, Karmacharya DB, et al. Strengthening adherence to Anti Retroviral Therapy (ART) monitoring and support: operation research to identify barriers and facilitators in Nepal. *BMC Health Serv Res*. 2015;15:188.
- [23] Wasti SP, Simkhada P, Randall J, et al. Factors influencing adherence to antiretroviral treatment in Nepal: a mixed-methods study. *PLoS One*. 2012;7:e35547.
- [24] Jhingan HP, Shyangwa P, Sharma A, et al. Prevalence of alcohol dependence in a town in Nepal as assessed by the CAGE questionnaire. *Addiction*. 2003;98:339–343.
- [25] National Center for HIV/AIDS and STD Control, Ministry of Health and Population, Nepal. HIV situation in Nepal. Kathmandu: Government of Nepal; 2015.
- [26] Muñoz M, Bayona J, Sanchez E, et al. Matching social support to individual needs: a community-based intervention to improve HIV treatment adherence in a resource-poor setting. *AIDS Behav*. 2011;15:1454–1464.
- [27] Basso CR, Helena ET, Caraciolo JM, et al. Exploring ART intake scenes in a human rights-based intervention to improve adherence: a randomized controlled trial. *AIDS Behav*. 2013;17:181–192.
- [28] Saunders JB, Aasland OG, Babor TF, et al. Development of the Alcohol Use Disorders Identification Test (AUDIT): WHO collaborative project on early detection of persons with harmful alcohol consumption—II. *Addiction*. 1993;88:791–804.
- [29] Pradhan B, Chappuis F, Baral D, et al. The alcohol use disorders identification test (AUDIT): validation of a Nepali version for the detection of alcohol use disorders and hazardous drinking in medical settings. *Subst Abuse Treat Prev Policy*. 2012;7:42.
- [30] Amberbir A, Woldemichael K, Getachew S, et al. Predictors of adherence to antiretroviral therapy among HIV-infected persons: a prospective study in Southwest Ethiopia. *BMC Public Health*. 2008;8:265.
- [31] Gordillo V, Del Amo J, Soriano V, et al. Sociodemographic and psychological variables influencing adherence to antiretroviral therapy. *Aids*. 1999;13:1763–1769.
- [32] World Health Organization. Department of Mental Health and Substance Dependence. WHOQOL-HIV BREF. Geneva (Switzerland): WHO; 2002.
- [33] Thapa R, Amatya A, Pahari DP, et al. Nutritional status and its association with quality of life among people living with HIV attending public anti-retroviral therapy sites of Kathmandu Valley, Nepal. *AIDS Res Ther*. 2015;12:14.
- [34] Adewuya AO, Afolabi MO, Ola BA, et al. Relationship between depression and quality of life in persons with HIV infection in Nigeria. *Int J Psychiatry Med*. 2008;38:43–51.
- [35] World Health Organization. WHO case definitions of HIV for surveillance and revised clinical staging and immunological classification of HIV-related disease in adults and children. Geneva (Switzerland): WHO; 2007.
- [36] Wolfe D, Carrieri MP, Shepard D. Treatment and care for injecting drug users with HIV infection: a review of barriers and ways forward. *Lancet*. 2010;376:355–366.
- [37] Altice FL, Kamarulzaman A, Soriano VV, et al. Treatment of medical, psychiatric, and substance-use comorbidities in people infected with HIV who use drugs. *Lancet*. 2010;376:367–387.
- [38] Hendershot CS, Stoner SA, Pantalone DW, et al. Alcohol use and antiretroviral adherence: review and meta-analysis. *J Acquir Immune Defic Syndr*. 2009;52:180–202.
- [39] Parsons JT, Rosof E, Mustanski B. Medication adherence mediates the relationship between adherence self-efficacy and biological assessments of HIV health among those with alcohol use disorders. *AIDS Behav*. 2008;12:95–103.
- [40] Samet JH, Cheng DM, Libman H, et al. Alcohol consumption and HIV disease progression. *J Acquir Immune Defic Syndr*. 2007;46:194–199.
- [41] Power R, Koopman C, Volk J, et al. Social support, substance use, and denial in relationship to antiretroviral treatment adherence among HIV-infected persons. *AIDS Patient Care STDS*. 2003;17:245–252.
- [42] Kalichman SC, Simbayi LC, Kaufman M, et al. Alcohol use and sexual risks for HIV/AIDS in sub-Saharan Africa: systematic review of empirical findings. *Prev Sci*. 2007;8:141–151.