



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

# Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection is likely to be androgen mediated



Carlos Gustavo Wambier, MD, PhD,<sup>a</sup> and Andy Goren, MD<sup>b</sup>  
*Providence, Rhode Island; and Irvine, California*

**Key words:** 5- $\alpha$  reductase; ACE2; androgen receptor; androgenetic alopecia; angiotensin converting enzyme 2; COVID-19; dutasteride; finasteride; human skin; retinoids; antiandrogen therapy; SARS-CoV-2; TMPRSS2; transmembrane protease serine 2.

*Abbreviations used:*

ACE2:	angiotensin converting enzyme 2
COVID-19:	coronavirus disease 2019
SARS-CoV-2:	severe acute respiratory syndrome coronavirus 2
TMPRSS2:	transmembrane protease, serine 2

Coronavirus disease 2019 (COVID-19) pandemic fatalities are rare before adrenarche/puberty (<10 years of age), and the vulnerability of males to severe disease<sup>1</sup> has been constantly reported over the past months of pandemic.

The first biologic step required for potential infectivity of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is the priming of the spike proteins by transmembrane protease, serine 2 (TMPRSS2). Although other proteases have been described to activate the spikes in vitro, only TMPRSS2 activity is regarded as essential for viral spread and pathogenesis in the infected hosts.<sup>2</sup> TMPRSS2 may also cleave angiotensin converting enzyme 2 (ACE2) for augmented viral entry.<sup>3</sup>

Androgen receptor activity has been considered a requirement for the transcription of the TMPRSS2 gene because no other known TMPRSS2 gene promoter has been described in humans to date.<sup>4,5</sup>

Male vulnerability may be further enhanced by X-linked inheritance of genetic polymorphisms (androgen receptor and ACE2 genes loci are in chromosome X). Obvious dermatologic signs of hyperactivation of androgen receptors are pattern reduction of density of scalp hair, increased density of facial and chest hair, acne, and oily skin. Theoretically, the hyperandrogenic phenotype might correlate with COVID-19 increased viral load, increased viral dissemination, and severity of lung involvement (Infographic 1).

Studies are still required for epidemiologic conclusions. Acknowledging the importance of androgens during the COVID-19 pandemic may offer another targeted therapy for trials, with androgen suppression to reduce host vulnerability when infection risk is high.

From the Department of Dermatology, The Warren Alpert Medical School of Brown University, Providence<sup>a</sup>; and Applied Biology, Inc, Irvine.<sup>b</sup>  
 Funding sources: None.

Conflicts of interest: None disclosed.

IRB approval status: Not applicable.


Correspondence and reprint requests to: Carlos Gustavo Wambier, MD, PhD, Rhode Island Hospital, 593 Eddy St, APC, 10th Floor, Providence, RI 02903. E-mail: [carlos\\_wambier@brown.edu](mailto:carlos_wambier@brown.edu).

J Am Acad Dermatol 2020;83:308-9.

0190-9622/\$36.00

© 2020 by the American Academy of Dermatology, Inc.

<https://doi.org/10.1016/j.jaad.2020.04.032>



**Infographic 1.** Androgen-mediated COVID-19.

**REFERENCES**

1. Shi Y, Yu X, Zhao H, Wang H, Zhao R, Sheng J. Host susceptibility to severe COVID-19 and establishment of a host risk score: findings of 487 cases outside Wuhan. *Crit Care*. 2020;24(1):108.
2. Hoffmann M, Kleine-Weber H, Schroeder S, et al. SARS-CoV-2 cell entry depends on ACE2 and TMPRSS2 and is blocked by a clinically proven protease inhibitor. *Cell*. 2020:1-10.
3. Heurich A, Hofmann-Winkler H, Gierer S, Liepold T, Jahn O, Pohlmann S. TMPRSS2 and ADAM17 cleave ACE2 differentially and only proteolysis by TMPRSS2 augments entry driven by the severe acute respiratory syndrome coronavirus spike protein. *J Virol*. 2014;88(2): 1293-1307.
4. Lucas JM, Heinlein C, Kim T, et al. The androgen-regulated protease TMPRSS2 activates a proteolytic cascade involving components of the tumor microenvironment and promotes prostate cancer metastasis. *Cancer Discov*. 2014;4(11):1310-1325.
5. National Institutes of Health. TMPRSS2 transmembrane serine protease 2 [Homo sapiens (human)] Gene ID: 7113. Updated March 13, 2020. Available at: <https://www.ncbi.nlm.nih.gov/gene/7113>. Accessed April 2, 2020.