

Research article

Open Access

Novel markers for differentiation of lobular and ductal invasive breast carcinomas by laser microdissection and microarray analysis

Gulisa Turashvili¹, Jan Bouchal*¹, Karl Baumforth², Wenbin Wei², Marta Dziechciarkova³, Jiri Ehrmann¹, Jiri Klein⁴, Eduard Fridman⁵, Jozef Skarda¹, Josef Srovnal³, Marian Hajduch³, Paul Murray² and Zdenek Kolar¹

Address: ¹Laboratory of Molecular Pathology, Institute of Pathology, Palacky University, Olomouc, Czech Republic, ²Cancer Research U.K. Institute of Cancer Studies, University of Birmingham, UK, ³Laboratory of Experimental Medicine, Department of Pediatrics, Palacky University, Olomouc, Czech Republic, ⁴Department of Surgery, Palacky University, Olomouc, Czech Republic and ⁵Department of Pathology, Tel Aviv University, Chaim Sheba Medical Center and Sackler School of Medicine, Tel Aviv, Israel

Email: Gulisa Turashvili - guliati@gmail.com; Jan Bouchal* - jan.bouchal@gmail.com; Karl Baumforth - k.r.baumforth@bham.ac.uk; Wenbin Wei - w.wei@bham.ac.uk; Marta Dziechciarkova - mdziechciarkova@seznam.cz; Jiri Ehrmann - jiri.ehrmann@tiscali.cz; Jiri Klein - kleinj@fnol.cz; Eduard Fridman - ef@sheba.health.gov.il; Jozef Skarda - jojos@email.cz; Josef Srovnal - josef.srovnal@seznam.cz; Marian Hajduch - hajduchm@atlas.cz; Paul Murray - p.g.murray@bham.ac.uk; Zdenek Kolar - kolarz@tunw.upol.cz

* Corresponding author

Published: 27 March 2007

Received: 27 November 2006

BMC Cancer 2007, 7:55 doi:10.1186/1471-2407-7-55

Accepted: 27 March 2007

This article is available from: <http://www.biomedcentral.com/1471-2407/7/55>

© 2007 Turashvili et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Invasive ductal and lobular carcinomas (IDC and ILC) are the most common histological types of breast cancer. Clinical follow-up data and metastatic patterns suggest that the development and progression of these tumors are different. The aim of our study was to identify gene expression profiles of IDC and ILC in relation to normal breast epithelial cells.

Methods: We examined 30 samples (normal ductal and lobular cells from 10 patients, IDC cells from 5 patients, ILC cells from 5 patients) microdissected from cryosections of ten mastectomy specimens from postmenopausal patients. Fifty nanograms of total RNA were amplified and labeled by PCR and *in vitro* transcription. Samples were analysed upon Affymetrix UI33 Plus 2.0 Arrays. The expression of seven differentially expressed genes (*CDH1*, *EMPI*, *DDR1*, *DVLI*, *KRT5*, *KRT6*, *KRT17*) was verified by immunohistochemistry on tissue microarrays. Expression of *ASPN* mRNA was validated by *in situ* hybridization on frozen sections, and *CTHRC1*, *ASPN* and *COL3A1* were tested by PCR.

Results: Using GCOS pairwise comparison algorithm and rank products we have identified 84 named genes common to ILC versus normal cell types, 74 named genes common to IDC versus normal cell types, 78 named genes differentially expressed between normal ductal and lobular cells, and 28 named genes between IDC and ILC. Genes distinguishing between IDC and ILC are involved in epithelial-mesenchymal transition, TGF-beta and Wnt signaling. These changes were present in both tumor types but appeared to be more prominent in ILC. Immunohistochemistry for several novel markers (*EMPI*, *DVLI*, *DDR1*) distinguished large sets of IDC from ILC.

Conclusion: IDC and ILC can be differentiated both at the gene and protein levels. In this study we report two candidate genes, asporin (*ASPN*) and collagen triple helix repeat containing 1 (*CTHRC1*) which might be significant in breast carcinogenesis. Besides E-cadherin, the proteins validated on tissue microarrays (*EMPI*, *DVLI*, *DDR1*) may represent novel immunohistochemical markers helpful in distinguishing between IDC and ILC. Further studies with larger sets of patients are needed to verify the gene expression profiles of various histological types of breast cancer in order to determine molecular subclassifications, prognosis and the optimum treatment strategies.

Background

Invasive ductal (IDC) and lobular carcinomas (ILC) are the most common histological types of breast cancer [1,2]. The terminology of ductal and lobular carcinomas is considered to be controversial as on purely anatomical grounds there is no justification for this nomenclature. Both carcinomas are derived from the terminal duct lobular unit (TDLU), and the differences in their morphology are likely to reflect the differences in mechanisms of carcinogenesis rather than the anatomical origin of the lesions. At presentation the clinical pathological parameters such as tumor site, size, grade and stage are similar for both tumor types [3], however, clinical follow-up data and the patterns of metastasis suggest that their development and progression are different [4,5]. Treatment for stage-matched tumors is similar [6], but ILCs are often resistant to neoadjuvant therapy [7]. Although patients with ILCs are older, have low grade tumor and less lymphatic invasion, they have no survival advantage compared with IDCs [8,9].

Expression profiling using microarrays is a powerful technology which enables the simultaneous study of the expression of thousands of genes and, in conjunction with laser capture microdissection, the high-throughput genetic analysis of morphologically distinct cell subpopulations within tumor tissue [10,11]. Microarray analysis has a number of applications, including tumor classification, molecular pathway modeling, functional genomics, and comparison of gene expression profiles between groups [12]. The study of gene expression in primary breast tumor tissues is complicated for two major reasons. First, breast cancer consists of many different cell types, including normal epithelial, stromal, adipose and endothelial cells. Second, tumor cells are morphologically and genetically diverse [13]. The recent development of laser capture microdissection has provided an opportunity to generate gene expression signatures from individual cell types [14-20].

Microarrays were used for analysis of breast tumor subclasses with clinical implication [21,22], for analysis of gene expression changes in single breast cancer cells from within the same tumor [14], for expression analysis of different gene families in breast cancer [23,24], and for analysis of gene expression in different cellular and tumor types [25-28]. ER status of the tumor was the most important discriminator of expression subtypes. Unsupervised hierarchical clustering segregated these tumors into two main clusters based on their basal (predominantly ER negative) and luminal (predominantly ER positive) characteristics [29]. Ductal breast cancer classes have been identified with aggressive phenotype and poor prognosis versus those with good prognosis [30-32]. Another study

reported distinct expression patterns based on BRCA1 and BRCA2 status [33].

To date, few papers have been published on gene expression profiles of normal cell populations in the mammary gland [34]. Several studies suggest differences in expression profiles of IDC and ILC. Inactivating mutations of E-cadherin gene are very frequent in ILC [35]. However, the loss of E-cadherin expression was shown to be an independent prognostic marker for recurrence, especially in node-negative breast cancer patients, irrespective of the histological type [36]. Abnormal cytoplasmic and nuclear localization of p120, a member of the E-cadherin/catenin adhesion complex, is mediated by E-cadherin loss and occurs in the early stages of ILC [37]. Non-microdissected IDC and ILC tissues have also been used to study specific gene expression profiles of lobular and ductal tumors [38,39]. We aimed to identify genes differentially expressed between normal cell types (ductal and lobular), between tumor cells and normal cells as well as between tumor types (ductal no special type and lobular). Despite examining limited number of patients, our study is the first full genome analysis of microdissected ductal and lobular tumor and normal cells which allowed us to detect normal mammary epithelium- and cancer-specific genes.

Methods

Laser capture microdissection and RNA isolation

Altogether ten surgical specimens with either invasive ductal carcinoma NST (no special type) (n = 5) or invasive lobular carcinoma (n = 5) were investigated. This research protocol was approved by the ethics committee at Palacky University. Tumor and normal tissues from the same mammary gland were identified by an experienced pathologist, snap-frozen in liquid nitrogen and stored at -80°C for further analysis. For microdissection, frozen tissues were embedded in TissueTek medium and cut on standard cryostat (Leica CM1850, Leica Microsystems GmbH, Wetzlar, Germany). Frozen sections (7 µm) were immediately fixed in acetone, stained by hematoxylin and dehydrated in alcohol and xylene. All solutions were prepared using diethyl pyrocarbonate-treated water. RNase free instruments and RnaseZap (Sigma, St Louis, MO, USA) were used throughout.

At least 1000 infiltrating ductal or lobular tumor cells together with normal lobular and ductal cells were microdissected from cryosections using the Veritas™ Laser Capture Microdissection System (Arcturus Bioscience, Inc., Mountain View, CA, USA) according to standard protocols. When microdissecting normal cells, no attempt has been made to exclude basal or myoepithelial cells for two reasons. First, both ducts and lobules are atrophic in menopause, and no clearly defined layers of luminal and

myoepithelial cells are preserved. Second, it is impossible to exclude contamination with or entrapment of myoepithelial cells during the microdissection procedure. Caps with captured cells were directly placed in 100 μ l lysis buffer (Qiagen, Hilden, Germany). Total cellular RNA was isolated (RNeasy[®] Micro Kit, Qiagen) according to manufacturer's recommendations and subsequently quantified on a Nanodrop spectrophotometer (Nanodrop Technologies, Wilmington, DE, USA).

RNA amplification, microarray target synthesis and hybridization

Fifty nanograms of total RNA were reverse transcribed and amplified by Microarray Target Amplification Kit (Roche diagnostics, Basel, Switzerland). In brief, total RNA (50 ng) was converted into cDNA using a modified oligo (dT) primer (TAS-T7 Oligo (dT)₂₄). A unique Target Amplification Sequence (TAS) with no homology to any known sequences in public databases generated the 3' anchor on the cDNA for subsequent PCR amplification. In order to include a 5' anchor sequence on the cDNA, the TAS-(dN)₁₀ primer was used for the initiation of the second strand cDNA synthesis. After purification using the Microarray Target Purification Kit (Roche), PCR was performed using the TAS primer and Expand PCR Enzyme Mix which is optimized for long (>1 kb) and unbiased PCR products. In order to ensure that messages were not amplified to saturation, the optimal number of PCR cycles was estimated by preliminary PCR and agarose electrophoresis of PCR products from cycles 21, 24, 27, 30 and 33. The optimal number of PCR cycles was set either to 27 cycles for patients 3, 4, 8, 9, and 10 or 29 cycles for patients 1, 2, 5, 6, and 7. All three populations from each patient were amplified by the same number of PCR cycles. If higher PCR cycling was needed for any cell population, new microdissection, RNA isolation, cDNA synthesis and PCR amplification were performed until the same PCR cycling was possible. After purification with the Microarray Target Purification Kit (Roche), the PCR products were labeled with biotin-14-CTP (Invitrogen, CarlsBad, CA, USA) and biotin-16-UTP (Roche) by *in vitro* transcription using Microarray RNA Target Synthesis Kit T7 (Roche). The labeled cRNA was purified using the Microarray Target Purification Kit (Roche), quantified by spectrophotometer and checked by agarose electrophoresis. The entire amplification and labeling process was monitored by GeneChip[®] Eukaryotic Poly-A RNA Control Kit (Affymetrix, Santa Clara, CA, USA) with exogenous positive controls which were spiked into the total RNA before cDNA synthesis. In all cases, 25 μ g of each biotinylated cRNA preparation was fragmented, assessed by gel electrophoresis, and placed in hybridization cocktail containing biotinylated hybridization controls (GeneChip[™] Eukaryotic Hybridization Control Kit, Affymetrix). Samples were first hybridized to Test3 Arrays for 16 hours, washed, stained

using antibody-mediated signal amplification and scanned. After passing this quality control stage, the samples were hybridized onto the large Human Genome U133 Plus 2.0 Arrays (Affymetrix).

Microarray data analysis

Scanned images of microarray chips were analysed by the GCOS (GeneChip Operating Software) from Affymetrix with the default settings except that the target signal was set to 100. Differentially expressed genes between cell types were identified using the GCOS change algorithm and Rank Products (RP) [40] following RMA (Robust Multiarray Analysis) [41]. GCOS pairwise analysis was performed to compare gene expression levels among the normal lobular, normal ductal and tumor cells within individual patients and between lobular and ductal carcinoma cells of different patients. For each comparison between two cell types, the number of increase and decrease calls of each probe set was calculated using MS Excel and probe sets with the highest number of consistent changes among all patients were identified. Probe level quantile normalization [40] and robust multiarray analysis [41] on the raw. CEL files were performed using the Affymetrix package of the Bioconductor [42]. The cut-off value of percentage of false-positives for RP analysis was set to 10%. Gene lists were uploaded to DAVID (Database for Annotation, Visualization and Integrated Discovery) [43], and functional annotation was performed. Further informations on genes were obtained from public databases, such as Gene Cards [44] and NETAFFX Analysis Center [45]. Hierarchical clustering was performed using dChip software [46]. The data discussed in this publication have been deposited in NCBI Gene Expression Omnibus (GEO) [47], and are accessible through GEO Series accession number GSE5764.

Validation by immunohistochemistry

Seven differentially expressed genes were verified by immunohistochemical staining on tissue microarrays (TMA), which were constructed from 119 breast cancer cases and contained 278 cores of 2.0 mm diameter. They consisted of 80 ductal carcinomas, 29 lobular carcinomas, one tubular carcinoma, 3 medullary carcinomas, 2 tubular-lobular carcinomas, 2 mixed ductal-lobular carcinomas, one mucinous and one papillary carcinoma. The construction of the tissue microarrays was done using a tissue arrayer (Beecher Instruments, Inc., Sun Prairie, WI, USA) [48]. The paraffin sections from 22 additional breast cancer samples containing normal mammary gland structures were used for comparison.

TMA sections were stained using standard immunohistochemical methods. Monoclonal mouse antibodies against cytokeratin 17 (dilution 1:100, clone 2D4-1G9, Abnova, Taipei, Taiwan), cytokeratin 5/6 (dilution 1:100, clone

D5/16 B4, Dako, Denmark) and E-cadherin (dilution 1:50, clone NCH-38, Dako), polyclonal mouse antibodies against EMP1 (epithelial membrane antigen 1; dilution 1:100, clone 2D4-1G9, Abnova), DDR1 (discoidin domain receptor 1; dilution 1:100, Abnova) and DVL1 (human homolog of the *Drosophila* dishevelled gene; dilution 1:100, Abnova) were used. Microwave antigen retrieval was performed in citrate buffer (pH 6.0). For antigen visualization, the EnVision/HRP system and DAB+ (Dako) were used, slides were subsequently counterstained with hematoxylin, dehydrated and mounted in Canadian balsam. Immunohistochemical procedure was optimized by testing different antigen retrieval methods and negative controls. We regarded cells as immunoreactive when an obvious membranous or submembranous (E-cadherin, DDR1, EMP1) and cytoplasmic (cytokeratins, DVL1) staining was seen. Immunoreactivity was scored as follows: retained (++) when more than 50% of membrane/cytoplasm were strongly positive, reduced (+) when 10–50% of the membrane/cytoplasm were positive, and absent (-) when 0–10% of the membrane/cytoplasm were positive. Frequencies of positive and negative staining in tumor and normal tissues were compared by Fisher's exact test (Statistica 6.0, StatSoft, Prague, Czech Republic).

Validation by in situ hybridization

Asporin mRNA expression was verified by chromogenic *in situ* hybridization (CISH) on frozen sections. Fluorescein labeled oligonucleotide (5'AAG TTG GTG GTA AGC CTT TAG GAA CTG AGG-FAM) was designed to the junction of exons five and six of the asporin mRNA (Generi Biotech, Hradec Kralove, Czech Republic). Sections were mounted on Superfrost Plus slides and stored at -80°C until further staining. Before CISH, the sections were heated at 50°C for 2 min, and then dried at room temperature for 30 min. Tissues were delipidized in chloroform at room temperature for 5 min and dried to evaporate the chloroform. Block of endogenous peroxidase activity was followed by incubation of tissue sections in 10% paraformaldehyde for 7 min, then washed in PBS for 3 min and in 2× SSC (1× SSC contains 150 mM NaCl, 15 mM sodium citrate; pH 7.2) twice for 5 min. The sections were covered with 100 µl hybridization buffer (60% formamide, 5× SSC) containing 150 ng/ml of probe and incubated in moist chamber at 37°C for 22 hours. After the hybridization, unbound probe was washed from the sections at 37°C by 2× SSC for 5 min, then three times with 60% formamide in 0.2× SSC at 37°C for 5 min, and twice with 2× SSC at room temperature for 5 min. Then polyclonal rabbit anti-FITC/HRP (dilution 1:40, Dako) was applied at room temperature for 45 min, followed by washing in 2× SSC twice for 10 min. For visualization, DAB+ (Dako) was used, slides were subsequently counterstained with hematoxylin, dehydrated and mounted in Canadian balsam.

Validation by PCR

Products after PCR amplification (see RNA amplification, microarray target synthesis and hybridization) were used for verification of *CTHRC1*, *ASPN* and *COL3A1* gene expression using gene specific PCR. The primers were as follows: *CTHRC1* forward 5'ACA AGT GCC AAC CCA GAT AGC AAC, reverse 5'ATC GCA CTT CTT CTG TGG AAG GAC (20 cycles of 95°C, 59°C and 72°C 1 min each, product length 79 bp, kindly provided by Dr. Danuta Radzioch, McGill University, Montreal, Canada), *ASPN* forward 5'GTT CAG CTT GGG AAC TTT GGA ATG TAA, reverse 5'ACT GCA ATA GAT GCTTGT TTC TCT CAA CCC (20 cycles of 95°C, 58.5°C and 72°C 1 min each, product length 243 bp [modified from [49]], and *COL3A1* forward 5'TTG TCA ACC AGT GCA AGT GAC CGA C, reverse 5'TGG TGA GCA CAG TCA TTG CTC TGC A (20 cycles of 95°C, 59°C and 72°C 1 min each, product length 276 bp) [50]. PCR products were separated on 10% polyacrylamide gels, stained with ethidium bromide and captured using a gel documentation system (DIANA II with cooled CCD camera, Raytest, Germany). Band intensities, expressed as "OD (optical density) × cm", were assessed by Multi-Analyst densitometric software (Bio-Rad, USA). The results were correlated with fluorescence signals from the relevant Affymetrix probe sets by Spearman coefficient using Statistica 6.0 software (StatSoft, Czech Republic).

Results

Unsupervised clustering

The clinical and histopathological characteristics including lymph node status, tumor grade, expression of estrogen/progesterone receptors (ER/PgR), c-erbB-B2/HER2/neu and accompanying lesions of ten breast cancer patients are shown in Table 1. We microdissected normal and tumor cells from these cases which resulted in a total of 30 samples for gene expression profiling (10 normal ductal, 10 normal lobular, 5 tumor ductal and 5 tumor lobular). Unsupervised clustering of all samples was performed using all probe sets without filtering. No tumor types were grouped together but two large clusters were evident: one mainly consisted of tumor cells and the other cluster mainly of normal cells. This clearly suggests differences in global gene expression profiles of tumor and normal cells (Figure 1).

Genes differentially expressed between normal ductal and lobular cells

In this comparison 367 probe sets were identified by pairwise comparison, and 501 probe sets by Rank Products (RP). There were 82 probe sets (78 named genes) identified by both methods. In pairwise comparisons, the differentially expressed genes were found by selecting those with count of changes more than 6 (out of 10, either increase or decrease). Multiple cytokeratins (5, 7, 14, 15, 17) were expressed in both cells, but none of them was

Table 1: Clinical and histopathological characteristics of breast cancer patients involved in microarray analysis

No	Diagnosis	Grade ¹	AC	LN	ER ³	PgR ³	HER-2	Bcl-2	E-CD
1	IDC	G3	IDH	N ₁	0	0	0	80%	>90%
2	IDC	G2	none	N ₁	40%	0	0	80%	>90%
3	IDC	G2	none	N ₀	10%	2%	0	70%	>90%
4	IDC	G3	none	N ₀	90%	50%	2+ ²	100%	>90%
5	IDC	G2	DCIS	N ₀	90%	1%	0	95%	>90%
6	ILC	G2	LCIS	N ₀	0	0	0	30%	0
7	ILC	G1	none	N ₀	80%	80%	0	100%	0
8	ILC	G1	none	N ₁	50%	0	0	80%	0
9	ILC	G2	LCIS	N ₁	0	0	0	<5%	0
10	ILC	G1	none	N ₀	50%	70%	0	80%	0

¹Tumors were graded using the Nottingham combined histologic grading system.

²No gene amplification was detected by fluorescent *in situ* hybridization.

³A complete H-score was calculated by summing the products of the percentage cells stained at a given staining intensity (0–100) and the staining intensity (0–3).

AC, accompanying changes; DCIS, ductal carcinoma *in situ*; E-CD, E-cadherin; ER, estrogen receptor; IDC, invasive ductal carcinoma, IDH, intraductal hyperplasia; ILC, invasive lobular carcinoma; LCIS, lobular carcinoma *in situ*; LN, lymph node; PgR, progesterone receptor.

useful to separate two cell types. Genes encoding proteins involved in proteolysis and metabolism (*USP25*, *TMPRSS3*, *ACACB*), protein biosynthesis and modification (*PDE4DIP*, *LOXL1*), ion transport (*CLCN3*, *GABRP*) and protein kinase cascade (*MAP4K5*) as well as genes regulating transcription (*REFAP*, *HSZFP36*) were upregulated in ductal cells. However, genes regulating cell growth, activation and motility (*TSPAN5*), genes encoding actin-binding proteins (*EHBP1*), small leucine-rich proteoglycan (*DCN*), and proteins with GTPase activity (*RHOBTB3*, *ARHGEF9*) were overexpressed in lobular cells (Table 2). Hierarchical clustering of normal cell types based on the 82 probe sets found by both RP and pairwise comparison showed that gene expression profiles of normal ductal and lobular cells from the same patients were similar, and they could not be well separated from each other (Figure 2).

Genes differentially expressed between tumor cell types and normal cells

A comparison of ductal carcinoma cells with normal ductal cells identified 1055 probe sets by pairwise analysis, 604 probe sets by RP and 326 probe sets by both methods. A comparison of ductal carcinoma cells with normal lobular cells identified 792 probe sets by pairwise analysis, 347 probe sets by RP and 171 probe sets by both methods. A comparison of lobular carcinoma cells with normal ductal cells identified 1022 probe sets by pairwise analysis, 350 probe sets by RP and 201 probe sets by both methods. A comparison of lobular carcinoma cells with normal lobular cells identified 983 probe sets by pairwise analysis, 344 probe sets by RP and 208 probe sets by both methods (see Additional file 1). In pairwise comparisons, the differentially expressed genes were identified by selecting those with count of changes more than 4 (out of 5, either increase or decrease).

Comparison between lobular carcinoma and normal cells

Combined pairwise analysis and RP have identified 106 differentially expressed probe sets (84 named genes) common to lobular carcinoma versus normal ductal and lobular cells. Of these probe sets, several genes encoding proteins involved in extracellular matrix (ECM)-tumor interaction and focal adhesion (*COL6A3*, *COL8A1*, *CTHRC1*, *THBS2*, *COMP*) were upregulated, whereas *ITGA2* gene with the same function was downregulated in tumor cells. *ASP* (asporin) was one of the upregulated genes with the highest fold change. It encodes a protein with porin activity which belongs to a family of leucine-rich repeat (LRR) proteins associated with the cartilage matrix.

Of the differentially expressed genes involved in Wnt signaling, *SFRP2* (secreted frizzled-related protein 2) and *WISP1* (WNT1 inducible signaling pathway protein 1) were upregulated in lobular carcinomas. Several genes encoding proteins involved in actin, calcium and metal ion binding were downregulated (*MYBPC1*, *DST*, *PIP*, *CA2*), and some genes with the same function (*BGN*, *ADAM12*) were upregulated in lobular carcinomas. *CDH1* (E-cadherin) was downregulated in lobular tumors accompanied by downregulation of *ITGA2* (integrin, alpha 2) which it is a ligand of. Several other genes were differentially expressed between lobular carcinomas and normal cells. *LYZ/LILRB1* was upregulated in lobular cancer cells. Other downregulated genes were as follows: *CGNL1* is a cellular myosin that appears to play a role in cytokinesis, cell shape, and specialized functions such as secretion and capping; *EHF* encodes a protein that belongs to an ETS transcription factor subfamily characterized by epithelial-specific expression (ESEs). This protein may be involved in epithelial differentiation and carcinogenesis; *CLIC6* encodes a member of the chloride




Figure 1
Unsupervised hierarchical clustering of all samples using all probe sets. No tumor types are grouped together, however, two large clusters are evident: one (blue) mainly consists of tumor cells and the other (black) mainly consists of normal cells. This suggests differences in global gene expression profiles of tumor and normal cells. 1–10, case numbers; D, ductal normal cells; L, lobular normal cells; Tduc, ductal tumor; Tlob, lobular tumor.

intracellular channel family of proteins and is involved in ion transport (Table 3).

Comparison between ductal carcinoma and normal cells

Combined pairwise analysis and RP have identified 90 differentially expressed probe sets (74 named genes) common to ductal carcinoma versus normal ductal and lobular cells. In particular, transcription regulators (ATF3, PIGR), genes encoding proteins with cytokine and growth factor activity (PTN, CX3CL1) and genes encoding proteins involved in ion transport and metabolism (ATP1A2, MMP7) were downregulated in ductal carcinomas when compared with normal cells (Table 3).

Comparison between both types of tumor cells and normal cells

Of 90 probe sets differentially expressed between ductal carcinoma and both normal cells, and 106 probe sets differentially expressed between lobular carcinoma and both normal cells, only 25 probe sets were common (Table 3). Hierarchical clustering based on those 25 probe sets showed that tumor samples were grouped together, however, gene expression profiles of normal cell types from cases 10 and 3 were different from other normal cells, and tumor cells from case 2 were different from other tumor cells (Figure 3).

Several upregulated genes (collagen type I, III, V, XI, fibronectin 1, versican) were related to tumor-ECM interactions and focal adhesion. However, another gene mediating focal adhesion, *MYLK*, was downregulated in tumor cells. This gene encodes a myosin light polypeptide kinase containing 1 fibronectin type-III domain. Genes encoding proteins involved in ion and electron transport (*CYB5R2*, *GABRP*), and genes encoding proteins with transcription factor and regulator activity (*ELF5*, *ID4*) were downregulated in both populations of tumor cells. They have also been implicated as regulators of cell proliferation, differentiation, and transformation. In addition, genes involved in cell differentiation and apoptosis (*PDZK1*) and genes encoding actin and actin-binding proteins (*CNN1*, *ACTA2*) were also downregulated in both types of tumor cells. Wnt signaling molecules were differentially expressed in our samples. Of these, *SFRP1* (secreted frizzled-related protein 1) was downregulated in tumor cells, and genes involved in calcium regulation pathway were not significantly changed. Structural constituents of cytoskeleton such as type I (14, 15, 17, 23) and type II (5) keratins were downregulated in tumor cells. Several other genes were also differentially expressed between tumor and normal cells. *PI15* (peptidase inhibitor 15) was downregulated, whereas *LRRRC15* (leucine rich repeat containing 15) was upregulated in both tumor cells.

Table 2: Genes differentially expressed between normal lobular versus normal ductal cells by both rank products and pairwise analysis

Probe Set ¹	Gene Symbol	Gene Title	Rank products	Pairwise comparison
			Fold-change	Incr/Decr
Regulation of transcription				
229431_at	RFXAP↓	Regulatory factor X-associated protein	-3.1	1/6
229732_at	HSZFP36↓	ZFP-36 for a zinc finger protein	-3.0	0/6
207117_at	H-plk↑	Krueppel-related zinc finger protein	2.4	6/1
Protein biosynthesis and modification				
214130_s_at	PDE4DIP↓	Phosphodiesterase 4D interacting protein (myomegalin)	-3.2	1/6
232752_at	— (LOXLI) ↓	CDNA FLJ10302 fis, clone NT2RM2000042 (lysyl oxidase-like 1)	-1.4	1/6
Proteolysis and metabolism				
49452_at	ACACB↓	Acetyl-Coenzyme A carboxylase beta	-18	0/8
220419_s_at	USP25↓	Ubiquitin specific peptidase 25	-2.9	1/6
220177_s_at	TMPRSS3↓	Transmembrane protease, serine 3	-2.8	1/6
Ion transport				
201735_s_at	CLCN3↓	Chloride channel 3	-2.8	2/6
205044_at	GABRP↓	Gamma-aminobutyric acid (GABA) A receptor, pi	-1.8	1/6
Actin binding				
212650_at	EHBP1↑	EH domain binding protein 1	2.6	6/0
GTPase activity				
202975_s_at	RHOBTB3↑	Rho-related BTB domain containing 3	2.5	7/1
203264_s_at	ARHGEF9↑	Cdc42 guanine nucleotide exchange factor (GEF) 9	2.6	6/0
Regulation of cell development, activation, growth and motility				
209890_at	TSPAN5↑	Tetraspanin 5	2.6	7/0
Small leucine-rich proteoglycan (slrp) family				
211896_s_at	DCN↑	Decorin	2.0	7/1
Protein kinase cascade				
211081_s_at	MAP4K5↓	Mitogen-activated protein kinase kinase kinase kinase 5	-2.3	1/7
Other				
206157_at	PTX3↓	Pentraxin-related gene, rapidly induced by IL-1 beta	-2.8	2/7
218431_at	Clorf133↓	Chromosome 14 open reading frame 133	-2.7	0/7
218514_at	FLJ10587↓	Hypothetical protein FLJ10587	-2.6	1/7
226760_at	LOC203411↑	Hypothetical protein LOC203411	3.0	6/2

↑ – upregulated in normal lobular cells; ↓ – downregulated in normal lobular cells

¹Affymetrix probe set number; If there was more than one probe set for the particular gene found among top 20 changed genes (according to the fold change), the probe set with the highest fold change is used and number of other probe set is indicated.

Genes differentially expressed between ductal and lobular carcinomas

In this comparison 208 probe sets were identified by pairwise comparison, 122 probe sets by RP, and 32 probe sets (28 named genes) by both methods (Table 4). In pairwise comparisons, the differentially expressed genes were identified by selecting those with count of changes more than 19 (out of 25 inter-patient comparison, every ductal carcinoma against every lobular carcinoma, either increase or decrease). These tumors were well separated by hierarchical clustering based on all 325 probe sets identified by RP and/or pairwise analysis (Figure 4). The expression of genes encoding proteins involved in cell adhesion was changed. Although CDH1 (E-cadherin), a classical member of the cadherin superfamily, was downregulated, THBS4 encoding calcium-binding adhesive glycoprotein, thrombospondin-4, was upregulated in all lobular carcinomas. DDR1 encoding receptor tyrosine kinase was overexpressed in ductal carcinomas.

The DVL1 gene encoding protein involved in Wnt signaling and the leucine-rich repeat protein ASPN (asporin) were upregulated in lobular carcinomas. Ductal carcinoma

showed upregulated genes which are involved in cell proliferation, signaling and cell cycle regulation, including RHOU, member of the Rho family of GTPases stimulating quiescent cells to reenter the cell cycle; PCSK6 encoding a calcium-dependent serine endoprotease; PRKCI encoding calcium-independent and phospholipid-dependent protein kinase C; PPP3CB encoding protein phosphatase 3; and CKS2 encoding a component of the CDC28 protein kinase. Epithelial membrane protein 1 (EMP1) was the only upregulated gene involved in cell growth and proliferation in lobular carcinomas. These changes were accompanied by the differential expression of transcription regulators. Majority of genes with this function were upregulated in ductal carcinomas such as AHCTF1, IRAK1, NRIP1, ADNP. Overexpressed genes in lobular carcinomas were the tumor suppressor FOXP1 and another transcription regulator MID1.

The genes encoding proteins involved in ubiquitin-mediated proteolysis, such as USP3, RKHD2 and TTC3, and nuclear components, such as DTL, GLCC11, TTC14, FAM54A, HIST1H3B, were upregulated in ductal carcinomas. Majority of genes encoding proteins with enzyme




Figure 2
Hierarchical clustering of normal ductal and lobular cells based on 82 probe sets found by both rank products and pairwise analysis. Ductal and lobular cells from the same patients tend to cluster together, and they can not be well separated from each other. This suggests expected similarities of expression signatures in normal epithelial (ductal and lobular) cells of mammary gland tree. 1–10, case numbers; D, ductal normal cells; L, lobular normal cells.

activity or implicated in metabolism were also upregulated in ductal carcinomas (*STK4*, *SLC1A2*, *B3GALT3*, *OSBPL10*, *CRBN*, *CHML*, *YWHAB*).

Two lobular and three ductal carcinomas were estrogen receptor-negative, whereas three lobular and two ductal carcinomas were estrogen receptor-positive. Hierarchical clustering using all probe sets was performed to determine whether receptor-positive and receptor-negative tumors could be separated. The tumors of the same histological type showed similar gene expression profiles without differences in relation to ER status as well as to other clinical parameters such as nodal status, stage and the expression of other immunohistochemical markers (data not shown).

Validation by immunohistochemistry on tissue microarrays

Seven differentially expressed genes (*KRT5*, *KRT6* and *KRT17* between tumor and normal cells, *CDH1*, *EMP1*, *DDR1* and *DVL1* between lobular and ductal carcinomas) were verified by immunohistochemical detection of proteins on TMA slides comprising of cores from 119 cases. The clinical and histopathological characteristics of these patients are shown in Table 5. The reduced expression or absence of cytokeratins 5/6 and 17 (*KRT5*, *KRT6*, *KRT17*) was found in both tumor tissues in comparison to terminal duct lobular units in 22 normal mammary tissues ($p < 0.0001$) (Table 6 and Figure 5). In a majority of ducts and lobules including TDLU, these cytokeratins were expressed in both basal and luminal cells, verifying the previously described variability of the expression of basal cytokeratins and their relationship to the cellular origin [51]. Cytokeratin 5 and 17 have also been found in a subset of breast cancer and identified patients with poor clinical outcome [31].

E-cadherin (*CDH1*) has successfully separated ductal and lobular invasive carcinomas. It was absent in 93.3% of lobular tumors compared with only 15% of ductal tumors ($p < 0.0001$). Epithelial membrane protein 1 (*EMP1*), discoidin domain receptor 1 (*DDR1*) and human homolog of the *Drosophila* dishevelled gene (*DVL1*) were found by pairwise comparison analysis to be differentially expressed between lobular and ductal carcinomas. Immunohistochemistry confirmed higher expression of *DVL1* and *EMP1* in lobular carcinomas and of *DDR1* in ductal carcinomas ($p < 0.0001$) (Table 7 and Figure 5). Of the special type carcinomas included on TMA slides, a papillary and two medullary carcinomas were positive for basal cytokeratins, one out of three medullary carcinomas was positive for *EMP1* and E-cadherin, a ductal-lobular carcinoma was positive for *DDR1*, all other special type carcinomas were negative for these markers, and finally none were positive for *DVL1* (data not shown).

Table 3: Genes differentially expressed between tumor versus normal cells by both rank products and pairwise analysis

Probe Set ¹	Gene Symbol	Gene Title	Fold-change			
			Tduc		Tlob	
			L	D	L	D
Lobular carcinoma vs normal cells (↑ – upregulated in Tlob, ↓ – downregulated in Tlob)						
Extracellular matrix-receptor interaction and focal adhesion (epithelial-mesenchymal transition)						
227314_at	ITGA2↓	Integrin, alpha 2	n	n	-4.7	-6.2
201438_at	COL6A3↑	Collagen, type VI, alpha 3	n	n	4.5	7.7
203083_at	THBS2↑	Thrombospondin 2	n	n	6.5	6.5
205713_s_at	COMP↑	Cartilage oligomeric matrix protein	n	n	6.7	8.1
226237_at	COL8A1↑	Collagen, type VIII, alpha 1	n	n	8.4	5.9
225681_at	CTHRC1↑	Collagen triple helix repeat containing 1	n	n	18.5	48.4
219087_at	ASPN↑	Asporin	n	n	23.3	22.1
WNT signalling						
223121_s_at 223122_s_at	SFRP2↑	Secreted frizzled-related protein 2	n	n	5.8 3.9	7.4 7.1
229802_at	WISP1↑	WNT1 inducible signaling pathway protein 1	n	n	9.1	12.4
Actin, calcium and metal ion binding						
204455_at	DST↓	Dystonin	n	n	-22.2	-18.3
214087_s_at	MYBPC1↓	Myosin binding protein C, slow type	n	n	-10.1	-12.2
209301_at	CA2↓	Carbonic anhydrase II	n	n	-9.2	-8.7
206509_at	PIP↓	Prolactin-induced protein	n	n	-7.0	-6.3
201131_s_at	CDH1↓	Cadherin 1, type I, E-cadherin	n	n	-6.2	-7.2
226777_at	ADAM12↑	ADAM metalloproteinase domain 12	n	n	5.7	5.3
213905_x_at	BGN/SDCCAG33↑	Biglycan/serologically defined colon cancer antigen 33	n	n	8.3	7.2
Other						
227742_at	CLIC6↓	Chloride intracellular channel 6	n	n	-9.3	-8.2
225817_at	CGNLI↓	Cingulin-like 1	n	n	-5.9	-7.8
219850_s_at	EHF↓	Ets homologous factor	n	n	-4.1	-6.3
213975_s_at	LYZ/ILRB1↑	Lysozyme	n	n	10.8	7.3
Ductal carcinoma vs normal cells (↓ – upregulated in Tduc, ↓ – downregulated in Tduc)						
Regulation of transcription						
202672_s_at	ATF3↓	Activating transcription factor 3	-4.5	-7.7	n	n
226147_s_at	PIGR↓	Polymeric immunoglobulin receptor	-4.8	-10	n	n
Cytokine and growth factor activity						
209466_x_at 211737_x_at	PTN↓	Pleiotrophin	-5.8 -7.1	-8.3 -12.5	n	n
823_at	CX3CL1↓	Chemokine (C-X3-C motif) ligand 1	-3.6	-8.3	n	n
Ion transport and metabolism						
203296_s_at	ATP1A2↓	ATPase, Na ⁺ /K ⁺ transporting, alpha 2	-8.2	-9.6	n	n
204259_at	MMP7↓	Matrix metalloproteinase 7	-4.5	-7.2	n	n

Table 3: Genes differentially expressed between tumor versus normal cells by both rank products and pairwise analysis (Continued)

212353_at	<i>SULF1</i> ↓	Sulfatase 1	4.6	4.0	n	n
Other						
1553602_at	<i>LOC118430</i> ↓	Small breast epithelial mucin	-7.1	-10.4	n	n
223623_at	<i>ECRG4</i> ↓	Esophageal cancer related gene 4 protein	-6.6	-5.6	n	n
206548_at	<i>FLJ23556</i> ↓	Hypothetical protein FLJ23556	-6.3	-11.2	n	n
218585_s_at	<i>DTL</i> ↑	Denticleless homolog (Drosophila)	3.6	4.8	n	n
Tumor vs normal cells (↑ – upregulated in tumor, ↓ – downregulated in tumor)						
Extracellular matrix-receptor interaction and focal adhesion (epithelial-mesenchymal transition)						
224823_at	<i>MYLK</i> ↓	Myosin, light polypeptide kinase	-3.5	-3.4	-5.9	-5.4
204620_s_at 221731_x_at	<i>CSPG2</i> ↑	Chondroitin sulfate proteoglycan 2 (versican)	3.0 3.5	4.9 5.3	4.5 4.0	5.8 6.1
221729_at	<i>COL5A2</i> ↑	Collagen, type V, alpha 2	3.5	4.4	9.3	11.3
211161_s_at	<i>COL3A1</i> ↑	Collagen, type III, alpha 1	4.1	3.3	9.8	14.9
212464_s_at	<i>FNI</i> ↑	Fibronectin 1	5.2	5.0	8.0	10.5
202311_s_at 202310_s_at	<i>COL1A1</i> ↑	Collagen, type I, alpha 1	5.6 5.1	6.0 5.1	12.7 10.4	13.0 10.8
37892_at	<i>COL11A1</i> ↑	Collagen, type XI, alpha 1	6.7	7.8	6.6	5.9
202404_s_at	<i>COL1A2</i> ↑	Collagen, type I, alpha 2	6.8	10.1	30.5	18.3
Ion and electron transport						
205044_at	<i>GABRP</i> ↓	Gamma-aminobutyric acid (GABA) A receptor, pi	-4.2	-8.7	-6.2	-8.6
220230_s_at	<i>CYB5R2</i> ↓	Cytochrome b5 reductase 2	-3.6	-3.8	-4.3	-4.2
Regulation of transcription						
220625_s_at	<i>ELF5</i> ↓	E74-like factor 5 (ets domain transcription factor)	-4.0	-4.8	-5.9	-8.4
209292_at	<i>ID4</i> ↓	Inhibitor of DNA binding 4	-2.9	-5.1	-10.8	-4.5
Actin and actin-binding proteins						
203951_at	<i>CNN1</i> ↓	Calponin 1, basic, smooth muscle	-6.7	-6.6	-6.9	-7.3
243140_at	<i>ACTA2</i> ↓	Actin, alpha 2, smooth muscle, aorta	-4.0	-4.1	-8.7	-4.0
WNT signaling						
202037_s_at	<i>SFRP1</i> ↓	Secreted frizzled-related protein 1	-7.1	-8.8	-8.5	-7.3
Intermediate filament proteins						
209351_at	<i>KRT14</i> ↓	Keratin 14	-22.9	-31.0	-21.9	-24.0
205157_s_at	<i>KRT17</i> ↓	Keratin 17	-17.7	-21.4	-9.9	-12.6
212730_at	<i>DMN</i> ↓	Desmuslin	-10.0	-10.4	-7.2	-7.5
201820_at	<i>KRT5</i> ↓	Keratin 5	-8.9	-12.7	-11.3	-12.1
Other						
229947_at	<i>PII5</i> ↓	Peptidase inhibitor 15	-6.5	-7.5	-18.5	-15.0
205380_at	<i>PDZK1</i> ↓	PDZ domain containing 1	-4.5	-5.8	-4.2	-6.4
213909_at	<i>LRRC15</i> ↑	Leucine rich repeat containing 15	6.2	5.7	7.3	11.8

Tduc – ductal tumor; Tlob – lobular tumor; D – normal ductal cells; L – normal lobular cells; n – not differentially expressed among top 20 genes. ¹Affymetrix probe set number; If there was more than one probe set for the particular gene found among top 20 changed genes (according to the fold change), the probe set with the highest fold change is used and number of other probe set is indicated.




Figure 3
Hierarchical clustering of tumor and normal samples based on 25 probe sets common to ductal and lobular carcinoma versus normal cells identified by both rank products and pairwise analysis. Tumor samples are grouped together, however, gene expression profiles of normal cell types from cases 10 and 3 are different from those of other normal cells, and tumor cells from case 2 are different from those of other tumor cells. 1–10, case numbers; D, ductal normal cells; L, lobular normal cells; Tduc, ductal tumor; Tlob, lobular tumor.

Validation of asporin expression by *in situ* hybridization

Combined RP and pairwise comparison showed that ASPN was one of the most upregulated genes in lobular carcinomas, when compared with normal ductal (fold change 23.3) and lobular (fold change 22.1) cells as well as with ductal carcinomas (fold change 3.9). None of the available antibodies against asporin worked on paraffin sections, therefore we utilised chromogenic *in situ* hybridization to detect asporin mRNA in frozen sections. All five lobular carcinomas appeared positive, whereas five cases of ductal carcinomas were negative or weakly and focally positive which is in agreement with the microarray data (Figure 6).

Validation of CTHRC1, ASPN and COL3A1 by PCR

Besides asporin, CTHRC1 was also upregulated in lobular cancer cells, when compared with normal ductal cells (fold change 48.4) and lobular cells (fold change 18.5), and COL3A1 was upregulated in both IDC and ILC cells, when compared with normal ductal cells (fold change 3.3 and 14.9, respectively) and lobular cells (fold change 4.1 and 9.8, respectively). As both RNA and cDNA were fully utilized for amplification, we used PCR amplification

products (see Methods) for validation by another PCR with specific primers. The results of semi-quantitative PCR correlated with the microarray data. Spearman coefficients were $r = 0.85$, $r = 0.73$ and $r = 0.70$ for CTHRC1, ASPN and COL3A1, respectively (all significant at $p < 0.0001$) (Figure 7).

Discussion

We examined 30 samples (microdissected tumor and normal ductal and lobular cells) from postmenopausal patients with lobular and ductal invasive breast carcinomas using Affymetrix arrays. Genes differentially expressed between the normal ductal and lobular cell types, which are less likely to be affected by fluctuating levels of female hormones as they were derived from postmenopausal women, are involved in ion transport and protein kinase cascade, protein biosynthesis and modification, proteolysis and metabolism, regulation of transcription and cell growth. However, hierarchical clustering of normal cell types based on the 82 probe sets identified by both RP and pairwise comparison showed that the gene expression profiles of normal ductal and lobular cells taken from the same patient are more similar to each other than ductal cells or lobular cells from different patients, and thus could not be well separated from each other. This may suggest expected similarities between expression signatures in normal epithelial (ductal and lobular) cells of the mammary gland. Since both IDC and ILC are believed to start in the terminal duct lobular unit (TDLU) of the breast [1,2], and these normal cell populations showed similar gene expression profiles, it is likely that the different morphological appearances of the two tumor types are mediated by differences in their mechanisms of carcinogenesis.

Combined pairwise comparison and RP analysis revealed that ductal and lobular carcinomas have a number of genes in common, however, they can be discriminated both at the gene and protein levels in our study. cDNA microarrays have previously been used to distinguish between IDC and ILC [38,39]. Unsupervised clustering of tumors failed to separate the two subtypes. There were 8 genes identified by MaxT permutation analysis using t tests, significance analysis for microarrays (SAM) and prediction analysis for microarrays (PAM) (E-cadherin, survivin, cathepsin B, TPI1, SPRY1, SCYA14, TFAP2B, and thrombospondin 4), and an additional 3 were identified by SAM and PAM (osteopontin, HLA-G, and CHC1) [38]. It has also been found that over half of ILCs differed from IDCs in global transcription programs, whereas the remaining ILCs closely resembled IDCs. Fifty two percent of the ILCs ("typical" ILCs) clustered together and displayed different gene expression profiles from the IDCs, whereas the other ILCs ("ductal-like" ILCs) were distributed between different IDC subtypes. Many of the differ-

Table 4: Genes differentially expressed in lobular versus ductal invasive carcinomas by rank products and/or pairwise analysis

Probe Set ¹	Gene Symbol	Gene Title	Rank products	Pairwise comparison
			Fold-change	Incr/Decr
Cell adhesion				
201131_s_at	CDH1↓	Cadherin 1, type 1, E-cadherin (epithelial)	-6.0	3/19
207169_x_a	DDR1↓	Discoidin domain receptor family, member 1	n	5/19
204776_at	THBS4↑	Thrombospondin 4	n	19/2
219087_at	ASPN↑	Asporin (LRR class 1)	3.9	23/0
Wnt signalling				
223168_at	RHOU↓	RAS homolog gene family, member U	n	2/20
203230_at	DVLI↑	Dishevelled, dsh homolog 1 (Drosophila)	n	20/0
Protein ubiquitination				
215392_at	USP3↓	Ubiquitin specific peptidase 3	-5.0	1/21
218247_s_at	RKHD2↓	Ring finger and KH domain containing 2	-4.6	0/19
1569472_s_at	TTC3↓	Tetratricopeptide repeat domain 3	-4.5	0/23
Regulation of transcription				
226115_at	AHCTF1↓	AT hook containing transcription factor 1	-4.0	0/22
201587_s_at	IRAK1↓	Interleukin-1 receptor-associated kinase 1	n	2/19
202599_s_at	NRIP1↓	Nuclear receptor interacting protein 1	n	0/19
201773_at	ADNP↓	Activity-dependent neuroprotector	n	0/20
203637_s_at	MIDI↑	Midline 1 (Opitz/BBB syndrome)	4.0	20/0
235444_at	FOXP1↑	Forkhead box P1	n	19/4
Electron and protein transport				
204351_at	S100P↓	S100 calcium binding protein P	-5.3	0/19
225177_at	RAB11FIP1↓	RAB11 family interacting protein 1 (class I)	-3.5	2/21
204041_at	MAOB↑	Monoamine oxidase B	3.5	20/4
212902_at	SEC24A↑	SEC24 related gene family, member A	3.5	19/2
Cell cycle regulation, cell proliferation and signaling				
204170_s_at	CKS2↓	CDC28 protein kinase regulatory subunit 2	-4.8	2/20
207414_s_at	PCSK6↓	Proprotein convertase subtilisin/kexin type 6	n	4/19
209678_s_at	PRKCI↓	Protein kinase C, iota	n	0/22
209817_at	PPP3CB↓	Protein phosphatase 3, catalytic subunit, beta isoform (calcineurin A beta)	n	0/24
201324_at	EMPI↑	Epithelial membrane protein 1	n	21/0
Metabolism and enzyme activity				
205509_at	CPB1↓	Carboxypeptidase B1 (tissue)	-11.1	0/19
223746_at	STK4↓	Serine/threonine kinase 4	-4.5	2/20
225491_at	SLC1A2↓	Solute carrier family 1 (glial high affinity glutamate transporter), member 2	-4.1	0/20
211379_x_at	B3GALT3↓	UDP-Gal:betaGlcNAc beta 1,3-galactosyltransferase, polypeptide 3	-4.0	0/20
219073_s_at	OSBPL10↓	Oxysterol binding protein-like 10	-4.0	1/21
229366_at	CRBN↓	Cereblon	-3.6	0/20
226350_at	CHML↓	Choroideremia-like (Rab escort protein 2)	n	0/24
217717_s_at	YWHA3↓	Tyrosine 3-monooxygenase/tryptophan 5-mono-oxygenase activation protein, beta polypeptide	n	1/20
49452_at	ACACB↑	Acetyl-Coenzyme A carboxylase beta	n	19/0
220770_s_at	LOC63920↑	Transposon-derived Buster3 transposase-like	5.3	24/0
Actin and cytoskeletal protein binding				
211776_s_at	EPB41L3↓	Erythrocyte membrane protein band 4.1-like 3	-5.3	0/21
201341_at	ENCI↓	Ectodermal-neural cortex (with BTB-like domain)	n	19/3
Nuclear components				
218585_s_at	DTL↓	Denticleless homolog (Drosophila)	-4.6	1/19
225706_at	GLCC1↓	Glucocorticoid induced transcript 1	-4.5	0/25
241863_x_at	TTC14↓	Tetratricopeptide repeat domain 14	-4.4	1/20
228069_at	FAM54A↓	Family with sequence similarity 54, member A	-3.8	0/20
208576_s_at	HIST1H3B↓	Histone 1, H3b	-3.3	1/19
Other				
1568838_at	---	Full-length cDNA clone CS0DF033YE17 of Fetal brain of Homo sapiens (human)	-5.0	1/20
226067_at	C20orf114↓	Chromosome 20 open reading frame 114	-4.9	0/19
1553979_at	---	Homo sapiens, clone IMAGE:3906992, mRNA	-4.1	0/19
230534_at	MGC15634↓	Hypothetical protein MGC15634	-3.6	2/20
226671_at	---	CDNA clone IMAGE:4797120	-3.8	2/20
219768_at	VTCN1↑	V-set domain containing T cell activation inhibitor 1	7.8	21/2
214657_s_at	TncRNA↑	Trophoblast-derived noncoding RNA	4.5	21/0
213004_at	ANGPTL2↑	Angiopoietin-like 2	n	19/4

↑ – upregulated in lobular carcinoma; ↓ – downregulated in lobular carcinoma; n – gene not found by rank products at the level of 10% of false positives, Incr/decr – increase or decrease in lobular versus ductal carcinoma by pairwise comparison.

¹All 32 probe sets identified by combined pairwise comparison and rank products are included, also a number of genes found by pairwise comparison were validated by immunohistochemistry (DVLI, DDR1, EMPI), therefore, functional categories also include named genes identified by pairwise analysis.




Figure 4
Hierarchical clustering of invasive ductal and lobular breast carcinomas based on 325 probe sets identified by rank products and/or pairwise analysis. Tumor types are well separated. 1–10, case numbers; Tduc, ductal tumor; Tlob, lobular tumor.

entially expressed genes encode for proteins involved in cell adhesion/motility, lipid/fatty acid transport and metabolism, immune/defense response, and electron transport. Many genes distinguishing between typical and ductal-like ILCs are involved in regulation of cell growth and immune response [39]. However, these two previous studies examined whole tumor tissues without microdissection, suggesting that expression of a number of genes could be related not only to tumor cells but also to other components of mammary tissue such as stromal, adipose, endothelial etc. Our study is the first full genome analysis

of microdissected ductal and lobular tumor and normal cells reporting both normal mammary epithelium- and cancer-specific genes expression profiles.

Importantly, *CDH1* (E-cadherin gene) was downregulated in our lobular carcinomas, and immunohistochemistry confirmed this loss at the protein level within tumors. E-cadherin is considered to be the most important cell adhesion molecule in the mammary gland. It acts as a tumor suppressor inhibiting invasion and metastasis. Mutations of this gene are correlated with gastric, breast, colorectal, thyroid and ovarian cancer. During tumor progression, E-cadherin can be functionally inactivated or silenced by different mechanisms such as post-translational control, somatic mutations, downregulation of gene expression through promoter hypermethylation, histone deacetylation, and transcriptional repression [52,53]. The latter induces cellular responses leading to the conversion of epithelial cells into invasive mesenchymal-like cells with increased motility and invasiveness, and this process is called an epithelial-mesenchymal transition (EMT) [52]. To date, it is believed that lost, non-polar or cytoplasmic expression of E-cadherin protein and/or transcriptional repression of its mRNA are hallmarks of EMT in cancer progression [53-55]. It has also been shown that several proteins such as fibronectin and integrin $\alpha\beta 6$ [54], Ets, TGF β , FGF-1,-2,-8, α -SMA, collagen type I, III and thrombospondins increase in abundance during EMT [56], conversely, amongst proteins that decrease in abundance are E-cadherin and cytokeratins [54]. According to our results, collagen type I and III, fibronectin and Ets domain transcription factor are upregulated and cytokeratins are downregulated in both tumor cell types. The expression of collagens and other mesenchyme-associated genes in microdissected breast cancer cells was also confirmed by Nishidate and co-workers [57]. In addition, pairwise comparisons revealed that thrombospondin 4 was upregulated only in lobular cancer cells, which agrees with the literature [38]. Thus we propose that the EMT plays a role in both tumor types but appears to be more important in lobular carcinomas. The EMT phenomenon seems to be promising because multiple molecules involved in EMT, such as receptor- and SRC-family tyrosine kinases, RAS and other small GTPases, can be envisioned as targets for anti-EMT therapy [58].

The tissue microenvironment, including the ECM-cell and cell-cell interactions, plays an important role in both normal mammary gland development and cancer. Neoplastic transformation of cells dramatically alters the synthesis of proteoglycans and other ECM proteins both in tumor and the surrounding matrix [59]. This can stimulate the growth and spread of tumor cells by decreasing the adhesive functions of the ECM [60]. Both tumor types examined show upregulated genes involved in tumor-ECM

Table 5: Clinical and histopathological characteristics of TMA cases

Features	IDC	ILC
G1	6 (7.5%)	15 (51.7%)
G2	57 (71.3%)	11 (37.9%)
G3	17 (21.2%)	3 (10.3%)
ER+	57 (71.3%)	19 (65.5%)
PgR+	50 (62.5%)	17 (58.6%)
HER-2+	12 (15%)	1 (3.4%)
Bcl-2+	13 (16.3%)	4 (13.8%)
N ₀	56 (70%)	21 (72.4%)
N ₁	16 (20%)	6 (20.7%)
N ₂	8 (10%)	2 (6.9%)
Total number of cases	80	29

ER, estrogen receptor; G1-3, grade 1-3; IDC, invasive ductal carcinoma; ILC, invasive lobular carcinoma; N₀₋₁, lymph node status; PgR, progesterone receptor; TMA, tissue microarray.

interactions, cell adhesion and migration processes including metastasis. Expression of majority of the proteins encoded by these genes is related to TGFβ or Wnt signaling, and both the TGFβ and Wnt pathways may affect ECM.

Asporin is a cartilage extracellular protein that has been reported to be associated with knee and hip osteoarthritis. This leucine-rich repeat protein was shown to interact with and inhibit TGFβ signaling which is thought to lead to insufficient quantities of aggrecan and type II collagen in osteoarthritis [61,62]. Asporin was more upregulated in our lobular carcinomas when compared with ductal tumors as well as with normal cell types. Overexpression of asporin mRNA in lobular carcinomas was then confirmed by chromogenic *in situ* hybridization and PCR. In support of our findings, upregulation of this gene has also been described in microdissected androgen-independent prostate cancer cells using Affymetrix Human Genome U133A GeneChips [63]. The authors did not discuss it and importantly, asporin has not been related to carcinogenesis to date.

We have found another candidate gene, collagen triple helix repeat containing 1 (*CTHRC1*), which was upregulated in ILC in comparison with normal cells, and its expression was also validated by PCR. Aberrant expression of *CTHRC1* has recently been reported in human solid tumors, including cancers of the gastrointestinal tract,

lung, breast, thyroid, ovarian, cervix, liver, and the pancreas. It is associated with cancer tissue invasion and metastasis and potentially plays important functional roles in cancer progression, perhaps by increasing cancer cell migration [64]. TGFβ upregulates *CTHRC1*, versican, *ADAM12*, and downregulates *SFRP1* and E-cadherin [65-67]. The loss of *SFRP1* is known to be associated with breast cancer progression and poor prognosis in early stages [68], and a similar expression profile is seen in our study. *SFRP1* is downregulated and versican is upregulated in both tumor types. Furthermore, pairwise comparison identified other overexpressed genes in lobular carcinoma such as *SFRP2* and *ADAM12*. E-cadherin, which is downregulated in lobular cancer cells, can also be repressed by TGFβ-induced expression of transcription factor complexes [69]. TGFβ signaling inhibitors have been shown to prevent EMT, to inhibit mammary tumor viability and to block metastasis in various murine models [70]. According to these results, we propose that deregulated TGFβ signaling is likely to be more important in lobular carcinogenesis.

Wnt signaling molecules are also expressed in our samples in several comparisons. The Wnt proteins are small secreted glycoproteins which are involved in the self-renewal of stem cells and may be responsible for the maintenance of mature tissues [71]. On binding to Frizzled receptors, Wnts can activate canonical (β-catenin-dependent) and/or non-canonical (β-catenin-independ-

Table 6: Immunohistochemical staining of cytokeratins in tumor and normal tissues

Antibodies	Tumors	Normal tissues (n = 22)	Fisher's p value
Cytokeratin 5/6	Ductal carcinomas (n = 80)	9 (11.2%)	<0.0001
	Lobular carcinomas (n = 29)	3 (10.3%)	<0.0001
Cytokeratin 17	Ductal carcinomas (n = 80)	8 (10%)	<0.0001
	Lobular carcinomas (n = 29)	2 (6.9%)	<0.0001




Figure 5
Representative immunohistochemical staining for the selected proteins: E-cadherin, DDR1, DVLI, cytokeratin 5/6, cytokeratin 17 and EMPI. 1.1. E-cadherin is negative in lobular carcinoma; 1.2. E-cadherin is positive in ductal carcinoma; 2.1. DDR1 is negative in lobular carcinoma; 2.2. DDR1 is positive in ductal carcinoma; 3.1. DVLI is positive in lobular carcinoma; 3.2. DVLI is negative in ductal carcinoma; 4.1. Cytokeratin 5/6 is negative in lobular carcinoma cells, but its expression is retained in normal ductal epithelial cells; 4.2. Cytokeratin 5/6 is negative in ductal carcinoma cells; 4.3. Duct lobular unit in normal mammary gland tissue is positive for cytokeratin 5/6; 5.1. Cytokeratin 17 is negative in lobular carcinoma cells, but its expression is retained in normal ductal epithelial cells; 5.2. Cytokeratin 17 is negative in ductal carcinoma cells, but its expression is retained in normal ductal epithelial cells; 5.3. Duct lobular unit in normal mammary gland tissue is positive for cytokeratin 17; 6.1. EMPI is positive in lobular carcinoma; 6.2. EMPI is negative in ductal carcinoma.

ent Wnt/planar cell polarity pathway and Wnt/Ca²⁺ pathway) Wnt signaling [72,73]. Downregulation of frizzled related proteins has been described in breast cancer [68]. Of these, *SFRP1* is downregulated in both types of our tumor cells, and genes involved in calcium regulation

pathway are not significantly changed. The promoter of fibronectin 1 contains LEF/TCF-binding sites, making it a direct target of canonical Wnt signaling [59,74]. Fibronectin 1 is also upregulated in both tumor types. However, there is a difference in Wnt signaling between ILC and

Table 7: Immunohistochemistry differentiates ductal and lobular carcinomas

Antibodies	Ductal carcinomas (n = 80)	Lobular carcinomas (n = 29)	Fisher's p value
E-cadherin	68 (85%)	2 (6.7%)	<0.0001
DVLI	20 (25%)	28 (96.5%)	<0.0001
EMPI	13 (16.3%)	27 (93.1%)	<0.0001
DDR1	77 (96.2%)	4 (13.8%)	<0.0001




Figure 6
Asporin mRNA detection by chromogenic *in situ* hybridization. 1–2. Various magnifications ($\times 100$, $\times 400$, $\times 400$) of lobular carcinoma; red arrows, lobular tumor cells are positive; blue arrows, stromal cells are negative; 3. Ductal carcinoma is negative ($\times 200$).

IDC. *SULF1* (sulfatase 1) is upregulated and *MMP7* is downregulated only in IDCs. *MMP7* is a confirmed Wnt target [75] and it has been shown to be activated in both canonical [76,77] and non-canonical Wnt signaling [78]. Binding Wnt ligands to frizzled receptors is regulated by the 6-O sulfation-desulfation of cell surface heparan sulfates (HSs) by sulfatase 1. Sulfated HSs bind to Wnt ligand with high affinity and inhibit Wnt signaling. Sulfatase 1 removes 0–6 sulfates from HSs and reduces their binding to Wnt ligands which in turn allows the formation of functional Wnt-Frizzled complexes and thus promotes Wnt signaling [79,80]. Frizzled-related proteins also have heparin-binding domains that promote the formation of Wnt-Frizzled complexes [81]. Although *SULF1* is upregulated, *SFRP1* and *MMP7* are downregulated in ductal cancer cells, whereas *SFRP2* and other Wnt molecules, such as *DVL1* and *WISP1*, are upregulated in lobular cancer cells. This suggests that Wnt signaling is activated in ILC cells but not in IDC cells. Since mRNA level of β -catenin was not changed between normal and tumor cells, the expression of Wnt molecules appears to be β -catenin-independent, favoring the non-canonical Wnt signaling in ILC. There is evidence that Wnts acting through the non-canonical pathway can promote tumor progression [82,83] which may also be true in ILC.

In addition to Wnt molecules such as *DVL1*, pairwise comparison revealed that *EMP1*, gene encoding a tumor-associated membrane protein involved in cell-cell interactions and proliferation control [84], was upregulated, whereas *DDR1*, epithelial-specific receptor kinase capable of binding Wnt5 and regulating the adhesion of mammary cells [85,86], was downregulated in ILCs. Immunohistochemistry has also confirmed the same differential expression of these three proteins in IDC and ILC on tis-

sue microarrays. Thus all the evidence suggests that the two tumor types can be distinguished both at the gene and protein levels. Specific changes in gene and protein expression are likely to reflect the differences in mechanisms of carcinogenesis as well as the specific histological and clinical characteristics of these tumors derived from the same anatomical compartment, TDLU.

Conclusion

Microdissection of normal and tumor cell types from the breast and full genome expression analysis by Affymetrix arrays allowed us to provide novel data on breast cancer. Invasive lobular and ductal breast carcinomas can be differentiated both at the gene and protein levels. Despite analyzing only thirty samples from ten patients, the results are in good accordance with previous literature [14,38,39,57,64]. Our data provide evidence for deregulated TGF β and Wnt signaling accompanied by the over-expression of mesenchyme-associated genes like the collagens, asporin and others which might be occurring in conjunction with an altered EMT. We propose that deregulated TGF β signaling and EMT phenomenon are involved in both tumor types, but they seem to be more important in lobular carcinomas which is in concordance with the loss of E-cadherin expression and their distinct morphology from ductal tumors. In this study we report two candidate genes, asporin (*ASPN*) and collagen triple helix repeat containing 1 (*CTHRC1*), which might be significant in mammary gland carcinogenesis and may also be important either in cancer diagnosis or therapy. Besides E-cadherin, the proteins validated on tissue microarrays by immunohistochemistry (*EMP1*, *DVL1*, *DDR1*) may represent novel tissue markers helpful in the differentiation of ductal and lobular cancers. Further studies with larger sets of patients are needed to verify the gene




Figure 7
PCR validation of microarray results for CTHRC1 (1), ASPN (2), and COL3A1 (3). Fluorescence signals from Affymetrix probe sets (225681_at, 219087_at and 211161_s_at, respectively) and optical density of PCR bands (OD × cm) were transformed to a percentage of the highest value. 1–10, case numbers; Tduc, ductal tumor; Tlob, lobular tumor; D, normal ductal cells; L, normal lobular cells.

expression profiles of various histological types of breast cancer in order to determine molecular subclassifications, prognosis, and the optimum treatment strategies.

Abbreviations

ADAM12 = ADAM metallopeptidase domain 12; ASPN = Asporin; CK = Cytokeratin; CTHRC1 = Collagen triple helix repeat containing 1; DAVID = Database for Annotation, Visualization and Integrated Discovery; DDR1 = Discoidin domain receptor 1; DVL1 = Human homolog of the *Drosophila* dishevelled gene; ECM = Extracellular matrix; EMP1 = Epithelial membrane antigen 1; EMT = Epithelial-mesenchymal transition; ER = Estrogen receptor; GCOS = GeneChip Operating Software; IDC = Invasive ductal carcinoma; ILC = Invasive lobular carcinoma; ISH = *In situ* hybridization; IVT = *In vitro* transcription; PCR = Polymerase chain reaction; PgR = Progesterone receptor; RMA = Robust multiarray analysis; RNA = Ribonucleic acid; RP = Rank products analysis; SFRP1 = Secreted frizzled-related protein 1; TDLU = Terminal duct lobular unit; TGF β = Transforming growth factor β ; TMA = Tissue microarray; WISP1 = WNT1 inducible signaling pathway protein 1.

Competing interests

The author(s) declare that they have no competing interests.

Authors' contributions

GT and JB contributed equally to this work. GT participated in collecting surgical specimens, cutting and evaluation of frozen sections, laser microdissection, immunohistochemistry, ISH, analysis of microarray results and drafting the manuscript. JB participated in the design of the study, collecting surgical specimens, RNA isolation, amplification and labeling, analysis of microarray results, ISH and PCR validation, drafting the manuscript. KB carried out the hybridization and scanning of the arrays and proofreading of the manuscript. WW performed data analysis and participated in drafting the manuscript. MD and MH were involved in laser microdissection. JE participated in collection and evaluation of frozen sections. JK participated in obtaining surgical specimens. EF and JS participated in tissue microarray construction. JS was involved in RNA isolation and PCR validation. PM participated in analysis of microarray results. ZK conceived of the study, participated in its design and coordination, as well as in collecting surgical specimens, evaluation of frozen sections and drafting the manuscript. All authors read and approved the final manuscript.

Additional material

Additional File 1

Supplementary tables. The data provided represent supplementary tables listing all differentially expressed probe sets found by both rank products and pairwise analysis between normal ductal and normal lobular cells, ductal carcinoma and normal ductal cells, ductal carcinoma and normal lobular cells, ductal carcinoma and lobular carcinoma cells, lobular carcinoma and normal ductal cells, and lobular carcinoma and normal lobular cells.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2407-7-55-S1.doc>]

Acknowledgements

The work was supported by grants NR 7844-3 and NR 8425-3 from the Czech Ministry of Health and MSM 6198959216 from the Czech Ministry of Education. The authors cordially thank to Jirina Zatloukalova, Jirina Prochazkova, Prof. George Burkadze, Dalimil Zurek, Veronika Krejci, Pravomila Abrahamova, Alena Lukasova, Renata Bilkova and Gabriela Korinkova for scientific discussions and kind technical assistance.

References

- Rosen PP: *Rosen's breast pathology* Lippincott-Raven Publishers; 1997.
- Weidner N, Cote R, Suster S, Weiss L: *Modern Surgical Pathology* Elsevier Science; 2003.
- Winchester DJ, Chang HR, Graves TA, Menck HR, Bland KI, Winchester DP: **A comparative analysis of lobular and ductal carcinoma of the breast: presentation, treatment, and outcomes.** *J Am Coll Surg* 1998, **186**:416-422.
- Silverstein MJ, Lewinsky BS, Waisman JR, Gierson ED, Colburn WJ, Senofsky GM, Gamagami P: **Infiltrating lobular carcinoma. Is it different from infiltrating duct carcinoma?** *Cancer* 1994, **73**:1673-1677.
- Toikkanen S, Pylkkanen L, Joensuu H: **Invasive lobular carcinoma of the breast has better short- and long-term survival than invasive ductal carcinoma.** *Br J Cancer* 1997, **76**:1234-40.
- Molland JG, Donnellan M, Janu NC, Carmalt HL, Kennedy CW, Gillett DJ: **Infiltrating lobular carcinoma – a comparison of diagnosis, management and outcome with infiltrating duct carcinoma.** *Breast* 2004, **13**:389-396.
- Newman LA, Buzdar AU, Singletary SE, Kuerer HM, Buchholz T, Ames FC, Ross MI, Hunt KK: **A prospective trial of preoperative chemotherapy in resectable breast cancer: predictors of breast-conservation therapy feasibility.** *Ann Surg Oncol* 2002, **9**:228-234.
- Mersin H, Yildirim E, Gulben K, Berberoglu U: **Is invasive lobular carcinoma different from invasive ductal carcinoma?** *Eur J Surg Oncol* 2003, **29**:390-395.
- Arpino G, Bardou VJ, Clark GM, Elledge RM: **Infiltrating lobular carcinoma of the breast: tumor characteristics and clinical outcome.** *Breast Cancer Res* 2004, **6**:R149-156.
- Perou CM, Jeffrey SS, van de Rijn M, Rees CA, Eisen MB, Ross DT, Pergamenschikov A, Williams CF, Zhu SX, Lee JC, Lashkari D, Shalon D, Brown PO, Botstein D: **Distinctive gene expression patterns in human mammary epithelial cells and breast cancers.** *Proc Natl Acad Sci USA* 1999, **96**:9212-9217.
- Schena M, Shalon D, Davis RW, Brown PO: **Quantitative monitoring of gene expression patterns with a complementary DNA microarray.** *Science* 1995, **270**:467-470.
- Bertucci F, Viens P, Hingamp P, Nasser V, Houlgatte R, Birnbaum D: **Breast cancer revisited using DNA array-based gene expression profiling.** *Int J Cancer* 2003, **103**:565-571.
- Tavassoli FA, Schnitt SJ: *Pathology of the breast* New York: Elsevier; 1992.
- Zhu G, Reynolds L, Crnogorac-Jurcevic T, Gillett CE, Dublin EA, Marshall JF, Barnes D, D'Arrigo C, Van Trappen PO, Lemoine NR, Hart

- IR: **Combination of microdissection and microarray analysis to identify gene expression changes between differentially located tumor cells in breast cancer.** *Oncogene* 2003, **22**:3742-3748.
15. Schuetz CS, Bonin M, Clare SE, Nieselt K, Sotlar K, Walter M, Fehm T, Solomayer E, Riess O, Wallwiener D, Kurek R, Neubauer HJ: **Progression-specific genes identified by expression profiling of matched ductal carcinomas in situ and invasive breast tumors, combining laser capture microdissection and oligonucleotide microarray analysis.** *Cancer Res* 2006, **66**:5278-5286.
 16. Yang F, Foekens JA, Yu J, Sieuwerts AM, Timmermans M, Klijn JG, Atkins D, Wang Y, Jiang Y: **Laser microdissection and microarray analysis of breast tumors reveal ER-alpha related genes and pathways.** *Oncogene* 2006, **25**:1413-1419.
 17. Aoyagi K, Tatsuta T, Nishigaki M, Akimoto S, Tanabe C, Omoto Y, Hayashi S, Sakamoto H, Sakamoto M, Yoshida T, Terada M, Sasaki H: **A faithful method for PCR-mediated global mRNA amplification and its integration into microarray analysis on laser-captured cells.** *Biochem Biophys Res Commun* 2003, **300**:915-920.
 18. Luzzi V, Mahadevappa M, Raja R, Warrington JA, Watson MA: **Accurate and reproducible gene expression profiles from laser capture microdissection, transcript amplification, and high density oligonucleotide microarray analysis.** *J Mol Diagn* 2003, **5**:9-14.
 19. Klur S, Toy K, Williams MP, Certa U: **Evaluation of procedures for amplification of small-size samples for hybridization on microarrays.** *Genomics* 2004, **83**:508-517.
 20. Ji W, Zhou W, Gregg K, Lindpaintner K, Davis S, Davis S: **A method for gene expression analysis by oligonucleotide arrays from minute biological materials.** *Anal Biochem* 2004, **331**:329-339.
 21. Sorlie T, Tibshirani R, Parker J, Hastie T, Marron JS, Nobel A, Deng S, Johnsen H, Pesich R, Geisler S, Demeter J, Perou CM, Lonning PE, Brown PO, Borresen-Dale AL, Botstein D: **Repeated observation of breast tumor subtypes in independent gene expression data sets.** *Proc Natl Acad Sci USA* 2003, **100**:8418-8423.
 22. Bertucci F, Houlgatte R, Granjeaud S, Nasser V, Lorioud B, Beaudoing E, Hingamp P, Jacquemier J, Viens P, Birnbaum D, Nguyen C: **Prognosis of breast cancer and gene expression profiling using DNA arrays.** *Ann N Y Acad Sci* 2002, **975**:217-231.
 23. Reinholz MM, Iturria SJ, Ingle JN, Roche PC: **Differential gene expression of TGF-beta family members and osteopontin in breast tumor tissue: analysis by real-time quantitative PCR.** *Breast Cancer Res Treat* 2002, **74**:255-269.
 24. Mackay A, Jones C, Dexter T, Silva RL, Bulmer K, Jones A, Simpson P, Harris RA, Jat PS, Neville AM, Reis LF, Lakhani SR, O'Hare MJ: **cDNA microarray analysis of genes associated with ERBB2 (HER2/neu) overexpression in human mammary luminal epithelial cells.** *Oncogene* 2003, **22**:2680-2688.
 25. Barsky SH: **Myoepithelial mRNA expression profiling reveals a common tumor-suppressor phenotype.** *Exp Mol Pathol* 2003, **74**:113-122.
 26. Pusztai L, Sotiriou C, Buchholz TA, Meric F, Symmans WF, Esteva FJ, Sahin A, Liu ET, Hortobagyi GN: **Molecular profiles of invasive mucinous and ductal carcinomas of the breast: a molecular case study.** *Cancer Genet Cytogenet* 2003, **141**:148-153.
 27. Dressman MA, Baras A, Malinowski R, Alvis LB, Kwon I, Walz TM, Polymeropoulos MH: **Gene expression profiling detects gene amplification and differentiates tumor types in breast cancer.** *Cancer Res* 2003, **63**:2194-2199.
 28. Nyante SJ, Devries S, Chen YY, Hwang ES: **Array-based comparative genomic hybridization of ductal carcinoma in situ and synchronous invasive lobular cancer.** *Hum Pathol* 2004, **35**:759-763.
 29. Sotiriou C, Neo SY, McShane LM, Korn EL, Long PM, Jazaeri A, Martiat P, Fox SB, Harris AL, Liu ET: **Breast cancer classification and prognosis based on gene expression profiles from a population-based study.** *Proc Natl Acad Sci USA* 2003, **100**:10393-10398.
 30. Sorlie T, Perou CM, Tibshirani R, Aas T, Geisler S, Johnsen H, Hastie T, Eisen MB, van de Rijn M, Jeffrey SS, Thorsen T, Quist H, Matese JC, Brown PO, Botstein D, Eystein Lonning P, Borresen-Dale AL: **Gene expression patterns of breast carcinomas distinguish tumor subclasses with clinical implications.** *Proc Natl Acad Sci USA* 2001, **98**:10869-10874.
 31. van de Rijn M, Perou CM, Tibshirani R, Haas P, Kallioniemi O, Kononen J, Torhorst J, Sauter G, Zuber M, Kochli OR, Mross F, Dieterich H, Seitz R, Ross D, Botstein D, Brown P: **Expression of cytokeratins 17 and 5 identifies a group of breast carcinomas with poor clinical outcome.** *Am J Pathol* 2002, **161**:1991-1996.
 32. van't Veer LJ, Dai H, van de Vijver MJ, He YD, Hart AA, Bernards R, Friend SH: **Expression profiling predicts outcome in breast cancer.** *Breast Cancer Res* 2003, **5**:57-58.
 33. Hedenfalk I, Duggan D, Chen Y, Radmacher M, Bittner M, Simon R, Meltzer P, Gusterson B, Esteller M, Kallioniemi OP, Wilfond B, Borg A, Trent J, Raffeld M, Yakhini Z, Ben-Dor A, Dougherty E, Kononen J, Bubendorf L, Fehrle W, Pittaluga S, Gruberberger S, Loman N, Johannsson O, Olsson H, Sauter G: **Gene-expression profiles in hereditary breast cancer.** *N Engl J Med* 2001, **344**:539-548.
 34. Grigoriadis A, Mackay A, Reis-Filho JS, Steele D, Iseli C, Stevenson BJ, Jongeneel CV, Valgeirsson H, Fenwick K, Iravani M, Leao M, Simpson AJ, Strausberg RL, Jat PS, Ashworth A, Neville AM, O'hare MJ: **Establishment of the epithelial-specific transcriptome of normal and malignant human breast cells based on MPSS and array expression data.** *Breast Cancer Res* 2006, **8**:R56.
 35. Huiping C, Sigurgeirsdottir JR, Jonasson JG, Eiriksdottir G, Johannsdottir JT, Egilsson V, Ingvarsson S: **Chromosome alterations and E-cadherin gene mutations in human lobular breast cancer.** *Br J Cancer* 1999, **81**:1103-1110.
 36. Asgeirsson KS, Jonasson JG, Tryggvadottir L, Olafsdottir K, Sigurgeirsdottir JR, Ingvarsson S, Ogmundsdottir HM: **Altered expression of E-cadherin in breast cancer: patterns, mechanisms and clinical significance.** *Eur J Cancer* 2000, **36**:1098-1106.
 37. Sarrio D, Perez-Mies B, Hardisson D, Moreno-Bueno G, Suarez A, Cano A, Martin-Perez J, Gamallo C, Palacios J: **Cytoplasmic localization of p120ctn and E-cadherin loss characterize lobular breast carcinoma from preinvasive to metastatic lesions.** *Oncogene* 2004, **23**:3272-3283.
 38. Korkola JE, DeVries S, Fridlyand J, Hwang ES, Estep AL, Chen YY, Chew KL, Dairkee SH, Jensen RM, Waldman FM: **Differentiation of lobular versus ductal breast carcinomas by expression microarray analysis.** *Cancer Res* 2003, **63**:7167-7175.
 39. Zhao H, Langerod A, Ji Y, Nowels KW, Nesland JM, Tibshirani R, Bukholm IK, Karesen R, Botstein D, Borresen-Dale AL, Jeffrey SS: **Different gene expression patterns in invasive lobular and ductal carcinomas of the breast.** *Mol Biol Cell* 2004, **15**:2523-2536.
 40. Breiting R, Armengaud P, Amtmann A, Herzyk P: **Rank products: a simple, yet powerful, new method to detect differentially regulated genes in replicated microarray experiments.** *FEBS Lett* 2004, **573**:83-92.
 41. Irizarry RA, Bolstad BM, Collin F, Cope LM, Hobbs B, Speed TP: **Summaries of Affymetrix GeneChip probe level data.** *Nucleic Acids Res* 2003, **31**:e15.
 42. **Affymetrix package of the Bioconductor** [<http://www.bioconductor.org>]
 43. **DAVID: Database for Annotation, Visualization and Integrated Discovery** [<http://david.abcc.ncifcrf.gov>]
 44. **Gene Cards** [<http://www.genecards.org>]
 45. **NETAFFX Analysis Center** [<http://www.affymetrix.com/analysis/>]
 46. **dChip software: Gene Expression Microarray and SNP Microarray Analysis** [<http://www.dchip.org>]
 47. **NCBI's Gene Expression Omnibus** [<http://www.ncbi.nlm.nih.gov/geo/>]
 48. Simon R, Sauter G: **Tissue microarrays for miniaturized high-throughput molecular profiling of tumors.** *Exp Hematol* 2002, **30**:1365-1372.
 49. Henry SP, Takanosu M, Boyd TC, Mayne PM, Eberspaecher H, Zhou W, de Crombrughe B, Hook M, Mayne R: **Expression pattern and gene characterization of asporin, a newly discovered member of the leucine-rich repeat protein family.** *J Biol Chem* 2001, **276**:12212-12221.
 50. Nakamura S, Terashima T, Yoshida T, Iseki S, Takano Y, Ishikawa I, Shinomura T: **Identification of genes preferentially expressed in periodontal ligament: specific expression of a novel secreted protein, FDC-SP.** *Biochem Biophys Res Commun* 2005, **338**:1197-1203.
 51. Gusterson BA, Ross DT, Heath VJ, Stein T: **Basal cytokeratins and their relationship to the cellular origin and functional classification of breast cancer.** *Breast Cancer Res* 2005, **7**:143-8.
 52. Peinado H, Portillo F, Cano A: **Transcriptional regulation of cadherins during development and carcinogenesis.** *Int J Dev Biol* 2004, **48**:365-375.

53. Grille SJ, Bellacosa A, Upson J, Klein-Szanto AJ, van Roy F, Lee-Kwon W, Donowitz M, Tschichl PN, Larue L: **The protein kinase Akt induces epithelial mesenchymal transition and promotes enhanced motility and invasiveness of squamous cell carcinoma lines.** *Cancer Res* 2003, **63**:2172-2178.
54. Thiery JP: **Epithelial-mesenchymal transitions in development and pathologies.** *Curr Opin Cell Biol* 2003, **15**:740-746.
55. Thiery JP: **Epithelial-mesenchymal transitions in tumour progression.** *Nat Rev Cancer* 2002, **2**:442-454.
56. Kalluri R, Neilson EG: **Epithelial-mesenchymal transition and its implications for fibrosis.** *J Clin Invest* 2003, **112**:1776-1784.
57. Nishidate T, Katagiri T, Lin ML, Mano Y, Miki Y, Kasumi F, Yoshimoto M, Tsunoda T, Hirata K, Nakamura Y: **Genome-wide gene-expression profiles of breast-cancer cells purified with laser microbeam microdissection: identification of genes associated with progression and metastasis.** *Int J Oncol* 2004, **25**:797-819.
58. Cheng JQ, Lindsley CW, Cheng GZ, Yang H, Nicosia SV: **The Akt/PKB pathway: molecular target for cancer drug discovery.** *Oncogene* 2005, **24**:7482-7492.
59. Schambony A, Kunz M, Gradl D: **Cross-regulation of Wnt signaling and cell adhesion.** *Differentiation* 2004, **72**:307-318.
60. Esko JD, Rostand KS, Weinke JL: **Tumor formation dependent on proteoglycan biosynthesis.** *Science* 1988, **241**:1092-1096.
61. Kizawa H, Kou I, Iida A, Sudo A, Miyamoto Y, Fukuda A, Mabuchi A, Kotani A, Kawakami A, Yamamoto S, Uchida A, Nakamura K, Notoya K, Nakamura Y, Ikegawa S: **An aspartic acid repeat polymorphism in asporin inhibits chondrogenesis and increases susceptibility to osteoarthritis.** *Nat Genet* 2005, **37**:138-144.
62. Loughlin J: **Polymorphism in signal transduction is a major route through which osteoarthritis susceptibility is acting.** *Curr Opin Rheumatol* 2005, **17**:629-633.
63. Best CJ, Gillespie JW, Yi Y, Chandramouli GV, Perlmutter MA, Gathright Y, Erickson HS, Georgevich L, Tangrea MA, Duray PH, Gonzalez S, Velasco A, Linehan WM, Matusik RJ, Price DK, Figg WD, Emmert-Buck MR, Chuaqui RF: **Molecular alterations in primary prostate cancer after androgen ablation therapy.** *Clin Cancer Res* 2005, **11**:6823-6834.
64. Tang L, Dai DL, Su M, Martinka M, Li G, Zhou Y: **Aberrant expression of collagen triple helix repeat containing I in human solid cancers.** *Clin Cancer Res* 2006, **12**:3716-3722.
65. Pygay P, Heroult M, Wang Q, Lehnert W, Belden J, Liaw L, Friesel RE, Lindner V: **Collagen triple helix repeat containing I, a novel secreted protein in injured and diseased arteries, inhibits collagen expression and promotes cell migration.** *Circ Res* 2005, **96**:261-268.
66. Zhao X, Ramsey KE, Stephan DA, Russell P: **Gene and protein expression changes in human trabecular meshwork cells treated with transforming growth factor-beta.** *Invest Ophthalmol Vis Sci* 2004, **45**:4023-4034.
67. Durmus T, Leclair RJ, Park KS, Terzic A, Yoon JK, Lindner V: **Expression analysis of the novel gene collagen triple helix repeat containing-I (Cthrc1).** *Gene Expr Patterns* 2006, **6**:935-940.
68. Klopocki E, Kristiansen G, Wild PJ, Klamann I, Castanos-Velez E, Singer G, Stohr R, Simon R, Sauter G, Leibiger L, Essers L, Weber B, Hermann K, Rosenthal A, Hartmann A, Dahl E: **Loss of SFRP1 is associated with breast cancer progression and poor prognosis in early stage tumors.** *Int J Oncol* 2004, **25**:641-649.
69. Eger A, Stockinger A, Park J, Langkopf E, Mikula M, Gotzmann J, Mikulits W, Beug H, Foisner R: **beta-Catenin and TGFbeta signalling cooperate to maintain a mesenchymal phenotype after FosER-induced epithelial to mesenchymal transition.** *Oncogene* 2004, **23**:2672-2680.
70. Huber MA, Azoitei N, Baumann B, Grunert S, Sommer A, Pehamberger H, Kraut N, Beug H, Wirth T: **NF-kappaB is essential for epithelial-mesenchymal transition and metastasis in a model of breast cancer progression.** *J Clin Invest* 2004, **114**:569-581.
71. Willert K, Brown JD, Danenberg E, Duncan AW, Weissman IL, Reya T, Yates JR 3rd, Nusse R: **Wnt proteins are lipidmodified and can act as stem cell growth factors.** *Nature* 2003, **423**:448-452.
72. Papkoff J, Rubinfeld B, Schryver B, Polakis P: **Wnt-1 regulates free pools of catenins and stabilizes APC-catenin complexes.** *Mol Cell Biol* 1996, **16**:2128-2134.
73. Veeman MT, Axelrod JD, Moon RT: **A second canon: functions and mechanisms of beta-catenin-independent wnt signaling.** *Dev Cell* 2003, **5**:367-377.
74. Gradl D, Kuhl M, Wedlich D: **The Wnt/Wg signal transducer beta-catenin controls fibronectin expression.** *Mol Cell Biol* 1999, **19**:5576-5587.
75. Marchenko GN, Marchenko ND, Leng J, Strongin AY: **Promoter characterization of the novel human matrix metalloproteinase-26 gene: regulation by the T-cell factor-4 implies specific expression of the gene in cancer cells of epithelial origin.** *Biochem J* 2002, **363**:253-262.
76. Brabletz T, Jung A, Dag S, Hlubek F, Kirchner T: **Beta-catenin regulates the expression of the matrix metalloproteinase-7 in human colorectal cancer.** *Am J Pathol* 1999, **155**:1033-1038.
77. Crawford HC, Fingleton BM, Rudolph-Owen LA, Goss KJ, Rubinfeld B, Polakis P, Matrisian LM: **The metalloproteinase matrilysin is a target of beta-catenin transactivation in intestinal tumors.** *Oncogene* 1999, **18**:2883-2891.
78. Rivat C, Le Floch N, Sabbah M, Teyrol I, Redeuilh G, Bruyneel E, Mareel M, Matrisian LM, Crawford HC, Gespach C, Attoub S: **Synergistic cooperation between the AP-1 and LEF-1 transcription factors in activation of the matrilysin promoter by the src oncogene: implications in cellular invasion.** *FASEB J* 2003, **17**:1721-1723.
79. Schlessinger J, Plotnikov AN, Ibrahim OA, Eliseenkova AV, Yeh BK, Yayon A, Linhardt RJ, Mohammadi M: **Crystal structure of a ternary FGF-FGFR-heparin complex reveals a dual role for heparin in FGFR binding and dimerization.** *Mol Cell* 2000, **6**:743-750.
80. Ai X, Do AT, Lozynska O, Kusche-Gullberg M, Lindahl U, Emerson CP Jr: **QSulf1 remodels the 6-O sulfation states of cell surface heparan sulfate proteoglycans to promote Wnt signaling.** *J Cell Biol* 2003, **162**:341-351.
81. Chong JM, Uren A, Rubin JS, Speicher DW: **Disulfide bond assignments of secreted Frizzled-related protein-1 provide insights about Frizzled homology and netrin modules.** *J Biol Chem* 2002, **277**:5134-5144.
82. Weeraratna AT, Jiang YA, Hostetter G, Rosenblatt K, Duray P, Bittner M, Trent JM: **Wnt5a signaling directly affects cell motility and invasion of metastatic melanoma.** *Cancer Cell* 2002, **1**:279-288.
83. Jonsson M, Dejmek J, Bendahl PO, Andersson T: **Loss of Wnt-5a protein is associated with early relapse in invasive ductal breast carcinomas.** *Cancer Res* 2002, **62**:409-416.
84. Chen Y, Medvedev A, Ruzanov P, Marvin KW, Jetten AM: **cDNA cloning, genomic structure, and chromosome mapping of the human epithelial membrane protein CL-20 gene (EMPI), a member of the PMP22 family.** *Genomics* 1997, **41**:40-48.
85. Dejmek J, Dib K, Jonsson M, Andersson T: **Wnt-5a and G-protein signaling are required for collagen-induced DDR1 receptor activation and normal mammary cell adhesion.** *Int J Cancer* 2003, **103**:344-351.
86. Jonsson M, Andersson T: **Repression of Wnt-5a impairs DDR1 phosphorylation and modifies adhesion and migration of mammary cells.** *J Cell Sci* 2001, **114**:2043-2053.

Pre-publication history

The pre-publication history for this paper can be accessed here:

<http://www.biomedcentral.com/1471-2407/7/55/prepub>