

Expression of a MHC Class II Transgene Determines Both Superantigenicity and Susceptibility to Mammary Tumor Virus Infection

By Carlo Pucillo, Rosemarie Cepeda, and Richard J. Hodes

From the Experimental Immunology Branch, National Cancer Institute, National Institutes of Health, Bethesda, Maryland 20892

Summary

Milk-borne mouse mammary tumor virus (MMTV) is a type B retrovirus that induces mammary carcinoma. Infectious MMTV, as well as genomically integrated mouse mammary proviruses, encode superantigens that are recognized by T cells that express appropriate T cell receptor $V\beta$ products. To determine the relationship between the superantigenic property of milk-borne MMTV and its in vivo infectivity, mice which were either positive or negative for expression of a transgene-encoded $E\alpha E\beta$ class II major histocompatibility complex (MHC) product were exposed to milk borne C3H MMTV. Superantigen-mediated deletion of $V\beta 14$ -expressing T cells occurred only in $E\alpha$ transgene-positive mice, indicating that the deletion was $E\alpha E\beta$ dependent. When mice were analyzed for viral infection by assaying viral p28 in the milk of recipient females, significant p28 levels were found only in $E\alpha E\beta$ transgene-positive mice. Similarly, the presence of C3H MMTV LTR mRNA in mammary glands, as detected by PCR, paralleled p28 levels. These findings indicate that $E\alpha$ expression or the $E\alpha$ -dependent T cell response to viral superantigen is causally related to susceptibility to MMTV infection, and that lack of a permissive class II product can protect mice from virus infection.

Mouse mammary tumor viruses (MMTV) are milk-borne retroviruses that are infectious and tumorigenic in mice (1). The natural history of these viruses, including the mechanism by which milk-borne transmission leads to host infection and the role of the immune system in this process, is poorly understood. Most mice also express a number of genomically integrated mammary tumor proviruses, the majority of which are defective and do not encode infectious virus (1). Recently, it was demonstrated that the open reading frame (ORF) of the MMTV 3' LTR encodes a superantigenic product which is recognized by T cells expressing an appropriate Tcr $V\beta$ product (2-7). This recognition can lead to $V\beta$ -specific T cell activation, anergy, or clonal deletion (2-8). It has been suggested that the ability of MMTV-encoded superantigens to induce T cell activation and consequent B cell activation in vivo may facilitate viral infection and replication in host lymphoid cells, and may thereby play a critical role in susceptibility to viral infection (9). Since T cell recognition of MMTV superantigens is dependent upon presentation by a permissive MHC class II product, a mechanism of viral infection which requires superantigenic stimulation might therefore be similarly dependent upon permissive class II expression. Indeed, early studies identified a significant MHC influence on susceptibility to MMTV-induced tumorigenesis, although precise identification of the relevant MHC-linked gene(s) was not possible (10, 11). To directly test the

relationship of MMTV susceptibility and class II-dependent superantigen recognition, susceptibility to MMTV infection was compared in mice that differ in their ability to respond to the MMTV superantigen as a result of expression or nonexpression of a transgenic class II $E\alpha$ product.

Materials and Methods

Animals. C57BL/6 (B6) mice were obtained from Frederick Cancer Research Center (Frederick, MD). C3H/OUJ mice were obtained from The Jackson Laboratory (Bar Harbor, ME). The MHC class II $E\alpha$ transgenic strains 107 (B6.107) and 36-5 (B6.36) have previously been characterized (12, 13) and were generously provided by David Lo (Scripps Institute, La Jolla, CA) and B. J. Fowlkes (NIH, Bethesda, MD). Inbred transgenic lines had been established by 10 generations of back-crossing to B6 mice and selection for transgene expression; transgenic mice used in experiments had been bred no more than three generations after establishing homozygosity.

Foster Nursing. MMTV-positive C3H/OUJ females were used as foster mothers. Newborn mice of the indicated strains were transferred to C3H/OUJ females for nursing within 3 d of birth.

Flow Cytometric Analysis. When mice were at least 4 mo of age (at which time maximal deletion of $V\beta 14^+$ cells had occurred), peripheral blood was analyzed for Tcr $V\beta$ expression. Lymphocytes were isolated from heparinized peripheral blood using Lymphocyte Separating Medium (Organon Teknica, Rockville, MD). Cells were stained using anti- $V\beta$ culture supernatant (14, 15) or control supernatant, followed by FITC-conjugated goat anti-rat

Ig, biotin-conjugated anti-CD4, and Texas red avidin. Two-parameter flow cytometric analysis was carried out as previously described (16).

Measurement of MMTV p28. Milk for assay of MMTV p28 was obtained from females of the appropriate strains. Pregnancies were induced, and following the birth of offspring, females were treated with 0.25 IU oxytocin i.p. and 5 min later milk was harvested by suction. ELISA was carried out by general procedures previously described (17). ELISA plates were coated with rabbit anti-MMTV p28 serum generously provided by Dr. Pierre Hainaut (University of York, York, England) (18). Plates were then washed and incubated with varying concentrations of milk or with a standard preparation of purified C3H milk virus protein. Plates were again washed and then incubated with biotin-conjugated rabbit anti-p28 antibody, washed, and incubated with avidin-conjugated enzyme followed by substrate. Plates were read spectrophotometrically, and virus protein was calculated with reference to a virus protein standard using software generously provided by Dr. Pierre Henkart (NIH, Bethesda, MD).

Polymerase Chain Reaction (PCR) Analysis of MMTV Expression. For analysis of MMTV LTR-ORF mRNA, total RNA was prepared from surgically removed mammary glands of lactating females as previously described (16, 19). cDNA synthesis was carried out as previously described (20). The cDNA samples were then amplified as previously described (20) in 100 μ l PCR reaction mixture consisting of: 5 μ l each of 20 μ M sense and antisense oligonucleotides in TE, 5 μ l of cDNA reaction, 1 \times PCR reaction buffer, 0.2 mM final concentration of each dNTP, and 2.5 U AmpliTaq (Perkin-Elmer Cetus Instrs., Norwalk, CT) per reaction. All reactions were overlaid with mineral oil and subjected to 30 cycles of amplification using a programmed thermal cycler (Perkin-Elmer Cetus Instrs.). Each cycle consisted of denaturation at 94°C for 1 min, annealing at 55°C for 1 min, and polymerization at 72°C for 2 min. 20- μ l samples were analyzed on 1.2% agarose ethidium bromide gels in 0.5 \times TBE run at 200 V, stained with ethidium bromide, and visualized under UV light.

Oligonucleotide primers capable of amplifying all known endogenous and milk-borne MMTV LTR ORF molecules were chosen on the basis of highly conserved sequences among characterized ORF molecules (21): 5' (sense) oligonucleotide: ATGCCG-GCGCCTGCAGCAGA (Mtv-C5); 3' (antisense) oligonucleotide: CCAAGTCAGGAAACCACTTG (Mtv-C3). To specifically detect milk-borne C3H MMTV, a 3' (antisense) oligonucleotide specific for the unique MMTV-C3H ORF carboxy terminus was used in combination with Mtv-C5: TCAGAGCTCAGATCAGAACCT (Mtv-C3H-3).

PCR products were transferred to a nylon membrane (Hybond-N⁺; Amersham, Bucks, UK) by alkali blotting. The membranes were prehybridized for 1 h in QuikHyb solution (Stratagene, La Jolla, CA). They were then hybridized with either ³²P-labeled cDNA probe or labeled oligonucleotide Mtv-C5. The cDNA probe was prepared from C3H milk virus LTR-ORF mRNA generously provided by Dr. R. Callahan (NIH) (22) and radiolabeled using the random hexamer priming method. The oligonucleotide probe Mtv-C3 was ³²P-5' end-labeled by Lofstrand Labs. Ltd. (Gaithersburg, MD). Membrane-bound cDNA was hybridized overnight at 42°C with the cDNA probe or at 45°C with Mtv-C3. cDNA-hybridized filters were washed twice with 2 \times SSC, 0.05% SDS at 45°C for 45 min and then once with 1 \times SSC, 0.05% SDS at 60°C for 30 min. Mtv-C3 oligonucleotide-hybridized filters were washed at room temperature for 20 min, then once with 1 \times SSC, 0.05% SDS at 45°C for 20 min. Autoradiography was performed using Phosphor Screens (Molecular Dynamics Inc., Sunnyvale, CA) and

exposed screens were scanned on a Phosphorimager (Molecular Dynamics Inc.).

Results

Influence of E α Transgene Expression on V β 14 Deletion. It was previously reported that the ability of the C3H milk virus superantigen to induce V β 14-specific deletion is MHC dependent, and it was suggested that this dependence is related to expression of an MHC class II E α E β product (2, 6). However, these previous studies did not directly test the role of class II E α E β expression in superantigen-mediated V β 14 deletion. Consistent with previous findings, it was found in the present study that exposure to milk-borne C3H MMTV resulted in significant deletion of V β 14⁺ T cells in B10.A (E α E β positive) but not B10 (E α E β negative) mice (data not shown). To directly examine the relationship of E α E β expression to V β 14 deletion, B6 mice, which fail to express an E α E β product due to the absence of expressed E α , were compared with the E α -transgenic B6 mouse line B6.107, that expresses E α E β in a normal cellular distribution (12). Newborn B6 or B6.107 mice were exposed to milk-borne MMTV by foster nursing on C3H/HeOJ females. The resulting adult mice were then analyzed for TCR V β expression. Analysis of CD4⁺ peripheral T cells demonstrated a marked decrease in the proportion of V β 14⁺ cells in B6.107 mice which had been exposed to milk-borne MMTV (1.8%), when compared to mice which had not been exposed to virus (9.7%) (Fig. 1). No differences were seen in expression of any other serologically detectable V β product (V β 's 2 through 13) (data not shown). In contrast, exposure of B6 mice (E α E β -negative) to milk-borne virus did not result in deletion of V β 14⁺ cells (7.1% in virus-exposed mice versus 7.0% in control mice). (B6 \times B6.107)_{F1} mice showed V β 14 deletion indistinguishable from that of homozygous B6.107 animals, indicating that the effect of transgene insertion and expression is dominant. Another E α transgenic line, B6.36, has been found to express E α E β molecules only on thymic epithelial cells (12, 13). Flow cytometric analysis confirmed the expression of E α E β product on peripheral lymphoid cells of B6.107 but not B6 or B6.36 mice (data not shown). To determine whether the restricted expression of E α E β in B6.36 mice is sufficient to support V β 14-specific deletion, B6.36 mice were also exposed to MMTV. These mice failed to undergo V β 14 deletion after MMTV exposure, indicating that the expression of E α E β limited to thymic epithelium is insufficient to support deletion. Together, these results demonstrate that appropriate expression of an E α E β class II product is necessary for the superantigen-mediated deletion of V β 14⁺ T cells in vivo.

Influence of E α Transgene Expression on Susceptibility to MMTV Infection. If susceptibility to MMTV infection were related to the host ability to respond to MMTV superantigen, it might be expected that E α expression would have an effect on viral infection that paralleled its effect on V β 14 deletion. To test this possibility, B6 and E α transgenic mice that either had or had not been exposed to milk-borne MMTV were




Figure 1. V β 14 deletion in MMTV-exposed mice. B6 mice (E α E β -negative), E α transgenic B6.107 mice (expressing E α E β in normal cellular distribution) and E α transgenic B6.36 mice (expressing E α E β only on thymic epithelium) were either nursed by MMTV-negative mothers or were foster-nursed by MMTV milk-virus positive C3H/HeOJ females. When mice were at least 4 mo of age (at which time maximal deletion of V β 14⁺ cells had occurred), they were bled, and peripheral blood was analyzed for the expression of V β 6⁺ and V β 14⁺ CD4⁺ cells. Exposure to MMTV did not induce changes in V β 6 expression (data not shown). Results for V β 14 expression are shown as means of 4 to 12 individual mice per group after subtraction of negative control staining. ■, Control; ▨, MMTV exposed.

examined for virus infection. Milk was harvested from female mice and analyzed using an ELISA specific for MMTV p28 gag product. No p28 was detected in control mice of any strain that had not been exposed to milk-borne virus. Virus-exposed E α -negative B6 mice had statistically in-




Figure 2. Assay of MMTV p28 in milk of virus-exposed or control mice. Female mice of the groups described in Fig. 1 were mated with virus-negative B6 males. Following delivery of offspring, milk was harvested by suction and analyzed for p28. The group designated as B6.107 is a pool of mice homozygous and heterozygous for the E α transgene. Results are expressed as ng of viral protein/ml of milk. The number of mice in virus-exposed groups was: B6 *n* = 11, B6.107 *n* = 11, B6.36 *n* = 4.

significant levels of p28 (Fig. 2). In contrast, the mean p28 content of virus-exposed B6.107 transgenic mice, which were E α E β positive and deleted V β 14⁺ CD4⁺ T cells, was 2.03 μ g viral protein/ml. Foster-nursed B6.36 mice, in which E α expression was confined to thymic epithelium and in which V β 14 deletion did not occur, had no detectable p28 in milk. Thus, p28 levels in these strains correlated with the ability to undergo superantigen-mediated deletion of V β 14⁺ cells.

MMTV infection was also analyzed by PCR. cDNA was




Figure 3. PCR analysis of C3H MMTV LTR-ORF mRNA in mammary tissue of virus-exposed or control mice. RNA was prepared from surgically removed mammary glands of female B6.107 or B6 mice which had been neonatally exposed to milk-borne C3H MMTV (+) or had not been exposed to milk-borne virus (-). RNA was reverse transcribed to prepare cDNA, and cDNA was subjected to PCR amplification using oligonucleotide primers that were either specific for the LTR of C3H milk-borne MMTV (A) or for common LTR sequences expressed by multiple endogenous as well as milk-borne MMTV's (B). Amplified product was analyzed by agarose gel electrophoresis and hybridization with ³²P-labeled MMTV oligonucleotide Mtv-C3.

prepared from the mammary glands of experimental mice and was amplified using oligonucleotide primers that were either specific for the LTR of milk-borne C3H MMTV, or were derived from common sequences expressed by multiple endogenous and milk-borne MMTV LTR's and were therefore capable of amplifying cDNA from all of these genes. Amplified products were analyzed by electrophoresis and hybridization with an MMTV oligonucleotide probe. Virus-exposed E α -transgenic B6.107 mice were positive for PCR-amplified C3H milk virus product, while transgene-negative B6 mice were negative (Fig. 3 A). Transgene-positive and negative mice, with or without exposure to milk virus, expressed products corresponding to common MMTV sequences shared by endogenous proviruses (Fig. 3 B). The observed products were of a size consistent with that expected for amplified MMTV product (21). The same results were seen after hybridization with a cDNA probe (data not shown). All mice also expressed similar quantities of PCR-amplified β 2-microglobulin product (data not shown). PCR analysis thus indicated that expression of the E α transgene was required for susceptibility to infection by milk-borne MMTV, in agreement with the results of viral p28 analysis.

Discussion

These findings demonstrate that host susceptibility to MMTV infection *in vivo* is strongly influenced by expression of a class II E α product. The mechanism underlying this effect is not certain. However, the observation that both the superantigenic effects of MMTV and the susceptibility of mice to viral infection are similarly dependent upon expression of a transgenic E α product suggests that the T cell response to viral superantigen is causally involved in viral infectivity. The expression of proviral *mtv* genes has been shown to be greatly enhanced in T and B lymphocytes following activation of these cells (9). Thus, it is possible that introduction of infectious MMTV into the host results in infection of cells including lymphoid cells, and that class II E α E β -positive cells such as B cells express and present the MMTV

superantigen to T cells expressing appropriate TCR V β products. The resulting activation of T cells and the consequent activation of B cells might result in enhanced viral expression and replication.

From such a model, it would be predicted that any maneuver which interferes with the superantigenic stimulation of T cells by MMTV-expressing cells would result in decreased susceptibility to virus infection and replication. The "knock-out" of E α E β expression in the wild type B6 strain is one such maneuver. In this situation, the failure of class II⁺ cells to express a permissive E α E β product results in failure of superantigenic stimulation of V β 14⁺ T cells and a consequent resistance to viral infection. A different approach was taken in the work of Golovkina et al. (23), who reported that the expression of a transgene encoding the C3H MMTV superantigen resulted in V β 14 deletion and in decreased susceptibility to infection by milk-borne MMTV. In this case, the interaction of V β 14⁺ T cells with E α E β ⁺ cells expressing an endogenous transgenic superantigen results in the deletion of V β 14⁺ T cells and a preempting of their ability to interact with cells bearing the infectious MMTV.

There thus appear to be two distinct mechanisms by which animals can be protected from susceptibility to infection with tumorigenic MMTV retrovirus. One mechanism involves the expression of *mtv* proviruses that lead to deletion of T cells that facilitate MMTV infection. The fact that multiple V β -specific endogenous *mtv* superantigens are conserved in the mouse suggests that this may be a biologically relevant mechanism of host protection. A second and distinct mechanism for host protection from MMTV infection is the failure to express class II E α E β . The fact that many wild as well as inbred mice have conserved genomic defects in class II E α E β expression is consistent with the biologic relevance of this mechanism as well. Both the expression of endogenous superantigens and the widespread deletion of a permissive class II product are phenomena which have to date been observed only in the mouse, suggesting that they may reflect a unique relationship between host and MMTV retroviruses in this species.

The authors wish to thank Ronald Gress and Robert Callahan for their comments and suggestions and Larry Granger for his assistance with flow cytometry.

Address correspondence to Carlo Pucillo, Bldg. 10, Rm. 4B17, National Institutes of Health, Bethesda, MD 20892.

Received for publication 20 May 1993 and in revised form 1 July 1993.

References

1. Slagle, B.L., and J.S. Butel. 1987. Exogenous and endogenous mouse mammary tumor viruses: replication and cell transformation. *In Cellular and Molecular Biology of Mammary Cancer*. D. Medina, W. Kidwell, G. Heppner, and E. Anderson, editors. Plenum Publishing Corp., New York. 275-306.
2. Marrack, P., E. Kushnir, and J.W. Kappler. 1991. A maternally inherited superantigen encoded by a mammary tumor virus. *Nature (Lond.)* 349:524.
3. Frankel, W.N., C. Rudy, J.M. Coffin, and B.T. Huber. 1991. Linkage of Mls genes to endogenous mammary tumor viruses

- of inbred mice. *Nature (Lond.)*. 349:526.
4. Woodland, D.L., M.P. Happ, K.J. Gollob, and E. Palmer. 1991. An endogenous retrovirus mediating deletion of $\alpha\beta$ T cells? *Nature (Lond.)*. 349:529.
 5. Dyson, P.J., A.M. Knight, S. Fairchild, E. Simpson, and K. Tomonari. 1991. Genes encoding ligands for deletion of V β 11 T cells cosegregate with mammary tumor virus genomes. *Nature (Lond.)*. 349:531.
 6. Acha-Orbea, H., A.N. Shakhov, L. Scarpellino, E. Kolb, V. Muller, A. Vessaz-Shaw, R. Fuchs, K. Blohlinger, P. Rollini, J. Billotte, et al. 1991. Clonal deletion of V β 14-bearing T cells in mice transgenic for mammary tumor virus. *Nature (Lond.)*. 350:207.
 7. Choi, Y., J.W. Kappler, and P. Marrack. 1991. A superantigen encoded in the open reading frame of the 3' long terminal repeat of mouse mammary tumor virus. *Nature (Lond.)*. 350:203.
 8. Speiser, D.E., Y. Chvatchko, R.M. Zinkernagel, and R.H. MacDonald. 1990. Distinct fates of self-specific T cells developing in irradiation bone marrow chimeras. Clonal deletion, clonal anergy, or in vitro responsiveness to self-Mls-1^a controlled by hematopoietic cells in the thymus. *J. Exp. Med.* 172:1305.
 9. Corley, R.B., F.E. Lund, T.D. Randall, L.B. King, S. Doerre, and D.L. Woodland. 1992. Mouse mammary tumor proviral gene expression in cells of the B lineage. *Semin. Immunol.* 4:287.
 10. Muhlbock, O., and A. Dux. 1981. Histocompatibility genes and mammary cancer. In *Mammary Tumors in the Mouse*, J. Hilgers and M. Sluysers, editors. Elsevier/North-Holland Biomedical Press, Amsterdam. 545-572.
 11. Demant, P., L.C.M.J. Oomen, and M. Oudshoorn-Snoek. 1989. Genetics of tumor susceptibility in the mouse: MHC and non-MHC genes. *Adv. Cancer Res.* 53:117.
 12. Marrack, P., D. Lo, R. Brinster, R. Palmiter, L. Burkly, R.H. Flavell, and J. Kappler. 1988. The effect of thymus environment on T cell development and tolerance. *Cell*. 53:627.
 13. Widera, G., L.C. Burkly, C.A. Pinkert, E.C. Bottger, C. Cowing, R.D. Palmiter, R.L. Brinster, and R.A. Flavell. 1987. Transgenic mice selectively lacking MHC class II (I-E) expression on B cells: an in vivo approach to investigate Ia gene function. *Cell*. 51:175.
 14. Liao, N., J. Maltzman, and D.H. Raulet. 1989. Positive selection determines T cell receptor V β 14 gene usage by CD8⁺ T cells. *J. Exp. Med.* 170:135.
 15. Karpati, R.M., S.M. Banks, B. Malissen, S.A. Rosenberg, M.A. Sheard, J.S. Weber, and R.J. Hodes. 1991. Phenotypic characterization of murine tumor infiltrating T lymphocytes. *J. Immunol.* 146:2043.
 16. Vacchio, M.S., J.J. Ryan, and R.J. Hodes. 1990. Characterization of the ligand(s) responsible for negative selection of V β 11- and V β 12-expressing T cells. Effects of a new Mls determinant. *J. Exp. Med.* 172:807.
 17. Vos, Q., and R.J. Hodes. 1992. Immunoglobulin (Ig) μ , κ transgenic mice express transgenic idiotype on endogenously rearranged IgM and IgA molecules by secretion of chimeric molecules. *J. Exp. Med.* 176:951.
 18. Hainaut, P., D.D. Vaira, C. Francois, C.M. Calberg-Bacq, and P.M. Osterrieth. 1985. Natural infection of Swiss mice with mouse mammary tumor virus (MMTV): viral expression in milk and transmission of infection. *Arch. of Virol.* 83:195.
 19. Chomczynski, P., and N. Sacchi. 1987. Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal. Biochem.* 162:156.
 20. Hathcock, K.S., H. Hirano, S. Murakami, and R.J. Hodes. 1992. Analysis of CD45 expression by B cells stimulated with IL5, anti-Ig, or LPS: expression of different CD45 isoforms by subpopulations of activated B cells. *J. Immunol.* 149:2286.
 21. Brandt-Carlson, C., J.S. Butel, and D. Wheeler. 1993. Phylogenetic and structural analyses of MMTV LTR ORF sequences of exogenous and endogenous origins. *Virology*. 193:171.
 22. Gallahan, D., and R. Callahan. 1987. Mammary tumorigenesis in feral mice: identification of a new *int* locus in mouse mammary tumor virus (Czech II)-induced mammary tumors. *J. Virol.* 61:66.
 23. Golovkina, T.V., A. Chervonsky, J.P. Dudley, and S.R. Ross. 1992. Transgenic MMTV superantigen expression prevents viral infection. *Cell*. 69:637.