

Cite this: *RSC Adv.*, 2018, 8, 26646

Hyperelatosides A–E, biphenyl ether glycosides from *Hypericum elatoides*, with neurotrophic activity†

Xi-Tao Yan,^a Zhen An,^a Dan Tang,^a Guang-Rui Peng,^a Chen-Yu Cao,^a Yuan-Zhen Xu,^a Chun-Huan Li,^a Pei-Liang Liu,^b Zai-Min Jiang^c and Jin-Ming Gao^{*,a}

Five new biphenyl ether glycosides, hyperelatosides A–E (1–5), one new benzoate glycoside, hyperelatoside F (6), along with nine known phenolic compounds (7–15), were isolated from the aerial parts of *Hypericum elatoides*. Their structures were elucidated by 1D and 2D NMR spectroscopy and HRESIMS, as well as chemical derivatization. This is the first report of the identification of biphenyl ether glycosides as plant metabolites and their possible biosynthetic pathway is proposed. Except for 3, the new phenolic metabolites exhibited significant neurotrophic activities to enhance nerve growth factor-induced neurite outgrowth in PC12 cells. In addition, the anti-neuroinflammatory and antioxidant activities of compounds 1–15 were preliminarily evaluated *in vitro*.

Received 21st June 2018

Accepted 5th July 2018

DOI: 10.1039/c8ra05322g

rsc.li/rsc-advances

Introduction

Plants belonging to the genus *Hypericum* (Hypericaceae) are well known for their therapeutic uses in neurological disorders.¹ In particular, the extract of *H. perforatum* (St. John's wort), the most known of the *Hypericum* species, has long been used for the treatment of mild to moderate depression.² Due to the continuous discovery of structurally novel natural products and their significant biological activities, plants of the genus *Hypericum* have attracted much attention from researchers in both chemistry and pharmacology. Previous chemical investigations on this genus have resulted in the isolation of phloroglucinol derivatives, xanthenes, benzophenones, naphthodianthrones, flavonoids, and coumarins as major constituents.^{3–6} These secondary metabolites possess extensive biological activities, such as antidepressant, neuroprotective, anti-neurodegenerative, antioxidant, memory-enhancing, anti-inflammatory, anticancer, anti-HIV, and antimicrobial activities.^{7–13}

Hypericum elatoides R. Keller is an herbaceous plant distributed in northwest China, especially in the western region of Shaanxi province.¹⁴ To date, there have been no phytochemical and biological studies on this plant. In the course of

our ongoing search for natural neurotrophic compounds with novel structures for potential neurodegenerative disease therapies,¹⁵ 15 phenolic compounds including five new uncommon biphenyl ether glycosides named hyperelatosides A–E (1–5), one new benzoate glycoside named hyperelatoside F (6), and nine known, structurally related phenolic derivatives (7–15) were isolated from the MeOH extract of the aerial parts of *H. elatoides* (Fig. 1). Herein, we described the isolation and structural elucidation of these new phenolic metabolites, as well as the biological evaluation of their promoting effects on nerve growth factor (NGF)-induced neurite outgrowth in PC12 cells, inhibitory effects on lipopolysaccharide (LPS)-induced nitric oxide (NO) production in BV-2 cells, and 1,1-diphenyl-2-picrylhydrazyl (DPPH) free radical-scavenging capacities.

Results and discussion

The MeOH extract of *H. elatoides* was subjected to liquid–liquid fractionation to afford a hexane-soluble fraction, an EtOAc-soluble fraction, and a BuOH-soluble fraction. The active EtOAc fraction was separated by semipreparative reversed phase (RP) HPLC and column chromatography (CC) using silica gel, RP-C₁₈, and Sephadex LH-20 to yield six new compounds (1–6) and nine known phenolic metabolites (7–15).

Compound 1 was isolated as a brown amorphous solid. Its molecular formula was determined to be C₂₂H₂₆O₁₃ by HRESIMS (*m/z* 521.1247 [M + Na]⁺, calcd for C₂₂H₂₆NaO₁₃, 521.1271). The IR spectrum showed absorption bands at 3418 and 1701 cm⁻¹ for hydroxyl and ester carbonyl groups, respectively. The low-field region of the ¹H NMR spectrum (measured in CD₃OD) exhibited two *para*-positional aromatic protons [δ_{H} 7.44 (1H, s, H-3) and 6.38 (1H, s, H-6)] and two *meta*-coupled

^aShaanxi Key Laboratory of Natural Products & Chemical Biology, College of Chemistry & Pharmacy, Northwest A&F University, Yangling 712100, China. E-mail: jinminggao@mwsuaf.edu.cn

^bKey Laboratory of Resource Biology and Biotechnology in Western China, College of Life Sciences, Northwest University, Xi'an 710069, China

^cCollege of Life Sciences, Northwest A&F University, Yangling 712100, China

† Electronic supplementary information (ESI) available: 1D NMR, 2D NMR, and HRESIMS of 1–6; chemical structures of 7–15. See DOI: 10.1039/c8ra05322g






Fig. 1 Chemical structures of compounds 1–6.

aromatic protons [δ_{H} 6.67 (1H, d, $J = 2.9$ Hz, H-4') and 6.10 (1H, d, $J = 2.9$ Hz, H-6')] (Table 1), indicating the presence of a 1,2,4,5-tetrasubstituted (A-ring) and a 1,2,3,5-tetrasubstituted (B-ring) benzene rings, respectively. Further, the aliphatic region of the ^1H NMR spectrum showed signals of three methoxy groups [δ_{H} 3.87 (3H, s), 3.78 (3H, s), and 3.65 (3H, s)] and a set of sugar protons (δ_{H} 3.41–4.84). The ^{13}C NMR spectrum (Table 2) supported by DEPT experiments displayed signals of three methoxy carbons (δ_{C} 57.1, 56.3, and 52.5), six characteristic glucopyranosyl carbons (δ_{C} 104.2, 78.6, 77.7, 75.0, 71.5, and 62.6), an ester carbonyl carbon (δ_{C} 167.9), and 12

aromatic carbons (δ_{C} 100.5–154.3), including eight quaternary carbons and four methine carbons, corroborating the presence of two tetrasubstituted aromatic rings detected in the ^1H NMR spectrum. Furthermore, the ^1H NMR spectrum of 1, measured in $\text{DMSO}-d_6$, displayed the downfield signals of two hydroxyl groups at δ_{H} 10.03 and 8.03 (Table 1), which were attached to C-5 position of A-ring and C-2' position of B-ring based on the HMBC correlations from the hydroxyl proton at δ_{H} 10.03 to C-4, C-5, and C-6 and from the hydroxyl proton at δ_{H} 8.03 to C-1', C-2', and C-3'. In the HMBC spectrum, the correlations from the methoxy protons at δ_{H} 3.78 to the carbonyl carbon at δ_{C} 167.9 (C-

Table 1 ^1H NMR spectroscopic data (500 MHz, δ in ppm, J in Hz) for compounds 1–5

| No. | 1 ^a | 1 ^b | 2 ^a | 3 ^a | 4 ^a | 4 ^b | 5 ^a |
|--------|----------------------|----------------------|----------------------|----------------------|----------------------|----------------------|----------------------|
| 3 | 7.44, s | 7.32, s | 7.35, s | 7.34, s | 6.85, d (8.8) | 6.84, d (8.8) | 6.84, d (8.8) |
| 4 | | | | | 6.59, dd (8.8, 2.9) | 6.58, dd (8.8, 3.0) | 6.56, dd (8.8, 2.9) |
| 6 | 6.38, s | 6.26, s | 6.40, s | 6.72, s | 6.46, d (2.9) | 6.45, d (3.0) | 6.39, d (2.9) |
| 4' | 6.67, d (2.9) | 6.64, d (2.9) | 6.68, d (2.8) | 6.72, d (2.9) | 6.24, d (2.9) | 6.19, d (2.9) | 6.67, d (2.9) |
| 6' | 6.10, d (2.9) | 6.14, d (2.9) | 6.33, d (2.8) | 6.54, d (2.9) | 5.91, d (2.9) | 5.74, d (2.9) | 6.17, d (2.9) |
| 1'' | 4.84, d (7.4) | 4.72, d (7.4) | 4.83, d (7.4) | 4.94, d (7.7) | 4.73, d (7.6) | 4.60, d (7.5) | 4.80, d (7.6) |
| 2'' | 3.52, m | 3.31, m | 3.49, m | 3.52, m | 3.46, m | 3.25, m | 3.52, m |
| 3'' | 3.50, m | 3.30, m | 3.48, m | 3.47, m | 3.41, m | 3.27, m | 3.49, m |
| 4'' | 3.41, m | 3.16, m | 3.39, m | 3.38, m | 3.42, m | 3.17, m | 3.41, m |
| 5'' | 3.45, m | 3.37, m | 3.44, m | 3.45, m | 3.25, m | 3.34, m | 3.44, m |
| 6'' | 3.71, dd (12.1, 5.8) | 3.47, dd (11.8, 5.9) | 3.70, dd (12.0, 5.9) | 3.67, dd (12.1, 5.9) | 3.69, dd (12.0, 4.7) | 3.45, dd (11.7, 5.8) | 3.71, dd (12.1, 5.7) |
| | 3.92, dd (12.1, 2.1) | 3.75, m | 3.91, dd (12.0, 2.2) | 3.86, dd (12.1, 2.2) | 3.77, dd (12.0, 2.4) | 3.62, m | 3.91, dd (12.1, 2.0) |
| 4-OMe | 3.87, s | 3.76, s | 3.85, s | 3.87, s | | | |
| 5-OMe | | | | | 3.68, s | 3.62, s | 3.66, s |
| 7-OMe | 3.78, s | 3.73, s | | | | | |
| 5'-OMe | 3.65, s | 3.62, s | 3.70, s | 3.77, s | 3.65, s | 3.59, s | 3.66, s |
| 2-OH | | | | | | 8.74, s | |
| 5-OH | | 10.03, s | | | | | |
| 2'-OH | | 8.03, s | | | | | |
| 3'-OH | | | | | | 9.08, br s | |
| 2''-OH | | 5.44, d (2.8) | | | | 6.05, br s | |
| 3''-OH | | 5.08, br s | | | | 5.12, d (4.4) | |
| 4''-OH | | 5.05, d (5.3) | | | | 4.99, d (4.6) | |
| 6''-OH | | 4.64, t (5.5) | | | | 4.31, t (5.7) | |

^a Data were recorded in CD_3OD . ^b Data were recorded in $\text{DMSO}-d_6$.

Table 2 ^{13}C NMR spectroscopic data (125 MHz, δ in ppm) for compounds 1–5

| No. | 1 ^a | 1 ^b | 2 ^a | 3 ^a | 4 ^a | 4 ^b | 5 ^a |
|--------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|
| 1 | 153.9 | 152.5 | 152.9 | 158.0 | 146.0 | 143.9 | 146.4 |
| 2 | 112.8 | 110.0 | 117.6 | 111.2 | 143.3 | 142.2 | 143.1 |
| 3 | 115.3 | 114.4 | 115.0 | 115.6 | 118.4 | 117.2 | 118.2 |
| 4 | 145.0 | 142.8 | 144.9 | 147.1 | 110.6 | 109.5 | 110.1 |
| 5 | 153.7 | 151.9 | 151.8 | 155.8 | 155.0 | 152.4 | 155.0 |
| 6 | 108.0 | 105.2 | 107.2 | 108.4 | 107.4 | 106.7 | 106.6 |
| 7 | 167.9 | 165.0 | 172.8 | 166.4 | | | |
| 1' | 147.0 | 143.9 | 146.8 | 154.0 | 152.1 | 150.9 | 146.8 |
| 2' | 133.7 | 132.5 | 134.7 | 130.0 | 131.4 | 129.7 | 133.7 |
| 3' | 148.4 | 147.0 | 148.6 | 150.8 | 153.1 | 151.4 | 148.5 |
| 4' | 100.5 | 99.1 | 101.2 | 101.8 | 98.2 | 96.3 | 100.4 |
| 5' | 154.3 | 151.8 | 154.0 | 159.2 | 158.8 | 156.2 | 154.4 |
| 6' | 101.1 | 100.2 | 102.7 | 101.0 | 97.3 | 95.4 | 101.2 |
| 1'' | 104.2 | 102.2 | 103.8 | 103.1 | 107.4 | 105.8 | 104.3 |
| 2'' | 75.0 | 73.3 | 75.0 | 74.9 | 75.5 | 73.8 | 75.0 |
| 3'' | 77.7 | 75.8 | 77.7 | 77.8 | 77.9 | 76.2 | 77.8 |
| 4'' | 71.5 | 70.0 | 71.6 | 71.5 | 71.0 | 69.6 | 71.6 |
| 5'' | 78.6 | 77.3 | 78.5 | 78.6 | 78.5 | 77.1 | 78.6 |
| 6'' | 62.6 | 60.8 | 62.6 | 62.6 | 62.2 | 60.8 | 62.6 |
| 4-OMe | 57.1 | 56.1 | 57.0 | 57.0 | | | |
| 5-OMe | | | | | 56.4 | 55.2 | 56.4 |
| 7-OMe | 52.5 | 51.5 | | | | | |
| 5'-OMe | 56.3 | 55.3 | 56.4 | 56.5 | 56.1 | 55.4 | 56.3 |

^a Data were recorded in CD_3OD . ^b Data were recorded in $\text{DMSO}-d_6$.

7), from H-3 (δ_{H} 7.44) to C-1 (δ_{C} 153.9), C-5 (δ_{C} 153.7), and C-7 (δ_{C} 167.9), from H-6 (δ_{H} 6.38) to C-2 (δ_{C} 112.8) and C-4 (δ_{C} 145.0), and from the methoxy protons at δ_{H} 3.87 to C-4 (δ_{C} 145.0) (Fig. 2) suggested the presence of a methoxycarbonyl group and a methoxy group located at the C-2 and C-4 positions of A-ring, respectively. In addition, the locations of a methoxy group at C-5' and a sugar moiety at C-3' of B-ring were established according to the HMBC correlations observed from the methoxy protons at δ_{H} 3.65 to C-5' (δ_{C} 154.3) and from the anomeric

proton at δ_{H} 4.84 (1H, d, $J = 7.4$ Hz, H-1'') to C-3' (δ_{C} 148.4), respectively. These two aromatic rings were bonded *via* an oxygen atom based on analysis of the chemical shifts of C-1 (δ_{C} 153.9) and C-1' (δ_{C} 147.0) together with the molecular formula. Acid hydrolysis followed by HPLC analysis after arylthiocarbamoyl-thiazolidine derivatization confirmed the characterization of a β -D-glucopyranosyl unit. Thus, the structure of **1** was elucidated as 2-(3-O- β -D-glucopyranosyl-2-hydroxy-5-methoxyphenoxy)-4-hydroxy-5-methoxy-methylbenzoate and was named hyperelatoside A.

Compound **2** was obtained as a brown amorphous solid. Its molecular formula was deduced to be $\text{C}_{21}\text{H}_{24}\text{O}_{13}$ by HRESIMS ion peak at m/z 483.1150 $[\text{M} - \text{H}]^-$ (calcd for $\text{C}_{21}\text{H}_{23}\text{O}_{13}$, 483.1139). The ^1H and ^{13}C NMR data of **2** (Tables 1 and 2) showed similarity to those of **1**, except for the lack of one methoxy signals as well as the chemical shift of C-7 (δ_{C} 172.8 in **2** and δ_{C} 167.9 in **1**). These observations indicated a carboxyl group in **2** instead of the methoxycarbonyl group in **1** at C-2 position, which was supported by the molecular formula. In addition, the HMBC correlations of **2** (Fig. 2) confirmed the same connectivity as in **1**. Therefore, the structure of **2** was determined as 2-(3-O- β -D-glucopyranosyl-2-hydroxy-5-methoxyphenoxy)-4-hydroxy-5-methoxy benzoic acid and was given the name hyperelatoside B.

Compound **3** was obtained as a brown amorphous solid. Its molecular formula was established as $\text{C}_{21}\text{H}_{22}\text{O}_{12}$ by HRESIMS (m/z 489.0998 $[\text{M} + \text{Na}]^+$, calcd for $\text{C}_{21}\text{H}_{22}\text{NaO}_{12}$, 489.1009), accounting for 11 degrees of unsaturation. The UV, IR, and NMR data of **3** were similar to those of **1** and **2** (Table 1), indicating that the structure of **3** is a biphenyl ether glycoside. The ^1H NMR spectrum of **3** revealed the presence of a 1,2,4,5-tetra-substituted benzene ring, a 1,2,3,5-tetrasubstituted benzene ring, two methoxy groups, as well as a sugar moiety, which were consistent with the constituent units of **2**. Similar HMBC correlations (Fig. 2) established that a sugar moiety, a carbonyl group, and two methoxy groups were located at the same




Fig. 2 Key HMBC (blue arrows) correlations of compounds 1–6.

positions as **2**. Compared with the chemical shifts of C-7 of **1** and **2** (Table 2), the appearance of an ester carbonyl carbon at δ_{C} 166.4 in **3** suggested that an ester group presented between these two aromatic rings *via* the linkage of C-7–O–C-2', consistent with the 11 degrees of unsaturation required by the molecular formula. Acid hydrolysis result confirmed the sugar to be a β -D-glucopyranose. Hence, the structure of **3** was characterized as 2',5-dihydroxy-4,5'-dimethoxy-3'-O- β -D-glucopyranosyl-2-carboxy-diphenyl ether 2,2'-lactone and was named hyperelatoside C.

Compound **4** was obtained as a brown amorphous solid with a molecular formula of $\text{C}_{20}\text{H}_{24}\text{O}_{11}$ based on HRESIMS (m/z 463.1205 $[\text{M} + \text{Na}]^+$, calcd for $\text{C}_{20}\text{H}_{24}\text{NaO}_{11}$, 463.1216). The ^1H NMR data of **4** (measured in CD_3OD , Table 1) exhibited a set of characteristic ABX aromatic protons at δ_{H} 6.85 (1H, d, $J = 8.8$ Hz, H-3), 6.59 (1H, dd, $J = 8.8, 2.9$ Hz, H-4), and 6.46 (1H, d, $J = 2.9$ Hz, H-6) and two *meta*-coupled aromatic protons at δ_{H} 6.24 (1H, d, $J = 2.9$ Hz, H-4') and 5.91 (1H, d, $J = 2.9$ Hz, H-6'), indicating the presence of a 1,2,4-trisubstituted (A-ring) and a 1,2,3,5-tetrasubstituted (B-ring) benzene rings, respectively. It also showed aliphatic signals for two methoxy groups [δ_{H} 3.68 (3H, s) and 3.65 (3H, s)] and a set of sugar protons (δ_{H} 3.25–4.73), including an anomeric proton at δ_{H} 4.73 (1H, d, $J = 7.6$ Hz, H-1'). The ^{13}C NMR spectrum displayed signals for 12 aromatic carbons (δ_{C} 97.3–158.8), two methoxy carbons (δ_{C} 56.4 and 56.1), and a characteristic glucopyranosyl moiety (δ_{C} 107.4, 78.5, 77.9, 75.5, 71.0, and 62.2) (Table 2). Acid hydrolysis result of **4** confirmed the sugar to be a β -D-glucopyranose. The assignment of signals corresponding to A-ring was based on the HMBC correlations from H-3 (δ_{H} 6.85) to C-1 (δ_{C} 146.0) and C-5 (δ_{C} 155.0), from H-4 (δ_{H} 6.59) to C-2 (δ_{C} 143.3) and C-6 (δ_{C} 107.4), and from H-6 (δ_{H} 6.46) to C-2 (δ_{C} 143.3) and C-4 (δ_{C} 110.6) (Fig. 2). The methoxy protons at δ_{H} 3.68 exhibited a HMBC correlation to C-5 (δ_{C} 155.0), which indicated the connection of a methoxy group at C-5 position of A-ring. Furthermore, the ^1H NMR spectrum of **4**, measured in $\text{DMSO}-d_6$, displayed a hydroxyl proton signal at δ_{H} 8.74 (1H, s, 2-OH), which showed HMBC correlations to C-1, C-2, and C-3, supporting that a hydroxyl group was attached to C-2 position of A-ring. Complete assignment of the ^1H and ^{13}C NMR data of B-ring was achieved by the HMBC correlations from H-4' (δ_{H} 6.24) to C-2' (δ_{C} 131.4), C-3' (δ_{C} 153.1), C-5' (δ_{C} 158.8), and C-6' (δ_{C} 97.3) and from H-6' (δ_{H} 5.91) to C-1' (δ_{C} 152.1), C-2' (δ_{C} 131.4), C-4' (δ_{C} 98.2), and C-5' (δ_{C} 158.8) (Fig. 2). The HMBC correlations from the methoxy protons at δ_{H} 3.65 to C-5' and from the anomeric proton at δ_{H} 4.73 to C-2' suggested the locations of a methoxy group at C-5' and a β -D-glucopyranosyl moiety at C-2', respectively. An ether linkage between C-1 (δ_{C} 146.0) of A-ring and C-1' (δ_{C} 152.1) of B-ring was then established due to their downfield chemical shifts and the molecular formula. Accordingly, compound **4** was identified as 2,3'-dihydroxy-5,5'-dimethoxy-2'-O- β -D-glucopyranosyl-diphenyl ether and was named hyperelatoside D.

Compound **5** was obtained as a brown amorphous solid. Its molecular formula was determined to be $\text{C}_{20}\text{H}_{24}\text{O}_{11}$ based on a deprotonated ion peak at m/z 439.1247 $[\text{M} - \text{H}]^-$ (calcd for $\text{C}_{20}\text{H}_{23}\text{O}_{11}$, 439.1240) in HRESIMS, identical to that found for

compound **4**. Moreover, the UV and IR spectra of **5** showed similar absorption bands to those of **4**, indicating their similar structures. The ^1H and ^{13}C NMR data (Tables 1 and 2) of A-ring of **5** were almost superimposable to those of **4**, but they differed in their B-ring, suggesting that the locations of the substituent groups on B-ring were different between **4** and **5**. The glycosylation site at C-3' position of B-ring of **5** was inferred from an upfield shift of δ_{C} 4.6 ppm for C-3' (*ipso*-C) and downfield shifts of δ_{C} 2.3 ppm for C-2' (*ortho*-C), 2.2 ppm for C-4' (*ortho*-C), and 3.9 ppm for C-6' (*para*-C) (Table 2). This was further supported by a HMBC cross-peak between the anomeric proton at δ_{H} 4.80 (H-1') and the corresponding aglycone carbon at δ_{C} 148.5 (C-3') (Fig. 2). The β -D-glucopyranosyl moiety was determined by the same method as described for **1**. Therefore, the structure of **5** was defined as 2,2'-dihydroxy-5,5'-dimethoxy-3'-O- β -D-glucopyranosyl-diphenyl ether and was named hyperelatoside E.

Compound **6** was purified as white needles with a melting point of 220–222 °C. The molecular formula, $\text{C}_{15}\text{H}_{20}\text{O}_{10}$, was deduced from HRESIMS (m/z 383.0946 $[\text{M} + \text{Na}]^+$, calcd for $\text{C}_{15}\text{H}_{20}\text{NaO}_{10}$, 383.0954). The IR spectrum showed absorption bands corresponding to a hydroxyl group (3439 cm^{-1}) and a conjugated carbonyl group (1666 cm^{-1}). The ^1H NMR spectrum exhibited signals attributable to one hydroxyl group [δ_{H} 10.46 (1H, s, 2-OH)], one 1,2,4,5-tetrasubstituted benzene ring [δ_{H} 7.22 (1H, s, H-6) and 6.71 (1H, s, H-3)], one anomeric proton at δ_{H} 5.04 (1H, d, $J = 7.2$ Hz, H-1'), and two methoxy groups [δ_{H} 3.88 (3H, s, H-8) and 3.73 (3H, s, H-9)] (Table 3). The ^{13}C NMR spectrum displayed 15 carbon resonances, which were classified as an ester carbonyl carbon (δ_{C} 169.3), six aromatic carbons (δ_{C} 103.5–156.6), a characteristic glucopyranosyl moiety (δ_{C} 99.4, 77.1, 76.7, 73.0, 69.5, and 60.6), and two methoxy carbons (δ_{C} 56.1 and 52.3) (Table 3). The β -D-glucopyranosyl unit was verified by the same method as described for **1**. In the HMBC spectrum of **6**, the methoxy protons at δ_{H} 3.88 (H-8) exhibited a correlation with the carbonyl carbon at δ_{C} 169.3 (C-7), which suggested that **6** is a derivative of methyl benzoate. The key HMBC correlations from H-6 (δ_{H} 7.22) to C-1 (δ_{C} 104.1), C-2 (δ_{C} 156.6), C-4 (δ_{C} 153.3), C-5 (δ_{C} 142.1), and C-7 (δ_{C} 169.3), from the hydroxyl proton at δ_{H} 10.46 to C-1 (δ_{C} 104.1), C-2 (δ_{C} 156.6), and C-3 (δ_{C} 103.5), and from the methoxy protons at δ_{H} 3.73 (H-9) to C-5 (δ_{C} 142.1) (Fig. 2), indicated the locations of a hydroxyl group and a methoxy group at C-2 and C-5, respectively. In addition, the anomeric proton at δ_{H} 5.04 (H-1') exhibited a HMBC correlation with C-4 (δ_{C} 153.3), indicating that the glucopyranosyl moiety was attached to C-4 position. Thus, compound **6** was assigned as 4-O- β -D-glucopyranosyl-2-hydroxy-5-methoxy-methylbenzoate and was named hyperelatoside F.

The remaining compounds were elucidated on the basis of NMR and ESIMS data analysis as well as by comparison with literature data. They were identified as 3,5-dihydroxy-4-methoxyxanthone (**7**),¹⁶ 1,5,6-trihydroxy-7-methoxyxanthone (**8**),¹⁷ 1,3,7-trihydroxy-2-(2-hydroxy-3-methyl-3-butenyl)-xanthone (**9**),¹⁸ 2,4-dihydroxy-3-methyl-6-O- β -D-glucopyranosyl benzophenone (**10**),¹⁹ garcimangosone D (**11**),²⁰ 1-(2-methylbutyl)-phloroglucinol-glucopyranoside (**12**),²¹ quercetin 3-O- α -L-rhamnopyranoside (**13**),²² quercetin 3-O- β -D-galactopyranoside (**14**),²³ and kaempferol 3-O- α -L-rhamnopyranoside (**15**)²⁴ (Fig. S54, ESI†).

Table 3 ^1H NMR (500 MHz) and ^{13}C NMR (125 MHz) spectroscopic data for compound **6** (DMSO- d_6)

| No. | δ_{H} (J in Hz) | δ_{C} |
|-------|-------------------------------|---------------------|
| 1 | | 104.1 |
| 2 | | 156.6 |
| 3 | 6.71, s | 103.5 |
| 4 | | 153.3 |
| 5 | | 142.1 |
| 6 | 7.22, s | 111.5 |
| 7 | | 169.3 |
| 8 | 3.88, s | 52.3 |
| 9 | 3.73, s | 56.1 |
| 4a | | |
| 4b | | |
| 8a | | |
| 9a | | |
| 1' | 5.04, d (7.2) | 99.4 |
| 2' | 3.26, m | 73.0 |
| 3' | 3.28, m | 76.7 |
| 4' | 3.16, m | 69.5 |
| 5' | 3.39, m | 77.1 |
| 6' | 3.45, dd (11.7, 5.2) | 60.6 |
| | 3.66, dd (11.7, 6.0) | |
| 1-OH | | |
| 2-OH | 10.46, s | |
| 7-OH | | |
| 2'-OH | 5.34, d (5.1) | |
| 3'-OH | 5.13, d (4.6) | |
| 4'-OH | 5.06, d (5.3) | |
| 6'-OH | 4.58, t (5.6) | |

Among them, compounds **7**, **10**, and **12** were isolated from the genus *Hypericum* for the first time.

Naturally occurring biphenyl ethers have been reported to be in some fungi in recent years, especially *Aspergillus* species,²⁵ but are rarely found in plants.²⁶ In fungi, these kinds of compounds are formed biosynthetically from the anthraquinone emodin, *via* sulochrin, and the grisandienes such as geodin was previously reported.²⁷ However, these intermediates of desmethylsulochrin, sulochrin, dihydrogeodin, and geodin are rarely as the plant metabolites. Given the isolation of benzoic acid derivatives (**6** and **12**), benzophenones (**10** and **11**), and xanthenes (**7**–**9**) from *H. elatoides*, the possible biosynthetic pathway of hyperelatosides A–E (**1**–**5**) is proposed in Scheme 1. The key step of this pathway is the xanthone ring cleavage reaction to form biphenyl ether skeleton, the immediate precursors of which may be 1,3,6,7- and 1,3,5,8-tetrahydroxyxanthenes. Although the formation of biphenyl ethers from xanthenes catalyzed by an enzyme system involving an oxygenase, has not yet detected in plants, the similar reaction has been observed in the biosynthetic studies of biphenyl ethers from anthraquinones in fungi. An anthraquinone ring cleavage enzyme, named questin oxygenase, was found in the cell-free extract of *Aspergillus terreus* and the reaction mechanism underlying the ring cleavage of anthraquinone is likely to be a chemical Baeyer–Villiger oxidation and hydrolysis of the formed lactone intermediate.^{27,28} This suggested that the same type of enzymes as questin oxygenase may be involved in the formation of biphenyl ethers from xanthenes in *H. elatoides*.

Xanthenes and benzophenones are two major constituents of the plants of the genus *Hypericum*.^{1c} The biosynthesis of xanthone products has been studied in *H. androsaemum*, *H. calycinum*, and *H. perforatum*.²⁹ 2,3',4,6-Tetrahydroxybenzophenone, which is formed from benzoic acid by the condensation of one benzoyl CoA and three malonyl CoA units catalyzed by benzophenone synthase, is cyclized regioselectively to yield either 1,3,7- or 1,3,5-trihydroxyxanthone, which is then regioselectively hydroxylated to 1,3,6,7- or 1,3,5,8-tetrahydroxyxanthone.³⁰ Subsequent xanthone ring cleavage reaction gives the biphenyl ether skeleton, which may generate the structures of **1**–**5** *via* enzymatic reactions of decarboxylation, *O*-methylation using *S*-adenosylmethionine (SAM), and glycosylation.

The neurotrophic activities of new compounds **1**–**6** were investigated in terms of their ability to enhance NGF-induced neurite outgrowth using rat pheochromocytoma (PC12) cells as a model system of neuronal differentiation. Fig. 3 showed that, relative to the NGF control (100%), compounds **1**, **2**, and **4**–**6** exerted a significant increase in neurite-bearing cells at 1 μM . In particular, the percentages of neurite-bearing cells for cells treated with hyperelatosides B (**2**) and D (**4**) in the presence of NGF (20 ng mL⁻¹) reached up to 192.78 \pm 17.71% and 228.70 \pm 12.51% relative to the NGF control, respectively. Interestingly, among the biphenyl ether glycosides (**1**–**5**), only **3** had no effect on NGF-mediated neurite outgrowth in PC12 cells at 1 μM , suggesting that the presence of a carboxyl or hydroxyl group at C-2 position of A-ring may be responsible for the neurotrophic activity. Compound **2** showed much stronger activity than **1**, indicating that attachment of an active hydrogen atom to the O-7 position of the carboxyl group could significantly improve this activity than attachment of a methyl group to O-7 position.

Neuroinflammation has been typically involved in the pathology of neurodegenerative diseases such as Alzheimer's disease and Parkinson's disease. Therefore, the inhibitory effects of compounds **1**–**15** on LPS-stimulated NO production in BV-2 microglial cells were investigated to establish their anti-neuroinflammatory activities. As shown in Fig. 4, compounds **5**, **8**, and **14** exhibited inhibitory effects against NO production in BV-2 cells at 25 μM . In order to investigate whether the inhibitory activities of the active compounds were due to their cytotoxicity towards BV-2 cells, the effects of compounds **5**, **8**, and **14** on cell proliferation/viability were measured by 3-(4,5-dimethylthiazole-2-yl)-2,5-diphenyltetrazolium bromide (MTT) method. Of these, compounds **5** and **8** showed no cytotoxicity with LPS treatment for 24 h.

In addition, severe oxidative stress has been also implicated in the pathogenesis of neurodegenerative diseases. The antioxidant activities of compounds **1**–**15** were preliminarily evaluated for their DPPH free radical scavenging properties at 1, 5, 10, 25, 50, and 100 μM using ascorbic acid as the positive control (IC₅₀ = 12.45 \pm 0.51 μM). The result showed that compound **14** exhibited the most significant DPPH radical scavenging activity with IC₅₀ value of 11.22 \pm 0.38 μM (Table 4), which showed stronger antioxidant activity than the positive control. Its potent activity may be due to the presence of two hydroxyl groups at C-3 and C-4 positions of the aromatic ring, which agrees with previous studies.³¹ Moreover, compounds **2**,




Fig. 3 Effects of compounds 1–6 on neurite outgrowth in PC12 cells. (A) Morphological changes of PC12 cells treated with 0.1% DMSO, NGF (20 ng mL⁻¹), and NGF (20 ng mL⁻¹) + 4 (1.0 μM) were visualized under light microphotography. The cells treated with 4 served as a representative example to demonstrate the enhancement of neurite outgrowth in the presence of NGF. (B) Neurite-bearing cells were quantitatively analyzed as described in measurement of neurite outgrowth. Data are expressed as percentages of the value of NGF-treated cells (means \pm standard deviations, $n = 3$) (D, 0.1% DMSO, control; N, 20 ng mL⁻¹ NGF, positive control; ### $p < 0.001$ vs. control; * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ vs. NGF control).




Fig. 4 Effects of compounds 1–15 on LPS-induced NO production in BV-2 microglial cells. Data are expressed as percentages of the value obtained with LPS-treated cells. (C, untreated control, 0.1% DMSO; L, LPS-treated control; Q, quercetin, 5 μM , positive control; ### $p < 0.001$ compared with control group, ** $p < 0.01$, *** $p < 0.001$ compared with LPS group).

Extraction and isolation

The aerial parts (1.3 kg) of *H. elatoides* were dried in shade, cut into small pieces, and extracted three times (each for 8 h) with 12 L of MeOH under reflux conditions (55 °C). After filtration and evaporation *in vacuo*, the obtained extract (334.8 g) was

suspended in distilled water (3 L) and partitioned successively with *n*-hexane (3 L \times 3), EtOAc (3 L \times 3), and *n*-BuOH (3 L \times 3) to yield a hexane-soluble fraction (55.6 g), an EtOAc-soluble fraction (49.2 g), and a BuOH-soluble fraction (71.0 g), respectively. The EtOAc fraction was subjected to silica gel CC and eluted with a gradient of CH₂Cl₂/MeOH (100 : 1 to 2 : 1) to yield fifteen fractions (E1–E15). Fr. E3 was further subjected to RP-C₁₈ CC (acetone–H₂O, 1 : 6) to give ten subfractions (E3.1–E3.10). Fr. E3.5 was subjected to silica gel CC (CHCl₃–MeOH, 100 : 1) to afford compound 7 (63.1 mg). Fr. E6 was subjected to silica gel CC (CH₂Cl₂–MeOH, 30 : 1 to 20 : 1) to yield eight subfractions (E6.1–E6.8). Fr. E6.3 was purified on a RP-C₁₈ column (MeOH–H₂O, 3 : 1) to yield compound 8 (26.5 mg). Fr. E6.4 was purified on a RP-C₁₈ column (acetone–MeOH, 1 : 5) to yield compound 9 (24.6 mg). Fr. E9 was subjected to silica gel CC and eluted with a gradient of CHCl₃–MeOH–H₂O (15 : 1 : 0.05 to 4 : 1 : 0.1) to yield eight fractions (E9.1–E9.8). Fr. E9.4 was further separated by RP-C₁₈ CC (MeOH–H₂O, 1 : 2 to 1 : 1) to yield seven subfractions (E9.4.1–E9.4.7). Compound 6 (5.3 mg) was obtained by the separation of Fr. E9.4.3 on a Sephadex LH-20 column (MeOH–H₂O, 1 : 2). Compounds 1 (34.2 mg), 2 (21.5 mg), 3 (19.4 mg), and 4 (10.0 mg) were obtained by the separation of Fr.

Table 4 Antioxidant activities of the isolated compounds by DPPH radical scavenging assay

| Compound ^a | IC ₅₀ (μM) |
|-----------------------|-----------------------|
| 2 | 85.50 ± 3.08 |
| 4 | 87.21 ± 1.64 |
| 5 | 49.28 ± 1.09 |
| 8 | 64.61 ± 2.99 |
| 13 | 16.44 ± 0.16 |
| 14 | 11.22 ± 0.38 |
| Ascorbic acid | 12.45 ± 0.51 |

^a Compounds 1, 3, 6, 7, 9–12, and 15 showed weak or no activities in the DPPH antioxidant assay and ascorbic acid was used as a positive control.

E9.4.5 on a RP-C₁₈ column (acetone–H₂O, 1 : 4 to 1 : 2.5). Fr. E9.4.6 was subjected to RP-C₁₈ CC (acetone–H₂O, 1 : 2.5) to yield compound **10** (35.9 mg). Fr. E9.5 was separated by RP-C₁₈ CC (acetone–H₂O, 1 : 6 to 1 : 4) to give seven subfractions (E9.5.1–E9.5.7). Fr. E9.5.2 was subjected to RP-C₁₈ CC (MeOH–H₂O, 1 : 3) to yield compound **11** (3.4 mg). Fr. E9.5.6 was subjected to RP-C₁₈ CC (MeOH–H₂O, 1 : 1.4) followed by RP-C₁₈ CC (acetone–H₂O, 1 : 2) to provide compound **12** (4.2 mg). Fr. E9.5.7 was purified on a RP-C₁₈ column (MeOH–H₂O, 1 : 1.5) to yield compound **15** (7.6 mg). Fr. E10 was separated by silica gel CC (CHCl₃–MeOH–H₂O, 15 : 1 : 0.05 to 3 : 1 : 0.1) to afford seven subfractions (E10.1–E10.7). Fr. E10.3 was subjected to RP-C₁₈ CC (acetone–H₂O, 1 : 5) to yield four fractions (E10.3.1–E10.3.4). Fr. E10.3.2 was subjected to RP-C₁₈ CC (MeOH–H₂O, 1 : 2.5 to 1 : 1.5) to afford eight fractions (E10.3.2.1–E10.3.2.8). Compound **5** (7.6 mg) was obtained by the separation of Fr. E10.3.2.2 on a RP-C₁₈ column (MeOH–H₂O, 1 : 2). Fr. E10.6 was separated on a silica gel column (CHCl₃–MeOH–H₂O, 4 : 1 : 0.1) to yield compound **13** (960.0 mg). Fr. E11 was subjected to silica gel CC and eluted with a gradient of CHCl₃–MeOH–H₂O (6 : 1 : 0.1 to 2 : 1 : 0.1) to afford six subfractions (E11.1–E11.6). Fr. E11.4 was further subjected to RP-C₁₈ CC (acetone–H₂O, 1 : 3 to 1 : 2) to give compound **14** (161.3 mg).

Hyperelatoside A (1). Brown amorphous solid; $[\alpha]_{\text{D}}^{20} -38.2$ (*c* 0.1, MeOH); UV (MeOH) λ_{max} (log ϵ) 226 (3.9), 260 (2.4), 292 (2.0) nm; IR (KBr) ν_{max} 3418, 2952, 1701, 1618, 1515, 1440, 1379, 1273, 1209, 1076, 1028, 630 cm⁻¹; ¹H and ¹³C NMR data (CD₃OD and DMSO-*d*₆), see Tables 1 and 2; HRESIMS (positive) *m/z* 521.1247 [M + Na]⁺ (calcd for C₂₂H₂₆NaO₁₃, 521.1271).

Hyperelatoside B (2). Brown amorphous solid; $[\alpha]_{\text{D}}^{20} -25.5$ (*c* 0.1, MeOH); UV (MeOH) λ_{max} (log ϵ) 225 (5.4), 260 (1.0) nm; IR (KBr) ν_{max} 3415, 1624, 1514, 1444, 1274, 1211, 1170, 1027, 630 cm⁻¹; ¹H and ¹³C NMR data (CD₃OD), see Tables 1 and 2; HRESIMS (negative) *m/z* 483.1150 [M – H]⁻ (calcd for C₂₁H₂₃O₁₃, 483.1139).

Hyperelatoside C (3). Brown amorphous solid; $[\alpha]_{\text{D}}^{20} = -21.2$ (*c* 0.05, MeOH); UV (MeOH) λ_{max} (log ϵ) 225 (5.4), 260 (2.4), 292 (1.0) nm; IR (KBr) ν_{max} 3434, 2954, 1640, 1514, 1454, 1019, 661 cm⁻¹; ¹H and ¹³C NMR data (CD₃OD), see Tables 1 and 2; HRESIMS (positive) *m/z* 489.0998 [M + Na]⁺ (calcd for C₂₁H₂₂NaO₁₂, 489.1009).

Hyperelatoside D (4). Brown amorphous solid; $[\alpha]_{\text{D}}^{20} -44.3$ (*c* 0.1, MeOH); UV (MeOH) λ_{max} (log ϵ) 290 (2.3) nm; IR (KBr) ν_{max} 3402, 2950, 2837, 1650, 1455, 1410, 1111, 1022, 671 cm⁻¹; ¹H and ¹³C NMR data (CD₃OD and DMSO-*d*₆), see Tables 1 and 2; HRESIMS (positive) *m/z* 463.1205 [M + Na]⁺ (calcd for C₂₀H₂₄NaO₁₁, 463.1216).

Hyperelatoside E (5). Brown amorphous solid; $[\alpha]_{\text{D}}^{20} -28.4$ (*c* 0.02, MeOH); UV (MeOH) λ_{max} (log ϵ) 290 (3.0) nm; IR (KBr) ν_{max} 3392, 2950, 2836, 1651, 1457, 1411, 1111, 1023, 671 cm⁻¹; ¹H and ¹³C NMR data (CD₃OD), see Tables 1 and 2; HRESIMS (negative) *m/z* 439.1247 [M – H]⁻ (calcd for C₂₀H₂₃O₁₁, 439.1240).

Hyperelatoside F (6). White needles; mp 220–222 °C; $[\alpha]_{\text{D}}^{20} -114.3$ (*c* 0.05, MeOH); UV (MeOH) λ_{max} (log ϵ) 222 (3.3), 256 (2.9), 319 (3.2) nm; IR (KBr) ν_{max} 3439, 2954, 1666, 1626, 1509, 1445, 1366, 1267, 1226, 1062, 694, 652 cm⁻¹; ¹H and ¹³C NMR data (DMSO-*d*₆), see Table 3; HRESIMS (positive) *m/z* 383.0946 [M + Na]⁺ (calcd for C₁₅H₂₀NaO₁₀, 383.0954).

Acid hydrolysis

The absolute configurations of the sugar moieties of **1–6** were determined by the acid hydrolysis method.³² Each compound (approximately 1.0 mg) was separately dissolved in 1 N HCl (0.5 mL), and then heated at 90 °C in a water bath for 2 h. After extraction with EtOAc two times, the H₂O-soluble fraction was evaporated to dryness under reduced pressure. The residue was dissolved in pyridine (0.1 mL) containing L-cysteine methyl ester hydrochloride (0.5 mg) and heated at 60 °C for 1 h. A 100 μL solution of *o*-tolylisothiocyanate (0.5 mg) in pyridine was added and the reaction mixture was heated at 60 °C for 1 h. Then each reaction mixture was analyzed by the Waters 1525 HPLC system using a Waters 2489 UV/vis detector (at 250 nm). Analytical HPLC was performed on the YMC Packed C₁₈ column (5 μm, 150 × 4.6 mm) with a linear gradient elution (CH₃CN–H₂O, 20 : 80 to 40 : 60) for 30 min. The derivative of D-glucose was identified in **1–6** by a comparison of the retention time with authentic D-glucose (*t*_R 16.3 min), which was subjected to the same derivatization procedure.

Measurement of neurite outgrowth

PC12 cells were purchased from the Cell Bank of Shanghai Institute of Biochemistry and Cell Biology, Chinese Academy of Sciences (Shanghai, China). PC12 cells were maintained in nutrient mixture F-12 medium supplemented with 10% inactivated HS, 5% inactivated FBS, penicillin G (100 U mL⁻¹), streptomycin (100 μg mL⁻¹), and sodium bicarbonate (2.5 g L⁻¹) at 37 °C in humidified atmosphere of 5% CO₂. Morphological analysis and quantification of neurite-bearing cells were performed using a phase-contrast microscope as previously described.¹⁵ Briefly, PC12 cells were seeded in poly-L-lysine-coated 24-well plates at a density of 2 × 10⁴ cells per mL with normal serum medium for 24 h. The F-12 medium containing low serum (2% HS and 1% FBS) was replaced prior to exposure to vehicle (0.1% DMSO) or indicated reagents. The cells were treated with tested compounds in the presence of NGF (20 ng mL⁻¹). Cells without treatment served as a negative control.

Cells treated with 20 ng mL⁻¹ of NGF served as a positive control. One concentration experiment was repeated in three wells. After an additional 48 h of incubation, neurite outgrowth of PC12 cells was observed under an inverted microscope using phase-contrast objectives and photographed by the digital camera. Eight images were selected randomly under a microscope for each well. At least 100 cells in each of eight randomly separated fields were scored. The cells with neurites greater than or equal to the length of one cell body were positive for neurite outgrowth and expressed as a percentage of the total cell number in eight fields. Experiments were repeated at least three times, and data are expressed as means \pm standard deviations.

Determination of NO production

BV-2 microglial cells were purchased from Peking Union Medical College Cell Bank (Beijing, China) and cultured in DMEM supplemented with 10% heat-inactivated fetal bovine serum, penicillin G (100 U mL⁻¹), and streptomycin (100 μ g mL⁻¹) at 37 °C in a humidified incubator with 5% CO₂. The NO concentration was detected by the Griess reagent.³³ Quercetin was used as a positive control. BV-2 cells were seeded at the density of 1.5 \times 10⁵ cells per mL in 96-well culture plate and treated with each compound and LPS (1.0 μ g mL⁻¹) for 24 h. After that, 50 μ L of cell-free supernatant was allowed to react with an equal volume of Griess reagent (Beyotime Institute of Biotechnology, Shanghai, China) for 15 min at room temperature in the dark. Then, the absorbance was measured at 540 nm using a microplate reader. The cell viability of the cultured cells was detected by MTT-based colorimetric method.

Antioxidant activity

The DPPH radical scavenging assay was performed according to a previously described method.³⁴ Ascorbic acid was used as a positive control. The absorbance of the resulting solution of each compound was measured at 517 nm with a spectrophotometer. Results were expressed as IC₅₀ values.

Conclusions

In summary, this study is the first report to describe the chemical constituents and their biological activities of *H. elatoides*. Chemical investigation of the aerial parts of *H. elatoides* led to the isolation of six new phenolic compounds including five unusual biphenyl ether glycosides, hyperelatosides A–E (1–5), and one benzoate glycoside, hyperelatoside F (6). To the best of our knowledge, this represents the first report of the isolation of biphenyl ether glycosides from plants and their putative biosynthetic pathway is proposed. Furthermore, except for 3, all the new isolates possessed potent ability to potentiate the activity of NGF to stimulate neurite outgrowth in PC12 cells. Compounds 5 and 8 exhibited moderate anti-neuroinflammatory activities and 14 showed significant antioxidant activity *in vitro*. Our findings show that the aerial parts of *H. elatoides* may be an excellent source of neurotrophic phytochemicals for the prevention and treatment of neurodegenerative diseases.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was financially supported by the Scientific Startup Foundation for Doctors of Northwest A&F University (No. 2452015356) and the Fundamental Research Funds for the Central Universities (No. 2452017173) and the Natural Science Foundation of China (No. 21572182).

Notes and references

- (a) C. C. Sánchez-Mateo, B. Prado and R. M. Rabanal, *J. Ethnopharmacol.*, 2002, **79**, 119; (b) M. F. Rieli, R. Mattei and C. E. L. de Araujo, *Fitoterapia*, 2002, **73**, 462; (c) A. Viana, J. C. do Rego, G. von Poser, A. Ferraz, A. P. Heckler, J. Costentin and S. M. K. Rates, *Neuropharmacology*, 2005, **49**, 1042; (d) O. Grundmann, Y. Lv, O. Kelber and V. Butterweck, *Neuropharmacology*, 2010, **58**, 767; (e) J. Zhao, W. Liu and J. C. Wang, *Chem. Biodiversity*, 2015, **12**, 309.
- (a) S. S. Chatterjee, S. K. Bhattacharya, M. Wonnemann, A. Singer and W. E. Müller, *Life Sci.*, 1998, **63**, 499; (b) J. Sarris, *Phytother. Res.*, 2007, **21**, 703; (c) S. Kasper, F. Caraci, B. Forti, F. Drago and E. Aguglia, *Eur. Neuropsychopharmacol.*, 2010, **20**, 747.
- (a) R. Ciochina and R. B. Grossman, *Chem. Rev.*, 2006, **106**, 3963; (b) I. P. Singh and S. B. Bharate, *Nat. Prod. Rep.*, 2006, **23**, 558; (c) J. J. Zhang, J. Yang, Y. Liao, X. W. Yang, J. Z. Ma, Q. L. Xiao, L. X. Yang and G. Xu, *Org. Lett.*, 2014, **16**, 4912; (d) X. W. Yang, R. B. Grossman and G. Xu, *Chem. Rev.*, 2018, **118**, 3508.
- (a) Z. Q. Yin, Y. Wang, W. C. Ye and S. X. Zhao, *Biochem. Syst. Ecol.*, 2004, **32**, 521; (b) N. Tanaka, Y. Kashiwada, S. Y. Kim, M. Sekiya, Y. Ikeshiro and Y. Takaishi, *Phytochemistry*, 2009, **70**, 1456; (c) W. J. Xu, R. J. Li, O. Quasie, M. H. Yang, L. Y. Kong and J. Luo, *J. Nat. Prod.*, 2016, **79**, 1971.
- (a) A. Hecka, B. Maunit, F. Aubriet and J. F. Muller, *Rapid Commun. Mass Spectrom.*, 2009, **23**, 885; (b) M. A. Farag and L. A. Wessjohann, *Planta Med.*, 2012, **78**, 488; (c) A. Porzel, M. A. Farag, J. Mülbradt and L. A. Wessjohann, *Metabolomics*, 2014, **10**, 574.
- (a) C. A. Ang'edu, J. Schmidt, A. Porzel, P. Gitu, J. O. Midiwo and G. Adam, *Pharmazie*, 1999, **54**, 235; (b) S. A. F. Tanemossu, K. Franke, N. Arnold, J. Schmidt, H. K. Wabo, P. Tane and L. A. Wessjohann, *Phytochemistry*, 2014, **105**, 171.
- (a) L. R. Jat, *Int. J. Pharm. Pharm. Sci.*, 2013, **5**, 9; (b) Z. B. Zhou, Z. R. Li, X. B. Wang, J. G. Luo and L. Y. Kong, *J. Nat. Prod.*, 2016, **79**, 1231.
- (a) M. A. G. del Rio, M. I. Sanchez-Reus, I. Iglesias, M. A. Pozo, M. Garcia-Arencibia, J. Fernandez-Ruiz, L. Garcia-Garcia, M. Delgado and J. Benedi, *CNS Neurol. Disord.: Drug Targets*, 2013, **12**, 665; (b) D. Zheleva-

- Dimitrova, P. Nedialkov and G. Momekov, *Pharmacogn. Mag.*, 2013, **9**, 1.
- 9 D. Ben-Eliezer and E. Yechiam, *Sci. Rep.*, 2016, **6**, 35700.
- 10 N. Huang, L. Rizshsky, C. Hauck, B. J. Nikolau, P. A. Murphy and D. F. Birt, *Phytochemistry*, 2011, **72**, 2015.
- 11 L. C. Brito, A. L. R. Berenger and M. R. Figueiredo, *Food Chem. Toxicol.*, 2017, **109**, 847.
- 12 H. Zhu, C. Chen, J. Yang, X. N. Li, J. Liu, B. Sun, S. X. Huang, D. Li, G. Yao, Z. Luo, Y. Li, J. Zhang, Y. Xue and Y. Zhang, *Org. Lett.*, 2014, **16**, 6322.
- 13 H. Jayasuriya, A. M. Clark and J. D. McChesney, *J. Nat. Prod.*, 1991, **54**, 1314.
- 14 (a) P. L. Liu, C. Du, Z. M. Jiang and J. Cai, *Xibei Zhiwu Xuebao*, 2011, **31**, 1268; (b) Q. Sun, P. L. Liu, Q. Zhang, L. X. Cao and Z. M. Jiang, *Xibei Zhiwu Xuebao*, 2012, **32**, 276.
- 15 (a) X. W. Shi, L. Liu, J. M. Gao and A. L. Zhang, *Eur. J. Med. Chem.*, 2011, **46**, 3112; (b) C. C. Zhang, X. Yin, C. Y. Cao, J. Wei, Q. Zhang and J. M. Gao, *Bioorg. Med. Chem. Lett.*, 2015, **25**, 5078; (c) R. Bai, C. C. Zhang, X. Yin, J. Wei and J. M. Gao, *J. Nat. Prod.*, 2015, **78**, 783; (d) C. C. Zhang, C. Y. Cao, M. Kubo, K. Harada, X. T. Yan, Y. Fukuyama and J. M. Gao, *Int. J. Mol. Sci.*, 2017, **18**, 1659.
- 16 Z. Zhang, H. N. ElSohly, M. R. Jacob, D. S. Pasco, L. A. Walker and A. M. Clark, *Planta Med.*, 2002, **68**, 49.
- 17 N. Tanaka and Y. Takaishi, *Phytochemistry*, 2006, **67**, 2146.
- 18 M. Iinuma, H. Tosa, T. Ito, T. Tanaka and M. Aqil, *Phytochemistry*, 1995, **40**, 267.
- 19 S. E. Ukwueze, P. O. Osadebe and F. B. C. Okoye, *Nat. Prod. Res.*, 2015, **29**, 1728.
- 20 Y. L. Huang, C. C. Chen, Y. J. Chen, R. L. Huang and B. J. Shieh, *J. Nat. Prod.*, 2001, **64**, 903.
- 21 G. Bohr, C. Gerhaeuser, J. Knauff, J. Zapp and H. Becker, *J. Nat. Prod.*, 2005, **68**, 1545.
- 22 T. Fukunaga, K. Nishiya, I. Kajikawa, Y. Watanabe, N. Suzuki, K. Takeya and H. Itokawa, *Chem. Pharm. Bull.*, 1988, **36**, 1180.
- 23 K. Yasukawa and M. Takido, *Phytochemistry*, 1987, **26**, 1224.
- 24 C. Y. Chen, S. L. Hsieh, M. M. Hsieh, S. F. Hsieh and T. J. Hsieh, *Chin. Pharm. J.*, 2004, **56**, 141.
- 25 (a) R. Liu, W. Zhu, Y. Zhang, T. Zhu, H. Liu, Y. Fang and Q. Gu, *J. Antibiot.*, 2006, **59**, 362; (b) T. Bunyapaiboonsri, S. Yoiprommarat, K. Intereya and K. Kocharin, *Chem. Pharm. Bull.*, 2007, **55**, 304; (c) M. J. Fang, H. Fang, W. J. Li, D. M. Huang, Z. Wu and Y. F. Zhao, *Nat. Prod. Res.*, 2012, **26**, 1224; (d) W. Peng, F. You, X. L. Li, M. Jia, C. J. Zheng, T. Han and L. P. Qin, *Zhongguo Tianran Yaowu*, 2013, **11**, 673; (e) L. L. Xu, C. C. Zhang, X. Y. Zhu, F. Cao and H. J. Zhu, *Nat. Prod. Res.*, 2017, **31**, 1875; (f) P. Phainuphong, V. Rukachaisirikul, S. Phongpaichit, S. Preedanon and J. Sakayaroj, *Tetrahedron*, 2017, **73**, 5920.
- 26 N. Li, H. S. Lee, N. Zhang, Y. N. Sun, J. L. Li, S. S. Xing, J. G. Chen and L. Cui, *Phytochem. Lett.*, 2015, **13**, 286.
- 27 (a) R. F. Curtis, C. H. Hassall, D. W. Jones and T. W. Williams, *J. Chem. Soc.*, 1960, 4838; (b) Z. G. Chen, I. Fujii, Y. Ebizuka and U. Sankawa, *Phytochemistry*, 1995, **38**, 299; (c) K. Huang, I. Fujii, Y. Ebizuka, K. Gomi and U. Sankawa, *J. Biol. Chem.*, 1995, **270**, 21495; (d) J. Hargreaves, J. Park, E. L. Ghisalberti, K. Sivasithamparam, B. W. Skelton and A. H. White, *J. Nat. Prod.*, 2002, **65**, 7.
- 28 I. Fujii, Y. Ebizuka and U. Sankawa, *J. Biochem.*, 1988, **103**, 878.
- 29 (a) W. Schmidt and L. Beerhues, *FEBS Lett.*, 1997, **420**, 143; (b) M. M. Gaid, D. Sircar, A. Müller, T. Beuerle, B. Liu, L. Ernst, R. Hänsch and L. Beerhues, *Plant Physiol.*, 2012, **160**, 1267; (c) I. El-Awaad, M. Bocola, T. Beuerle, B. Liu and L. Beerhues, *Nat. Commun.*, 2016, **7**, 11472.
- 30 (a) W. Schmidt, S. Peters and L. Beerhues, *Phytochemistry*, 2000, **53**, 427; (b) H. R. El-Seedi, M. A. El-Barbary, D. M. H. El-Ghorab, L. Bohlin, A. K. Borg-Karlson, U. Göransson and R. Verpoorte, *Curr. Med. Chem.*, 2010, **17**, 854.
- 31 (a) P. V. Kiem, N. X. Cuong, N. X. Nhiem, V. K. Thu, N. K. Ban, C. V. Minh, B. H. Tai, T. N. Hai, S. H. Lee, H. D. Jang and Y. H. Kim, *Bioorg. Med. Chem. Lett.*, 2011, **21**, 633; (b) X. T. Yan, S. H. Lee, W. Li, Y. N. Sun, S. Y. Yang, H. D. Jang and Y. H. Kim, *Food Chem.*, 2014, **156**, 408.
- 32 T. Tanaka, T. Nakashima, T. Ueda, K. Tomii and I. Kouno, *Chem. Pharm. Bull.*, 2007, **55**, 899.
- 33 (a) Z. Huang, Z. X. Zhu, Y. T. Li, D. R. Pang, J. Zheng, Q. Zhang, Y. F. Zhao, D. Ferreira, J. K. Zjawiony, P. F. Tu and J. Li, *J. Nat. Prod.*, 2015, **78**, 2276; (b) J. Li, N. Li, X. Li, G. Chen, C. Wang, B. Lin and Y. Hou, *J. Nat. Prod.*, 2017, **80**, 3081; (c) J. Wei, W. H. Guo, C. Y. Cao, R. W. Kou, Y. Z. Xu, M. Górecki, L. D. Bari, G. Pescitelli and J. M. Gao, *Sci. Rep.*, 2018, **8**, 2175.
- 34 (a) O. P. Sharma and T. K. Bhat, *Food Chem.*, 2009, **113**, 1202; (b) C. D. Zheng, G. Li, H. Q. Li, X. J. Xu, J. M. Gao and A. L. Zhang, *Nat. Prod. Commun.*, 2010, **5**, 1759.