



Whole-Genome Sequence of the *Mycoplasma* (*Mesomycoplasma*) *hyorhinis* DSM 25591 Type Strain

Lisa Käbisch,^{a,b} Anne-Kathrin Schink,^a Dennis Hanke,^a  Torsten Semmler,^c Corinna Kehrenberg,^b  Stefan Schwarz^a

^aInstitute of Microbiology and Epizootics, Centre for Infection Medicine, Department of Veterinary Medicine, Freie Universität Berlin, Berlin, Germany

^bInstitute for Veterinary Food Science, Justus-Liebig-Universität Gießen, Gießen, Germany

^cNG1-Microbial Genomics, Robert Koch Institute, Berlin, Germany

ABSTRACT The whole-genome sequence of the type strain *Mycoplasma* (*Mesomycoplasma*) *hyorhinis* DSM 25591 is reported and compared to the available sequences of the corresponding type strains from other strain collections to ascertain conformity. Knowledge of the identity of type strains is of importance for their application in standardized test systems.

Due to reclassification of the *Mollicutes*, the assignment of *Mycoplasma hyorhinis* to the novel genus *Mesomycoplasma* is under discussion (1–3). The American Type Culture Collection (ATCC), the National Collection of Type Cultures (NCTC), and the German Collection of Microorganisms and Cell Cultures GmbH (DSMZ) provide defined type strains. Investigating the degree of identity between type strains of different collections, we subjected *Mycoplasma* (*Mesomycoplasma*) *hyorhinis* DSM 25591 from the DSMZ to whole-genome sequencing. The obtained sequence was compared to the complete genome sequence of *M. hyorhinis* NCTC 10130 (GenBank accession number [LS991950.1](https://doi.org/10.1093/nar/33/1/LS991950.1)) and the partially available sequence of *M. hyorhinis* ATCC 17981 ([NZ_KB911485.1](https://doi.org/10.1093/nar/33/1/NZ_KB911485.1)).

Prior to DNA isolation with the MasterPure complete DNA and RNA purification kit (Epicentre, Madison, WI, USA), *M. hyorhinis* DSM 25591 was cultured using modified Friis broth for 5 days at 37°C in a humidified, 7.5% CO₂ atmosphere (4, 5). The DNA was quantified using a Qubit 3.0 fluorometer (double-stranded DNA broad-range assay kit; Thermo Fisher Scientific, Waltham, MA, USA), and its quality was assessed by spectral analysis (NanoDrop spectrophotometer; Thermo Fisher Scientific). Short-read sequencing was performed on an Illumina MiSeq sequencer with the MiSeq reagent kit v3 (Illumina, Inc., San Diego, CA, USA) and the Nextera XT library preparation kit (Illumina), resulting in 300-bp paired-end reads and a >90-fold coverage. In addition, long-read sequencing was performed using Oxford Nanopore MinION (Oxford Nanopore Technologies [ONT], Oxford, UK). MinION one-dimensional (1D) libraries were constructed using the SQK-RBK004 kit (ONT) and loaded according to the manufacturer's instructions onto an R9.4 flow cell. Sequencing data were collected for 48 h. More detailed information on the sequencing parameters is provided in Table 1. A closed genome sequence was generated by a *de novo* hybrid assembly using a combination of short and long reads with Unicycler v0.4.7 (9). A draft genome assembly created with SPAdes v3.12 and consecutive connection of contigs, using the long reads from MinION, resulted in a single circular chromosomal DNA of 839,538 bp with a G+C content of 25.9% (10). Default parameters were used for the applied software.

Comparison with the sequence of *M. hyorhinis* NCTC 10130 showed an identity of 99.8%. The incomplete sequence of *M. hyorhinis* ATCC 17981 matched our sequence with an identity of 99.1 to 100%, depending on the respective contig. Two open reading frames (ORFs) within the regions encoding variable lipoproteins (Vlps) were further analyzed by PCR and Sanger sequencing (LGC Genomics, Berlin, Germany), as they

Citation Käbisch L, Schink A-K, Hanke D, Semmler T, Kehrenberg C, Schwarz S. 2021. Whole-genome sequence of the *Mycoplasma* (*Mesomycoplasma*) *hyorhinis* DSM 25591 type strain. *Microbiol Resour Announc* 10:e00164-21. <https://doi.org/10.1128/MRA.00164-21>.

Editor Irene L. G. Newton, Indiana University, Bloomington

Copyright © 2021 Käbisch et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Stefan Schwarz, stefan.schwarz@fu-berlin.de.

Received 13 February 2021

Accepted 24 March 2021

Published 22 April 2021

TABLE 1 Sequencing characteristics of *M. hyorhinis* DSM 25591

Technology	Total no. of reads	Mean read length (bp)	Quality control	Adapter trimming	Error correction	Base calling	Avg genome coverage (x)	Amt of DNA used (ng)	SRA accession no.
Illumina MiSeq	1,500,074	285	FastQC (6)	Flexbar v3.0.3 (7)	BayesHammer (8)	NA	437	1	SRX9929725
Nanopore MinION	112,219	5,324	FastQC (6)	NA ^a	NA	Guppy	711	400	SRX9929726

^aNA, not applicable.




FIG 1 Presentation of the total genome sequence of *Mycoplasma* (*Mesomycoplasma*) *hyorhinis* DSM 25591 as a circular closed chromosomal DNA molecule of 839,538 bp, deposited in GenBank under the accession number [CP064323](#). (a) Comparison of the *vlpG* and *vlpB* genes of *M. hyorhinis* DSM 25591 ([CP064323](#)) with those of *M. hyorhinis* NCTC 10130 ([LS991950.1](#)). The genes are shown as arrows, with the arrowheads indicating the direction of transcription. The individual repeats are displayed as boxes. Sequence similarities of the genes are indicated by gray shading. Compared to NCTC 10130, strain DSM 25591 showed 13 instead of 12 repeats of 66 bp (ORF position, 440934 to 442004) in *vlpG* and 24 repeats of 36 bp (ORF position, 442427 to 443614) instead of 6 complete and 1 partial repeat in *vlpB*. (b) Multisequence alignment of the 23S rRNA regions involved in macrolide resistance of *M. hyopneumoniae* J ([NC_007295.1](#)), *M. hyorhinis* DSM 25591 ([CP064323](#)), *M. hyorhinis* NCTC 10130 ([LS991950.1](#)), and *M. hyorhinis* ATCC 17981 ([NZ_KB911485.1](#)) compared to that of *E. coli* K-12 MG1655 ([NC_000913.3](#)). The nucleotide transition presumably conferring resistance to 14-membered macrolides is marked in bold and underlined.

differed from those in the *M. hyorhinis* NCTC 10130 sequence and were not included in the available sequence of *M. hyorhinis* ATCC 17981 (Fig. 1a). The number of repeats is known to potentially differ from generation to generation (11–13).

Analysis of the 23S rRNA of *M. hyorhinis* DSM 25591 identified a G2057A (*Escherichia coli* numbering) transition. In *Mycoplasma (Mesomycoplasma) hyopneumoniae*, transitions at nucleotide G2057 confer resistance to 14-membered macrolides (14). Therefore, the intrinsic resistance to erythromycin reported in the literature in *M. hyorhinis* might be explained by the G2057A transition, which is also seen in other available sequences deposited in the NCBI database (Fig. 1b) (14–17).

Data availability. The complete genome sequence of *Mycoplasma hyorhinis* DSM 25591 has been deposited in GenBank under the accession number [CP064323](https://ncbi.nlm.nih.gov/nucl/CP064323). The version described in this paper is the first version, [CP064323.1](https://ncbi.nlm.nih.gov/nucl/CP064323.1), annotated using the NCBI Prokaryotic Genome Annotation Pipeline (PGAP). The associated BioProject accession number is [PRJNA674728](https://ncbi.nlm.nih.gov/bioproject/PRJNA674728); the SRA numbers are [SRX9929726](https://ncbi.nlm.nih.gov/sra/SRX9929726) and [SRX9929725](https://ncbi.nlm.nih.gov/sra/SRX9929725).

ACKNOWLEDGMENTS

We thank Petra Krienke (Institute of Microbiology and Epizootics, Freie Universität Berlin, Berlin, Germany) for the Illumina MiSeq sequencing.

This work was supported by the German Federal Ministry of Food and Agriculture (BMEL) and the Federal Office for Agriculture and Food (BLE), grant number FKZ 2818HS015. Funding for T.S. was provided by the German Federal Ministry of Health (BMG) within the ZooSeq MinION-Zoo project, grant number ZMV11-2519NIK704. We acknowledge support from the Open Access Publication Initiative of Freie Universität Berlin.

REFERENCES

- Gupta RS, Sawnani S, Adeolu M, Alnajjar S, Oren A. 2018. Phylogenetic framework for the phylum Tenericutes based on genome sequence data: proposal for the creation of a new order *Mycoplasmoidales* ord. nov., containing two new families *Mycoplasmoidaceae* fam. nov. and *Metamycoplasmataceae* fam. nov. harbouring *Eperythrozoon*, *Ureaplasma* and five novel genera. *Antonie Van Leeuwenhoek* 111:1583–1630. <https://doi.org/10.1007/s10482-018-1047-3>.
- Balish M, Bertaccini A, Blanchard A, Brown D, Browning G, Chalker V, Frey J, Gasparich G, Hoelzle L, Knight T, Jr, Knox C, Kuo C-H, Manso-Silvan L, May M, Pollack JD, Ramirez AS, Spargser J, Taylor-Robinson D, Volokhov D, Zhao Y. 2019. Recommended rejection of the names *Malacoplasma* gen. nov., *Mesomycoplasma* gen. nov., *Metamycoplasma* gen. nov., *Metamycoplasmataceae* fam. nov., *Mycoplasmoidaceae* fam. nov., *Mycoplasmoidales* ord. nov., *Mycoplasmoides* gen. nov., *Mycoplasmopsis* gen. nov. [Gupta, Sawnani, Adeolu, Alnajjar and Oren 2018] and all proposed species comb. nov. placed therein. *Int J Syst Evol Microbiol* 69:3650–3653. <https://doi.org/10.1099/ijsem.0.003632>.
- Gupta RS, Oren A. 2020. Necessity and rationale for the proposed name changes in the classification of *Mollicutes* species. Reply to: 'Recommended rejection of the names *Malacoplasma* gen. nov., *Mesomycoplasma* gen. nov., *Metamycoplasma* gen. nov., *Metamycoplasmataceae* fam. nov., *Mycoplasmoidaceae* fam. nov., *Mycoplasmoidales* ord. nov., *Mycoplasmoides* gen. nov., *Mycoplasmopsis* gen. nov. [Gupta, Sawnani, Adeolu, Alnajjar and Oren 2018] and all proposed species comb. nov. placed therein', by M. Balish et al. (*Int J Syst Evol Microbiol*, 2019;69:3650–3653). *Int J Syst Evol Microbiol* 70:1431–1438. <https://doi.org/10.1099/ijsem.0.003869>.
- Friis NF. 1975. Some recommendations concerning primary isolation of *Mycoplasma suis* pneumoniae and *Mycoplasma flocculare*: a survey. *Nord Vet Med* 27:337–339.
- Leibniz Institut DSMZ-Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH. 2007. DSMZ medium 1078 Friis medium. DSMZ, Braunschweig, Germany. https://www.dsmz.de/microorganisms/medium/pdf/DSMZ_Medium1078.pdf.
- Andrews S. 2010. FastQC: a quality control tool for high throughput sequence data. <https://www.bioinformatics.babraham.ac.uk/projects/fastqc/>.
- Roehr JT, Dieterich C, Reinert K. 2017. Flexbar 3.0—SIMD and multicore parallelization. *Bioinformatics* 33:2941–2942. <https://doi.org/10.1093/bioinformatics/btx330>.
- Nikolenko SI, Korobeynikov AI, Alekseyev MA. 2013. BayesHammer: Bayesian clustering for error correction in single-cell sequencing. *BMC Genomics* 14:S7. <https://doi.org/10.1186/1471-2164-14-S1-S7>.
- Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Unicycler: resolving bacterial genome assemblies from short and long sequencing reads. *PLoS Comput Biol* 13:e1005595. <https://doi.org/10.1371/journal.pcbi.1005595>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Citti C, Watson-McKown R, Droesse M, Wise KS. 2000. Gene families encoding phase- and size-variable surface lipoproteins of *Mycoplasma hyorhinis*. *J Bacteriol* 182:1356–1363. <https://doi.org/10.1128/jb.182.5.1356-1363.2000>.
- Rosengarten R, Wise KS. 1990. Phenotypic switching in mycoplasmas: phase variation of diverse surface lipoproteins. *Science* 247:315–318. <https://doi.org/10.1126/science.1688663>.
- Citti C, Wise KS. 1995. *Mycoplasma hyorhinis* *vlp* gene transcription: critical role in phase variation and expression of surface lipoproteins. *Mol Microbiol* 18:649–660. https://doi.org/10.1111/j.1365-2958.1995.mmi_18040649.x.
- Gautier-Bouchardon AV. 2018. Antimicrobial resistance in *Mycoplasma* spp. *Microbiol Spectr* 6:ARBA-0030-2018. https://doi.org/10.1128/microbiolspec_ARBA-0030-2018.
- Wu CC, Shryock TR, Lin TL, Faderan M, Veenhuizen MF. 2000. Antimicrobial susceptibility of *Mycoplasma hyorhinis*. *Vet Microbiol* 76:25–30. [https://doi.org/10.1016/s0378-1135\(00\)00221-2](https://doi.org/10.1016/s0378-1135(00)00221-2).
- Ter Laak EA, Pijpers A, Noordergraaf JH, Schoevers EC, Verheijden JH. 1991. Comparison of methods for in vitro testing of susceptibility of porcine *Mycoplasma* species to antimicrobial agents. *Antimicrob Agents Chemother* 35:228–233. <https://doi.org/10.1128/aac.35.2.228>.
- Kobayashi H, Nakajima H, Shimizu Y, Eguchi M, Hata E, Yamamoto K. 2005. Macrolides and lincomycin susceptibility of *Mycoplasma hyorhinis* and variable mutation of domain II and V in 23S ribosomal RNA. *J Vet Med Sci* 67:795–800. <https://doi.org/10.1292/jvms.67.795>.