



Bioactive molecules from plants: a prospective approach to combat SARS-CoV-2


Gagan Kumar Panigrahi¹ · Shraban Kumar Sahoo¹ · Annapurna Sahoo¹ · Shibasish Behera¹ · Snigdharani Sahu¹ · Archana Dash¹ · Kunja Bihari Satapathy¹

Received: 18 May 2021 / Accepted: 8 July 2021
© Institute of Korean Medicine, Kyung Hee University 2021

Abstract

The emergence of Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) or 2019 Novel Coronavirus (2019-nCoV) has put the entire globe into unrest, primarily due to unavailability of specific drug against the viral proteins. In the last two decades the world has withstood many contagious disease crashes. SARS-CoV-2 has put the world and the mankind in danger. It is spreading unstopably all over the world. The virus is evolving and thus the pathogenicity of SARS-CoV-2 strains has been different and making it difficult to develop a broad-spectrum anti-viral molecule that would be effective against all the SARS-CoV-2 variants. This imperative situation demands development of molecules for effective treatment against SARS-CoV-2. The phytomolecules or the bioactive molecules of plants could be a great alternative to combat SARS-CoV-2. The bioactive molecules with their antiviral properties and the secondary metabolites may effectively deactivate the functioning of viral proteins. The structural configuration of 2019-nCoV proteins and genomic information are available, thus contributing immensely for fast molecular docking studies and hence, enables screening of numerous accessible phytomolecules. In the current study, we have essentially highlighted common phytomolecules against the known viral proteins and described the mode of action of few plant-derived molecules which have the potential to suppress the activity of the viral proteins.

Graphic abstract



Keywords SARS-CoV-2 · Pathogenicity · Antiviral · Phytomolecules · Secondary metabolites · Molecular docking

Gagan Kumar Panigrahi, Shraban Kumar Sahoo, and Annapurna Sahoo have contributed equally.

Extended author information available on the last page of the article

Introduction

The outburst situation caused due to the Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) represents a serious public health crisis across the globe. Since, December 2019, the whole world is suffering from the crisis of Corona virus. The virus is supposed to be originated from bats and latter on transmitted to humans. The outbreak of this human pathogen emerged in the city of Wuhan in China, and resulted to human disease, termed as COVID-19 (Arti and Bhatnagar 2020; Burki 2020; Chen et al. 2020; Huang et al. 2020). World Health Organization (WHO) declared Public Health Emergency of International Concern (PHEIC) owing to its fast rate of transmission within the humans (Chan et al. 2020; Chen et al. 2020; Li et al. 2020a, b; Sun et al. 2020). Over 14.57 million cases have been detected in 213 countries till now. The virus possesses crown like spikes on its outermost layer, so it was named as coronavirus (Fig. 1). The SARS-CoV-2 belongs to the beta (β), Coronavirus genus, closely related to the previously identified severe acute respiratory syndrome Coronavirus (SARS-CoV), family Coronaviridae, order Nidovirales and the sub family Orthocoronaviridae (Lu et al. 2020; Sher-een et al. 2020; Wu et al. 2020; Zhou et al. 2020). Middle East Respiratory Syndrome (MERS) and severe acute respiratory syndrome (SARS) are commonly caused by viruses belonging to the Coronaviridae. Coronaviridae has two sub families; Torovirinae and Coronavirinae and the members of Coronaviridae family affects the mammals and aves (Zhou et al. 2020). The virus is 65–125 nm in diameter and the genomic content is estimated to be 26–32 kilobases. SARS-CoV-2 is an RNA virus and it contains single-stranded RNA genome. The virus has four subgroups i.e. Alpha (α), Beta (β), Gamma (γ) and Delta (δ). The Alpha (α) and Beta (β) corona virus infect mammals. The Beta (β) corona virus causes respiratory illnesses to humans and the Gamma (γ) and Delta (δ) corona virus affect aves and some selective mammals (Woo et al. 2012). SARS-CoV-2 makes use of

its own proteins to safely harbour in the host cells (Fig. 1). Reportedly, a densely glycosylated spike (S) protein, SARS-CoV-2 main protease (M^{pro}) and RNA-dependent RNA polymerase (RdRp) are defining features paving the path of virus from entry to infection in the host cell (Cui et al. 2019; Lung et al. 2020; Ton et al. 2020; Wrapp et al. 2020). Essentially, the S protein undergoes extensive structural reorganization in order to fuse with the host membranes, thus establishing a physical link between the virus and the host cell (Booth et al. 2003; Li et al. 2020a). Eventually, a stable conformation is established (Wrapp et al. 2020). RdRp catalyzes the replication event, resulting in the synthesis of viral RNA. Remarkably, SARS-CoV and SARS-CoV-2 share similar nucleotide sequences and resulting RNA-dependent RNA polymerase (Liu et al. 2020). The SARS-CoV-2 proteins, metabolites and host cell factors can be targeted to reduce the viral replications within the host cells as for instance; the viral protein M^{pro} facilitates the synthesis of functional viral proteins out of the precursors (Fig. 2). M^{pro} , S and RdRp proteins can be targeted for developing diagnostics, antibodies and vaccines. Human corona virus (HCoV), HCoV OC43, MERS-CoV and SARS-CoV are some of the viruses which causes respiratory problems in human and are being transmitted from one individual to another. This viral infection is transmitted from human to human through the coughed, sneezed droplets or by the contact of infected surface. The solved three dimensional structures of the viral proteins provide an outstanding ground for discovering specific ligands (Liu et al. 2020). In South-east countries like South Korea, China, Japan and India, many traditional medications are used to treat SARS-CoV-2, but the efficiency of the compounds are limiting and molecular mechanisms are also unclear. Plants possess amazing defense competence towards diseases (Panigrahi and Satapathy 2021; Panigrahi et al. 2021). The phytomolecules are also known as the bioactive compounds which have the tendency to modify the cellular physiological processes. Here, we highlight several phytomolecules having the ability to restrict the activity of SARS-CoV-2 and thus future studies may reveal the underlying molecular mechanism(s) and the efficacy of these phytomolecules.





Fig. 1 A typical schematic view of SARS-CoV-2. The RNA virus possesses 4 structural proteins known to be spike protein, membrane protein, nucleocapsid protein and envelope protein

Covid-19 and its transmission

People are worstly affected by Covid-19 in different ways. The most common symptoms of Covid-19 include fever, dry cough, and tiredness. It also shows symptoms like diarrhoea, headache, loss of taste or smell, aches and pain, rash on skin, discoloration of finger on toes etc. The symptoms of Covid-19 are same as the symptoms of corona virus (CoV) which had appeared in 2003 as SARS. It was named as SARS-CoV-2 by WHO on 11th February 2020 and the disease was termed as CoV disease-19 (Covid-19) (Jiang

Fig. 2 Possible potential targets to restrict the activity of SARS-CoV-2. There are number of promising targets which can be exploited to inhibit the viral infection. Metabolites including nucleoside analogues and anti-metabolites can be targeted to prevent viral replication. Likewise, host cell proteins such as ACE-2 (Angiotensin Converting Enzyme-2), Importin α 1- β 1 inhibitors and Signal transducers and activators of transcription 1 (STAT1) can also be targeted. Three categories of proteins; structural, non-structural and accessory can also be targeted to repress the activity of the virus



et al. 2020; Velvan and Meyer 2020; Wang et al. 2020a,b,c). Transmission of SARS-CoV-2 is severe as compared to the SARS-CoV. Severe lung damage pneumonia like symptoms were seen in patient suffering from SARS-CoV-2 (Zhu et al. 2020). The Chinese center for disease control and prevention sent a medical emergency team to assist the health authorities of Hubai province and Wuhan city to inspect about the disease, but later WHO gave confirmation about the outbreak of the disease. According to WHO the disease was not caused by any particular animal in Hunan South China sea food market place. Subsequently the transmission of the disease occurred rapidly, the situation became out of control when the whole world was affected badly. The genomic evidences related to SARS-CoV-2 were published (Ferh et al. 2017; Zheng 2020). All possible ways including vaccines, monoclonal antibodies, oligo nucleotides, interferons and small molecules, drugs were tried to be developed for the control of SARS-CoV-2. WHO recommended to practise social distancing among people so as to deaccelerate the higher rate of infections.

In countries including China, United States of America, Italy, India and many other countries, SARS CoV-2 had spread rapidly (Giovanetti et al. 2020; Paraskevis et al. 2020). Several studies suggested that bat may be the cause of SARS- CoV-19, but it is not confirmed that bat is the stockpile of SARS-CoV-2. CoVs acts as nonfatal pathogen, which cause only common cold in case of humans (Li et al. 2005a,b; Hampton 2005; Banerjee et al. 2019; Giovanetti et al. 2020; Paraskevis et al 2020; Paules et al. 2020).

Covid-19 mainly spreads and enters into human body in 3 different ways; firstly by droplets transmission-by sneezing, coughing; secondly by the patient it transmits through contact—by touching mouth, nose and eyes in contaminated area; thirdly, by aerosol transmission (Guan et al. 2020; Jin et al. 2020). In case of humans, symptoms such as abdominal discomfort and diarrhoea are also prominent. It was found that the Angiotensin-Converting Enzyme 2 gene (ACE2) was highly expressed in Covid-19 patients (Zhang et al. 2020a, b). The most general symptoms of this disease are cold, fever, myalgia or fatigue, common pneumonia and complicated dyspnoea and less common symptoms are headache, diarrhoea, hemoptysis, runny nose and phlegm producing cough. The affected individuals showing mild symptoms usually get recovered within one week whereas individuals showing acute symptoms are prone to damage in alveolar tissues and subsequently leading towards severe respiratory failure. Mainly the old aged patients already suffering from diseases such as cirrhosis, tumour, hypertension, diabetes and coronary heart damage get very severe problems and proven to be highly fatal (Arti et al. 2020; Huang et al. 2020; Li and Clercq 2020; Li et al. 2020a, b). As compared to other countries, the proportion of infection in India was low i.e. 1.9%. In India lockdown started from 25th March 2020, all over the country, after the rapid transmission of Covid-19 and situation was back to normal around the start of 2021. But, unusually India is witnessing the dreadful nature of this pandemic now, where it is believed to be because of a variant strain of SARS-CoV-2. Currently, nation-wide crisis

has again led to lockdown situations and speculations suggest that the rise in Covid infections would reach around the mid of May, 2021.

Conceivable drugs and vaccines against Covid-19

On December 2019, an unknown outburst of viral Pneumonia in Wuhan city made clinicians across the globe to analyze different therapeutics to defy Covid-19. Since definite vaccine for Covid-19 was not discovered, the disease got escalated all over the world and forced the clinicians to use former convenient drugs for the treatment. Covid-19 treatment may involve targeting of Autophagy and endocytic signaling events (Yang et al. 2020). Primarily, virus and host cell interaction is critical and there are several check-points where suitable drugs may be targeted (Figs. 3 and 4). Essentially, SARS-CoV-2 mainly affected the lungs and majority of fatality was due to the collapse of respiratory system. Consequently, patients were first aided with oxygen therapy, extra corporeal membrane oxygenation (ECMO) and invasive mechanical ventilation. Many drugs such as remdesivir, ribavirin, nitazoxanide, penciclovir, chloroquine, nafamostat and favipiravir were tested in counter to this lethal disease on Vero E6 cells and was monitored that chloroquine and remdesivir were the most suitable drugs with less cytotoxic effects (Wang et al. 2020a, b, c). In addition to these drugs, antibiotics such as hydroxychloroquine and azithromycin were able to control the viral growth. Lopinavir, baricitinb, interferon- α and ribavirin may be given to the patients with acute

respiratory system, but addition of lopinavir has side effects like diarrhoea, nausea or liver damage (Hirsch et al. 2013). Use of steroids can check the inflammatory damage, but a heavy dose has several harmful effects (Booth et al. 2003; Griffith et al. 2005; Minneci et al. 2009; Clark and Baillie 2020). Plasma therapy was also under observation for the treatment of Covid-19. This therapy essentially involved in deriving antibodies from B-lymphocytes of the recovered patients. It can help in reducing inflammation and viral growth. Traditional Chinese Medicine (TCM) therapy was another choice for the treatment of Covid-19 as it was a successful therapy curing disease from ancient time. It was highlighted that Lian Hua Qing Wen and Shu Feng Jie Du capsules gave positive response in treatment of SARS-CoV-2 (Lu 2020). The western medicines and combination of western and traditional Chinese medicines were experimented separately for the effectiveness towards the disease and was found that the combination of western and traditional Chinese medicine was more effective. The combined medicine stabilized the body temperature in less time and also lowered the mortality rate. Vaccination was necessary to cease the transmission and develop immunity against Covid-19. Recombinant protein subunits for MERS, DNA plasmids, viral vectors, virus like particles and nanoparticles were used to develop the first vaccine. The mRNA vaccine was first tested in humans on March 16, 2020 in United States of America. The vaccine mimics mRNA 1273 which encodes spike protein of SARS-CoV-2 and is primarily enclosed within nanomaterials. However, the understanding of the SARS-CoV-2-host interaction(s) and the underlying molecular mechanism(s) still remains blurred.




Fig. 3 Possible virus replication check-points to restrict the activity of SARS-CoV-2. Virus interacts with host cell by associating with specific cell surface receptors and thereby establishes stable interaction

followed by various downstream events allowing the virus to proliferate and infect the host system




Fig. 4 Few antiviral drugs having the potential to restrict virus replication in the host cells. Various synthetic antiviral drugs have proven activity of inhibiting viral propagation by strategically interfering in several check points related to host-virus interaction

Bioactive molecules against coronavirus


Plant metabolites

The primary and secondary metabolites have different group of compounds (Fig. 5). Improvised analytical techniques would modulate in generating informations regarding bioactive molecules derived from various natural sources including plants. Thus, sophisticated analytical techniques and choosing right natural source may help in overcoming many diseases, particularly for those having no specific treatment. The major compounds that are categorized as primary metabolites and found in all the living systems include proteins, lipids, carbohydrates, nucleic acid and chlorophyll (Fig. 5). Likewise, secondary metabolites include flavonoids, phenol acid, alkaloids, volatile oils, terpenes, saponins, tannins, and many more (Dias et al. 2012; Azmir et al. 2013; Woolley 2001; Panda et al. 2020). Different species of plants accumulate specific secondary metabolites and also plays an imperative role during survival and adaptability process of the plant primarily by modulating the cellular homeostasis (Panigrahi and Satapathy 2020a, b, c; Sahoo and Satapathy 2020).

Bioactive molecules

The modern science is exploring new natural derivatives present in variety of living organisms and subsequently these natural products display highly efficient therapeutic properties for treating various diseases (Dash et al. 2020). These natural products have diverse application and unique properties which are useful for medicine production processes (Kang et al. 2013). From ancient time, the traditional natural medicines are playing vital role in disease control and ancient medicine practitioners were expert in this field with or without knowledge of bioactive molecules. Now-a-days researchers are focusing more on plant-baed bioactive molecules to overcome dreadful disease as these products have negligible side effects on humans (Kang et al. 2013). The therapeutic capacity of medicinal plants is due to their chemical composition which primarily comprises of vitamins, minerals and bioactive molecules (Singh et al. 2020a). Many scientific studies suggest that the synthetic vitamins and minerals cannot give the benefits as much as natural products. The medicinal plants contain many minerals and vitamins that easily get assimilated by human body. Natural products contain more bioactive molecules as it has complementary action between vitamins, enzymes and

Fig. 5 A variety of plant metabolites involved in regulating plant physiology. The primary and secondary metabolites have different group of compounds. The major compounds that are categorized as primary metabolites and found in all the living systems include proteins, lipids, carbohydrates, nucleic acid and chlorophyll. Likewise, secondary metabolites include flavonoids, phenol acid, alkaloids, volatile oils, terpenes, saponins, tannins, and many more



minerals (Singh et al. 2020b). Synthetic drugs possess more side effects and have more disadvantages towards human body as compared to natural medicines (Singh et al. 2020b). The plant products have phytomolecules containing active ingredients with therapeutic properties (Dias et al. 2012). The secondary metabolites are specific from species to species. Bioactive molecules harbour therapeutic, toxicological and immune stimulation properties and could prove to be an effective alternative against viral diseases (Panigrahi and Satapathy 2020d; Sahoo and Satapathy 2020).

Across the globe, the mankind is facing the deadly effect of Covid-19. Every vaccine is under the trial basis and it has been reported that the drugs that were generally used for the Human Immunodeficiency Virus (HIV) such as lopinavir/ritonavir can be used for treating Covid-19 patients (Wang et al. 2020a, b, c). Some other drugs like pitavastatin, nelfinavir, perampanel and praziquantel can also be used against covid-19 (Xu et al. 2020). Plants are repositories of several types of natural bioactive molecules which might play a critical role in addressing the current pandemic. Essentially the various secondary metabolites which are secreted by the plant cells can be utilized for developing anti-viral drugs using plant biotechnological approaches (Fig. 6). Usually secondary metabolites such as flavonoids, alkaloids, terpenoids and polyphenols possess antiviral properties (Yang et al. 2018; Khaerunnisa et al. 2020; Singh et al. 2020c). Medicinal plants are abundantly rich in phenolic metabolites (Yang et al. 2018). Molecular docking studies revealed that

medicinal plants-derived phytomolecules such as quercetin, curcumin, kaempferol, catechin, naringenin, butein-7-glucoside, apigenin-7-glucoside, demethoxy-curcumin, oboeuropein and epigallocatechin that have the potential for combating against SARS-CoV-2 (Singh et al. 2020c). These bioactive molecules display similar pharmacophore properties as nelfinavir as revealed in in silico analysis. The bioactive phytomolecules released from a range of plant species bears therapeutic properties against SARS-CoV-2. Many phytomolecules such as rutin, diacetyl curcumin, diosmin, (E)-1-(2-Hydroxy-4-methoxyphenyl)-3-[3-[(E)-3-(2-hydroxy-4-methoxyphenyl)-3-oxoprop-1-enyl]phenyl]prop-2-en-1-one, beta'-(4-Methoxy-1,3-phenylene)bis(2'-hydroxy-4',6'-dimethoxyacrylophenone) and apigenin are also effective anti-virals (Adem et al. 2020; Singh et al. 2020c; Table 1). Lupane-type triterpenes, R-cadinol, labdane-type and abietane-type diterpenes, lignoids, curcumin and sesquiterpenes are few other identified bioactive compounds which would also play a vital role in defending the host cells against SARS-CoV-2 (Wen et al. 2007; Yang et al. 2020; Gong et al. 2008; Nguyen et al. 2012). In the last decade, many numbers of traditional Chinese herbs have been identified which were effective against SARS-CoV (Chang et al. 2012). Herbal extracts from *Houttuynia cordata*, Chinese Rhubarb extracts and beta-sitosterol from roots of *Isatis indigotica* that have the potential to impede the enzymatic activity of SARS-CoV (Chang et al. 2012; Singh et al. 2020c). Many more phytomolecules like epigallocatechin




Fig. 6 Developmental routes for plant-based diagnostics and therapeutics against SARS-CoV-2. Plant-based secondary metabolites can be substantially utilized for developing anti-viral drugs using plant biotechnological approaches

gallate, aloe-emodin, quercetin, rhoifolin, hesperetin, sinigrin, 3β -Friedelanol from *Euphorbia neriifolia*, blaucoxanthone from the roots of plant species such as *Calophyllum blancoi*, *Lycoris radiata*, *Pyrrhosia lingua*, *Artemisia annua* and *Lindera aggregata* possess anti-viral activity against SARS-CoV-2 (Shen et al. 2005; Singh et al. 2020c). Molecular docking simulations have revealed that the phytochemicals from *Nilavembu kudineer* have the potential to bind to the host cell viral receptors; ACE2 and thus may mediate the restriction of virus to attach to the host cells (Li et al. 2005a,b; Singh et al. 2020c; Walter et al. 2020).

Mode of action of phytochemicals against viral pathogens

Viral diseases are causing a distressing threat for human beings across the globe. A number of viral diseases are continuously reported which predominantly display severe health related issues and unfortunately unavailability of effective antiviral cure makes these diseases more severe (Kapoor et al. 2017). Essentially, flaviviruses and alphaviruses are the causative agents for several types of viral diseases such as chikungunya, influenza and HIV which are still considered as potential threats. Similarly, the COVID-19 which resulted in a pandemic situation has badly affected a large number of population across the globe. As the rate of genomic mutation is faster in case of coronavirus, it has become difficult to develop effective synthetic drug-based treatment strategies (Irwin et al. 2016). One of the drawbacks of synthetic antiviral drugs is that they display adverse side effects which would affect the human well-being. On the contrary, plant-derived drugs can be considered as a

suitable alternative since herbal treatments exhibit minimal side effects with maximal health benefits (Biswas et al. 2019). For several emerging viral diseases, plant-derived compounds exhibiting antiviral properties have been evaluated and found to be effective and most importantly have been proposed to be applied along with the pre-existing therapies so as to enhance the efficacy (Kapoor et al. 2017; Lillehoj et al. 2018). The antiviral properties of the plant-based compounds are governed by a variety of mechanisms. It has been reported that few phytochemicals have the potential to restrict the entry of viruses primarily by ensuring unavailability of specific sites for attachment of viruses to specific host cells as the phytochemicals can bind to the carbohydrate moiety of the target host cell (Idris et al. 2016). Also, few phytochemicals inhibit viral replication and propagation and thus prevents the infectious action of viruses (Kapoor et al. 2017).

Phytochemicals can be strategically employed against viral infection by targeting any possible viral replication check-points. For example, the phytochemical; epigallocatechin gallate (EGCG) specifically inactivates the viral as well as host enzymes (reverse transcriptase, protease, RNA polymerase) and ultimately restricts the viral growth (Lipson et al. 2017). Another way the flavonoids restrict the viral RNA synthesis is by preventing protein phosphorylation resulting in inhibition of several viruses such as herpes simplex virus, influenza virus and HIV (Kumar and Pandey 2013). Several phytochemicals including fiesitin, baicalein and quercetagenin have the potential to inhibit the chikungunya virus essentially by restricting the viral replication events (Lani et al. 2016). Similarly luteolin also possess antiviral replication activity during the post entry process as this phytochemical cannot inhibit the viral replication

Table 1 Potent phytomolecules against coronavirus

S.I. No	Name of plants	Phytomolecules	Mechanism of Action	References
1	<i>Allium sativum</i>	Allicin	Inhibit SARS-3CLpro activity	Shang et al. (2019)
2	<i>Amaranthus tricolor</i>	Amaranthin	Inhibit 3CLpro interacting through Thr26, Glu166, Cys145, His41, Thr24 H-bonding	ul Qamar et al. (2020)
3	<i>Astragalus membranaceus</i>	Flavonoids, lectin, isoflavones	Inhibit SARS-3CLpro activity	Jo et al. (2020)
4	<i>Capsicum annum</i>	Quercetin, Luteolin-7-glucoside	Inhibit the cellular entry of SARS CoV	Miean et al. (2001), Chen et al. (2008)
5	<i>Citrus sinensis</i>	Naringenin	Potential Inhibitor of COVID-19 Main Protease	Salehi et al. (2019), Khaerunnisa et al. (2020)
6	<i>Dioscorea rhizoma</i>	Saponins	Activates macrophages and inflammatory responses	Wen et al. (2011), Cheng et al. (2016)
7	<i>Dryopteris crassirhizoma</i>	Sutchuenoside A	Play significant role in regulation of inflammatory process	Cheng et al. (2016)
8	<i>Ephedra sinica</i>	(3R)-3-O- β -D-glucopyranosyl-3-phenyl propanoic acid	Inhibits viral replication and 3CLpro	Zhang et al. (2020a, b)
9	<i>Flos lonicerae</i>	Chlorogenic acid	Inhibits viral replication	Chen et al. (2004), Zhang et al. (2020a, b)
10	<i>Fraxinus sieboldiana</i>	Calceolarioside B	Inhibit 3CLpro interacting through His41, Gly143, Cys145, Glu166, Thr24, Thr25 H-bonding	ul Qamar et al. (2020)
11	<i>Galla chinensis</i>	Luteolin and tetra-O-gallol- β -D glucose gall, tannin	Avidly binds with surface spike protein of SARS-CoV	Yi et al. (2004)
12	<i>Glycyrrhizae uralensis</i>	Glycyrrhizin	Inhibit 3CLpro	Chen et al. (2004)
13	<i>Glycyrrhiza radix</i>	β -sistosterol, Liquiritin	Inhibit viral adsorption and penetration	Cinatl et al. (2003)
14	<i>Glycyrrhiza uralensis</i>	Flavonoids, glycyrrhetic acid and Licoleafol	Inhibit SARS-3CLpro activity	Jo et al. (2020), ul Qamar et al. (2020)
15	<i>Houttuynia cordata</i>	Flavonoids, polysaccharides and Quercetin	Inhibit the viral SARS-3CLpro activity and Block viral RNAdependent RNA polymerase activity (RdRp) Immunomodulation	Lin et al. (2005), Lau et al. (2008)
16	<i>Hyptis atrorubens</i>	Methyl rosmarinat	Inhibit 3CLpro interacting through Cys145, His41, Thr24, Thr25, Thr26, H-bonding	ul Qamar et al. (2020)
17	<i>Isatis indigotica</i>	5-hydroxyoxyindole, isaindigo- tone, indole-3-carboxaldehyde	Inhibit the cleavage activity of SARS-3CLpro enzyme	Lin et al. (2005)
18	<i>Lonicera japonica</i>	Iridoids	Inhibits corona virus replication at non-toxic concentration	Xiong et al. (2013)
19	<i>Lycium chinense</i>	Polyphenols, flavonoids, carotenoids	Inhibit the viral SAES-3CL pro activity	Miean et al. (2001)
20	<i>Lycoris radiata</i>	Lycorine	Posses anticorona virus	Chen et al. (2004)
21	<i>Myrica cerifera</i>	Myricitrin	Inhibit 3CLpro interacting through Gly143, Cys145, His41, Thr24, Thr25, Thr26 H-bonding	ul Qamar et al. (2020)
22	<i>Nigella sativa</i>	Thymoquinones, sterols and saponins	Increased production of interferons	Yimer et al. (2019)
23	<i>Olea europaea</i>	Oleuropein	Exhibits high antioxidant activity	Nicoli et al. (2019)
24	<i>Phaseolus vulgaris</i>	3,5,7,3',4',5'hexahydroxy flavanone-3-Obeta-Dglucopyranoside	Inhibit 3CLpro interacting through His41, Cys145, Thr24, Thr25, Thr26, Gln189 H-bonding	ul Qamar et al. (2020)

Table 1 (continued)

S.I. No	Name of plants	Phytomolecules	Mechanism of Action	References
25	<i>Phragmites australis</i>	lectins	Inhibits corona virus by interfering with two targets in the viral replication cycle	Keyaerts et al. (2007)
26	<i>Phyllanthus emblica</i>	(2S)Eriodictyol 7O-(6''-Ogalloyl)-beta D glucopyranoside	Inhibit 3CLpro interacting through Thr24, Thr25, Gly143, Met49, Cys145, His41, Thr26 H-bonding	ul Qamar et al. (2020)
27	<i>Psoralea argyrea</i>	5,7,3',4'Tetrahydroxy2'-(3,3dimethylallyl) isoflavones	Inhibit 3CLpro interacting through His41, Cys145, Thr24, Thr25, Thr26, H-bonding	ul Qamar et al. (2020)
28	<i>Rheum palmatum</i>	Emodin, Rhein	Inhibit interaction of SARS-CoV spike protein	Ho et al. (2007), Cheng et al. (2016)
29	<i>Saposhnikovia radix</i>	Cimifugin	Inhibitory effect against corona virus	Zheng et al. (2011)
30	<i>Scutellaria baicalensis</i>	Kaempferol	Prevent the early stage of HCoV-22E9 infection, including viral attachment and penetration	Deng et al. (2012)
31	<i>Scutellariae radix</i>	Baicalin	Antiviral activity	Islam et al. (2012)
32	<i>Spinacia oleracea</i>	Kaempferol	Prevent the early stage of infection, including viral attachment and penetration	Dabeek et al. (2019)
33	<i>Tonina sinensis</i>	Gallic acid, Quercetin	Inhibit the cellular entry of SARSCoV	Chen et al. (2008)
34	<i>Veronica linariifolia</i>	Iridoid	Binds with surface spike protein of corona virus	Yi et al. (2004)

during the entry of virus into the host cell (Fan et al. 2016). It is also evident from recent studies that the pinocembrin specifically inhibits Zika virus replication during late early stages (Lee et al. 2019). Also, genistein effectively inhibits the post-replication events mediated by the entry of herpes B virus (Lee et al. 2019). Interestingly, combinatorial effect of genistein and synthetic drugs such as ganciclovir and acyclovir has been found to be more effective (LeCher et al. 2019). Similarly, around 43 alkaloids have been reported which are effective against influenza virus. Majorly, the alkaloids are believed to induce interferons and macrophages which enhances the immunogenic response against the virus (Moradi et al. 2018). Likewise, alkaloids also play an important role in diminishing the effect of influenza virus by inhibiting the viral replication or by preventing the synthesis of viral proteins (Moradi et al. 2018). *Commelina communis* contains homonojirimycin (HNJ), an alkaloid which strongly inhibits the infectious activity of influenza virus (Zhang et al. 2013). Alkaloid such as loliolide extracted from *Phyllanthus urinaria* impairs the entry of Hepatitis C virus (HCV) into the host cell (Chung et al. 2016). Also plant extract from *Aglaiia* sp. possess antiviral activity owing to the presence of terpenes which can act against HIV-1 primarily by restricting the viral proliferation because of phytochemical-generated cytotoxic environment. In similar fashion, the transition phase between S and G2/M of the cell

cycle is inhibited by the 3, 4-secodammarane triterpenoid, which ultimately prohibits the viral dissemination (Esimone et al. 2010). *Marrubium vulgare* contains terpenes which are effective against HSV-1 (Fayyad et al. 2014). Honokiol, one of the lignin extracted from *Magnolia* tree are mostly used in Chinese medications (Fang et al. 2015). Honokiol is effective against dengue virus as it can stall the synthesis of dsRNA and replication event. Similarly, 3-hydroxy caruilignan C (3-HCL-C), a lignin prevents the viral transcriptional and translational phases (Wu et al. 2012). One of the Coumarin (C4) is known to suppress the IHNV infection, essentially by promoting cellular damage and preventing apoptosis and thus can be one of the potential alternative measures for generating anti-IHNV drug/vaccine (Hu et al. 2019). A detailed study on the underlying molecular mechanism(s) which mediates silencing of viral activities inside the host cell is yet to be fully revealed.

Conclusion and future prospective

Traditional plants have been widely explored for the treatment of various illness including viral infections. The major advantage of phytochemicals is that they exhibit minimal toxicity and side effects. Now-a-days researchers are keen to reveal more plant-based molecules having tremendous

potential against various diseases. Thousands of plant-derived secondary metabolites have been used as antiviral, antioxidant and antibacterial chemicals as revealed in several *in silico* and *in vitro* studies. Interestingly, phytochemicals along with other synthetic compounds including drugs can also be treated as an effective approach against viral infections. But, at the same time detailed molecular mechanism including the dosage potency of phytochemicals also need to be elucidated. Since most of the prevailing therapies against viral infections are synthetic drug-based which display adverse side effects, it is most important to approach for developing strategies which would be cost effective and less harmful for the human beings. In this context, phytochemicals can be considered as a suitable alternative.

The year 2020 and till now it has been a disastrous year for the world population due to sudden outbreak of

Covid-19. Though the vaccines are given to the patients, but the recovery and disease infection rate is still at the peak, so the new drugs or vaccines with high efficiency is yet to be discovered. WHO strongly advocates in maintaining several preventive measures such as washing hands frequently, maintaining social distance, by avoiding crowded places and wearing mask frequently. Different lockdown plans are made by the government authorities to protect the public from this pandemic. Plant-derived bioactive molecules may play a significant role in developing traditional medications which would be a boon for the entire mankind. Many plants contain the natural derivative compounds having the antiviral properties, consequently effective drugs can be formulated and would be a stepping stone for combating against new strains of virus (Fig. 7). Phytomolecules have tremendous potential to act as anti-viral or anti-pathogen agents. Now-a-days,




Fig. 7 Strategy to restrict the synthesis of SARS-CoV-2 proteins inside the host cell. Plant-derived molecules can be effectively recruited in the host cell specifically to prevent viral proliferation

the brutal spread of the coronavirus is a wham to the mankind. Combined application of phytochemicals and FDA-approved synthetic drugs can be proved to be highly effective and thus related studies could be beneficial for a long-term preventive measure against viral pathogens. Additionally, repurposing of known phytochemicals which are effective against virus can be practised. Furthermore, nanotechnological interventions such as encapsulation of antiviral phytochemicals in nanoparticles can enhance the stability and delivery within the host cells. In addition, exploring biodiversity rich regions can also be advantageous in isolating antiviral phytochemicals. It is absolutely important to explore the bioactive molecules-based therapeutic measures and get ourselves prepared for future encounter with virus.

Acknowledgements Authors thank the administration and management of Centurion University of Technology and Management, Odisha, India for their heartfelt support. We apologize to all colleagues whose work could not be discussed owing to space limitations.

Author contribution All the authors have substantial contribution for the preparation of the manuscript. KBS and GKP conceived the idea. Data curation and writing: GKP, SKS, AS, SB, SS, AM and KBS. Review and editing: KBS, GKP, SKS and AS. All the authors have read and approved the final manuscript before submission.

Funding The present study was supported by the Centurion University of Technology and Management, Odisha, India. The financial support was provided to Gagan Kumar Panigrahi (GKP) and Shraban Kumar Sahoo (SKS) (Reference: CSREM TRUST BHUBANESWAR:UTRP UNBH21195785007).

Declarations

Ethical statement This article does not contain any studies involving animals performed by any of the authors. This article does not contain any studies involving human participants performed by any of the authors.

Conflict of interest Gagan Kumar Panigrahi has no conflict of interest. Shraban Kumar Sahoo has no conflict of interest. Annapurna Sahoo has no conflict of interest. Shibashish Behera has no conflict of interest. Snigdha Rani Sahu has no conflict of interest. Archana Dash has no conflict of interest. Kunja Bihari Satapathy has no conflict of interest.

References

Adem S, Eyupoglu V, Sarfraz I, Rasul A, Ali M (2020) Identification of potent COVID-19 main protease (Mpro) inhibitors from natural polyphenols: an *in silico* strategy unveils a hope against CORONA. Preprint 2020030333. <https://doi.org/10.20944/preprints202003.0333.v1>

Arti MK, Bhatnagar K (2020) Modeling and predictions for Covid 19 Spread in India. Preprint <https://doi.org/10.13140/RG.2.2.11427.81444>

Azmir J, Zaidul ISM, Rahman MM, Sharif KM, Mohamed A, Sahena F, Jahurul MHA, Ghafoor K, Norulaini NAN, Omar AKM

(2013) Techniques for extraction of bioactive compounds from plant materials: a review. *J Food Eng* 117(4):426–436

Banerjee A, Kulcsar K, Misra V, Frieman M, Mossman K (2019) Bats and Coronaviruses. *Viruses* 11(1):E41. <https://doi.org/10.3390/v11010041>

Biswas D, Nandy S, Mukherjee A, Pandey DK, Dey A (2019) *Moringa oleifera* Lam and derived phytochemicals as promising antiviral agents: a review. *S Afr J Bot*. <https://doi.org/10.1016/j.sajb.2019.07.049>

Booth CM, Matukas LM, Tomlinson GA, Rachlis AR, Rose DB, Dwosh HA (2003) Clinical features and short term outcomes of 144 patients with SARS in the greater Toronto area. *J Am Med Assoc* 289:2801–2809

Burki TK (2020) Coronavirus in China. *Lancet Respir Med* 8(3):238

Chan JFW, Yuan S, Kok KH et al (2020) A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster. *Lancet* 395:514–523

Chang FR, Yen CT, Ei-Shazly M, Lin WH, Yen MH, Lin KH, Wu YC (2012) Anti-human coronavirus (anti-HCoV) triterpenoids from the leaves of *Euphorbia nerifolia*. *Nat Prod Commun* 7(11)

Chen F, Chan KH, Jiang Y, Kao RYT, Lu HT, Fan KW, Cheng VCC, Tsui WHW, Hung IFN, Lee TSW, Guan Y (2004) In vitro susceptibility of 10 clinical isolates of SARS coronavirus to selected antiviral compounds. *J Clin Virol* 31(1):69–75

Chen CJ, Michaelis M, Hsu HK, Tsai CC, Yang KD, Wu YC, Cinatl J Jr, Doerr HW (2008) *Toona sinensis* Roem tender leaf extract inhibits SARS coronavirus replication. *J Ethnopharmacol* 120(1):108–111

Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, Qiu Y, Wang J, Liu Y, Wei Y, Yu T (2020) Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet* 395(10223):507–513

Cheng C, Giri SS, Jun WJ, Kim HJ et al (2016) Immunomodulatory effects of a bioactive compound isolated from *Dryopteris crassirhizoma* on the Grass Carp *Ctenopharyngodon idella*. *J Immunol Res* 3068913:1–12

Chung CY, Liu CH, Burnouf T, Wang GH, Chang SP, Jassey A, Tai CJ, Tai CJ, Huang CJ, Richardson CD, Yen MH (2016) Activity-based and fraction-guided analysis of *Phyllanthus urinaria* identifies loliolide as a potent inhibitor of hepatitis C virus entry. *Antiviral Res* 130:58–68

Cinatl J, Morgenstern B, Bauer G, Chandra P, Rabenau H, Doerr HW (2003) Glycyrrhizin, an active component of liquorice roots, and replication of SARS-associated coronavirus. *Lancet* 361(9374):2045–2046

Clark DRJEM, Baillie JK (2020) Clinical evidence does not support corticosteroid treatment for 2019-nCoV lungs injury. *Lancet* 6736(20):30317–30322

Cui J, Li F, Shi ZL (2019) Origin and evolution of pathogenic coronaviruses. *Nat Rev Microbiol* 17(3):181–192

Dabeek WM, Marra MV (2019) Dietary Quercetin and Kaempferol: bioavailability and potential cardiovascular-related bioactivity in humans. *Nutrients* 11(10):1–15

Dash S, Panda MK, Singh MC, Jit BP, Singh YD, Patra JK (2020) Bioactive molecules from *alpinia* genus: a comprehensive review. *Curr Pharma Biotechnol* 21:1

Deng YF, Aluko RE, Jin Q, Zhang Y, Yuan LJ (2012) Inhibitory activities of baicalin against renin and angiotensin-converting enzyme. *Pharmaceut Biol* 50(4):401–406

Dias DA, Urban S, Roessner U (2012) A historical overview of natural products in drug discovery. *Metabolites* 2(2):303–336

Esimone CO, Eck G, Nworu CS, Hoffmann D, Überla K, Proksch P (2010) Dammarenolic acid, a secodammarane triterpenoid from

- Aglaia* sp. shows potent anti-retroviral activity in vitro. *Phytochemistry* 17(7):540–547
- Fan W, Qian S, Qian P, Li X (2016) Antiviral activity of luteolin against Japanese encephalitis virus. *Virus Res* 220:112–116
- Fang CY, Chen SJ, Wu HN, Ping YH, Lin CY, Shiuan D, Chen CL, Lee YR, Huang KJ (2015) Honokiol, a lignan biphenol derived from the magnolia tree, inhibits dengue virus type 2 infection. *Viruses* 7(9):4894–4910
- Fayyad AG, Ibrahim N, Yaakob WA (2014) Phytochemical screening and antiviral activity of *Marrubium vulgare*. *Malays J Microbiol* 10(2):106–111
- Fehr AR, Channappanavar R, Perlman S (2017) Middle East respiratory syndrome: emergence of a pathogenic human coronavirus. *Annu Rev Med* 68:387–399
- Giovanetti M, Benvenuto D, Angeletti S, Ciccozzi M (2020) The first two cases of 2019-nCoV in Italy: Where they come from? *J Med Virol*. <https://doi.org/10.1002/jmv.25699>
- Gong SJ, Su XJ, Wu HP, Li J, Qin YJ, Xu Q, Luo WS (2008) A study on anti-SARS-CoV 3CL protein of flavonoids from *Litchi Chinensis*. *Chin Pharma Bull* 24(5):699–700
- Griffith JF, Antonio GE, Kumta SM, Hui DSC, Wong JKT, Joynt GM (2005) Osteonecrosis of hip and knee in patients with severe acute respiratory syndrome treatment with steroids. *Radiol* 235:168–175
- Guan W-J, Ni Z-Y, Hu Y, Liang W-H, Ou C-Q, He J-X et al (2020) Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med*. <https://doi.org/10.1056/NEJMoa2002032>
- Hampton T (2005) Bats may be SARS reservoir. *JAMA* 294(18):2291
- Hirsch HH, Martino R, Ward KN, Boeckh M, Einsele H, Ljungman P (2013) Guidelines for diagnosis and treatment of human respiratory syncytial virus parainfluenza virus, metapneumovirus, rhinovirus and coronavirus. *Clin Infect Dis* 56:258–266
- Ho TY, Wu SL, Chen JC, Li CC, Hsiang CY (2007) Emodin blocks the SARS coronavirus spike protein and angiotensin-converting enzyme 2 interaction. *Antiviral Res* 74(2):92–101
- Hu Y, Chen W, Shen Y, Zhu B, Wang GX (2019) Synthesis and antiviral activity of Coumarin derivatives against infectious hematopoietic necrosis virus. *Bioorg Med Chem Lett* 29(14):1749–1755
- Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y et al (2020) Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 395:497–506. [https://doi.org/10.1016/S0140-6736\(20\)30183-5](https://doi.org/10.1016/S0140-6736(20)30183-5)
- Idris F, Muharram SH, Diah S (2016) Glycosylation of dengue virus glycoproteins and their interactions with carbohydrate receptors: possible targets for antiviral therapy. *Arch Virol* 161(7):1751–1760
- Irwin KK, Renzette N, Kowalik TF, Jensen JD (2016) Antiviral drug resistance as an adaptive process. *Virus Evol* 2(1):vew14
- Islam MN, Chung HJ, Kim DH, Yoo HH (2012) A simple isocratic HPLC method for the simultaneous determination of bioactive components of *Scutellariae radix* extract. *Nat Prod Res* 26(21):1957–1962
- Jiang S, Xia S, Ying T, Lu L (2020) A novel coronavirus (2019-nCoV) causing pneumonia-associated respiratory syndrome. *Cell Mol Immunol*. <https://doi.org/10.1038/s41423-020-0372-4>
- Jin YH, Cai L, Cheng ZS, Cheng H, Deng T, Fan YP, Fang C, Huang D, Huang LQ, Huang Q, Han Y (2020) A rapid advice guideline for the diagnosis and treatment of 2019 novel coronavirus (2019-nCoV) infected pneumonia (standard version). *Military Med Res* 7(1):4
- Jo S, Kim S, Shin DJ, Kim MS (2020) Inhibition of SARS-CoV 3CL protease by flavonoids. *J Enzyme Inhib Med Chem* 35(1):145–151
- Kang YB, Mallikarjuna PR, Fabian DA, Gorajana A, Lim CL, Tan EL (2013) Bioactive molecules: current trends in discovery, synthesis, delivery and testing. *IeJSME* 7(Suppl 1):S32–S46
- Kapoor R, Sharma B, Kanwar SS (2017) Antiviral phytochemicals: an overview. *Biochem Physiol* 6(2):7
- Keyaerts E, Vijgen L, Pannecouque C, Van Damme E, Peumans W, Egberink H, Balzarini J, Van Ranst M (2007) Plant lectins are potent inhibitors of coronaviruses by interfering with two targets in the viral replication cycle. *Antiviral Res* 75(3):179–187
- Khaerunnisa S, Kurniawan H, Awaluddin R, Suhartati S, Soetjipto S (2020) Potential inhibitor of COVID-19 main protease (M pro) from several medicinal plant compounds by molecular docking study. Preprint <https://doi.org/10.20944/preprints202003.0226>
- Kumar S, Pandey AK (2013) Chemistry and biological activities of flavonoids: an overview. *Sci World J* 2013:162750
- Lani R, Hassandarvish P, Shu MH, Phoon WH, Chu JJ, Higgs S, Vanlandingham D, Bakar SA, Zandi K (2016) Antiviral activity of selected flavonoids against Chikungunya virus. *Antiviral Res* 133:50–61
- Lau KM, Lee KM, Koon CM, Cheung CSF, Lau CP, Ho HM, Lee MYH, Au SWN, Cheng CHK, Bik-San Lau C, Tsui SKW (2008) Immunomodulatory and anti-SARS activities of *Houttuynia cordata*. *J Ethnopharmacol* 118(1):79–85
- LeCher JC, Diep N, Krug PW, Hilliard JK (2019) Genistein has antiviral activity against herpes B virus and acts synergistically with antiviral treatments to reduce effective dose. *Viruses* 11(6):499
- Lee LJ, Loe MW, Lee RC, Chu JJ (2019) Antiviral activity of pinocebrin against Zika virus replication. *Antiviral Res* 167:13–24
- Li G, Clercq E (2020) Therapeutic options for the 2019 novel coronavirus (2019-nCoV). *Nat Rev Drug Discov*. <https://doi.org/10.1038/d41573-020-00016-0>
- Li SY, Chen C, Zhang HQ, Guo HY, Wang H, Wang L, Zhang X, Hua SN, Yu J, Xiao PG, Li RS (2005a) Identification of natural compounds with antiviral activities against SARS-associated coronavirus. *Antiviral Res* 67(1):18–23
- Li W, Shi Z, Yu M, Ren W, Smith C, Epstein JH et al (2005b) Bats are natural reservoirs of SARS-like coronaviruses. *Science* 310(5748):676–679
- Li Q, Guan X, Wu P, et al (2020a) Early transmission dynamics in Wuhan, China, of Novel Coronavirus–Infected Pneumonia New England. *J Med*, 1–9
- Li T, Wei C, Li W, Hongwei F, Shi J (2020b) Beijing union medical college hospital on "pneumonia of novel coronavirus infection" diagnosis and treatment proposal. *Med J Peking Union Med Coll Hosp* <https://kns.cnki.net/kcms/detail/11.5882.r.20200130.1430.002.html>
- Lillehoj H, Liu Y, Calsamiglia S, Fernandez-Miyakawa ME, Chi F, Cravens RL, Oh S, Gay CG (2018) Phytochemicals as antibiotic alternatives to promote growth and enhance host health. *Vet Res* 49(1):76
- Lin CW, Tsai FJ, Tsai CH, Lai CC, Wan L, Ho TY, Hsieh CC, Chao PDL (2005) Anti-SARS coronavirus 3C-like protease effects of *Isatis indigotica* root and plant-derived phenolic compounds. *Antiviral Res* 68(1):36–42
- Lipson SM, Karalis G, Karthikeyan L, Ozen FS, Gordon RE, Ponnala S, Bao J, Samarrai W, Wolfe E (2017) Mechanism of antirotavirus synergistic activity by epigallocatechin gallate and a proanthocyanidin-containing nutraceutical. *Food Environ Virol* 9(4):434–443
- Liu W, Morse JS, Lalonde T, Xu S (2020) Learning from the Past: Possible urgent prevention and treatment options for severe acute respiratory infections caused by 2019-nCoV. *Chem Bio Chem*, 730–738
- Lu, H (2020) Drug treatment options for the 2019-new coronavirus (2019-nCoV). *Biosci Trends*, 10–12
- Lu R, Zhao X, Li J et al (2020) Genomic characterisation and epidemiology of 2019 novel coronavirus: implications for virus origins and receptor binding. *Lancet* 395:565–574

- Lung JH, Yu-Shih Lin Yao-Hsu Yang et al (2020) The potential chemical structure of anti-SARS-CoV-2 RNA-dependent RNA polymerase. *J Med Virol*, 10–19
- Miean KH, Mohamed S (2001) Flavonoid (myricetin, quercetin, kaempferol, luteolin, and apigenin) content of edible tropical plants. *J Agric Food Chem* 49(6):3106–3112
- Minneci PC, Deans KJ, Eichacker PQ, Natanson C (2009) The effects of steroids during sepsis depend on dose and severity of illness: an updated meta-analysis. *Clin Microbiol Infect* 15(4):308–318. <https://doi.org/10.1111/j.1469-0691.2009.02752.x>
- Moradi MT, Karimi A, Lorigooini Z (2018) Alkaloids as the natural anti-influenza virus agents: a systematic review. *Toxin Rev* 37(1):11–18
- Nguyen TTH, Woo HJ, Kang HK, Kim YM, Kim DW, Ahn SA, Xia Y, Kim D (2012) Flavonoid-mediated inhibition of SARS coronavirus 3C-like protease expressed in *Pichia pastoris*. *Biotech Lett* 34(5):831–838
- Nicoli F, Negro C, Vergine M, Aprile A, Nutricati E, Sabella E, Miceli A, Luvisi A, De Bellis L (2019) Evaluation of phytochemical and antioxidant properties of 15 Italian *Olea europaea* L. *Culti Leaves Mol* 24(10):1–10
- Panda S, Sahoo S, Tripathy K, Singh YD, Sarma MK, Babu PJ, Singh MC (2020) Essential oils and their pharmacotherapeutics applications in human diseases. *Adv Tradit Med*, 1–15
- Panigrahi GK, Satapathy KB (2020a) Sacrificed surveillance process favours plant defense: a review. *Plant Archives* 20(1):2551–2559
- Panigrahi GK, Satapathy KB (2020b) *Arabidopsis* DCP5, a decapping complex protein interacts with Ubiquitin-5 in the processing bodies. *Plant Archives* 20(1):2243–2247
- Panigrahi GK, Satapathy KB (2020c) Formation of *Arabidopsis* Poly(A)-Specific Ribonuclease associated processing bodies in response to pathogenic infection. *Plant Archives* 20(2):4907–4912
- Panigrahi GK, Satapathy KB (2020d) Potent Phytomolecules against the RNA dependent RNA polymerase of the SARS-COV-2. *Biosci Biotech Res Comm* 13(12):127–130
- Panigrahi GK, Satapathy KB (2021) *Pseudomonas syringae* pv. *syringae* Infection orchestrates the fate of the *Arabidopsis* J domain containing cochaperone and decapping protein factor 5. *Physiol Mol Plant Pathol* 101598:1–9
- Panigrahi GK, Sahoo A, Satapathy KB (2021) Insights to plant immunity: defense signaling to epigenetics. *Physiol Mol Plant Pathol* 101568:1–7
- Paraskevis D, Kostaki EG, Magiorkinis G, Panayiotakopoulos G, Sourvinos G, Tsiodras S (2020) Full-genome evolutionary analysis of the novel corona virus (2019-nCoV) rejects the hypothesis of emergence as a result of a recent recombination event. *Infect Genet Evol* 79:104212
- Paules CI, Marston HD, Fauci AS (2020) Coronavirus infections more than just the common cold. *JAMA*. <https://doi.org/10.1001/jama.2020.0757>
- Sahoo AS, Satapathy KB (2020) *In Silico* molecular docking-based screening reveals phytomolecules against SARS-COV-2 main protease. *Biosci Biotech Res Comm* 13(12):131–134
- Salehi B, Fokou PVT, Sharifi-Rad M, Zucca P, Pezzani R, Martins N, Sharifi-Rad J (2019) The therapeutic potential of naringenin: a review of clinical trials. *Pharmaceuticals* 12(1):11
- Shang A, Cao SY, Xu XY, Gan RY, Tang GY, Corke H, Mavumengwana V, Li HB (2019) Bioactive compounds and biological functions of garlic (*Allium sativum* L.). *Foods* 8(7):246
- Shen YC, Wang LT, Khalil AT, Chiang LC, Cheng PW (2005) Bioactive pyranoxanthones from the roots of *Calophyllum blancoi*. *Chem Pharm Bull* 53(2):244–247
- Shereen MA, Khan S, Kazmi A, Bashir N, Siddique R (2020) COVID-19 infection: origin, transmission, and characteristics of human coronaviruses. *J Adv Res* 24:91–98
- Singh AK, Singh A, Shaikh A, Singh R, Misra A (2020a) Chloroquine and hydroxychloroquine in the treatment of covid-19 with or without diabetes: a systematic search and a narrative review with special reference to India and other developing countries. *Diabet Metab Syndrome Clin Res Rev* 14:241–246
- Singh YD, Jena B, Ningthoujam R et al (2020b) Potential bioactive molecules from natural products to combat against coronavirus. *Adv Tradit Med (ADTM)*. <https://doi.org/10.1007/s13596-020-00496-w>
- Singh YD, Panda MK, Satapathy KB (2020c) Ethnomedicine for drug discovery. *Advances in pharmaceutical biotechnology*. Springer, Singapore, pp 15–28
- Sun J, He WT, Wang L, Lai A, Ji X, Zhai X, Li G, Suchard MA, Tian J, Zhou J, Veit M (2020) COVID-19: epidemiology, evolution, and cross-disciplinary perspectives. *Trends Mol Med* 26(5):483–495
- Ton A-T, Gentile F, Hsing M et al. (2020) Rapid Identification of Potential Inhibitors of SARS-CoV-2 Main Protease by Deep Docking of 1.3 Billion Compounds. *Mol Inform*, 1–18
- ul Qamar MT, Alqahtani SM, Alamri MA, Chen LL (2020) Structural basis of SARS-CoV-2 3CLpro and anti-COVID-19 drug discovery from medicinal plants. *J Pharmaceut Anal* 10(4):313–319
- Velavan TP, Meyer CG (2020) The COVID-19 epidemic. *Trop Med Int Health*. <https://doi.org/10.1111/tmi.13383>
- Walter TM, Justinraj CS, Nandini VS (2020) Effect of *Nilavembu kudineer* in the Prevention and Management of COVID-19 by inhibiting ACE2 Receptor. *Siddha Papers* 15(2):1–9
- Wang C, Horby PW, Hayden FG, Gao GF (2020a) A novel coronavirus outbreak of global health concern. *Lancet* 395:470–473
- Wang M, Cao R, Zhang L, Yang X, Liu J, Xu M (2020b) Remdesivir and Chloroquine effectively inhibit the recently emerged coronavirus (2019-nCoV) in vitro. *Cell Res* 30:269–271
- Wang Z, Chen X, Lu Y, Chen F, Zhang W (2020c) Clinical characteristics and therapeutic procedure for four cases with 2019 novel coronavirus pneumonia receiving combined Chinese and Western medicine treatment. *Biosci Trends* 14(1):64–68. <https://doi.org/10.5582/bst.2020.01030>
- Wen CC, Kuo YH, Jan JT, Liang PH, Wang SY, Liu HG, Lee CK, Chang ST, Kuo CJ, Lee SS, Hou CC (2007) Specific plant terpenoids and lignoids possess potent antiviral activities against severe acute respiratory syndrome coronavirus. *J Med Chem* 50(17):4087–4095
- Wen CC, Shyur LF, Jan JT, Liang PH, Kuo CJ, Arulselvan P, Wu JB, Kuo SC, Yang NS (2011) Traditional Chinese medicine herbal extracts of *Cibotium barometz*, *Gentiana scabra*, *Dioscorea batatas*, *Cassia tora*, and *Taxillus chinensis* inhibit SARS-CoV replication. *J Tradit Complement Med* 1(1):41–50
- Woolley JG (2001) Plant alkaloids. De Montfort University Leicester, Nature Publishing Group, London, Encyclopedia of life sciences
- Wrapp D, Wang N, Corbett KS et al (2020) Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. *Science* 1263:1260–1263
- Wu SF, Lin CK, Chuang YS, Chang FR, Tseng CK, Wu YC, Lee JC (2012) Anti-hepatitis C svirus activity of 3-hydroxy caruilignan C from *Swietenia macrophylla* stems. *J Viral Hepat* 19(5):364–370
- Wu F, Zhao S, Yu B et al (2020) A new coronavirus associated with human respiratory disease in China. *Nature* 579:265–269
- Xiong J, Li S, Wang W, Hong Y, Tang K, Luo Q (2013) Screening and identification of the antibacterial bioactive compounds from *Lonicera japonica* leaves. *Food Chem* 138(1):327–333
- Xu Z, Peng C, Shi Y, Zhu Z, Mu K, Wang X, Zhu W (2020) Nelfinavir was predicted to be a potential inhibitor of 2019-nCoV main protease by an integrative approach combining homology modelling, molecular docking and binding free energy calculation. *BioRxiv*. <https://doi.org/10.1101/2020.01.27.921627>

- Yang L, Wen KS, Ruan X, Zhao YX, Wei F, Wang Q (2018) Response of plant secondary metabolites to environmental factors. *Molecules* 23(4):762
- Yang Y, Islam MS, Wang J, Li Y, Chen X (2020) Traditional Chinese medicine in the treatment of patients infected with 2019-new coronavirus (SARS-CoV-2): a review and perspective. *Int J Biol Sci* 16(10):1708
- Yi L, Li Z, Yuan K, Qu X, Chen J, Wang G, Zhang H, Luo H, Zhu L, Jiang P, Chen L (2004) Small molecules blocking the entry of severe acute respiratory syndrome coronavirus into host cells. *J Virol* 78(20):11334–11339
- Yimer EM, Tuem KB, Karim A, Ur-Rehman N, Anwar F (2019) *Nigella sativa* L. (Black Cumin): a promising natural remedy for wide range of illnesses. *Evid Based Complement Alternat Med* 2019:1528635
- Zhang G, Zhang B, Zhang X, Bing F (2013) Homonijirimycin, an alkaloid from dayflower inhibits the growth of influenza a virus in vitro. *Acta Virol* 57(1):85–86
- Zhang DH, Wu KL, Zhang X, Deng SQ, Peng B (2020a) *In silico* screening of Chinese herbal medicines with the potential to directly inhibit 2019 novel coronavirus. *J Integr Med* 18(2):152–158
- Zhang H, Kang Z, Gong H, Xu D, Wang J, Li Z, et al (2020b) The digestive system is a potential route of 2019-nCov infection: a bioinformatics analysis based on single-cell transcriptomes. bioRxiv <https://doi.org/10.1101/2020.01.30.927806>
- Zheng J (2020) SARS-CoV-2: an emerging coronavirus that causes a global threat. *Int J Biol Sci* 16(10):1678–1685
- Zheng ZG, Wang RS, Cheng HQ, Duan TT, He B, Tang D, Gu F, Zhu Q (2011) Isolated perfused lung extraction and HPLC–ESI–MSn analysis for predicting bioactive components of *Saposhnikovia Radix*. *J Pharm Biomed Anal* 54(3):614–618
- Zhou P, Yang XL, Wang XG, Hu B, Zhang L, Zhang W, Si HR, Zhu Y, Li B, Huang CL, Chen HD (2020) A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* 579(7798):270–273
- Zhu N, Zhang D, Wang W, Li X, Yang B, Song J et al (2020) A novel coronavirus from patients with pneumonia in China, 2019. *N Engl J Med*. <https://doi.org/10.1056/NEJMoa2001017>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Authors and Affiliations

Gagan Kumar Panigrahi¹ · Shraban Kumar Sahoo¹ · Annapurna Sahoo¹ · Shibasish Behera¹ · Snigdharani Sahu¹ · Archana Dash¹ · Kunja Bihari Satapathy¹

✉ Kunja Bihari Satapathy
kunjabihari.satapathy@cutm.ac.in

¹ School of Applied Sciences, Centurion University of Technology and Management, Khurda, Odisha, India