

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

4-Acetyl-1H-pyrrole-2-carbaldehyde

Tao Sun, Jia-Wei Xie, Ren-Ying Zhao, Ai-Guo Zhu and Yan-Qing Ge*

 Taishan Medical University, Tai'an 271016, People's Republic of China
 Correspondence e-mail: yqge@yahoo.cn


Received 20 August 2012; accepted 13 September 2012

 Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.042; wR factor = 0.128; data-to-parameter ratio = 14.5.

The title compound, $\text{C}_7\text{H}_7\text{NO}_2$, was synthesized *via* a one-pot Vilsmeier–Haack and subsequent Friedel–Crafts reaction. The pyrazole ring makes dihedral angles of 4.50 (9) and 2.06 (8)°, respectively, with the aldehyde and acetyl groups. In the crystal, classical $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds and weak $\text{C}-\text{H}\cdots\text{O}$ interactions assemble the molecules into a chain along the b axis.

Related literature

For the synthetic procedure, see: Ge *et al.* (2009). For related structures, see: Ge *et al.* (2011); Hao *et al.* (2012).



Experimental

Crystal data

$\text{C}_7\text{H}_7\text{NO}_2$
 $M_r = 137.14$
 Monoclinic, $P2_1/n$
 $a = 3.811$ (5) Å
 $b = 13.219$ (5) Å
 $c = 13.167$ (5) Å
 $\beta = 95.602$ (5)°

$V = 660.2$ (9) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.10$ mm⁻¹
 $T = 293$ K
 $0.15 \times 0.13 \times 0.10$ mm

Data collection

Bruker APEXII CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2005)
 $T_{\min} = 0.541$, $T_{\max} = 0.556$
 3752 measured reflections
 1348 independent reflections
 1010 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.056$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.042$
 $wR(F^2) = 0.128$
 $S = 1.04$
 1348 reflections
 93 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.20$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.14$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1A}\cdots\text{O2}^{\text{i}}$	0.86	2.11	2.876 (2)	148
$\text{C7}-\text{H7A}\cdots\text{O1}^{\text{ii}}$	0.96	2.54	3.453 (5)	159

Symmetry codes: (i) $x - \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$; (ii) $x + \frac{1}{2}, -y + \frac{1}{2}, z - \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2005); cell refinement: SAINT (Bruker, 2005); data reduction: SAINT; program(s) used to solve structure: SIR97 (Altomare *et al.*, 1999); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: XP in SHELXTL (Sheldrick, 2008); software used to prepare material for publication: WinGX (Farrugia, 1999).

This study was supported by the Shandong Natural Science Foundation (No. ZR2012BL04).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IM2396).

References

- Altomare, A., Burla, M. C., Camalli, M., Casciarano, G. L., Giacovazzo, C., Guagliardi, A., Moliterni, A. G. G., Polidori, G. & Spagna, R. (1999). *J. Appl. Cryst.* **32**, 115–119.
- Bruker (2005). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Ge, Y. Q., Hao, B. Q., Duan, G. Y. & Wang, J. W. (2011). *J. Lumin.* **131**, 1070–1076.
- Ge, Y. Q., Jia, J., Yang, H., Zhao, G. L., Zhan, F. X. & Wang, J. W. (2009). *Heterocycles*, **78**, 725–736.
- Hao, B.-Q., Xu, W.-R., Meng, F.-C. & Duan, G.-Y. (2012). *Acta Cryst.* **E68**, o877.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2012). E68, o2947 [doi:10.1107/S1600536812039153]

4-Acetyl-1H-pyrrole-2-carbaldehyde

Tao Sun, Jia-Wei Xie, Ren-Ying Zhao, Ai-Guo Zhu and Yan-Qing Ge

Comment

Synthesis of nitrogen-containing heterocyclic compounds has been a subject of great interest due to the wide application in agrochemical and pharmaceutical fields (Ge *et al.*; 2009, 2011). Some pyrrole derivatives which belong to this category exhibit interesting biological properties, such as anti-bacterial, anti-inflammatory, anti-oxidant, anti-tumor, anti-fungal, and immune suppressant activities. The title pyrazole (I) (Fig. 1) was synthesized in order to study its biological properties. (I) was screened for anticancer activities and found to be inactive. We report here the crystal structure of the title compound. In the title compound, C₇H₇NO₂, the pyrazole ring makes dihedral angles of 4.50 (9)° and 2.06 (8)°, respectively, with the aldehyde group and acetyl group. The crystal structure is determined by classical intermolecular N—H···O (H1A···O2 = 2.11 Å) hydrogen bonding and weak C—H···O (H7A···O1 = 2.54 Å) interactions, which assemble the molecules into a one-dimensional chain structure.

Experimental

A solution of dimethylformamide (5.5 mmol) in 1,2-dichloroethane (10 ml) in a 3-necked flask was cooled in an ice bath. To the stirred and cooled solution was added a solution of oxalyl chloride (5.5 mmol) in 1,2-dichloroethane (10 ml) over a period of 10 min. The suspension of white solid was then allowed to stir at room temperature for 15 min. The suspension was cooled in ice and a solution of pyrrole (5 mmol) in 1,2-dichloroethane (10 ml) was added over 10 min. The light orange solution obtained was allowed to stir 15 min at room temperature. To this was added aluminium chloride (11 mmol), followed by acetyl chloride (5 mmol), rapidly and at room temperature. The mixture was stirred for 3 h. The mixture was then poured onto about 50 ml of ice and water, 50% aqueous sodium hydroxide (4 ml) was added, and the mixture was stirred rapidly for about 10 min. The mixture was then made slightly acidic with concentrated hydrochloric acid and the organic and aqueous layers were separated. The aqueous layer was extracted with ethyl ether. The organic extracts were washed with water, dried, filtered and concentrated. The final product was isolated by column chromatography on silica gel (yield 76%). Crystals of (I) suitable for X-ray diffraction were obtained by allowing a refluxed solution of the product in ethyl acetate (0.10 M) to cool slowly to room temperature (without temperature control) and allowing the solvent to evaporate for 3 d.

Refinement

All H atoms were placed in geometrically calculated positions and refined using a riding model with C—H = 0.93 Å (for aldehyde group and pyrrole ring), C—H = 0.96 Å (for CH₃) and N—H = 0.86 Å with U_{iso}(H) = 1.2 U_{eq}(C,N) for the aldehyde group and pyrrole hydrogen atoms and U_{iso}(H) = 1.5 U_{eq}(C) for CH₃.

Computing details

Data collection: *APEX2* (Bruker, 2005); cell refinement: *SAINTE* (Bruker, 2005); data reduction: *SAINTE* (Bruker, 2005); program(s) used to solve structure: *SIR97* (Altomare *et al.*, 1999); program(s) used to refine structure: *SHELXL97*

(Sheldrick, 2008); molecular graphics: *XP* in *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *WinGX* (Farrugia, 1999).




Figure 1

The molecular structure of compound, showing displacement ellipsoids drawn at the 30% probability level.




Figure 2

The one-dimensional infinite chain structure, which is formed by intermolecular N—H \cdots O hydrogen bonding and weak C—H \cdots O interactions.

4-Acetyl-1*H*-pyrrole-2-carbaldehyde

Crystal data

$C_7H_7NO_2$

$M_r = 137.14$

Monoclinic, $P2_1/n$

Hall symbol: $-P 2_1n$

$a = 3.811 (5) \text{ \AA}$

$b = 13.219 (5) \text{ \AA}$

$c = 13.167 (5) \text{ \AA}$

$\beta = 95.602 (5)^\circ$

$V = 660.2 (9) \text{ \AA}^3$

$Z = 4$

$F(000) = 288$
 $D_x = 1.380 \text{ Mg m}^{-3}$
 Mo $K\alpha$ radiation, $\lambda = 0.71069 \text{ \AA}$
 Cell parameters from 1324 reflections
 $\theta = 3.1\text{--}25.7^\circ$

$\mu = 0.10 \text{ mm}^{-1}$
 $T = 293 \text{ K}$
 Block, colorless
 $0.15 \times 0.13 \times 0.10 \text{ mm}$

Data collection

Bruker APEXII CCD area-detector
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 phi and ω scans
 Absorption correction: multi-scan
 (SADABS; Bruker, 2005)
 $T_{\min} = 0.541$, $T_{\max} = 0.556$

3752 measured reflections
 1348 independent reflections
 1010 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.056$
 $\theta_{\max} = 26.4^\circ$, $\theta_{\min} = 2.2^\circ$
 $h = -4 \rightarrow 4$
 $k = -15 \rightarrow 16$
 $l = -12 \rightarrow 16$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.042$
 $wR(F^2) = 0.128$
 $S = 1.04$
 1348 reflections
 93 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods
 Secondary atom site location: difference Fourier
 map

Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0703P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.20 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.13 \text{ e \AA}^{-3}$
 Extinction correction: SHELXL97 (Sheldrick,
 2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
 Extinction coefficient: 0.031 (9)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.4828 (4)	0.16971 (9)	0.21257 (10)	0.0472 (4)
H1A	0.4131	0.1739	0.2726	0.057*
O2	0.7983 (4)	0.23425 (9)	-0.10389 (9)	0.0634 (4)
C4	0.5018 (4)	0.24663 (12)	0.14876 (11)	0.0448 (4)
H4	0.4423	0.3133	0.1619	0.054*
C6	0.6799 (4)	0.27169 (12)	-0.02967 (12)	0.0459 (4)
C2	0.5910 (4)	0.08234 (11)	0.16858 (11)	0.0452 (4)
O1	0.4742 (4)	-0.03023 (10)	0.29831 (10)	0.0777 (5)
C3	0.6811 (4)	0.10776 (11)	0.07349 (12)	0.0437 (4)
H3	0.7652	0.0638	0.0264	0.052*

C5	0.6236 (4)	0.21205 (11)	0.05991 (11)	0.0413 (4)
C7	0.5885 (6)	0.38131 (13)	-0.02795 (14)	0.0606 (5)
H7A	0.6548	0.4132	-0.0887	0.091*
H7B	0.7126	0.4126	0.0308	0.091*
H7C	0.3392	0.3887	-0.0248	0.091*
C1	0.5825 (5)	-0.01457 (13)	0.21628 (14)	0.0582 (5)
H1	0.6661	-0.0696	0.1819	0.070*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0623 (9)	0.0475 (8)	0.0331 (7)	-0.0014 (6)	0.0119 (6)	-0.0040 (6)
O2	0.0926 (10)	0.0596 (8)	0.0412 (7)	-0.0007 (6)	0.0225 (6)	-0.0005 (6)
C4	0.0576 (10)	0.0385 (8)	0.0390 (9)	0.0007 (7)	0.0082 (7)	-0.0040 (7)
C6	0.0535 (9)	0.0477 (9)	0.0366 (9)	-0.0042 (7)	0.0046 (7)	-0.0013 (7)
C2	0.0538 (10)	0.0432 (9)	0.0383 (9)	-0.0001 (7)	0.0026 (7)	-0.0017 (7)
O1	0.1213 (12)	0.0589 (9)	0.0546 (9)	-0.0035 (7)	0.0173 (8)	0.0133 (7)
C3	0.0503 (9)	0.0423 (9)	0.0389 (9)	0.0017 (7)	0.0065 (7)	-0.0067 (7)
C5	0.0467 (9)	0.0420 (9)	0.0356 (8)	-0.0020 (6)	0.0053 (6)	-0.0024 (6)
C7	0.0764 (12)	0.0480 (10)	0.0585 (11)	0.0044 (9)	0.0112 (9)	0.0073 (8)
C1	0.0797 (13)	0.0467 (10)	0.0480 (10)	-0.0001 (8)	0.0052 (9)	0.0011 (8)

Geometric parameters (\AA , $^\circ$)

N1—C4	1.325 (2)	C2—C1	1.429 (2)
N1—C2	1.373 (2)	O1—C1	1.211 (2)
N1—H1A	0.8600	C3—C5	1.405 (2)
O2—C6	1.2207 (18)	C3—H3	0.9300
C4—C5	1.378 (2)	C7—H7A	0.9600
C4—H4	0.9300	C7—H7B	0.9600
C6—C5	1.452 (2)	C7—H7C	0.9600
C6—C7	1.491 (2)	C1—H1	0.9300
C2—C3	1.372 (2)		
C4—N1—C2	109.91 (13)	C5—C3—H3	126.1
C4—N1—H1A	125.0	C4—C5—C3	106.19 (13)
C2—N1—H1A	125.0	C4—C5—C6	126.74 (14)
N1—C4—C5	109.11 (13)	C3—C5—C6	127.07 (13)
N1—C4—H4	125.4	C6—C7—H7A	109.5
C5—C4—H4	125.4	C6—C7—H7B	109.5
O2—C6—C5	121.72 (16)	H7A—C7—H7B	109.5
O2—C6—C7	120.76 (14)	C6—C7—H7C	109.5
C5—C6—C7	117.52 (14)	H7A—C7—H7C	109.5
C3—C2—N1	106.92 (14)	H7B—C7—H7C	109.5
C3—C2—C1	129.75 (15)	O1—C1—C2	124.74 (17)
N1—C2—C1	123.23 (15)	O1—C1—H1	117.6
C2—C3—C5	107.87 (13)	C2—C1—H1	117.6
C2—C3—H3	126.1		
C2—N1—C4—C5	0.08 (18)	C2—C3—C5—C6	-179.38 (15)

C4—N1—C2—C3	0.20 (18)	O2—C6—C5—C4	177.98 (16)
C4—N1—C2—C1	-176.51 (16)	C7—C6—C5—C4	-1.8 (2)
N1—C2—C3—C5	-0.40 (17)	O2—C6—C5—C3	-2.2 (2)
C1—C2—C3—C5	176.02 (16)	C7—C6—C5—C3	177.96 (15)
N1—C4—C5—C3	-0.33 (17)	C3—C2—C1—O1	-174.36 (18)
N1—C4—C5—C6	179.50 (14)	N1—C2—C1—O1	1.5 (3)
C2—C3—C5—C4	0.45 (17)		

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N1—H1 <i>A</i> \cdots O2 ⁱ	0.86	2.11	2.876 (2)	148
C7—H7 <i>A</i> \cdots O1 ⁱⁱ	0.96	2.54	3.453 (5)	159

Symmetry codes: (i) $x-1/2, -y+1/2, z+1/2$; (ii) $x+1/2, -y+1/2, z-1/2$.