

# Effects of the Percutaneous Carbon Dioxide Therapy on Post-surgical and Post-traumatic Hematoma, Edema and Pain

Salvatore Ratano<sup>1</sup>, Biljana Jovanovic<sup>2</sup>, Eric Choudja Ouabo<sup>2</sup>

## Learning Point of the Article:

These findings encourage the development of specific treatment protocols with CDT to heal hematoma and edema, enhance tissue regeneration, and improve functional recovery after trauma or surgical interventions.

## Abstract

**Introduction:** Carbon dioxide therapy (CDT) has been used since the Roman Empire to improve the microcirculation and oxygenation of tissues altered. The classical indications are vasculopathies and ischemic diseases.

**Hypothesis:** The CDT could be effective in favoring joint mobility recovery and pain reduction in patients with post-surgical or post-traumatic hematoma, edema, and pain.

**Study Design:** Case report.

**Material and Methods:** Eight patients were treated once a day for 5–10 days; a single session lasted 50 min. For post-surgical cases, the treatment began the day or the day after they were discharged from the hospital.

**Result:** For all patients in this series, the CDT has brought clinical benefits in terms of decreasing pain and improving of joint mobility.

**Conclusion:** The CDT is a safe and effective treatment to provide a greater amount of oxygen to the injured tissues. It favors the healing of post-surgical and post-traumatic hematoma and edema, promoting the mobility recovery of patients.

**Keywords:** Carbon dioxide therapy, hematoma, edema, pain, mobility recovery, case report.

## Introduction

The term “carboxytherapy” has appeared for the first time in the scientific community in 1995 by Parassoni during the XVI National Assembly of Aesthetic Medicine in Rome, and it refers to the transcutaneous or percutaneous administration of CO<sub>2</sub> with therapeutic purposes [1].

Carbon dioxide therapy (CDT) finds its origins in Royat, France, where the first Roman spa called “Rubiaccum” (because of rosy water contenting ferrum) was found, dating to 20 BC during the empire of Augustus. It arouses the interest of people even in the

following eras, like the Celts, who worshiped this place as miraculous, but, however, after the invasion of the Barbarians in the 5th century, the site was forgotten [2]. Later, in 1932, Barrieu first treated patients with arterial and venous disorders with CO<sub>2</sub> spring gas injections [3]. In 1946, in France, the Institute of Research of Royat was inaugurated, managing to produce during the second half of the century nearly 400 publications about carboxytherapy, also called carbocrenotherapy [2]. Over the years, CDT has found considerable development until a consensus regarding the effects of CO<sub>2</sub> was formulated during

Access this article online

Website:  
www.jocr.co.in

DOI:  
<https://doi.org/10.13107/jocr.2023.v13.i12.4058>

## Author's Photo Gallery



Dr. Salvatore Ratano



Dr. Biljana Jovanovic



Dr. Eric Choudja Ouabo

<sup>1</sup>Department of Orthopaedic Surgery, University of Palermo, Palermo, Italy,

<sup>2</sup>Private Practice, Center of Orthopedic Surgery, Clinique Bois-Cerf, 1006 Lausanne, Switzerland.

### Address of Correspondence:

Dr. Eric Choudja Ouabo,  
Private Practice, Center of Orthopedic Surgery, Clinique Bois-Cerf, 1006 Lausanne, Switzerland.  
E-mail: e\_choudja@hotmail.com

Submitted: 18/09/2023; Review: 08/10/2023; Accepted: November 2023; Published: December 2023

DOI: <https://doi.org/10.13107/jocr.2023.v13.i12.4058>

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License <https://creativecommons.org/licenses/by-nc-sa/4.0/>, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms



**Figure 1:** (a and b) patient positioned supine on the table during CDT

the International Conference in Freiburg-en-Breisgau, Germany, in 1989 and 1999.

The following therapeutic effects were highlighted in consensus: (a) local increase of blood supply and opening of functionally closed capillaries; (b) dilatation of precapillary segments; (c) improvement of tissue oxygenation by arterial increased liberation of oxygen; (d) improved deformability of erythrocytes; (e) modification of threshold of thermoreceptors; (f) antiseptic effect [4,5]. The physiopathological bases of these therapeutic effects reside in the administration of CO<sub>2</sub>, which improves the arterial blood flow, the microcirculation, and increase the tcPO<sub>2</sub> in ischemic tissues, which is explained by the Bohr effect. In fact, the presence of CO<sub>2</sub> causes a rightward shift of the O<sub>2</sub>-Hb dissociation curve with a consequent increase in pCO<sub>2</sub> or decrease in pH. In lower pH and higher pCO<sub>2</sub>, the affinity of hemoglobin to oxygen is decreased [6].

The traditional therapeutic indication is vasculopathy or ischemic diseases, including diabetic periphery syndrome, Morbus Buerger, Reynaud's syndrome, chronic venous insufficiency, chronic venous-lymphatic insufficiency, and recently also erectile dysfunction associated with

microangiopathy. The CDT has found indications in other branches of medicine, such as dermatology, where this method is very useful in wound healing, including leg ulcers and hair disorders, and helps to improve psoriasis and scleroderma [6].

Depending on the method of administration of CO<sub>2</sub>, different types of treatment are available: (a) use of spa therapy like in Royat Center; (b) use of artificial CO<sub>2</sub>-enriched water for bathing to improve ischemic limb symptoms; (c) subcutaneous injection of CO<sub>2</sub> in plastic surgery; (d) percutaneous administration of CO<sub>2</sub>-rich spa gas [7].

Considering all these indications, the different methods, and the effects of CDT, the authors have assumed clinical benefits in other ischemic disorders like hematoma: A blood collection inside a tissue following a breaking blood vessel involved a reduction of oxygen in the affected tissues, which showed signs of adaptation to hypoxia [8].

The aim of the present article was to expose the effects of percutaneous CDT on the healing of wounded tissues in patients with post-surgical and post-traumatic hematoma, edema, and pain. Better oxygenation has an anti-inflammatory effect and decreases pain.

### Materials and Methods

Between the patients treated from 2019 to 2022, we selected 8 patients, 2 men and 6 women, with a mean age of 56.8 years (range 36–72 years), who were treated at the Center of Surgery, Hand, and Aesthetics in Lausanne, Switzerland. Informed consent was obtained from all patients, and the rights of the subjects were protected. The device used was the CARBO® Fit system produced by Derma Art, Slovenia.

Their selection is based on the presence of hematoma, edema,



**Figure 2:** (a) post-traumatic hematoma throughout the leg, ankle, and foot; (b) MRI showing myotendinous tear of both medial and lateral gastrocnemius muscles with edema; (c, d). After 5 days, hematoma had disappeared.

Indications	Contraindications
Haematoma, edema and post-traumatic and post-surgical pain	Pregnancy
Wound healings	Acute myocardial infarction
Muscle injury	Severe respiratory failure
Morel-Lavallée syndrome	Severe renal failure
Vasculopathies	Severe anaemia
Diabetic ulcer	Chronic liver failure
CRPS	Skin infection
Buerger's disease	Gaseous gangrene
Reynaud's syndrome	Cancer
Polyneuropathy	Deep venous thrombosis
Chronic venous insufficiency	Vascular surgery of the lower limb during the 6 months before
Chronic venous-lymphatic insufficiency	
Leg ulcers	
Skin disorder (to improve psoriasis, scleroderma, Angiectactic rosacea)	
<b>CDT: Carbon dioxide therapy</b>	

**Table 1: Indications and contraindications of the CDT.**

pain, and joint functional reduction after a trauma (3 patients) or a surgical intervention (5 patients). The patients were treated once a day for 5–10 days. For post-surgical cases, the treatment began the day or the day after they were discharged from the hospital. A clinical examination was conducted to check the correct indication for the CDT and to exclude any contraindications during the first day of therapy (Table 1). In view of the fact that the effect of CDT depends on many effects (its concentration, surface area exposure, contact time, skin temperature, number, and frequency of sessions), the number of

sessions is variable depending on the patient's clinical condition and his evolution after the first or second CDT treatment. In some acute cases, double-daily administration may be indicated. During the treatment, the patient is positioned supine on the table, and the body region to be treated is covered with a sealed envelope (Fig. 1a and b). The ambient air is removed and replaced with 99.5% medical carbon dioxide at a pressure of 49.5 bars at 15°C. The quantity of CO<sub>2</sub> used depends on the body surface to be treated; generally, we use between 0.25 and 0.3 L of CO<sub>2</sub> per session. A single session lasted 50 min. The patient is comfortable and can relax by reading a book, listening to music, or sleeping. If necessary, during the wintertime, the skin is heated for a few minutes stay in an infrared sauna, which increases the passage rate of carbon dioxide.

## Cases

### Post-traumatic cases

Case No. 1: A 76-year-old male patient (70 kg, 172 cm) was evaluated for strong leg pain (visual analog scale [VAS]: 8/10) after a fall on skiing. He walked with crutches without leaning his foot on the ground, with a limited extension of 30° of the left knee. The pain was refractory to conservative treatment with non-steroidal anti-inflammatory drugs (NSAIDs). The initial clinical presentation was a hardening of the medial gastrocnemius region associated with hematoma throughout the leg, ankle, and foot (Fig. 2a). Magnetic resonance imaging demonstrated a myotendinous tear of both medial and lateral gastrocnemius muscles with edema over the entire leg (Fig. 2b). The patient experienced immediate relief of the pain (VAS: 4/10); after 3 days, he stopped taking NSAIDs. After 5 days, he walked independently without limping, and the pain had disappeared (VAS: 0/10) (Fig. 2c and d). At 5 months of follow-

	Cases	sex	Age	weight	Height (m)	IMC (Kg/m <sup>2</sup> )	Initial VAS	Final VAS	Sessions of CDT
Post-Traumatic	n.1	Male	76	70	1.72	23.7	8	0	5
	n.2	Female	57	58	1.73	19.4	5	0-1	5
	n.3	Female	47	60	1.68	21.3	7-8	1-2	10
Post-Surgical	n.4	Male	36	72	1.78	22.7	8-9	1-2	7
	n.5	Female	59	60	1.74	19.8	7-8	1-2	7
	n.6	Female	62	50	1.65	18.4	6-7	1-2	7
	n.7	Female	60	65	1.70	22.5	9	2-3	10
	n.8	Female	58	79	1.57	32.0	9	2	8

**Table 2: Demographic data and results of reported cases.**





**Figure 3:** (a) edema and hematoma after ankle sprain; (b,c) resorption of edema and hematoma after CDT

up, the patient had no disorder: complete range of motion of the knee, normal muscle contractions, and without endurance.

**Case No. 2:** A 57-year-old female patient (58 kg, 173 cm) had shown up for a severe ankle sprain (initial VAS: 5/10); she walked with crutches and presented edema and hematoma on the peroneal side (Fig. 3a). The indication was given for carboxytherapy treatment: 1 time a day for 5 days. After 1 day of therapy, the malleolar swelling was greatly decreased (Fig. 3b); after 4 days, the malleolar edema disappeared, and the hematoma was almost completely reduced (Fig. 3c). After 6 days from her accident, the patient returned to sport, and she resumed swimming (final VAS: 0–1/10).

**Case No. 3:** A 47-year-old female patient (60 kg, 168 cm) was evaluated after a scooter accident. Clinically, she presented right zygomatico-maxillary fractures and an extended hematoma and edema on the whole lower limb with almost full limitation of knee mobility (Fig. 4a). The diagnosis of Morel-Lavallée syndrome was posed (initial VAS: 7–8/10). The knee blending was surgically drained, with post-operative improvement of knee flexion up to 80° (Fig. 4b). A few days later, treatment for CDT was initiated. After 10 days of treatment, the patient had no pain anymore, and the hematoma and edema were greatly reduced. Moreover, the knee flexion was improved by up to 120° (final

VAS: 1–2/10) (Fig. 4c and d).

#### Post-surgical cases

**Case No. 4:** A 36-year-old male patient (72 kg, 178 cm) consulted 3 days after arthroscopic combined anterior cruciate ligament and anterolateral reconstruction with hamstring autograft complaining pain and knee limited extension at 25° and flexion at 60° (VAS: 8–9/10) (Fig. 5a and b) despite taking strong doses of analgic drugs. After a cycle of seven sessions of carboxytherapy, the patient had no pain anymore and an improved knee range of motion: a complete extension and flexion of 90° (final VAS: 1–2/10) (Fig. 5c). He was now able to begin post-surgical rehabilitation.

surgical rehabilitation.


**Case No. 5:** A 59-year-old female patient (60 kg, 174 cm) was evaluated 4 days after a bilateral Reviel osteotomy of the 1st metatarsus and Akin osteotomy of the 1st phalanx foot and percutaneous pinning of the 2nd toe of the left foot for bilateral hallux valgus and claw toe (Fig. 6a and b). She took a triple dose of NSAIDs and paracetamol for pain control (initial VAS: 7–8/10). After 3 carboxytherapy sessions, the patient was able to stop any medication, and the edema was sharply reduced. After 7 sessions of therapy, she had no pain anymore (VAS: 1–2/10) (Fig. 6c).

**Case No. 6:** A 62-year-old female patient (50 kg, 165 cm). The patient had on the left foot a plate removal of the second metatarsal bone and a resection of an exostosis of the navicular bone. In the same operation, she also had on the right foot a Reviel osteotomy of the 1st metatarsal bone, an Akin osteotomy of the first phalanx of the great toe, and an exostosis resection (Fig. 7a). She complained of important post-operative pain and edema of both feet and a slight dehiscence of the osteotomy scar (initial VAS: 6–7/10) (Fig. 7b). Treatment with carboxytherapy was indicated. After 7 sessions of CDT, the patient had no pain anymore, and the edema was clearly reduced (VAS: 1–2/10) (Fig. 7c).



**Figure 4:** (a) Post-traumatic hematoma in Morel-Lavallée syndrome; (b) After surgical drainage the knee blending; (c,d) hematoma and edema greatly reduced after 10 days of CDT.





**Figure 5:** (a,b) Edema and hematoma 3 days after ACL reconstruction; (c) Knee joint mobility improvement after CDT.

Case No. 7: A 60-year-old female patient (65 kg, 170 cm) was evaluated after total knee arthroplasty with anterior tibial tuberosity osteotomy of the right knee for tricompartmental knee osteoarthritis with patella baja. She is complaining of pain, edema, and hematoma of the entire inferior leg (initial VAS: 9/10) (Fig. 8a and b). A clear clinical improvement was obtained after 10 sessions of CDT: almost complete pain disappeared and there was sharp reduction of hematoma and edema (final VAS: 2–3/10) (Fig. 8c).

Case No. 8: A 58-year-old female patient (79 kg, 157 cm) presented with hematoma of the leg after a tibial open wedge high tibia osteotomy left knee for varus medial femoro-tibial osteoarthritis, complaining of pain and reduced knee mobility (40° of knee flexion). She takes NSAIDs and morphine derivatives (tramadol chlorhydrate) (VAS: 9/10) (Fig. 9a and b). She was treated for 8 days with CDT and reported favorable clinical results: improvement of knee range of motion (knee flexion at 85°) and reduction of hematoma. After 4 days, she gave up tramadol chlorhydrate, and after 8 days, she only takes NSAIDs if needed (VAS: 2/10) (Fig. 9c and d).

For all patients in this series, the CDT has brought clinical benefits in terms of decreasing edema, pain, and improving joint mobility (Table 2). The reason is the effects of CDT on the microcirculation and oxygenation of the tissues altered by surgery or trauma. The administration of CO<sub>2</sub> improves arterial blood flow and increases the tcPO<sub>2</sub> in ischemic tissues, thanks to the Bohr effect [6].

These patients recovered their joint mobility in a mean time of 7 days. This is very significant data if we consider the clinical results of Jackson and Feagin, who studied the prognosis of quadriceps contusions in young athletes treated with rest and early passive pain-free motion. They found an average disability of 7 days for mild contusions, 56 days for moderate contusions, and 72 days for severe contusions [9].

To our knowledge, in the literature, no study has addressed the effects of the CDT on the mechanisms of healing of hematoma, edema, and pain. Some studies have addressed different pathological frameworks relating to the skin and subcutaneous layers.

Brandi et al. studied the role of CDT on 70 patients with chronic wounds; their conclusion was that this treatment is simple to use, without important side-effects, and provides a positive

## Discussion



**Figure 6:** (a,b) Post-operative X-rays and foot hematoma; (c) Hematoma reduction after 7 sessions of CDT.




**Figure 7:** (a,b) Post-operative foot X-ray, edema and hematoma; (c) Edema and hematoma reduction after 7 sessions of CDT.


contribution to promote wound healing [8]. Pinheiro et al. observed that CDT used with radiofrequency improved abdominal skin elasticity after abdominoplasty [10]. Pianez et al. (2016), in their studies, assessed the effectiveness of CDT in

the reduction of cellulite on buttocks and thighs [11].

Carboxytherapy was also studied on different types of alopecia with five sessions at 1–3-weeks intervals. In cases of androgenic baldness, hair regrowth has been observed within thinning sites



**Figure 8:** (a,b) Post-operative Total knee arthroplasty X-ray, Leg edema and hematoma; (c) Leg, edema and hematoma reduction after 10 sessions of CDT.



**Figure 9:** (a,b) Post-operative edema and hematoma, X-ray after High tibial osteotomy; (c,d) Edema and hematoma reduction after 8 days of CDT.



following CDT [12].

The present article has several limitations due to the small number of patients, the shortness of the follow-up, and the absence of case controls. Therefore, despite the encouraging results, long-term controlled trials are required to better define specific treatment protocols for CDT in patients with hematoma, edema, and pain.

### Conclusion

CDT is a safe and effective treatment to provide a greater amount of oxygen to the injured tissues. This series highlights his favorable clinical benefits for patients with

post-surgical and post-traumatic hematoma, edema, and pain. Further studies are needed to confirm these promising results.

### Clinical Message

- The use of percutaneous coroxytherapy, started between D0 and D7, has a clear analgesic effect and allows faster functional recovery.
- Some contraindications to CTP treatment must be respected, as listed in Table 1.

**Declaration of patient consent:** The authors certify that they have obtained all appropriate patient consent forms. In the form, the patient has given the consent for his/ her images and other clinical information to be reported in the journal. The patient understands that his/ her names and initials will not be published and due efforts will be made to conceal their identity, but anonymity cannot be guaranteed.

**Conflict of interest:** Nil **Source of support:** None

### References

1. Varlaro V, Manzo G, Mugnaini F, Bisacci C, Fiorucci P, De Rango P, et al. Carboxytherapy: Effects on microcirculation and its use in the treatment of severe lymphedema. *Acta Phlebol* 2007;8:79-91.
2. Fabry R, Dubost JJ, Schmidt J, Body J, Schaff G, Baguet JC. Thermal treatment in arterial diseases: An expensive placebo or an effective therapy? *Therapie* 1995;50:113-22.
3. Barrieu M. Royat thermal gas subcutaneous injections. *Maroc Med* 1949;28:265.
4. Body J, Morel F, Schaff G. Effets vaso-actifs du CO2 thermal. *Angeiologie* 2000;52:71-5.
5. Finzgar M, Melik Z, Cankar K. Effect of transcutaneous application of gaseous carbon dioxide on cutaneous microcirculation. *Clin Hemorheol Microcirc* 2015;60:423-35.
6. Sakai Y, Miwa M, Oe K, Ueha T, Koh A, Niikura T, et al. A novel system for transcutaneous application of carbon dioxide causing an "artificial bohr effect" in the human body. *Plos One* 2011;6:e24137.
7. Ito T, Moore JI, Koss MC. Topical application of CO2 increases skin blood flow. *J Invest Dermatol* 1989;93:259-62.
8. Brandi C, Grimaldi L, Nisi G, Brafa A, Campa A, Calabrò M, et al. The role of carbon dioxide therapy in the treatment of chronic wounds. *In Vivo* 2010;24:223-6.
9. Jackson DW, Feagin JA. Quadriceps contusions in young athletes. Relation of severity of injury to treatment and prognosis. *J Bone Joint Surg Am* 1973;55:95-105.
10. Pinheiro NM, Crema VO, Millan BM, Carvalho FA, Mendonca AC. Comparison of the effects of carboxytherapy and radiofrequency on skin rejuvenation. *J Cosmet Laser Ther* 2015;17:156-61.
11. Pianez LR, Custódio FS, Guidi RM, De Freitas JN, Sant'Ana E. Effectiveness of carboxytherapy in the treatment of cellulite in healthy women: A pilot study. *Clin Cosmet Investig Dermatol* 2016;9:183-90.
12. King S, King M. Using carboxytherapy as an adjuvant aesthetic treatment for patients with alopecia. *J Aesthetic Nurs* 2013;2:434-5.

**Conflict of Interest:** Nil  
**Source of Support:** Nil

**Consent:** The authors confirm that informed consent was obtained from the patient for publication of this case report

### How to Cite this Article

Ratano S, Jovanovic B, Choudja Ouabo E. Effects of the Percutaneous Carbon Dioxide Therapy on Post-surgical and Post-traumatic Hematoma, Edema and Pain. *Journal of Orthopaedic Case Reports* 2023 December;13(12): 11-17.

