

SCIENTIFIC REPORTS

OPEN

Relationship between auxiliary gamma subunits and mallotoxin on BK channel modulation

Xin Guan, Qin Li & Jiusheng Yan

Received: 28 October 2016
Accepted: 03 January 2017
Published: 06 February 2017

The large-conductance, calcium- and voltage-activated K^+ (BK) channel consists of the pore-forming α subunits (BK α) and auxiliary subunits. The auxiliary γ 1-3 subunits potently modulate the BK channel by shifting its voltage-dependence of channel activation toward the hyperpolarizing direction by approximately 145 mV (γ 1), 100 mV (γ 2), and 50 mV (γ 3). Mallotoxin is a potent small-molecule BK channel activator. We analyzed the relationship between mallotoxin and the γ subunits in their BK channel-activating effects in membrane patches excised from HEK-293 cells. We found that mallotoxin, when applied extracellularly, shifted the half-activation voltage ($V_{1/2}$) of BK α channels by -72 mV. The channel-activating effect of mallotoxin was greatly attenuated in the presence of the γ 1, γ 2, or γ 3 subunit, with resultant $\Delta V_{1/2(+/-\text{ mallotoxin})}$ values of -9 , -28 , or -15 mV, respectively. Most examined γ 1 mutant subunits antagonized mallotoxin's channel-activating effect in a manner that was largely dependent on its own modulatory function. However, mallotoxin caused an irreversible functional and structural disengagement of the γ 1-F273S mutant from BK channels. We infer that the auxiliary γ subunit effectively interferes with mallotoxin on BK channel modulation via either a direct steric competition or an indirect allosteric influence on mallotoxin's binding and action on BK α .

The mammalian big/large-conductance, voltage- and calcium-activated potassium (BK) channel is widely expressed in various tissues and cell types and plays important roles in many physiological processes, including contractile activity of smooth muscles¹, regulation of neurotransmitter release and neuronal firing²⁻⁵, and frequency tuning of auditory hair cells⁶. The BK channel features large single-channel conductance and dual activation by membrane depolarization and elevation of intracellular free calcium ($[Ca^{2+}]_i$). BK channels consist of the pore-forming, voltage- and Ca^{2+} -sensing α subunits (BK α), either alone or together with tissue-specific auxiliary β subunits (β 1- β 4) or γ subunits (γ 1- γ 4)⁷⁻⁹. The BK channel γ subunits (BK γ) exhibit different tissue-specific mRNA expression and may broadly modulate BK channels in different tissues⁹. The few published studies have shown that the γ 1 subunit regulates BK channels in prostate cancer cells^{7,8}, salivary gland cells¹⁰, airway epithelial cells^{11,12}, and also probably in arterial smooth muscle cells¹³.

The auxiliary γ subunits are a group of four leucine-rich repeat (LRR)-containing membrane proteins, γ 1 (LRRC26), γ 2 (LRRC52), γ 3 (LRRC55), and γ 4 (LRRC38). They are structurally distinct from the double membrane-spanning BK channel β subunits in possessing an N-terminal signal peptide, an extracellular LRR domain, a single transmembrane segment, and a short intracellular C-terminus^{8,9,14}. The γ 1, γ 2 and γ 3 subunits, all ~ 35 kDa in size, facilitate BK channel activation by shifting voltage-dependence of channel activation in the hyperpolarizing direction over an exceptionally large range, by approximately 145 mV (γ 1), 100 mV (γ 2), and 50 mV (γ 3) in the absence of calcium^{8,15}.

In addition to their regulation by auxiliary proteins, BK channels are modulated by a variety of endogenous or exogenous small peptide or chemical molecules¹⁶⁻¹⁹. The pharmacological properties of BK channels are extensively altered in the presence of the auxiliary β subunits²⁰. The interactions between the β subunits and small molecule modulators of BK channels have been a subject of intensive investigations^{20,21}. However, currently little is known about the influence of the newly identified γ subunits on the BK channel's pharmacological properties.

Mallotoxin, also called rottlerin, is the principal phloroglucinol constituent of kamala, a red powder on the surface of the fruit of the kamala tree (*Mallotus philippinensis*) that has been used as a folk medicine in treating tapeworm infection, eye diseases, bronchitis, abdominal disease, spleen enlargement, and other illnesses²². At

Department of Anesthesiology and Perioperative Medicine, The University of Texas MD Anderson Cancer Center, Houston, Texas, USA. Correspondence and requests for materials should be addressed to J.Y. (email: jyan1@mdanderson.org)




Figure 1. Effects of mallotoxin on BK channels formed by BK α alone. (a,b) Voltage protocols (left) and the representative activation (a) and deactivation (b) currents of the BK α channel at different voltages in the absence (-MTX) and presence of mallotoxin (+MTX). (c) Voltage-dependence of BK channel activation. (d,e) Voltage-dependence of the kinetics of channel activation (d) and deactivation (e).

micromolar concentrations, mallotoxin potently shifted the voltage-dependence of BK channel activation toward the hyperpolarizing potential direction when applied extracellularly on whole cells²³. The BK channel-activating effect of mallotoxin was largely abolished when the $\gamma 1$ subunit was either endogenously expressed in salivary gland cells or heterologously expressed in human embryonic kidney (HEK)-293 cells¹⁰. Despite the potent activating effects of the $\gamma 1$ subunit and mallotoxin on BK channels, the mechanisms underlying their individual actions and their interactions within BK channel complexes are largely unknown. In this study, we analyzed the relationships between the BK channel auxiliary γ subunits and mallotoxin and probed the mechanisms underlying the apparent antagonizing effect of the γ subunits on mallotoxin's channel-activating function.

Results

Effects of mallotoxin on BK α channels in excised membrane patches. We observed, in excised patches of HEK-293 cell membrane, that extracellular exposure of the heterologously expressed BK α channels to mallotoxin potently shifted the voltage-dependence of channel activation. Mallotoxin was included in pipette solution under an inside-out configuration and data was collected after the mallotoxin achieved its maximal effect in shifting the BK channel's conductance-voltage (G-V) relationship towards hyperpolarizing direction. The results from mallotoxin treatment were compared to those without mallotoxin that were obtained in separate patches. Mallotoxin at 2 μ M resulted in a large shift of 72 mV in the half-activation voltages ($V_{1/2}$) of BK channels toward the hyperpolarizing direction (to 92 ± 6 mV, $n = 4$), compared to the untreated BK α channels ($V_{1/2} = 164 \pm 3$ mV, $n = 11$) in the virtual absence of $[Ca^{2+}]_i$ (Fig. 1a and c; Table 1). The mallotoxin-induced left shift in the conductance-voltage (G-V) relationship was accompanied by a great deceleration in channel deactivation (e.g., ~ 10 times decrease in rate at -160 mV), while the rate of channel activation at tested voltages (120–200 mV) was largely unaffected (Fig. 1b,d and e). Increase in mallotoxin concentration from 2 to 10 and 20 μ M showed no significant increase in modulatory effect on BK channels. We also observed that mallotoxin at a high concentration, e.g., 10 or 20 μ M, precipitated in pipette solution. Thus, 2 μ M mallotoxin was used throughout this study except where another concentration is specified.

Effects of mallotoxin on BK channels in the presence of auxiliary γ subunits. To test whether the auxiliary γ subunits affect mallotoxin's channel-activating effect on BK channels, we co-expressed $\gamma 1$, $\gamma 2$, $\gamma 3$, and

Expression	No mallotoxin			+2 μ M mallotoxin		
	V _{1/2} (mV)	z	n ^a	V _{1/2} (mV)	z	n
BK α alone	164 \pm 3	1.28 \pm 0.05	11	92 \pm 6	1.32 \pm 0.05	4
+ γ 1	23 \pm 3	1.57 \pm 0.11	10	14 \pm 2	1.29 \pm 0.10	6
+ γ 1 [*]	27 \pm 6	1.10 \pm 0.06	4			
+ γ 2	61 \pm 3	1.17 \pm 0.07	8	33 \pm 2	1.10 \pm 0.12	3
+ γ 3	115 \pm 2	1.36 \pm 0.05	6	100 \pm 5	1.13 \pm 0.03	4
+ γ 4	154 \pm 3	1.27 \pm 0.07	7	85 \pm 8	1.22 \pm 0.09	4
+ γ 1/ γ 4-linker&tail	97 \pm 3	1.03 \pm 0.03	5	53 \pm 6	1.00 \pm 0.11	5
+ γ 1/ γ 4-TMa	69 \pm 3	1.33 \pm 0.05	3	35 \pm 5	1.15 \pm 0.15	4
+ γ 1/ γ 4-TMc	59 \pm 3	1.11 \pm 0.09	4	29 \pm 4	1.19 \pm 0.06	5
+ γ 1-S272V	64 \pm 5	1.02 \pm 0.05	5	24 \pm 1	1.25 \pm 0.15	3
+ γ 1-L274A	25 \pm 4	1.43 \pm 0.13	3	20 \pm 2	1.35 \pm 0.10	4
+ γ 1-V275A	26 \pm 2	1.60 \pm 0.13	3	23 \pm 4	1.00 \pm 0.18	3
+ γ 1-P270V/F273S	155 \pm 2	1.48 \pm 0.14	5	107 \pm 8	0.98 \pm 0.05	4
+ γ 1-F273S/L274A	157 \pm 2	1.25 \pm 0.19	3	81 \pm 3	1.05 \pm 0.06	5
+ γ 1-F273S/V275A	152 \pm 2	1.14 \pm 0.07	4	74 \pm 7	0.76 \pm 0.06	5
+ γ 1- Δ 3R	40 \pm 1	1.34 \pm 0.05	4	38 \pm 4	1.07 \pm 0.05	6
+ γ 1- Δ 4R	146 \pm 4	1.15 \pm 0.05	11	94 \pm 4	0.93 \pm 0.15	3
+ γ 1-F273S/ Δ 3R	163 \pm 7	1.13 \pm 0.04	5	134 \pm 6	1.00 \pm 0.03	5
+ γ 1- Δ tailN ²⁹¹⁻²⁹⁸	168 \pm 3	1.22 \pm 0.11	5	66 \pm 3	0.97 \pm 0.07	4
+ γ 1-F273S	18 \pm 1 (79%) ^b	1.55 \pm 0.07 ^b	13	96 \pm 5	1.12 \pm 0.05	5
	126 \pm 8 (21%) ^{bc}	0.99 \pm 0.16 ^{bc}				
+ γ 1-F273S [*]	147 \pm 3	1.14 \pm 0.08	6			

Table 1. Boltzmann fit parameters of voltage-dependent BK channel activation. ^{*}Condition of mallotoxin withdrawal in which cells were preincubated with 2 μ M mallotoxin for 5 min and then transferred to a mallotoxin-free solution. ^an values are the number of recorded excised inside-out patches from different HEK-293 cells. ^bThe indicated percentage in parentheses here refers to the portion of the channels' subpopulation that was obtained from a double Boltzmann function fit. ^cBecause of the difficulty in obtaining reliable parameter values from a double Boltzmann function fit for the minor portion (e.g., \leq 35%), the estimated values of the V_{1/2} and errors provided here are considered less reliable and used for reference only.

γ 4 individually with BK α in HEK-293 cells. In the presence of the γ 1 subunit, application of mallotoxin caused no significant change in BK channel V_{1/2} (Δ V_{1/2} = -9 mV). The V_{1/2} values of mallotoxin-treated and -untreated BK α γ 1 channels in the virtual absence of [Ca²⁺] were 14 \pm 2 mV (n = 5, fitted by single Boltzmann function) and 23 \pm 3 mV (n = 10), respectively (Fig. 2a,c; Table 1). The BK α γ 1 channel deactivation was moderately slower in the presence of mallotoxin (e.g., ~3 times slower at -160 mV) than in the absence of mallotoxin, while the channel activation was less affected (Fig. 2b,d and e). Application of mallotoxin on the membrane patches over an extended time (e.g., more than 15 minutes) had little further effect on the G-V relationship, but the membrane patches appeared to be less stable and more leaky, presumably due to some non-specific effects of the hydrophobic mallotoxin on cell membranes. Therefore, all electrophysiological data presented in this study were collected within 5–10 minutes of mallotoxin application.

The γ 2 subunit also exerted a significant attenuating effect on mallotoxin. Mallotoxin produced a shift of only -28 mV in BK channel voltage-dependence of activation, from 61 \pm 3 mV (n = 8) for the untreated BK α γ 2 channel to 33 \pm 2 mV (n = 3) for the treated channels (Fig. 3a; Table 1). The presence of the γ 3 subunit, like the γ 1 subunit, abolished most of the channel-activating effect of mallotoxin, with a difference of only -15 mV in the BK α γ 3 channel V_{1/2} values in the presence (100 \pm 5 mV, n = 4) and absence (115 \pm 2 mV, n = 6) of mallotoxin (Fig. 3b; Table 1). The γ 4 subunit, which was barely effective in BK modulation, had no significant effect on mallotoxin's channel-activating effect. Mallotoxin induced a significant shift of -69 mV in BK channel V_{1/2}, from 154 \pm 3 mV (n = 7) for the untreated BK α γ 4 channels to 85 \pm 8 mV (n = 4) for the treated channels (Fig. 3c; Table 1), a result similar to that with BK α channels in the absence of γ subunits.

Inverse correlation of the activating effects of mallotoxin on BK channels and the modulatory function of most BK γ 1 mutants.

Our group recently reported that a peptide region of the γ subunit encompassing the single transmembrane segment and its neighboring intracellular polybasic charged cluster are responsible for the difference in modulatory functions of the four γ subunits²⁴ and also are the key determinants for the γ 1 subunit's modulatory function²⁵. To define the relationship between the γ 1 subunit's modulatory function and the channel-activating effect of mallotoxin, we analyzed the modulatory effects of mallotoxin in the presence of various γ 1 mutants, which displayed a range of different capacities for shifting the BK channel V_{1/2} toward the hyperpolarizing direction via mutations in the transmembrane or C-tail region²⁵. It is notable that, for most of the 13 γ 1 mutants examined, their V_{1/2}-shifting capacities were inversely correlated with the mallotoxin's V_{1/2}-shifting effect in the mutant's presence (Fig. 4; Table 1). The BK channel-activating effect of mallotoxin was not significantly attenuated by most of the loss-of-function mutants. The F273S/P270V, F273S/L274A, F273S/




Figure 2. Effects of mallotoxin on BK channels in the presence of the auxiliary $\gamma 1$ subunit. (a,b) Voltage protocols (left) and the representative activation (a) and deactivation (b) currents of BK $\alpha\gamma 1$ channel complexes at different voltages in the absence (–MTX) and presence of mallotoxin (+MTX). (c) Voltage-dependence of BK channel activation. (d,e) Voltage-dependence of the kinetics of channel activation (d) and deactivation (e).

V275A, $\Delta 4R$, and $\Delta tailN^{291-298}$ mutants lost $\geq 87\%$ of the $V_{1/2}$ -shifting capacity of the wild-type (WT) $\gamma 1$ subunit. Consequently, they were largely unable to antagonize the mallotoxin-induced shift in BK channel $V_{1/2}$ ($\Delta V_{1/2} = -52$ to -102 mV). The L274A, V275A, and $\Delta 3R$ mutants retained more than 85% of the $V_{1/2}$ -shifting capacity of the $\gamma 1$ -WT subunit. Like the $\gamma 1$ -WT subunit, they also largely suppressed the BK channel-activating effect of mallotoxin, with nearly no noticeable shift in $V_{1/2}$ ($|\Delta V_{1/2}| \leq 5$ mV) between the treated and untreated membrane patches. The four partially functional mutants, $\gamma 1/\gamma 4$ -linker&tail, $\gamma 1/\gamma 4$ -TMa, $\gamma 1/\gamma 4$ -TMc, and S272V, shifted BK channel $V_{1/2}$ in a range of -67 to -100 mV (48–75% of the capacity of the $\gamma 1$ -WT subunit) in the absence of mallotoxin. Their presence only partially antagonized the BK channel-activating effect of mallotoxin, and the resultant $\Delta V_{1/2} (+/- \text{ mallotoxin})$ values were -44 mV, -34 mV, -30 mV, and -40 mV, respectively (Fig. 4b; Table 1), which are close to a line drawn between the effects of mallotoxin on the BK α channels and on the BK $\alpha\gamma 1^{WT}$ channels (Fig. 4c; Table 1). These results showed that the $\gamma 1$ subunit antagonized the mallotoxin's channel-activating effect in a manner that is largely dependent on its own modulatory function. However, the F273S/ $\Delta 3R$ mutant appeared to be exceptional which was inactive in BK channel modulation but still effective in attenuating the BK channel-activating effect of mallotoxin, with a resultant $\Delta V_{1/2} (+/- \text{ mallotoxin})$ value of only 29 mV.

Mallotoxin irreversibly disengaged the $\gamma 1$ -F273S mutant from BK channels. We recently reported that the F273S mutant of the $\gamma 1$ subunit ($\gamma 1$ -F273S) resulted in weakened association between the $\gamma 1$ and BK α subunits, which caused a small fraction ($\sim 20\%$) of the channels to be high voltage-activated channels with a $V_{1/2} \geq 100$ mV²⁵. We found that the total expression of F273S was not significant different from WT but overexpression of the F273S mutant enhanced its modulatory function by increasing the portion of low $V_{1/2}$ channel from 79 to 91%, suggesting a likely decreased binding affinity of the F273S mutant to BK α ²⁵. Interestingly, in contrast to the slightly activating effect of mallotoxin on the BK channels in the presence of $\gamma 1$ -WT (Fig. 5a; Table 1), mallotoxin exerted a significant inhibitory effect on BK channels initially complexed with $\gamma 1$ -F273S by shifting the G-V relationship toward to the positive voltage direction with a $V_{1/2}$ value of 96 ± 5 mV which is close to that of the mallotoxin-modulated BK α alone channels (Fig. 5b; Table 1). We further found that the




Figure 3. Effects of mallotoxin (MTX) on voltage-dependence of BK channel activation in the presence of the auxiliary subunits $\gamma 2$ (a), $\gamma 3$ (b), or $\gamma 4$ (c). The voltage protocol used for data recording is the same as in Figs 1a and 2a. Representative activation currents of the BK $\alpha\gamma 1$ -3 channels are in the absence and presence of mallotoxin and the plotted voltage-dependence of BK channel activation are shown on the left, middle, and right sides, respectively.

modulatory function of the F273S mutant was, indeed, largely irreversibly lost upon pre-application (5 min) and then withdrawal of mallotoxin, with a resultant $V_{1/2}$ of 147 ± 3 mV, which is close to that of the BK α channel alone (Fig. 5a; Table 1). In contrast, there was no drastic difference for the BK channel's G-V relationship before mallotoxin treatment and after mallotoxin withdrawal in the presence of $\gamma 1$ -WT (Fig. 5a; Table 1). As expected from a competition of mallotoxin with $\gamma 1$ -F273S for BK channel modulation, we found that a reduction in the concentration of mallotoxin to 0.5 or 0.25 μM during the pre-treatment resulted in a great decrease in the loss of the channel modulatory function of $\gamma 1$ -F273S. After mallotoxin withdrawal, the channel modulatory effect of the $\gamma 1$ -F273S mutant was largely retained in 3 out of 8 patches for cells pretreated with 0.5 μM mallotoxin (Fig. 5c) and fully retained in 4 patches and only partially lost in 1 patch (totally 5 patches) for cells pretreated with 0.25 μM mallotoxin (Fig. 5d).

To determine whether mallotoxin can affect the physical association between the $\gamma 1$ and BK α subunits, we performed immunoprecipitation and immunoblot analyses of the BK $\alpha\gamma 1^{\text{WT}}$ and BK $\alpha\gamma 1^{\text{F273S}}$ channel complexes in the absence and presence of mallotoxin (Fig. 5e and f). The BK channel complex was immunoprecipitated by immobilized anti-BK α antibody that was covalently crosslinked to the agarose beads and then immunoblotted with anti-BK α and anti-V5 antibodies for the BK α subunit and the V5-tagged $\gamma 1$ subunit, respectively. Similar to the recently reported weakened association between the $\gamma 1$ -F273S mutant and BK α ²⁵, only ~20% of the $\gamma 1$ -F273S mutant protein, as compared to $\gamma 1$ -WT, remained associated with BK α in the isolated channel complexes. Mallotoxin treatment caused a further >60% reduction in the BK α -associated $\gamma 1$ -F273S mutant protein




Figure 4. The relationship between the $\gamma 1$ subunit and mallotoxin (MTX) in determining their efficacies in shifting the BK channel $V_{1/2}$. (a) The schematic structures of the used chimeric $\gamma 1/\gamma 4$ mutants and the location of mutations in amino acid sequence. (b) The shifts in BK channel $V_{1/2}$ ($\Delta V_{1/2}$) induced by wild-type (WT) and mutant $\gamma 1$ subunits in the absence of mallotoxin (left) and by mallotoxin in the presence of the corresponding $\gamma 1$ proteins (right). (c) The relationship between the values of $\Delta V_{1/2}$ induced by $\gamma 1$ and mallotoxin. For reference, a straight line was drawn between the two data points of BK channels in the absence of $\gamma 1$ and in the presence of $\gamma 1$ -WT.

in the isolated channel complexes. However, no significant difference in the association of the $\gamma 1$ -WT subunit to BK α was observed in the isolated channel complexes in the absence or presence of mallotoxin treatment.

Discussion

In this study, we found that mallotoxin, when applied extracellularly on excised membrane patches, shifted the half-activation voltage ($V_{1/2}$) of BK channels by -72 mV in the absence of γ subunits. The channel-activating effect of mallotoxin was largely abolished or greatly attenuated in the presence of subunits $\gamma 1$, $\gamma 2$, and $\gamma 3$, with resultant $\Delta V_{1/2}$ (+/- mallotoxin) values of -9 , -28 , and -15 mV, respectively. Most of the examined $\gamma 1$ mutant subunits antagonized mallotoxin's channel-activating effect in a manner that was largely dependent on its own modulatory function. However, mallotoxin caused an irreversible functional and structural disengagement of the $\gamma 1$ -F273S mutant from BK channels.

Mallotoxin has been found to affect numerous cellular events, including ion channel activation^{23,26}, many protein kinase signaling pathways^{27,28}, induction of mitochondrial uncoupling²⁹, and autophagy³⁰. The previous electrophysiological studies on the effects of mallotoxin on BK channels were mostly recorded in whole cell configuration^{10,23}, which could be complicated by mallotoxin's effect on intracellular events, such as many kinase activities^{27,28}. The present study relied on patch-clamp recording on excised membrane patches to minimize interference from mallotoxin-induced intracellular events. The observed ~ 70 mV shift in the voltage-dependence of BK α channel gating in HEK-293 cells in the presence of $2 \mu\text{M}$ mallotoxin is close to that observed in whole cell recording in CHO-K1 cells (70 mV at $5 \mu\text{M}$)¹⁰ but smaller than that originally reported through whole cell recording in HEK-293 cells (>100 mV at $0.5 \mu\text{M}$)²³. The discrepancy could be due to a difference in cell recording configurations. We also confirmed that, in excised membrane patches, the $\gamma 1$ subunit largely attenuated the BK channel-activating effect of mallotoxin ($\Delta V_{1/2} = 9$ mV at $2 \mu\text{M}$), a result similar to that previously observed in whole cell recording of native channels in parotid acinar cells ($\Delta V_{1/2} = 6$ mV at $5 \mu\text{M}$) and recombinant BK α - $\gamma 1$ channels in CHO-K1 cells ($\Delta V_{1/2} = 17$ mV at $5 \mu\text{M}$)¹⁰. Our observation that the $\gamma 2$ and $\gamma 3$ subunits also largely attenuated the BK channel-activating effect of mallotoxin suggests that the pharmacological properties of BK channels exerted by the three different auxiliary γ subunits are similar. Therefore, the results we obtained on excised membrane patches confirmed previous findings on whole cell recording^{10,23} and further demonstrated that mallotoxin directly activated BK channels in the absence of the auxiliary γ subunit but became largely ineffective in the presence of the auxiliary $\gamma 1$, $\gamma 2$, and $\gamma 3$ subunits.

To probe the mechanism underlying the drastic difference in the activating efficacy of mallotoxin on the BK α and BK α $\gamma 1$ channels, we examined the effects of mallotoxin on BK channels complexed with various $\gamma 1$ mutants. Our finding that, for most $\gamma 1$ subunit mutants tested, the BK channel $V_{1/2}$ -shifting capacity was correlated with their antagonizing effects on the mallotoxin's channel-activating function. One may argue that channel-activating function of mallotoxin is simply dependent on the extent of channel-stimulation or -activation by any other activator through an uncompetitive mechanism which involves two independent binding/active sites. For example, the $\gamma 1$ subunit might affect the efficacy of mallotoxin by maximally affecting the same gating parameter, e.g.,




Figure 5. Mallotoxin disengaged γ 1-F273S mutant from BK channel modulation. (a) Mallotoxin (MTX) at 2 μ M had little effect on the G-V relationship of BK channels formed by BK α and the wild-type γ 1 subunit (γ 1-WT) during treatment and after withdrawal of mallotoxin. (b) Mallotoxin at 2 μ M induced an irreversible drastic loss of the channel-modulatory function of the γ 1-F273S mutant. (c,d) Treatment of cells with a low concentration of mallotoxin produced less loss of the channel-modulatory function of the γ 1-F273S mutant. A double Boltzmann function was used to generate G-V fitting curves for the data of no mallotoxin treatment in (b) and of mallotoxin withdrawal for 3 patches in (c) and for all patches in (d). (e) Immunoblot analysis of the BK α γ 1 channel complex immunoprecipitated with anti-BK α antibody in the absence or presence of mallotoxin. BK α and γ 1-V5 were immunoblotted with anti-BK α and anti-V5 antibody, respectively. (f) Densitometric analysis of the relative content of γ 1-V5. Data were collected from six independent experiments. For each experiment, the intensity of the γ 1-V5 band was first normalized to that of the BK α band and then to the mean of untreated and treated WT samples. * $p < 0.05$.

the allosteric coupling factor between the voltage sensors and channel pore gate^{8,10}. However, an uncompetitive mechanism can be ruled out by three lines of evidence. First, the effect of mallotoxin on the BK α channel was largely unaffected by another activator, the ligand Ca²⁺²³. Second, the γ 3 subunit and the γ 1-F273S/ Δ 3R mutant,

which are a much weaker modulator than $\gamma 1$ -WT or a nearly loss-of-function modulator, respectively, were still effective in suppressing mallotoxin's channel-activating effect. Third, as discussed next, mallotoxin can structurally compete off the $\gamma 1$ -F273S mutant from BK channels.

We observed that the $\gamma 1$ subunit and mallotoxin mutually affected each other on their modulatory effects on BK channels. The mallotoxin-induced shifts in the G-V curves toward the depolarizing direction in the initial presence of the F273S mutant can be explained by mallotoxin-induced displacement of the F273S mutant protein from the BK channels, evidenced by the irreversible loss of BK channel modulation by the F273S mutant upon withdrawal of mallotoxin and also the mallotoxin treatment-induced dissociation of the F273S mutant protein in the immunoprecipitated channel complexes. The F273 residue was predicted to be near or in the middle of the single transmembrane segment of the $\gamma 1$ subunit and was found to play an important role in the $\gamma 1$ subunit's association with and modulatory function on BK channels²⁵. Therefore, the $\gamma 1$ subunit and mallotoxin can mutually affect their modulatory efficacy on BK channels in a specific $\gamma 1$ transmembrane residue-dependent manner. This result can be explained by either a competitive or a noncompetitive mechanism between the γ subunit and mallotoxin for their mutual influence on BK channel modulation. For a competitive mechanism, the γ subunit and mallotoxin sterically compete on overlapping binding or active sites on BK α . The F273S mutation in the middle of the $\gamma 1$ subunit's transmembrane domain might create room and/or additional interaction to allow the mallotoxin to bind and thus to irreversibly disengage the $\gamma 1$ subunit from BK channels. For a noncompetitive mechanism, the binding/active sites of the $\gamma 1$ subunit and mallotoxin on BK α are distinct but allosterically and negatively coupled. The ability of mallotoxin to functionally and structurally disengage the $\gamma 1$ -F273S mutant could be simply due to the latter's weakened association to BK α . For either of these two mechanisms, these $\gamma 1$ mutants with partially reduced G-V shifting capacities likely exert less influence on the mallotoxin's binding on BK α than the WT and thus allow mallotoxin to regain influence on BK channel gating. More studies will be needed to distinguish these two different mechanisms.

The mechanism for the decreased association of the F273S mutant to BK α remains unclear. We found that overexpression of the F273S mutant enhanced its modulatory function by increasing the portion of low $V_{1/2}$ channel from 79 to 91%, suggesting the possibility of either a decreased expression or a comprised binding affinity to BK α .

Materials and Methods

Heterologous expression of BK α and γ proteins in HEK-293 cells. Recombinant cDNA constructs of human BK α (hSlo), $\gamma 1$ -4 subunits, and $\gamma 1$ mutants were used for heterologous expression in HEK-293 cells. As described previously^{8,9,24}, fusion cDNA constructs that encode precursor fusion proteins of human BK α and C-terminal-tagged BK γ proteins were generated with the pCDNA6 vector and used to facilitate the co-translational assembly of BK $\alpha\gamma$ protein complexes after endogenous cleavage by peptidases at the linker (signal peptide) region in the mature proteins. HEK-293 cells were obtained from ATCC. The cells were transfected with plasmids using Lipofectamine 2000 (Invitrogen) and subjected to electrophysiological assays 16–24 h after transfection.

Electrophysiology. Inside-out BK channel currents were acquired at room temperature using EPC-10 (HEKA). Symmetric internal and external patch-clamp recording solutions contained 136 mM KMeSO₃, 4 mM KCl, and 20 mM HEPES (pH = 7.20). The external solution was supplemented with 2 mM MgCl₂, and the internal solution was supplemented with 5 mM HEDTA without Ca²⁺ to achieve virtual 0 [Ca²⁺]_i. All patch-clamp data were collected at virtual 0 [Ca²⁺]_i. Steady-state activation was expressed as the normalized conductance (G/G_{max}) calculated from the relative amplitude of the tail currents (deactivation at -120 mV). The voltage of half-maximal activation ($V_{1/2}$) and the equivalent gating charge (z) were obtained by fitting the relations of G/G_{max} vs. voltage with the single Boltzmann function $G/G_{max} = 1/(1 + e^{-zF(V-V_{1/2})/RT})$ or with the double Boltzmann function $G/G_{max} = a/(1 + e^{-zaF(V-V_{a1/2})/RT}) + (1 - a)/(1 + e^{-zbF(V-V_{b1/2})/RT})$. Mallotoxin was purchased from Sigma-Aldrich, stored at -20 °C in aliquots in DMSO, and freshly diluted and applied in the external recording solution at a final concentration of 2 μ M. For mallotoxin withdrawal experiments, the cells were pretreated with 2 μ M mallotoxin for 5 minutes in phosphate-buffered saline (PBS) buffer (pH 7.4) and then transferred to mallotoxin-free solution and subjected immediately to patch-clamp recording.

Immunoprecipitation and immunoblotting. HEK-293 cells expressing the BK $\alpha\gamma 1^{WT}$ or BK $\alpha\gamma 1^{F273S}$ channel complexes were preincubated with 2 μ M mallotoxin in PBS buffer (pH 7.4) for 5 minutes. The channel complexes were then solubilized from cell membranes with 1% dodecyl maltoside (DDM) in Tris-buffered saline (TBS) buffer (50 mM Tris, 150 mM NaCl, pH 7.6) supplemented with 2 μ M mallotoxin. After centrifugation at 17,000 g for 10 minutes, the solubilized channel complexes in the supernatant were incubated for 2 hours with mouse monoclonal anti-BK α antibody (University of California-Davis/NIH Neuromab facility) that was covalently crosslinked to protein-A/G agarose beads (Thermo Fisher Scientific). The captured protein complexes were washed three times with TBS buffer supplemented with 1% DDM and 2 μ M mallotoxin, eluted with SDS-PAGE sample buffer, and then loaded directly to 12% SDS-PAGE gel to be separated by electrophoresis. Resolved proteins were transferred to PVDF membranes (Thermo Fisher Scientific) and probed by a mouse monoclonal anti-V5 antibody (1:10000, Invitrogen) for V5-tagged γ subunits and a mouse monoclonal anti-BK α antibody (1:1000, University of California-Davis) for BK α . The intensities of the protein bands were analyzed with ImageJ software (US National Institutes of Health).

Statistical analyses. Experimental values are reported as means \pm standard error of the mean. The Student *t*-test was used for comparison of two groups. Differences were considered statistically significant at $p < 0.05$.

References

- Brayden, J. E. & Nelson, M. T. Regulation of arterial tone by activation of calcium-dependent potassium channels. *Science* **256**, 532–5 (1992).
- Shao, L. R., Halvorsrud, R., Borg-Graham, L. & Storm, J. F. The role of BK-type Ca^{2+} -dependent K^+ channels in spike broadening during repetitive firing in rat hippocampal pyramidal cells. *J Physiol* **521** Pt 1, 135–46 (1999).
- Golding, N. L., Jung, H. Y., Mickus, T. & Spruston, N. Dendritic calcium spike initiation and repolarization are controlled by distinct potassium channel subtypes in CA1 pyramidal neurons. *J Neurosci* **19**, 8789–98 (1999).
- Hu, H. *et al.* Presynaptic Ca^{2+} -activated K^+ channels in glutamatergic hippocampal terminals and their role in spike repolarization and regulation of transmitter release. *J Neurosci* **21**, 9585–97 (2001).
- Raffaelli, G., Saviane, C., Mohajerani, M. H., Pedarzani, P. & Cherubini, E. BK potassium channels control transmitter release at CA3-CA3 synapses in the rat hippocampus. *J Physiol* **557**, 147–57 (2004).
- Ramanathan, K., Michael, T. H., Jiang, G. J., Hiel, H. & Fuchs, P. A. A molecular mechanism for electrical tuning of cochlear hair cells. *Science* **283**, 215–7 (1999).
- Gessner, G. *et al.* BKCa channels activating at resting potential without calcium in LNCaP prostate cancer cells. *J Membr Biol* **208**, 229–40 (2005).
- Yan, J. & Aldrich, R. W. LRRC26 auxiliary protein allows BK channel activation at resting voltage without calcium. *Nature* **466**, 513–6 (2010).
- Yan, J. & Aldrich, R. W. BK potassium channel modulation by leucine-rich repeat-containing proteins. *Proc Natl Acad Sci USA* **109**, 7917–22 (2012).
- Almassy, J. & Begenisich, T. The LRRC26 protein selectively alters the efficacy of BK channel activators. *Mol. Pharmacol.* **81**, 21–30 (2012).
- Manzanares, D. *et al.* IFN-gamma-mediated reduction of large-conductance, Ca^{2+} -activated, voltage-dependent K^+ (BK) channel activity in airway epithelial cells leads to mucociliary dysfunction. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **306**, L453–62 (2014).
- Manzanares, D. *et al.* Airway Surface Dehydration by Transforming Growth Factor beta (TGF-beta) in Cystic Fibrosis Is Due to Decreased Function of a Voltage-dependent Potassium Channel and Can Be Rescued by the Drug Pirfenidone. *J. Biol. Chem.* **290**, 25710–6 (2015).
- Evanson, K. W., Bannister, J. P., Leo, M. D. & Jaggari, J. H. LRRC26 Is a Functional BK Channel Auxiliary gamma Subunit in Arterial Smooth Muscle Cells. *Circ. Res.* **115**, 423–31 (2014).
- Zhang, J. & Yan, J. Regulation of BK channels by auxiliary gamma subunits. *Front Physiol* **5**, 401 (2014).
- Yan, J. & Aldrich, R. W. BK potassium channel modulation by leucine-rich repeat-containing proteins. *Proc. Natl. Acad. Sci. USA* **109**, 7917–22 (2012).
- Shi, J. *et al.* Mechanism of magnesium activation of calcium-activated potassium channels. *Nature* **418**, 876–80 (2002).
- Tang, X. D., Santarelli, L. C., Heinemann, S. H. & Hoshi, T. Metabolic regulation of potassium channels. *Annu Rev Physiol* **66**, 131–59 (2004).
- Bentzen, B. H., Olesen, S. P., Ronn, L. C. & Grunnet, M. BK channel activators and their therapeutic perspectives. *Front Physiol* **5**, 389 (2014).
- Nardi, A. & Olesen, S. P. BK channel modulators: a comprehensive overview. *Curr Med Chem* **15**, 1126–46 (2008).
- Torres, Y. P., Granados, S. T. & Latorre, R. Pharmacological consequences of the coexpression of BK channel alpha and auxiliary beta subunits. *Front Physiol* **5**, 383 (2014).
- Li, Q. & Yan, J. Modulation of BK Channel Function by Auxiliary Beta and Gamma Subunits. *Int Rev Neurobiol* **128**, 51–90 (2016).
- Mitra, R. & Kapoor, L. D. Kamala—the national flower of India—its ancient history and uses in Indian medicine. *Indian J Hist Sci* **11**, 125–32 (1976).
- Zakharov, S. I., Morrow, J. P., Liu, G., Yang, L. & Marx, S. O. Activation of the BK (SLO1) potassium channel by mallotoxin. *J Biol Chem* **280**, 30882–7 (2005).
- Li, Q., Fan, F., Kwak, H. R. & Yan, J. Molecular basis for differential modulation of BK channel voltage-dependent gating by auxiliary gamma subunits. *J Gen Physiol* **145**, 543–54 (2015).
- Li, Q., Guan, X., Yen, K., Zhang, J. & Yan, J. The single transmembrane segment determines the modulatory function of the BK channel auxiliary gamma subunit. *J Gen Physiol* **147**, 337–51 (2016).
- Zeng, H. *et al.* Mallotoxin is a novel human ether-a-go-go-related gene (hERG) potassium channel activator. *J Pharmacol Exp Ther* **319**, 957–62 (2006).
- Bain, J. *et al.* The selectivity of protein kinase inhibitors: a further update. *Biochem J* **408**, 297–315 (2007).
- Davies, S. P., Reddy, H., Caivano, M. & Cohen, P. Specificity and mechanism of action of some commonly used protein kinase inhibitors. *Biochem J* **351**, 95–105 (2000).
- Soltoff, S. P. Rotlerin is a mitochondrial uncoupler that decreases cellular ATP levels and indirectly blocks protein kinase C delta tyrosine phosphorylation. *J Biol Chem* **276**, 37986–92 (2001).
- Hasima, N. & Ozpolat, B. Regulation of autophagy by polyphenolic compounds as a potential therapeutic strategy for cancer. *Cell Death Dis* **5**, e1509 (2014).

Acknowledgements

This work was supported by the National Institutes of Health National Institute of Neurological Disorders and Stroke (Grant NS078152).

Author Contributions

J.Y. and X.G. designed the study, analyzed data and wrote the paper. X.G. performed electrophysiological and biochemical experiments. Q.L. constructed plasmids and mutants.

Additional Information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Guan, X. *et al.* Relationship between auxiliary gamma subunits and mallotoxin on BK channel modulation. *Sci. Rep.* **7**, 42240; doi: 10.1038/srep42240 (2017).

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2017