

Inhibitory activity of stilbenes against filamentous fungi

Luce Mattio,¹ Giorgia Catinella,¹

Marcello Iriti,² Lisa Vallone³

¹Department of Food, Environmental and Nutritional Sciences, University of Milan; ²Department of Agricultural and Environmental Sciences, University of Milan; ³Department of Health, Animal Science and Food Safety “Carlo Cantoni”, University of Milan (VESPA), Italy

Abstract

Stilbenoids (resveratrol and its derivatives) are secondary metabolites produced by plants as defence mechanism to microbial infection. These compounds are known for their anti-inflammatory action and health benefits in preventing a wide range of disorders (e.g. cancer and cardiovascular diseases). However, their antimicrobial properties are less investigated. A series of 8 stilbenoid compounds were synthesized and their antifungal activity against 19 wild strains of filamentous fungi and yeasts (isolated from the environment and food) was tested *in vitro*. Using an agar diffusion assay, compounds were tested at the concentration of 100 µg/ml on filamentous fungi and yeasts at 10⁴ CFU/ml. The results showed that tested derivatives possess moderate antifungal activity: in particular, monomeric stilbenoids 3'-hydroxy-pterostilbene and piceatannol, and dimeric stilbenoids (±)-trans- δ -viniferin and pallidol were active against mycotoxigenic fungi.

Introduction

Stilbenoids are among the most important classes of phytoalexins produced by 72 plant species belonging to 31 genera, with a particular emphasis on stilbenes from the *Vitaceae* (Valletta *et al.*, 2021; Jaillon *et al.*, 2007). Over 400 different stilbenoids are currently known (El Khawand *et al.*, 2018), mostly derived from *trans*-resveratrol (3,5,4'-trihydroxy-*trans*-stilbene), although different structures can be found in some plant families (Chong *et al.*, 2009).

Stilbenoids are mainly involved in constitutive and inducible protection of the plant against biotic (phytopathogenic microorganisms and herbivores) and abiotic (e.g. UV radiation and tropospheric ozone) stresses (Chong *et al.*, 2009; Jeandet *et al.*,

2010), due to their antibiotic and antioxidant activities (Valletta *et al.*, 2021). Among stilbenoids, resveratrol is the most investigated and its health benefits as anti-inflammatory, anticancer, estrogenic, neuroprotective, cardio protective, anti-atherosclerotic, anti-aging, anti-diabetic, anti-osteoporosis and anti-obesity agent have been documented in several preclinical (*in vitro/in vivo*) studies (Valletta *et al.*, 2021).

In the last decades, the interest in resveratrol has been amplified since its presence in wine was indicated as a possible explanation for the “French paradox”, i.e. the reduced risk of cardiovascular disease associated to moderate, regular consumption of red wine at main meals for people consuming a diet rich in saturated fats (Catalgol *et al.*, 2012; Jeandet *et al.*, 2021).

Currently, resveratrol has been exploited in pharmaceutical, cosmetic and food industries. In this latter area, possible applications as antimicrobial agent in the conservation of food are under evaluation (Oh *et al.*, 2018; Ma *et al.*, 2018). The *in vitro* antimicrobial properties of resveratrol are widely known (Albert *et al.*, 2011; Chalal *et al.*, 2014), although the mechanism of action on pathogenic and food-borne bacteria, fungi and yeasts is not yet fully understood (Lee *et al.*, 2015).

Resveratrol exhibited inhibitory activity against yeasts and filamentous fungi (Seppanen *et al.*, 2019) such as *Botrytis cinerea* (Adrian *et al.*, 1997; Hoss *et al.*, 1990; Paul *et al.*, 1998; Sarig *et al.*, 1997), *Rhizopus stolonifer* (Valletta *et al.*, 2021), *Phomopsis viticola* (Hoss *et al.*, 1990), *Fusarium nivale* (Bala *et al.*, 1999), *Saccharomyces cerevisiae*, *Penicillium expansum* and *Aspergillus niger* (Seppanen *et al.*, 2019; Valletta *et al.*, 2021; Adrian *et al.*, 1997). Conversely, Weber *et al.* (Weber *et al.*, 2011), showed that resveratrol was not active on several *Candida* species (Weber *et al.*, 2011; Shevelev *et al.*, 2018), though recent reports suggested that resveratrol derivatives could inhibit *Candida* species as well (Lee *et al.*, 2015; Shevelev *et al.*, 2020).

Recently, other monomeric and oligomeric stilbenoid derivatives including pterostilbene, pinosylvin, piceatannol and viniferins have attracted the attention of researchers. Pterostilbene and *trans*- ϵ -viniferin have been found to be 5-fold more active than resveratrol as antifungal agents, indicating their high antimicrobial potential (Chalal *et al.*, 2014; Houillé *et al.*, 2014). These compounds were tested *in vitro* against *B. cinerea*, and pterostilbene, in particular, inhibited conidial germination and *in vitro* mycelium growth more effectively than resveratrol, indicating that methylation

Correspondence: Lisa Vallone, Dipartimento VESPA, Via dell'Università n. 9, Lodi (LO), Italy.

Tel. +39.250334312.

E-mail: lisa.vallone@unimi.it

Key words: Stilbenes, Resveratrol, Antimicrobial activity, Food-borne fungi.

Acknowledgments: The work of Giorgia Catinella has been partially funded by Fondazione F. Ili Confalonieri (PhD Scholarship).

Contributions: The authors contributed equally.

Conflict of interest: The authors declared no potential conflict of interest.

Received for publication: 31 July 2020.

Revision received: 15 November 2020.

Accepted for publication: 18 November 2020.

This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0).

©Copyright: the Author(s), 2021

Licensee PAGEPress, Italy

Italian Journal of Food Safety 2021; 10:8461

doi:10.4081/ijfs.2021.8461

of -OH groups could have a role in the antifungal activity. Schouten *et al.* (Schouten *et al.*, 2002) demonstrated that resveratrol, though not toxic to *B. cinerea*, is converted into a fungitoxic derivative by a specific fungal laccase (Caruso *et al.*, 2011).

A possible mode of action of stilbenoids may involve membrane peroxidation (Lee *et al.*, 2017). Pterostilbene caused destruction of the endoplasmic reticulum, and the nuclear and mitochondrial membranes in *B. cinerea* dormant conidia. A positive correlation between antifungal activity of natural and synthetic stilbenoids and their hydrophobicity was found, suggesting that pterostilbene is more active than the less hydrophobic resveratrol due to its increased diffusion through the cytoplasmic membrane (Caruso *et al.*, 2011).

Owing to the need of alternative compounds to control food spoilage and contamination, the aim of this study was to assay *in vitro* the antimicrobial activity of a small collection of stilbenoid monomers and dimers against filamentous fungi and yeasts isolated from the environment and food of animal origin.

Materials and methods

Sixteen fungal strains, *Alternaria alter-*

nata, *Aspergillus flavus*, *Aspergillus niger*, *Aspergillus ochraceus*, *Aspergillus terreus*, *Byssoschlamys nivea*, *Botrytis cinerea*, *Fusarium graminearum*

Fusarium verticillioides, *Geotrichum candidum*, *Mucor circinelloides*, *Penicillium expansum*

Penicillium italicum, *Penicillium roqueforti*, *Rhizopus nigricans* and 3 yeast strains *Candida albicans*, *Candida parapsilosis*, *Malassezia pachydermatis* were grown to assess the inhibitory activity of 4 monomeric stilbenoids (i.e. resveratrol, piceatannol, pterostilbene, 3'-hydroxy-pterostilbene) and 4 dimers (i.e. (±)-trans- δ -viniferin, (±)-trans- ϵ -viniferin, pallidol (±)-pterostilbene-trans-dihydrodimer) (Figure 1). Stilbenoids were dissolved in DMSO to obtain a stock solution of 10 mg/ml. Wild strains were used instead of ATCC strain because more aggressive and less tamed to tested compounds.

The bioassay was carried out on Sabouraud agar medium, by paper disk diffusion assay. From microbanks, all the mycetes were first suspended into M₂ broth and incubated for 5 days at 25°C. A spectrophotometer was used to adjust the final

cell concentration at 10⁴ cfu/ml by reading the OD at 600 nm.

Then, 100 μ l of the microbial suspensions were spread on Sabouraud agar medium. The 6-mm-diameter, sterile disks impregnated with 10 μ l of stilbenoid compounds at the concentration of 100 μ g/ml were placed on the inoculated agar. The inoculated plates were incubated at 25°C for 7 days. As positive controls, cyclopiroxolamine and tebuconazole (1 μ g/disk) were used. Inhibitory activity was determined by measuring the zone of inhibition.

Results

The inhibitory activity of the tested stilbenoids varied according to the different strains. Piceatannol showed an antifungal activity against *P. roqueforti*, *A. flavus*, *P. italicum*, *A. terreus*, *G. candidum*, *F. verticillioides* and *C. parapsilosis*. 3'-Hydroxy-pterostilbene affected the growth of *C. parapsilosis*, *P. roqueforti*, *A. ochraceus*, *A. flavus* and *C. albicans*. Pallidol inhibited the growth of *P. italicum* and *F. verticil-*

loide. Finally, (±)-trans- δ -viniferin and (±)-pterostilbene-trans-dihydrodimer were effective only on one strain, *A. flavus* and *F. verticillioides*, respectively. Overall, *A. flavus* was sensitive to piceatannol, δ -viniferin and 3'-hydroxy-pterostilbene; *F. verticillioides* was sensitive to (±)-pterostilbene-trans-dihydrodimer, pallidol and piceatannol; *P. roqueforti* was sensitive to piceatannol and 3'-hydroxy-pterostilbene; *P. italicum* was sensitive to pallidol and piceatannol and *C. parapsilosis* was sensitive to 3'-hydroxy-pterostilbene and piceatannol; *A. terreus* and *G. candidum* were sensitive to piceatannol; *A. ochraceus* and *C. albicans* were sensitive to 3'-hydroxy-pterostilbene (Table 1). Values are mean inhibition zone (mm) \pm S.D. of three replicates; positive controls (tebuconazole and cyclopiroxolamine) > 20 mm.

Discussion

Results indicated that the antifungal activity of selected stilbenoids was strictly related to the chemical structure of the




Figure 1. Structure of selected stilbenoids, tebuconazole and cyclopiroxolamine.

Table 1. Inhibitory activity of selected stilbenoids against filamentous fungi and yeasts.

Fungi and yeasts	piceatannol	3'-hydroxy-pterostilbene	(±)-trans-δ-viniferin	(±)-pterostilbene-trans-dihydrodimer	pallidol
<i>Alternaria alternata</i>	n.i.	n.i.	n.i.	n.i.	n.i.
<i>Aspergillus flavus</i>	20 ± 0.0	3 ± 0.0	5 ± 0.2	n.i.	n.i.
<i>Aspergillus niger</i>	n.i.	n.i.	n.i.	n.i.	n.i.
<i>Aspergillus ochraceus</i>	n.i.	5 ± 0.1	n.i.	n.i.	n.i.
<i>Aspergillus terreus</i>	5 ± 0.1	n.i.	n.i.	n.i.	n.i.
<i>Byssosclamyces nivea</i>	n.i.	n.i.	n.i.	n.i.	n.i.
<i>Botrytis cinerea</i>	n.i.	n.i.	n.i.	n.i.	n.i.
<i>Fusarium graminearum</i>	n.i.	n.i.	n.i.	n.i.	n.i.
<i>Fusarium verticillioides</i>	2 ± 0.2	n.i.	n.i.	10 ± 0.0	5 ± 0.1
<i>Geotrichum candidum</i>	5 ± 0.1	n.i.	n.i.	n.i.	n.i.
<i>Mucor circinelloides</i>	n.i.	n.i.	n.i.	n.i.	n.i.
<i>Penicillium expansum</i>	n.i.	n.i.	n.i.	n.i.	n.i.
<i>Penicillium italicum</i>	n.i.	n.i.	n.i.	n.i.	20 ± 0.2
<i>Penicillium roqueforti</i>	20 ± 0.1	6 ± 0.2	n.i.	n.i.	n.i.
<i>Rhizopus nigricans</i>	n.i.	n.i.	n.i.	n.i.	n.i.
<i>Candida albicans</i>	n.i.	2 ± 0.2	n.i.	n.i.	n.i.
<i>Candida parapsilosis</i>	2 ± 0.0	10 ± 0.1	n.i.	n.i.	n.i.
<i>Malassezia pachydermatis</i>	n.i.	n.i.	n.i.	n.i.	n.i.

n.i.: no inhibition zone formation.

molecule, although it is not possible to define a relationship between the structure of a given compound (number/position of -OH/-OMe groups) and its antimicrobial activity (Chalal *et al.*, 2014). In general, the monomeric species (in particular piceatannol) were more active against the tested fungi compared to the dimeric derivatives. Interestingly, the tested stilbenoids were more active against phytopathogenic fungi (i.e., *Aspergillus* spp., *Penicillium* spp. and *F. verticillioides*). We can speculate that this behaviour could be related to the coevolution between pathogen and host plant (and its phytoalexins). In this sense, phytopathogenic fungi would be more sensitive to the fungitoxic effects of stilbenoids.

Interestingly, on dermatophyte fungi, inhibitory activity has been demonstrated at concentrations of 25-50 µg/ml, while on *Candida albicans* it appeared evident at lower concentrations (10-20 µg/ml) (Vestergaard *et al.*, 2019).

Conclusions

In this study, some stilbene derivatives were assayed and exhibited a weak to moderate antifungal activity against a panel of fungal strains and yeasts. Monomeric stilbenoids, in particular 3'-hydroxy-pterostilbene and piceatannol, and dimeric stilbenoids (±)-trans-δ-viniferin and pallidol were active against mycotoxigenic fungi,

thus showing a promising potential as food preservatives. In further studies, these compounds could be tested at higher concentrations, in combination with other natural compounds or low-dose conventional antimicrobials. In this view, stilbenoids could contribute to reduce the risk of selecting resistant fungal strains, a relevant issue due to the global burden of antimicrobial resistance. Finally, the efficacy of these compounds could be improved by formulation, including their functionalization with nanostructures or incorporation in active packaging.

References

- Adrian M, Jeandet P, Veneau J, Weston LA, Bessis R, 1997. Biological activity of resveratrol, a stilbenic compound from grapevines, against *Botrytis cinerea*, the causal agent for gray mold. *J Chem Ecol* 23:1689-702.
- Albert S, Horbach R, Deising HB, Siewert B, Csuk R, 2011. Synthesis and antimicrobial activity of (E) stilbene derivatives. *Bioorg Med Chem* 19:5155-66.
- Bala AE, Kollmann A, Ducrot PH, Majira A, Kerhoas L, Delorme R, Einhorn J, 1999. Antifungal activity of resveratrol oligomers from *Cyphostemma crotalaroides*. *Pestic Sci* 55:206-8.
- Caruso F, Mendoza L, Castro P, Cotoras M, Aguirre M, Matsushiro B, Isaacs M, Rossi M, Viglianti A, Antonioletti R, 2011. Antifungal Activity of Resveratrol against *Botrytis cinerea* is improved using 2-Furyl Derivatives. *PLoS One* 6:10.
- Catalgol B, Batirel S, Taga Y, Ozer NK, 2012. Resveratrol: French paradox revisited. *Front Pharmacol* 3:141.
- Chalal M, Klinguer A, Echairi A, Meunier P, Vervandier-Fasseur D, Adrian M, 2014. Antimicrobial activity of resveratrol analogues. *Molecules* 19:7679-88.
- Chan MMY, 2002. Antimicrobial effect of resveratrol on dermatophytes and bacterial pathogens of the skin. *Biochem Pharmacol* 63:99-104.
- Chong J, Poutaraud A, Huguency P, 2009. Metabolism and roles of stilbenes in plants. *Plant Sci* 177:143-55.
- El Khawand T, Courtois A, Valls J, Richard T, Krisa S, 2018. A review of dietary stilbenes: sources and bioavailability. *Phytochem Rev* 17:1007-29.
- Hoss G, Blaich R, 1990. Influence of Resveratrol on Germination of Conidia and Mycelial Growth of *Botrytis cinerea* and *Phomopsis viticola*. *J Phytopathol* 129:102-10.
- Houillé B, Papon N, Boudesocque L, Bourdeaud E, Besseu S, Courdavault V, Enguehard-Gueiffier C, Delanoue G, Guerin L, Bouchara JP, Clastre M, Giglioli-Guivarch N, Guillard J, Lanoue A, 2014. Antifungal Activity of Resveratrol Derivatives against *Candida* Species. *J Nat Prod* 77:1658-

- 62.
- Jaillon O, Aury JM, Noel B, Policriti A, Clepet C, 2007. French-Italian Public Consort. Grapevine Genome Charact. The grapevine genome sequence suggests ancestral hexaploidization in major angiosperm phyla. *Nature* 449:463–73.
- Jeandet P, Delaunois B, Conreux A, Donnez D, Nuzzo V, Cordelier S, Clément C, Courot E, 2010. Biosynthesis, metabolism, molecular engineering, and biological functions of stilbene phytoalexins in plants. *Biofactors* 36:331-41.
- Jeandet P, Vannozzi A, Sobarzo-Sanchez E, Uddin MdS, Bru R, Martinez-Marquez A, Clement C, Cordelier S, Manayi A, Nabavi SF, Rasekhian M, El-Saber Batiha G, Khan H, Morkunas I, Belwal T, Jiang J, Koffaso M, Nabavi SM, 2021. Phytostilbenes as agrochemicals: biosynthesis, bioactivity, metabolic engineering and biotechnology. *Nat Prod Rep* 1:1-48.
- Lee J, Lee DG, 2015. Intracellular Mechanism of Resveratrol in *C. albicans*. *Curr Microbiol* 70:383-9.
- Lee W, Lee DG, 2017. Resveratrol induces membrane and DNA disruption via pro-oxidant activity against *Salmonella typhimurium*. *Biochem Biophys Res Commun* 489:228-34.
- Ma DSL, Tan LTH, Chan KG, Yap WH, Pusparajah P, Chuah LH, Ming LC, Khan TM, Lee LH, Goh BH, 2018. Resveratrol - Potential Antibacterial Agent against Foodborne Pathogens. *Front Pharmacol* 9:102.
- Oh WY, Shahidi F, 2018. Antioxidant activity of resveratrol ester derivatives in food and biological model systems. *Food Chem* 261:267-73.
- Paul B, Chereyathmanjiyil A, Masih I, Chapuis L, Benoît A, 1998. Biological control of *Botrytis cinerea* causing grey mould disease of grapevine and elicitation of stilbene phytoalexin (resveratrol) by a soil bacterium. *FEMS Microbiol Lett* 165:65-70.
- Sarig P, Zutkhi Y, Monjauze A, Lisker N, Ben-Arie R, 1997. Phytoalexin elicitation in grape berries and their susceptibility to *Rhizopus stolonifera*. *Physiol Mol Plant Pathol* 50:337-47.
- Schouten A, Wagemakers L, Stefanato FL, Van der Kaaij RM, Van Kan JAL, 2002. Resveratrol acts as a natural antifungal and induces self-intoxication by a specific laccase. *Mol Microbiol* 43:883–94.
- Seppanen VF, Vestergaard M, Ingmer H, 2019. Antibacterial and antifungal properties of resveratrol. *Int J Antimicrob Agents* 53:716-23.
- Shevelev AB, Isakova EP, Trubnikova EV, La Porta N, Martens S, Medvedeva OA, Trubnikov DV, Akbaev RM, Biryukova YK, Zylkova MV, Lebedeva AA, Smirnova MS, Deryabina YI, 2018. A study of antimicrobial activity of polyphenols derived from wood. *Bull Russ State Med Univ* 7:46-9.
- Shevelev AB, La Porta N, Isakova EP, Martens S, Biryukova YK, Belous AS, Sivokhin DA, Trubnikova EV, Zylkova MV, Belyakova AV, Smirnova MS, Yulia I, Deryabina YI, 2020. In Vivo Antimicrobial and Wound-Healing Activity of Resveratrol, Dihydroquercetin and Dihydromyricetin against *Staphylococcus aureus*, *Pseudomonas aeruginosa* and *Candida albicans*. *Pathogens* 9:296.
- Valletta A, Iozia LM, Leonelli F, 2021. Impact of Environmental Factors on Stilbene. *Biosynthesis. Plants* 10:90.
- Vestergaard M, Ingmer H, 2019. Antibacterial and antifungal properties of resveratrol. *Int J Antimicrob Agents* 53:716-23.
- Weber K, Schulz B, Ruhnke M, 2011. Resveratrol and its antifungal activity against *Candida* species. *Mycoses* 54:30-3.