

LETTER TO THE EDITOR

Open Access

First description of *Trypanosoma cruzi* human infection in Esmeraldas province, Ecuador

Ángel Guevara^{1*}, Juan Moreira^{1,2^}, Hipatia Criollo¹, Sandra Vivero¹, Marcia Racines¹, Varsovia Cevallos³, Rosanna Prandi², Cynthia Caicedo², Francisco Robinzon² and Mariella Anselmi^{2,4}

Abstract

Chagas disease was described in Ecuador in 1930 in the province of Guayas and thereafter in various provinces. Triatomine were reported in the province of Esmeraldas but no human infection has been described. Here we report the first evidence that the disease does exist in the province of Esmeraldas. In indigenous Awá communities located in the northwest jungle of the Esmeraldas province, 144 individuals were tested using ELISA and PCR for *T.cruzi* of which 5 (3.47%) were positive. Twenty eight triatomine were collected, 27 were *Triatoma dispar* and 1 *Pastronylus rufotuberculatus*, *T.cruzi* was detected in 11 (42.3%) of 26 insects.

Keywords: *Triatoma dispar* L, *Trypanosoma cruzi*, Ecuador, Esmeraldas, Awá population, Chagas disease

Letter

Chagas disease or American trypanosomiasis, is caused by an infection by the protozoan hemoflagellate *Trypanosoma cruzi*, which is transmitted to humans through bites of infected triatomine insects. The infection is widespread throughout Latin America although an increasing number of cases in non-endemic countries have been described [1]. In Ecuador, human *T.cruzi* infections have been observed in different provinces since 1930 [2,3]. However, in the Esmeraldas province, although triatomine insects were reported, autochthonous cases of *T.cruzi* human infection have not been documented [2]. The province of Esmeraldas, bordering the southern regions of Colombia, is located in the northwest of the country. Most of the inhabitants are of African descendant with dispersed indigenous populations such as the Epera, Chachi and Awá. The latter are also called *kwaiker*, speak their own language (*Awá pit*) and live in remote areas, isolated from any urban areas. Reports from individuals from three Awá's villages: Mataje Alto (17 N 0772280, UTM 0134144, 221 m), Pambilar (17 N 0766542, UTM 0124494, 144 m) and Balsareño (17 N 0761275, UTM

0128009, 44 m) revealed the presence of triatomine insects in their homes and in peridomestic areas. Therefore, a protocol to study insects and human blood of Awá population was prepared and approved by the Bioethics Committee COBI-ASFORUM (Federalwide Assurance FWZ00002482, IEC IORG0001932, IRB00002438, IEPID NPI 125754-12/132854-13) in Quito-Ecuador.

We collected twenty-eight insects and sent them to the laboratory for identification and detection of *T.cruzi* infection. Twenty-seven of the insects were *Triatoma dispar* Lent and one was *Pastronylus rufotuberculatus* according to conventional taxonomic keys. Since the insects were not optimally preserved, it was impossible to detect living forms of trypanosomes using microscopic techniques. Specific PCR for *T.cruzi* DNA was performed according to previously described procedures [4,5] in 26 of 28 specimens collected and 11 (42.3%) were positive (Table 1, Figure 1).

In two Awá's villages (Pambilar and Balsareño), 144 blood samples were obtained with informed consent and sera were analyzed for antibodies against *T.cruzi* utilizing three different serological tests according to manufacturer's instructions. The three different commercial ELISA tests showed high absorbance values indicating a strong anti-*T.cruzi* IgG response in 5 (3.47%) of 144

* Correspondence: angel_gustavoguevara@hotmail.com

[^] Deceased

¹Laboratorio de Parasitología Molecular y Medicina Tropical, Centro de Biomedicina, Carrera de Medicina, Universidad Central del Ecuador, Sodiro N14121 e Iquique, Quito, Ecuador

Full list of author information is available at the end of the article

Table 1 Triatomine insects collected in Awá's communities of northwest jungle of Esmeraldas province, Ecuador

Community	No	Species	Habitat
Pambilar	17	<i>Triatoma dispar</i>	Peridomestic
Balsareño	10	<i>Triatoma dispar</i>	Domestic
Balsareño	1	<i>Pastrongylus rufotuberculatus</i>	Domestic





Table 2 Results expressed in absorbance values from human sera samples with three different ELISA tests

Sample code	Age	Gender	Biokit (ELISA) [®]	Bioschile (ELISA) [®]	Chagatest (ELISA) [®]
Pamb 10	21	Male	1.82	1.50	0.62
Pamb 12	29	Female	> 2.0	> 2.0	2.0
Pamb 18	48	Female	> 2.0	> 2.0	> 2.0
Pamb 59	62	Male	1.84	1.96	2.0
Bals 37	32	Male	> 2.0	> 2.0	> 2.0

Cut-off value: Biokit negative <0.9; Bioschile negative <0.9; Chagatest negative < 0.230.



samples tested. *T.cruzi* specific PCR [6,7] was also positive in all 5 *T.cruzi* ELISA positive individuals (Table 2, Figure 2).

The present study demonstrates human *T.cruzi* infection in the northwest jungle of Ecuador. The extent of the infection as well as the associated Chagas disease pathology, if any, in the Awá population remains to be determined. In any case, our findings should alert Ecuadorian health authorities to start an integrated strategy to provide treatment and prevention measures to avoid further transmission. *Triatoma dispar* Lent is considered as a sylvatic species and has been reported in the northeast jungle of Ecuador [8] but not in the northwest jungle, in this letter 27 (96%) of 28 triatomine insects collected in northwest province of Esmeraldas were *T.dispar* L and many of them were collected in domestic areas. Studies related to vector biology and *T.cruzi* genotyping in vectors, human beings and wild reservoirs are required to understand the dynamics of *T.cruzi* transmission in this particular area since triatomine vectors had been shown an ease move from wild to domestic areas [9].

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

AG designed the study and wrote the manuscript. HC carried out PCR assays with insects. SV and MR did the serological and PCR tests with humans. RP, CC, FR collected data and samples and socialized the project with communities. VC did the triatominae identification. MA and JM did the protocol for the study, revised the manuscript and search for funding. All authors read and approved the final version of the manuscript. Dr. Juan Moreira passed away on July 14th, this article was read and approved by him on early June 2014.

Acknowledgements

We thank the Federación de Pueblos Awá for their support in the field. Thanks to Ronald Guderian and Thomas Unnasch for critical reviewing of the manuscript.

Author details

¹Laboratorio de Parasitología Molecular y Medicina Tropical, Centro de Biomedicina, Carrera de Medicina, Universidad Central del Ecuador, Sodiro N14121 e Iquique, Quito, Ecuador. ²CECOMET, Nelson Estupiñán Bass 210 y Luis Tello, Esmeraldas, Ecuador. ³Instituto Nacional de Investigación en Salud Pública, Entomología, Ministerio de Salud del Ecuador, Sodiro y Yaguachi, Quito, Ecuador. ⁴Centro de Enfermedades Tropicales, Hospital Sacro Cuore, Verona, Negrar, Italia.

Received: 12 June 2014 Accepted: 16 July 2014

Published: 6 August 2014

References

1. Bonney K: Chagas disease in the 21st century: a public health success or an emerging threat? *Parasite* 2014, **21**:11.
2. Aguilar M, Abad-Franch F, Racines J, Paucar A: Epidemiology of Chagas disease in Ecuador. A brief review. *Mem Inst Oswaldo Cruz* 1999, **94**:387–393.
3. Arteaga C: Investigaciones sobre la existencia de la enfermedad de Chagas en la zona del ferrocarril de la Costa (Provincia del Guayas). *Rev Univ Guayaquil* 1930, **1**:89–101.
4. Guevara A, Garzón E, Bowen C, Córdova X, Gómez E, Ouaiissi A: High infection rates of *Triatoma dimidiata* are associated with low levels of *Trypanosoma cruzi* seroprevalence in Pedro Carbo, Ecuador: Use of a *tc24* gene-based PCR approach. *Parasite* 2005, **12**:65–68.
5. Smith B, Conlan C, Hwang W, Weirauch C: Polymerase Chain Reaction Detection of *Trypanosoma cruzi* in Suboptimally Preserved Vectors and Comparative Infection Rates 2007–2010 in Escondido Southern California. *Vector-Borne Zoonotic Dis* 2011, **11**(12):1603–1604.
6. Guevara A, Eras J, Recalde M, Vinuesa L, Cooper P, Ouassi A, Guderian R: Severe digestive pathology associated with chronic Chagas disease in Ecuador: report of two cases. *Rev Soc Bras Med Trop* 1997, **30**:389–392.
7. Kirchoff L, Votava J, Ochs D, Moser D: Comparison of PCR and microscopic methods for detecting *Trypanosoma cruzi*. *J Clin Microb* 1996, **34**(5):1171–1175.
8. Zeledon R, Ugalde J, Paniagua L: Entomological and Ecological Aspects of Six Sylvatic Species of Triatomines (Hemiptera, Reduviidae) from the Collection of the National Biodiversity Institute of Costa Rica, Central America. *Mem Inst Oswaldo Cruz Rio de Janeiro* 2001, **96**(6):757–764.
9. Grijalva M, Villacís A, Ocaña S, Yumiseva C, Baus E: Limitations of selective deltamethrin application for triatomine control in central coastal Ecuador. *Parasites & Vectors* 2011, **4**:20.

doi:10.1186/1756-3305-7-358

Cite this article as: Guevara et al.: First description of *Trypanosoma cruzi* human infection in Esmeraldas province, Ecuador. *Parasites & Vectors* 2014 **7**:358.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

