



Effects of a Trans-Theoretical Model-Based Health Education Program on the Management of Cognitive Dysfunction in Older Adults With Mild Cognitive Impairment: Study Rationale and Protocol Design for a Randomized Controlled Trial

OPEN ACCESS

Edited by:

Yilong Wang,
Capital Medical University, China

Reviewed by:

Anna Rita Atti,
University of Bologna, Italy
Maddia Lozupone,
University of Bari Aldo Moro, Italy

*Correspondence:

Lina Wang
aring2000@163.com

†These authors have contributed
equally to this work and share first
authorship

Specialty section:

This article was submitted to
Aging Psychiatry,
a section of the journal
Frontiers in Psychiatry

Received: 12 October 2020

Accepted: 10 December 2020

Published: 13 January 2021

Citation:

Liu X, Wang L, Tao H, Ge C, Zhen X,
Sun X, Wang S and Su L (2021)
Effects of a Trans-Theoretical
Model-Based Health Education
Program on the Management of
Cognitive Dysfunction in Older Adults
With Mild Cognitive Impairment: Study
Rationale and Protocol Design for a
Randomized Controlled Trial.
Front. Psychiatry 11:616420.
doi: 10.3389/fpsy.2020.616420

Xiaoshen Liu^{1†}, Lina Wang^{2*†}, Hong Tao³, Chenxi Ge¹, Xueting Zhen¹, Xue Sun¹,
Simeng Wang¹ and Liming Su¹

¹ School of Medicine, Huzhou University, Huzhou, China, ² School of Medicine, Huzhou University, Huzhou Central Hospital, Huzhou, China, ³ AdventHealth Whole-Person Research, Orlando, FL, United States

Introduction: Recent studies have confirmed that the management of cognitive dysfunction produces considerable positive effects in individuals with mild cognitive impairment (MCI), however, compliance with participation in various cognitive dysfunction management strategies remains scant in older adults with MCI. Health education programs can improve the level of knowledge of the disease effectively, though it remains unclear as to whether health education programs are sufficient to promote behavior changes of older adults with MCI in the community.

Objective: The study aims to provide insight into the effect of a trans-theoretical Model (TTM)-based health education program on increasing knowledge about mild cognitive impairment (MCI), compliance in cognitive dysfunction management, and other cognition-related health outcomes (general cognitive function, sleep quality, depression symptoms, apathy symptoms) for older adults with MCI.

Methods: This study is a single-blinded, randomized, prospective clinical trial. We will recruit 132 participants with MCI who will be randomly assigned to a TTM-based health education group and a standard health education group in a ratio of 1:1. The intervention group will receive a TTM-based health education program (1 session/week, 4–560 min/session for 8 weeks), while the control group will receive standard health education. Assessors blinded to participant allocation will conduct baseline, post-intervention, and 3-month follow-up assessments. Statistical analyses will consist of the Wilcoxon test, the Chi-square test, the *T*-test, and 2 (group) × 3 (time) ANOVA with a 5% cut-off for significance.

Discussion: Supposing the TTM-based health education program will provide validated community-based cognitive dysfunction management strategies for older adults with

MCI, this would be a feasible approach to improve the compliance of participation in cognitive dysfunction management and the cognition-related health outcomes.

Clinical Trial Registration: ChiCTR1900028351. Registered on December 19, 2019. <http://www.chictr.org.cn/edit.aspx?pid=47223&htm=4>.

Keywords: cognitive dysfunction, self-management, trans-theoretical model, health education, mild cognitive impairment

BACKGROUND

Dementia, which remains incurable using existing technologies, has become a global health crisis. Recently, clinical research and basic research have focused on the preclinical stage of dementia—mild cognitive impairment (MCI) (1). MCI denotes problems with memory, attention, or other cognition, which are greater than expected for healthy aging but insufficient to interfere with daily activities (2). The clinical outcomes of MCI may remain stable, progress to dementia, or return to neurological integrity. The annual conversion rates from MCI to dementia are estimated from 10 to 59.4%, whereas 14.4 to 55.6% of those with MCI revert to normal cognition (NC) (3). The higher rates of MCI reversion emphasize the importance of early prevention and treatment at the MCI stage to forestall further cognitive decline.

Drug treatments resulted in no effect in any of the MCI studies ($N = 53,693$ patients, 95% CI: 0.000 to 0.005) (4), meaning that recent efforts have focused on non-pharmacological cognitive dysfunction management strategies, which have been proved promising for delaying cognitive decline, including physical exercise, cognitive training, musical therapy, mental/social activity, eating habits, and multi-domain interventions (5, 6).

In fact, older adults with MCI have poor compliance with participation in these cognitive dysfunction management strategies, with an average non-adherence rate of ~25% (7). There are many factors that influence compliance, including patients' lack of understanding of their disease, lack of social support, low sense of self-efficacy, and financial constraints (8). A promising intervention to improve compliance and educational strategies would help participants understand health information, and this may be essential to help individuals generate the motivations, beliefs and improve their overall compliance behavior (7, 9).

Health education can improve individual's level of disease knowledge, and it is an essential component of chronic disease management strategies. Health education helps individuals to understand the essence of disease management and encourages individuals to participate in their own disease management (10, 11). Nevertheless, previous studies have shown that older adults in the community still lack knowledge about the cause, diagnosis, and treatment of cognitive impairment. A total of 55.5% of community-dwelling older adults did not perceive themselves as very knowledgeable about dementia; more than 65% thought dementia was caused by healthy aging (12, 13). Meanwhile, older adults with MCI had lower scores with knowledge of

dementia than those with normal cognition, and the scores would decrease as cognitive impairment worsened gradually (14). In addition, some limitations also occurred in health education activities aimed at the population with cognitive dysfunction, educators placing more emphasis on patients with dementia and their caregiver rather than those with MCI; the lack of health education programs for older adults with MCI in the community setting (15); and the failure to assess the extent of behavioral change caused by a health education (16). Anghel et al. found that better compliance behaviors were related to increased knowledge of the disease (17). Nevertheless, whether health education can improve the compliance of participation in cognitive dysfunction management for older adults with MCI is not clear.

The trans-theoretical model (TTM) is an integrative model of behavior change that has been applied successfully to improve compliance, self-management, and health behavior in older adults and patients with chronic diseases (18–20). The effectiveness of behavioral changes may increase, framing them with the TTM (18). TTM emphasizes behavior change as a process rather than a single event, which posits that individuals move through five stages of adoption health-promoting behaviors (i.e., pre-contemplation, contemplation, preparation, action, and maintenance) (21). Nevertheless, no education program based on TTM has been proven to promote behavior change and other cognition-related health outcomes on older adults with MCI.

To address this gap, we developed a TTM-based health education program, which was targeted to improve the disease knowledge and the compliance of participation in cognitive dysfunction management for older adults with MCI.

Our study seeks to answer two primary questions:

(1) What are the impacts of a TTM-based health education program on older adults with MCI? We hypothesize that this program would positively affect the level of compliance of participation in cognitive dysfunction management, disease knowledge, and other cognition-related health outcomes (general cognitive function, sleep quality, depression symptoms, apathy symptoms) for older adults with MCI compared to the control group.

(2) Is it feasible to use the TTM-based health education program for recruitment and compliance in the community setting? We hypothesize that older adults with MCI would be unlikely to refuse to participate in this program and would have high compliance, and no adverse events would occur during the intervention.

MATERIALS AND METHODS

Study Design and Participants

This study is a randomized controlled trial (RCT, ClinicalTrials.gov number: ChiCTR1900 028351) with a 3-month follow up. This protocol has been approved by an independent ethics committee (Third People's Hospital of Huzhou, Zhejiang Province). It will be conducted in accordance with the ethical principles of the Declaration of Helsinki. The participants will be recruited from the community's health care center in Huzhou, Zhejiang Province, China. This program will consist of two arms: the intervention group (TTM-based health education program) and the control group (standard health education).

Our staff will distribute recruitment leaflets at community health care centers and local senior centers. Local health care providers will also help to make referrals to our study. Individuals who show interest will be invited for an in-person interview to screen for eligibility by three trained staff and a trained neurologist-psychiatrist will examine the potential participants to provide the final diagnosis of MCI. In the first round of eligibility assessments, potential participants complaining of memory impairment will be recruited. The second round of eligibility assessments will include demographic data, sensory function evaluation, cognitive function, and functional evaluation of activities of daily and medical history.

Participants will be eligible for the trial if they are 60 years of age or older with Petersen criteria for MCI (22). Participants who have severe physical disorders or mental disorders at present, or who have unfortunate events during the trial or follow-up period will not continue to receive the health education program and will be withdrawn from the study. Detailed inclusion and exclusion criteria are listed in **Table 1**. All participants will be fully informed of the program protocol, and only those who sign the informed consent before baseline will be included in this study. They will receive a gift for the completion of each session of the TTM-based health education program.

Sample Size

Based on our pilot study, Cohen's $d = 0.684$, effect size $= 0.324$. Cohen's d and the effect sizes were calculated for between-group

differences of means of the level compliance of participation in cognitive dysfunction management using an independent-groups pre-test and post-test design (IGPP). Sample size estimation was conducted using statistical software G*power. The analysis, with an independent t -test using $\beta = 0.95$ and $\alpha = 0.05$, requires a total sample size of 114 participants. Considering a 15% dropout rate, our total sample size comes to 132 participants (66 cases for both the intervention and control groups).

Randomization, Allocation Concealment, and Blinding

After completing the screening, eligible participants will be allocated randomly to one of two groups: TTM-based health education program group (intervention group) or standard health education group (control group) in a ratio of 1:1 using a computer-generated random number list. The random allocation sequence will be sealed in opaque envelopes by a staff member who will be not involved in study intervention, screening assessments, data collection, or data analysis. Participants will be informed that they will learn facts and management strategies about MCI, and that they will be blinded as to whether they receive health education based on TTM or standard health education. The data collectors and the data analyst will be blinded to group allocations.

Intervention

Intervention Group

The intervention group participants will attend one TTM-based health education session/week, 45–60 min/session for 8 weeks. The TTM encourages an assessment of an individual's current stage of behavior change and provides ten strategies to facilitate the construction of health education content according to the specific behavior stage (27). Therefore, before the intervention, the behavior stage of the participants' cognitive dysfunction management will be assessed by the researcher according to some questions (21) which will guide the researcher to match the appropriate health education contents (see **Figure 1**). After completing each health education session that matches the current behavioral characteristics, the participant's behavior


TABLE 1 | Inclusion criteria and exclusion criteria.

Inclusion criteria

- Aged 60 years or over
- Meet diagnostic criteria of MCI proposed by Petersen (22)
- Chief complaint of memory impairment confirmed by self-report/relatives;
- Objective cognitive impairment inconsistent with age and education Montreal Cognitive Assessment-Basic (MoCA-B) scores 18 for those with elementary education level, scores 21 for those with secondary education level, scores 23 for those with tertiary education level (23);
- Preserved global cognitive function as evidenced by Mini-mental State Examination (MMSE) score ≥ 24 (24), and absence of dementia [diagnosis by a psychiatrist according to DSM-V (25)];
- Intact Activities of Daily Living (ADL) [ADL score was ≤ 16 (26)]
- Absence of self-reported visual or auditory impairment
- Capacity to provide written informed consent
- Willingness to participate in the entirety of the study

Exclusion criteria

- History of severe disease (e.g., severe cardiovascular and cerebrovascular diseases, cancer and other severe physical disease, psychiatric disorders (neurodevelopmental disorders, schizophrenia spectrum, major depression disorder, obsessive-compulsive and related disorders and other major psychotic disorders), terminal illness)
- Serious events occurring during the intervention period (e.g., acute stroke, fall)
- Lost to follow-up
- Participation in another training related to cognitive function within 1 year



stage of cognitive dysfunction management will be reassessed, and health education content will be adjusted accordingly. With any behavior change, relapses are common (28), and we will still re-assess the behavior stage and provide stage-matched health education content for participants. The content of health education sessions will be derived from evidence-based knowledge and recommendations will be founded on mild cognitive impairment guidelines (3). The details of this program are displayed in **Supplementary Table 1**.

Control Group

The control group participants will attend one health promotion session/2 weeks, 45 min/session for 8 weeks that will be delivered by the researcher in another community health care center. The health promotion session will also be derived from evidence-based knowledge and recommendations (3). Each health education session covers different subjects, including basic knowledge of MCI, e.g., the prevalence of MCI, risk factors of MCI, symptoms of MCI, consequences and impacts of MCI, development, and diagnosis of MCI; management strategies of MCI, e.g., taking part in regular exercise (twice/week), encouraging participation in reading, puzzle games, and other cognitive activities, learning basic knowledge of a healthy diet and recommending a Mediterranean diet. For any questions asked by the control participants, general advice, without giving information relating to the behavior change stage and corresponding health education, will be given for ethical consideration.

The staff will record attendance for each participant each session over the 8-week intervention period and ask about the

presence of any adverse effects, such as pain or discomfort at each session. The interventions will be delivered at the different sites to minimize experimental contamination among the participants. The study procedure is shown in **Figure 2**.

Outcomes and Measuring Instruments


Socio-Demographic Profile

A self-designed sheet will be used to collect the demographic data, including age, gender, education, job, financial status, marital status, living condition, family history of dementia, amount of smoking, amount of alcohol consumption, eating habits, physical activities, cognitive activities, physical health status, and clinical subtype of MCI [tested by the Auditory Verbal Learning Test (Chinese Version) (29) according to the Winblad criteria (30)].

Primary Outcomes

Level of Compliance of Participation in Cognitive Dysfunction Management

The compliance of participation in cognitive dysfunction management will be determined using the Cognitive Dysfunction Management Compliance Scale (31). This scale was developed by our research team. It is composed of six dimensions: (1) perceived susceptibility (seven items); (2) perceived seriousness (four items); (3) perceived benefits (three items); (4) perceived barriers (seven items); (5) cues to action (three items); and (6) self-efficacy (eight items). Each item is scored based on a five-point Likert scale (1=not at all agree; 2=disagree; 3=not concerned; 4=agree; 5=very agree), (-) for reverse counting. A higher score means more adherence to cognitive management.



The Cronbach's α coefficient of the overall questionnaire was 0.904, and content validity I-CVI was 0.92.

Level of Knowledge of MCI

The scores of MCI knowledge will be determined using the MCI Knowledge Questionnaire. The questionnaire, designed by HY, Pan of Fudan University (32), has three domains: (1) basic knowledge of MCI (four items); (2) risk factors of MCI (eight items); and (3) prevention, treatment, and cognition of MCI (eight items). The answer for each item is true (1) or false (0). Total scores on the questionnaire range from 0 to 20, with higher scores indicating more knowledge. The Cronbach's α coefficient of the overall questionnaire was 0.85, the content validity I-CVI was 0.93 (32).

Secondary Outcomes General Cognitive Function

The cognitive function will be assessed using the Chinese version of the Montreal Cognitive Assessment Basic (MoCA-B). It was developed as a collaborative project between Canada and Thailand to differentiate between older adults with and without MCI regardless of level of literacy. MoCA-B includes nine domains: executive functioning, memory, language, orientation, counting, abstract, visual perception, naming, and attention (33). According to the Chinese cultural background, the optimal cutoff scores for MCI screening were 19 for individuals with no more than 6 years of education, 22 for individuals with 7 to 12 years' education, and 24 for individuals with more than 12 years of

education. The Cronbach's α was reported at 0.807, and the criterion-related validity was 0.787 (25).

Theory of Mind

Affective and cognitive Theory of mind (ToM) will be assessed using the "Yoni task" (Chinese version) (34). A total of 103 pictures can be subdivided into three types that correspond to affective ToM, cognitive ToM, and control conditions. A face named "Yoni" is shown in the middle with all pictures. The participant will be asked to complete special tasks depending on Yoni's gaze direction, facial expression and the information contained in the sentence. Correct answers received one point and incorrect answers received zero points (35).

Depression Symptoms

Depressive symptoms will be assessed using the Chinese version of the 15-item Geriatric Depression Scale (GDS). The total score ranges between 0 and 15, using seven as the cutoff point, with higher scores indicating severe depression. It has been validated in Chinese populations with reliability (Cronbach's α was 0.89) and validity was 0.96 (36).

Apathy Symptoms

Apathy symptoms will be assessed using the 18-item Apathy Evaluation Scale (AES), which involves cognitive, behavioral, emotional, and "other." Scores range from 18 to 72 with higher scores corresponding to more severe apathy (37). It has been validated in the Chinese older adult population with Cronbach's α 0.89, test retest reliability 0.88, and inter-tater reliability 0.86 (38).

Sleep Quality

The Pittsburgh Sleep Quality Index (PSQI) has been widely used to measure subjective sleep quality in older adults with MCI (39, 40). It involves seven components: sleep quality, sleep latency, sleep duration, habitual sleep efficiency, sleep disturbances, use of sleep medication and daytime dysfunction (41). Each component is reflected by a score ranging from 0 to 3, total scores range from 0 to 21. Higher scores indicate worse subjective sleep quality. It has been validated in a Chinese population with Cronbach's α 0.82–0.83, and the test-retest 0.85 (42).

Data Collection

Before randomized allocation, the baseline data (T0) will be collected at a community health care center by the staff blinded to group allocation, including socio-demographic profile, primary outcomes and secondary outcomes; primary outcomes, and secondary outcomes will be measured again at the end of the intervention (T1) and 3 months after intervention (T2). **Table 2** shows the assessment and timeline of the study.

Retention Strategies

Several safeguards will be put in place to decrease the dropout rate. (1) collecting multiple contact information, including contact information of a secondary contact person; (2) distributing a gift for time spent at baseline and follow up interviews; (3) enrolling in WeChat Group (WeChat, a mobile application, which is an instant messaging application) and receiving reminder messages of upcoming intervention; (4) making multiple contact attempts to complete each intervention; and (5) providing a quiet and comfortable environment for intervention.

Statistical Analysis

Data collection will be conducted in a single room and kept in a password protected computer folder. Collected data will be encoded in Excel and double-checked. Data processing and statistical analysis will be conducted using SPSS 22.0. Descriptive statistics will summarize demographic data and observable variables using frequencies and percentages for categorical variables, means \pm standard deviations for continuous variables. We will employ exploratory analysis (normality testing and distribution curves) to examine the distribution of data. The proportionality of baseline data for both groups will be compared using the non-parametric Wilcoxon test, the Chi-square test, and the *T*-test. A 2 (group) \times 3 (time) ANOVA will be used to determine the effect of the group factors, the intervention group and control group differences within-group over time, and interaction effects. Simple effect analysis will be added when there is an interaction between group factors and time factors. The intention-to-treat principle will follow data analysis. A two-tail *P*-value of < 0.05 will be considered statistically significant.

DISCUSSION

Although older adults are at risk of dementia, older adults with MCI progress to dementia more quickly than do those who are cognitively intact (43). The symptoms of early MCI are not visible, and most individuals are hidden in the community (44).

One primary challenge in implementing health education in the community is the lack of cognitive health education programs for older adults with MCI. Given limited community health care resources in China, only a few large cities can provide professional cognitive training and mental health therapists (45). It is essential, therefore, to design a cognitive health education program for older adults with MCI in the community.

Health education, as the non-pharmacological method, can improve health literacy; nevertheless, whether it can facilitate behavior change remains to be further investigated. This study innovatively uses the compliance of participation in cognitive dysfunction management as a primary indicator to assess of the effectiveness of health education interventions. This study will determine the effectiveness of a TTM-based cognitive health education program on increasing the compliance of participation in cognitive dysfunction management and disease knowledge for older adults with MCI.

TTM explains how behavior change occurs and offers explicit suggestions on how to help individuals change their behavior (21). In recent years, TTM also played an essential role in health education that promotes behavior change. Mary et al. (46) performed a randomized controlled trial of a TTM-based exercise stage-matched intervention with exercise behavior in sedentary older women. They found that, compared to the control group, the TTM-based intervention group demonstrated more willingness to start exercising, with longer adherence to an exercise, and higher exercise self-efficacy. Additionally, the authors claimed that health education based on TTM was considerably significant for future behavior changes. However, the effectiveness of a health education based on TTM for older adults with MCI is not well-understood.

Recent longitudinal studies identified depression and sleep complaints as remarkable risk factors to predict cognitive decline (39). Depressive symptoms, apathy symptoms and poor sleep are highly prevalent among older adults with MCI, with prevalence of 31.8, 44.8, and 63%, respectively (47–49). These conditions greatly interfere with the normal life of older adults with MCI. Some studies have shown that cognitive dysfunction management can significantly improve the neuropsychiatric symptoms (depression and apathy) and sleep quality (39). Considering the potential positive effect of TTM-based cognitive health education on cognitive dysfunction management behavior, further verification is needed to demonstrate whether health education can create favorable mental and physiological conditions to improve neuropsychiatric symptoms and poor sleep in older adults with MCI.

Retention and dropout are crucial issues for the intervention study. Since the TTM-based health education program will last for 8 weeks, we might have difficulty in recruiting participants and sustaining their attendance throughout the intervention period. The researcher will adopt retention strategies and work closely with community health care center staff.

There are some limitations to the current study. First, the sample population may not represent the overall characteristics of older adults with MCI, thereby limiting the generalizability of the TTM-based health education program. Second, self-reported subjective sleep quality will be measured in this study; therefore, the results may be influenced by the subjective factors of the

TABLE 2 | Summary of study assessments and time-lines.

	Questionnaire items	Time point				
		Baseline	Randomized allocation	Intervention	Post intervention	Follow-up 3 months
		T0			T1	T2
Enrollment						
Eligibility screening						
Informed consent						
Screening	General information					
	Demographic information	✓				
	Daily functions					
	Activities of Daily Living (ADL)					
	Neuro psychological examination					
	Mini-Mental State Examination (MMSE)					
	Montreal Cognitive Assessment Basic (MoCA-B)					
	Auditory Verbal Learning Test (AVLT)					
Measurements						
Primary outcomes	Compliance of cognitive dysfunction management					
	Cognitive Dysfunction Management Compliance Scale	✓			✓	✓
	Level of knowledge of MCI					
	MCI Knowledge Questionnaire	✓			✓	✓
Secondary outcomes	General cognitive function					
	Montreal Cognitive Assessment-Basic (MoCA-B)	✓			✓	✓
	Theory of Mind (ToM)	✓			✓	✓
	Neuropsychiatric symptoms					
	Depression Scale Short-form (GDS-15)	✓			✓	✓
	Apathy Evaluation Scale (AES)	✓			✓	✓
	Sleep quality					
	The Pittsburgh Sleep Quality Index (PSQI)	✓			✓	✓

participants. Third, the follow-up of health education in this study will be limited to 3 months. Hence, the more long-term effect of health education on cognitive dysfunction management compliance needs to be further explored. Future research is needed to explore the above research gaps. Finally, there are several limitations of TTM, which would not be avoided in the high level of social ecological effects. TTM ignores the social context in which change occurs, including socioeconomic

status (SES) and income. In addition, this model assumes that individuals make coherent and logical plans in their decision-making process when this is not always true. In this study, the participants might be still in the initial stage of behavioral change after the end of the intervention period (8 weeks), however, we hope that the older adults will get some knowledge of MCI, and their self-efficacy to participate in the management of cognitive functions can be evoked by the TTM education program.

In summary, this study will provide insight into the effect of a TTM-based health education program on increasing knowledge about MCI and compliance in management of cognitive dysfunction. If proven effective, health care professionals and practitioners in the geriatric field will be provided with a validated community-based cognitive health education program and will be prompted to increase its application in older adults with MCI. The benefits of this program will be perceived over time with a trend toward increasing the knowledge of MCI, which may lead to higher compliance with cognitive dysfunction management and improvements in global cognitive function.

TRIAL STATUS

The study is currently ongoing. Recruitment began in July 2020 and will conclude in May 2021.

ETHICS STATEMENT

Medical Ethics Committee of the Third People's Hospital of Huzhou, Zhejiang Province has approved and regulates the ethical execution of this research (number 2019-080). The patients/participants provided their written informed consent to participate in this study.

AUTHOR CONTRIBUTIONS

LW: research concept, research design, and critical revision of the manuscript for intellectual content. XL: drafting the first version of the manuscript, submitting the manuscript for

publication, and assisting in the development of the research design. HT: contributing to the design of the study. CG and XZ: assisting with statistical analytic planning. XS, SW, and LS: assisting with participant enrolment, consenting, and data acquisition plan. All authors contributed to the design and drafting of the manuscript, read, and approved the final manuscript.

FUNDING

This work was supported by the National Natural Science Foundation of China (No. 71704053), China Scholarship Council foundation (No. 201908330251), Zhejiang Provincial Natural Science Foundation of China (No. LQ17G030002), and General Scientific Research Project of Zhejiang Education Department (Y201942543 and Y202044216). The funding bodies participated neither in the study's design nor in the data analysis and manuscript elaboration.

ACKNOWLEDGMENTS

All authors thank all participants in the health education program and members of the health education program group for their cooperation.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fpsy.2020.616420/full#supplementary-material>

REFERENCES

- Wang J, Wang L, Zhou X, Wen X, Zhen X. Risk factors for predicting progression from normal cognition to mild cognitive impairment: protocol for a systematic review and meta-analysis of cohort studies. *BMJ Open*. (2019) 9:e027313. doi: 10.1136/bmjopen-2018-027313
- Qarni T, Salardini A. A multifactor approach to mild cognitive impairment. *Semin Neurol*. (2019) 39:179–87. doi: 10.1055/s-0039-1678585
- Petersen RC, Lopez O, Armstrong MJ, Getchius TSD, Ganguli M, Gloss D, et al. Practice guideline update summary: mild cognitive impairment: report of the guideline development, dissemination, and implementation subcommittee of the American academy of neurology. *Neurology*. (2018) 90:126–35. doi: 10.1212/WNL.0000000000004826
- Ströhle A, Schmidt DK, Schultz F, Fricke N, Staden T, Hellweg R, et al. Drug and exercise treatment of Alzheimer disease and mild cognitive impairment: a systematic review and meta-analysis of effects on cognition in randomized controlled trials. *Am J Geriatr Psychiatry*. (2015) 23:1234–49. doi: 10.1016/j.jagp.2015.07.007
- Langa KM, Levine DA. The diagnosis and management of mild cognitive impairment: a clinical review. *JAMA*. (2014) 312:2551–61. doi: 10.1001/jama.2014.13806
- Couch E, Lawrence V, Co M, Prina M. Outcomes tested in non-pharmacological interventions in mild cognitive impairment and mild dementia: a scoping review. *BMJ Open*. (2020) 10:e035980. doi: 10.1136/bmjopen-2019-035980
- Miller TA. Health literacy and adherence to medical treatment in chronic and acute illness: a meta-analysis. *Patient Educ Couns*. (2016) 99:1079–86. doi: 10.1016/j.pcc.2016.01.020
- Nafradi L, Nakamoto K, Schulz PJ. Is patient empowerment the key to promote adherence? A systematic review of the relationship between self-efficacy, health locus of control and medication adherence. *PLoS ONE*. (2017) 12:e0186458. doi: 10.1371/journal.pone.0186458
- Kröger E, Tatar O, Vedel I, Giguère AMC, Voyer P, Guillaumie L, et al. Improving medication adherence among community-dwelling seniors with cognitive impairment: a systematic review of interventions. *Int J Clin Pharm*. (2017) 39:641–56. doi: 10.1007/s11096-017-0487-6
- Liao P, Lin T, Wang T, Ting M, Wu I, Huang H, et al. Long-term and interactive effects of pay-for-performance interventions among diabetic nephropathy patients at the early chronic kidney disease stage. *Medicine*. (2016) 95:e3282. doi: 10.1097/MD.0000000000003282
- Nunes S, Rego G, Nunes R. Difficulties of Portuguese patients following acute myocardial infarction: predictors of readmissions and unchanged lifestyles. *Asian Nurs Res*. (2016) 10:150–7. doi: 10.1016/j.anr.2016.03.005
- Panelli RJ. SUDEP: a global perspective. *Epilepsy Behav*. (2020) 103:106417. doi: 10.1016/j.yebeh.2019.07.018
- Zhang H, Loi SM, Zhou SA, Zhao M, Lv X, Wang J, et al. Dementia literacy among community-dwelling older adults in urban china: a cross-sectional study. *Front Public Health*. (2017) 5:124. doi: 10.3389/fpubh.2017.00124
- Lee J, Park S, Kim KW, Kwon JE, Park JH, Kim MD, et al. Differences in knowledge of dementia among older adults with normal cognition, mild cognitive impairment, and dementia: a representative nationwide sample of Korean elders. *Arch Gerontol Geriatr*. (2016) 66:82–88. doi: 10.1016/j.archger.2016.04.013
- Du L, Hu J. The effects of health education on knowledge about Alzheimer's disease and health-promoting behaviours of older Chinese adults in a nursing home: a pilot study. *Int J Nurs Pract*. (2016) 22:31–42. doi: 10.1111/ijn.12349

16. Casemiro FG, Quirino DM, Diniz MAA, Rodrigues RAP, Pavarini SCI, Gratão ACM. Effects of health education in the elderly with mild cognitive impairment. *Rev Bras Enferm.* (2018) 71:801–10. doi: 10.1590/0034-7167-2017-0032
17. Anghel LA, Farcaș AM, Oprean RN. Medication adherence and persistence in patients with autoimmune rheumatic diseases: a narrative review. *Patient Prefer Adherence.* (2018) 12:1151–66. doi: 10.2147/PPA.S165101
18. Peleg M, Michalowski W, Wilk S, Parimbelli E, Bonaccio S, O Sullivan D, et al. Ideating mobile health behavioral support for compliance to therapy for patients with chronic disease: a case study of atrial fibrillation management. *J Med Syst.* (2018) 42:234. doi: 10.1007/s10916-018-1077-4
19. Allmer H, Allmer M, Euskirchen J, Froböse I, Wallmann B, Walter T, et al. Zugangswege zu bewegungs- und sportinaktiven Älteren [Approaching physically inactive elderly for physical activity]. *Gesundheitswesen.* (2015) 77(Suppl. 1):S43–4. doi: 10.1055/s-0033-1333715
20. Ren Q, Lian M, Liu Y, Thomas Hawkins C, Zhu L, Shen Q. Effects of a transtheoretical model-based WeChat health education programme on self-management among haemodialysis patients: a longitudinal experimental intervention study. *J Adv Nurs.* (2019) 75:3554–65. doi: 10.1111/jan.14182
21. Adams J, White M. Are activity promotion interventions based on the transtheoretical model effective? A critical review. *Br J Sports Med.* (2003) 37:106–14. doi: 10.1136/bjsm.37.2.106
22. Petersen RC. Mild cognitive impairment as a diagnostic entity. *J Intern Med.* (2004) 256:183–94. doi: 10.1111/j.1365-2796.2004.01388.x
23. Chen K, Xu Y, Chu A, Ding D, Liang X, Nasreddine ZS, et al. Validation of the Chinese version of montreal cognitive assessment basic for screening mild cognitive impairment. *J Am Geriatr Soc.* (2016) 64:e285–90. doi: 10.1111/jgs.14530
24. Li G, Shen YC, Chen CH, Zhao YW, Li SR, Lu M. An epidemiological survey of age-related dementia in an urban area of Beijing. *Acta Psychiatr Scand.* (1989) 79:557–63. doi: 10.1111/j.1600-0447.1989.tb10303.x
25. Sullivan D, Brady N, Manning E, O'Shea E, O'Grady S, O'Regan N, et al. Validation of the 6-item cognitive impairment test and the 4AT test for combined delirium and dementia screening in older emergency department attendees. *Age Ageing.* (2018) 47:61–8. doi: 10.1093/ageing/afx149
26. Lawton MP, Brody EM. Assessment of older people: self-maintaining and instrumental activities of daily living. *Gerontologist.* (1969) 9:179–86.
27. Mundorf N, Redding C, Paiva A. Sustainable transportation attitudes and health behavior change: evaluation of a brief stage-targeted video intervention. *Int J Environ Res Public Health.* (2018) 15:150. doi: 10.3390/ijerph15010150
28. The 6 Stages of Behavior Change. Available online at: <https://www.verywellmind.com/the-stages-of-change-2794868#citation-3> (accessed November 14, 2020).
29. Guo QH, Sun YT, Yu PM, Hong, ZL. Norm of auditory verbal learning test in the normal aged in China community. *Chin J Clin Psychol.* (2007) 15:132–4. doi: 10.3969/j.issn.1005-3611.2007.02.007
30. Winblad B, Palmer K, Kivipelto M, Jelic V, Fratiglioni L, Wahlund LO, et al. Mild cognitive impairment—beyond controversies, towards a consensus: report of the international working group on mild cognitive impairment. *J Intern Med.* (2004) 256:240–6. doi: 10.1111/j.1365-2796.2004.01380.x
31. Zhen XT. *Development reliability validity test of compliance prediction scale in cognitive function management for older adults with MCI.* (dissertation/master's thesis). Huzhou University, Zhejiang (2020).
32. Pan HY. *The elderly people with MCI in the community of Jinhua: cross-sectional study and intervention study.* (dissertation/master's thesis). Fudan University, Shanghai (2012).
33. Julayanont P, Tangwongchai S, Hemrungronj S, Tunvirachaisakul C, Phanthumchinda K, Hongsawat J, et al. The montreal cognitive assessment-basic: a screening tool for mild cognitive impairment in illiterate and low-educated elderly adults. *J Am Geriatr Soc.* (2015) 63:2550–4. doi: 10.1111/jgs.13820
34. Yang LQ, Wang K, Zhu CY, Zhang YF, Lv XY, Cao H, et al. Theory of mind abilities between autistic children's and the normal children's parents. *Chin J Behav Med Brain Sci.* (2011) 20:812–14. doi: 10.3760/cma.j.issn.1674-6554.2011.09.015
35. Shamy-Tsoory SG, Aharon-Peretz J. Dissociable prefrontal networks for cognitive and affective theory of mind: a lesion study. *Neuropsychologia.* (2007) 45:3054–67. doi: 10.1016/j.neuropsychologia.2007.05.021
36. Lee CM, Chang CF, Pan MY, Hsu TH, Chen MY. Depression and its associated factors among rural diabetic residents. *J Nurs Res.* (2017) 25:31–40. doi: 10.1097/jnr.000000000000143
37. Marin RS, Biedrzycki RC, Firinciogullari S. Reliability and validity of the apathy evaluation scale. *Psychiatry Res.* (1991) 38:143–62. doi: 10.1016/0165-1781(91)90040-v
38. Lee SH, Wen MC, Chao CC, Chen YJ, Yen CF. Apathy in late-life depression among Taiwanese patients. *Int Psychogeriatr.* (2008) 20:328–37. doi: 10.1017/S1041610207005698
39. Song D, Yu DSF. Effects of a moderate-intensity aerobic exercise programme on the cognitive function and quality of life of community-dwelling elderly people with mild cognitive impairment: a randomised controlled trial. *Int J Nurs Stud.* (2019) 93:97–105. doi: 10.1016/j.ijnurstu.2019.02.019
40. Alperin N, Wiltshire J, Lee SH, Ramos AR, Cardenache RH, Rundek T, et al. Effect of sleep quality on amnesic mild cognitive impairment vulnerable brain regions in cognitively normal elderly individuals. *Sleep.* (2019) 42:zsy254. doi: 10.1093/sleep/zsy254
41. Buysse DJ, Reynolds CF, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh sleep quality index: a new instrument for psychiatric practice and research. *Psychiat Res.* (1989) 28:193–213. doi: 10.1016/0165-1781(89)90047-4
42. Tsai P, Wang S, Wang M, Su C, Yang T, Huang C, et al. Psychometric evaluation of the Chinese version of the Pittsburgh sleep quality index (CPSQI) in primary insomnia and control subjects. *Qual Life Res.* (2005) 14:1943–52. doi: 10.1007/s11136-005-4346-x
43. Kasper S, Bancher C, Eckert A, Forstl H, Frolich L, Hort J, et al. Management of mild cognitive impairment (MCI): the need for national and international guidelines. *World J Biol Psychiatry.* (2020) 21:579–94. doi: 10.1080/15622975.2019.1696473
44. Hu C, Yu D, Sun X, Zhang M, Wang L, Qin H. The prevalence and progression of mild cognitive impairment among clinic and community populations: a systematic review and meta-analysis. *Int Psychogeriatr.* (2017) 29:1595–608. doi: 10.1017/S1041610217000473
45. Zhang H, Wang J, Sun T, Wang Z, Lyu X, Yu X, et al. A randomized controlled trial of combined executive function and memory training on the cognitive and noncognitive function of individuals with mild cognitive impairment: study rationale and protocol design. *Alzheimers Dement.* (2018) 4:556–64. doi: 10.1016/j.trci.2018.09.004
46. Findorff MJ, Stock HH, Gross CR, Wyman JF. Does the transtheoretical model (TTM) explain exercise behavior in a community-based sample of older women? *J Aging Health.* (2007) 19:985–1003. doi: 10.1177/0898264307308341
47. Song D, Li PWC, Yu DSF. The association between depression and mild cognitive impairment: a cross-sectional study. *Int J Geriatr Psych.* (2017) 33:672–4. doi: 10.1002/gps.4798
48. Ma L. Depression, anxiety, and apathy in mild cognitive impairment: current perspectives. *Front Aging Neurosci.* (2020) 12:9. doi: 10.3389/fnagi.2020.00009
49. McKinnon A, Terpening Z, Hickie IB, Batchelor J, Grunstein R, Lewis SJG, et al. Prevalence and predictors of poor sleep quality in mild cognitive impairment. *J Geriatr Psychol.* (2014) 27:204–11. doi: 10.1177/0891988714527516

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 Liu, Wang, Tao, Ge, Zhen, Sun, Wang and Su. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.