



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Review

The autonomic aspects of the post-COVID19 syndrome

Arad Dotan^{a,b,c,*}, Paula David^{b,d}, Dana Arnheim^{b,c}, Yehuda Shoenfeld^{a,b,e}^a Ariel University, Israel^b Zabłudowicz Center for Autoimmune Diseases, Sheba Medical Center, Tel-Hashomer, Ramat-Gan 52621, Israel^c Sackler Faculty of Medicine, Tel Aviv University, Tel Aviv, Israel^d Department of Medicine B, Sheba Medical Center, Tel Hashomer, Israel^e Laboratory of the Mosaics of Autoimmunity, Saint Petersburg University, 199034, Russia

ARTICLE INFO

Keywords:

COVID-19
 Post-COVID19 syndrome
 Long COVID-19
 Dysautonomia
 Autonomic nervous system dysfunction
 Fibromyalgia
 Chronic fatigue syndrome
 Autoimmunity
 Autoimmune disease

ABSTRACT

The SARS-CoV-2 outbreak, responsible for the widespread COVID-19, led to one of the most rogue pandemics in modern time, yet the major effects of the pandemic may still be ahead of us. SARS-CoV-2 had been found to possess autoimmune properties. Close to 20 distinct autoantibodies which target GPCR of the nervous system and renin-angiotensin system-related molecules were found significantly associated with the clinical severity of COVID-19. The new on-set of more than 10 various autoimmune disorders were documented as well. Additionally, clinical presentations of persisted symptoms were triggered in numerous recently recovered COVID-19 patients, which led to the formulation of the novel term “post-COVID19 syndrome”. Manifestations related to post-COVID-19 syndrome exist among approximately 50–80% of symptomatic COVID-19 patients who recovered, and among patients reported more than 50 different long-term effects of the SARS-CoV-2 infection. Many of the common symptoms of the post-COVID19 syndrome are not explained by the virus-related injury alone. Similarly to chronic fatigue syndrome and fibromyalgia, autoimmune-mediated autonomic nervous system dysfunction may play a significant part in the pathogenesis of such symptoms, including chronic fatigue, cognitive impairment, mood related disorders, and numerous more. Importantly, therapeutic options such as immunomodulatory and immunosuppressive therapy may favor some post-COVID19 patients, while plasmapheresis and IVIG could be considered in severe cases. Nevertheless, as physical exercise has been found to stabilize the autonomic nervous system, exercise therapy might be a safer and more effective remedy for the post-COVID19 syndrome.

1. The acute phase of COVID-19

The severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) outbreak, responsible for the widespread coronavirus disease of 2019 (COVID-19), led to one of the most rogue pandemics in modern time. This novel disease had caused morbidity and mortality at an unanticipated measure on a global scale. Nowadays, COVID-19 is perhaps most known for its acute phase, commonly presented with fever, respiratory tract symptoms [1] and, in some patients, extra-respiratory manifestations [2]. Although most individuals infected by SARS-CoV-2 are asymptomatic or present merely mild symptoms [3,4], severally-ill patients have extremely high mortality rates and overwhelm local health

authorities [5]. The high inflammatory state was frequently referred to as “cytokine storm syndrome” (SCC) to describe a vicious cycle of the hyperstimulated immune system [6], leading to an uncontrolled inflammatory response. This disorder was recognized in the ‘graft versus host disease’ and was later adopted for the severe inflammatory state of extensive viral and bacterial infections [7]. The SCC involves numerous circulating cytokines and chemokines such as IL-1 α , IL-1 β , IL-2, IL-7, IL-6, IL-8, IL-10, tumor necrosis factor, interferon γ and high concentrations of ferritin, C-reactive protein can be detected [5,8–11]. SCC had been associated with extremely high mortality due to the extreme hyperstimulation of the immune system, leading to severe self-damage of tissue, often in the form of acute respiratory distress syndrome (ARDS)

Abbreviations: SARS-CoV-2, Syndrome coronavirus 2; COVID-19, Coronavirus disease of 2019; SCC, Cytokine storm syndrome; ARSD, Acute respiratory distress syndrome; aPL, Antiphospholipid antibodies; anti-TPO, Anti-thyroid peroxidase; WHO, World Health Organization; GPCRs, G-coupled receptors; CFS, Chronic fatigue syndrome; POTS, Postural orthostatic tachycardia syndrome; IVIG, Intravenous immunoglobulin.

* Corresponding author at: Ariel University, Israel.

E-mail address: araddotan@mail.tau.ac.il (A. Dotan).

<https://doi.org/10.1016/j.autrev.2022.103071>

Received 7 February 2022; Accepted 14 February 2022

Available online 16 February 2022

1568-9972/© 2022 Elsevier B.V. All rights reserved.

and multiple organ failure [12]. The state of hyperstimulation of the immune system that occurs in severely ill patients contributes to the wide range of autoimmune manifestations [6,13]. Autoimmunity in such patients is believed to be contributed to dysregulation of immunological components and loss of self and non-self description in individuals with a genetic predisposition. An additional mechanism that is not less important is molecular mimicry between viral components and human peptides, which had been described relating to the spike protein of SARS-CoV-2 [8,14,15].

In order to have a deep understanding of the post-COVID19 syndrome, the fundamental properties that initiate at the acute COVID-19 phase are established. For that matter, close to 20 distinct functionally active autoantibodies which target GPCR of the nervous system and renin-angiotensin system-related molecules that are all significantly associated with the clinical severity of COVID-19. The new on-set of more than 10 various autoimmune disorders were documented as well. Some of the frequently found manifestations include the presents of antiphospholipid antibodies (aPL); according to aPL data in COVID-19 patients from different cohorts, the positive rate of aPL and its subtypes are considerably high, although broad: ranging from 24.2% to 57.1%, significantly higher than ill non-COVID patients [16]. In contrast, the prevalence of aPL in the general population is much lower, estimated as 50 (95% CI 42–58) per 100,000 population [17].

Development of some autoantibodies seems to be acquired in the acute phase of COVID-19. At the same time, many autoimmune manifestations presented by COVID-19 patients were developed weeks or months after their recovery [8,18]. For instance, among 159 acute COVID-19 patients, anti-thyroid peroxidase (anti-TPO) and anti-thyroglobulin were present in 20% and 11%, respectively. Follow-up (after approximately three months) was done for 127 patients. 7 (5.5%) patients which were initially negative anti-TPO were later found positive, whereas only one patient, originally positive for anti-TPO, was found negative in the follow-up [18]. The same group also found 20.1% (41/204) with “long COVID-19”, defined as the presence or persistence of symptoms upon follow-up [18]. These results are worrisome because they may indicate the development of autoimmunity in COVID-19 patients in the long term. Importantly, additional autoantibodies share a similar long-term development and persistency [19].

Autoantibodies are essential laboratory findings because some of them can represent a pre-clinical phase of autoimmunity, while others have well-known pathogenetic outcomes. Moreover, new onset of autoimmune manifestations and diseases were also developed following an acute phase of COVID-19 [8]. Furthermore, new-onset of immune-mediated neuropathies, such as Guillain-Barré syndrome, Miller Fisher syndrome, and myasthenia gravis were recognized as well [20–23]. Some of the autoimmunity following COVID-19 were organ/system specific, such as numerous cases of Kawasaki-like disease secondary to COVID-19 cases [24], immune-mediated neuropathies, and immune thrombocytopenic purpura [25], while others were systemic diseases [8].

Some immune-mediated disease flares and new-onset disease following COVID-19 vaccination were documented [26]. These manifestations could be related to the vaccination and are important recognition, yet should not use to create fear of vaccination, mainly because the COVID-19 is associated with much higher risks [26,27].

2. Introduction to post-COVID19 syndrome

In this review, we discuss the most current and vital discoveries that have been made, which have revealed the syndrome as a multi-organ disease with a broad spectrum of symptoms. Additionally, we will review the possible mechanisms that might play a crucial role in developing the chronic post-COVID19 syndrome and which treatments might be suitable for these patients. Nonetheless, and maybe most importantly, is our discussion on how individuals can prevent such chronic syndrome. It should be raised that the possible ability of SARS-CoV-2 to trigger

autoimmunity in the acute disease may contribute to the development of the chronic symptoms of the post-COVID19 syndrome in multiple mechanisms, which will be further discussed.

Recent studies have focused on analyzing the clinical presentation and epidemiology of such symptoms, leading to meaningful conclusions that formulated the novel term “post-COVID19 syndrome”, “post-acute COVID19 syndrome”, “Persistent COVID-19 Symptoms”, “long COVID-19”, etc. In October 2021, the World Health Organization (WHO) had released a formal clinical definition for the post-COVID19 syndrome, which includes: “*Illness that occurs in people who have a history of probable or confirmed SARS-CoV-2 infection; usually within three months from the onset of COVID-19, with symptoms and effects that last for at least two months*” [28]. Importantly, for these symptoms to be diagnosed as a post-COVID19 syndrome, they should not be explained by an alternative disorder.

Significant amount of time has passed since the pandemic outbreak, and many recovered patients are experiencing non-specific symptoms that substantially impact the everyday functioning of COVID-19 recovered individuals. These symptoms are correlated with the timing of the SARS-CoV-2 infection, do not have an alternative cause, and last for at least two months. The common morbidity of such symptoms includes substantial neurological and psychiatric (e.g., cognitive dysfunction, chronic fatigue, shortness of breath, sleep disturbances, anosmia, and paraneesthesia) [29,30]. Other symptoms, also included in the post-COVID19 syndrome manifestations, are likely connected to the organ-specific injury due to the viral damage and inflammation, such as pulmonary fibrosis, cardiomyopathy, and thromboembolic events [29]. Notably, the morbidity of the post-COVID19 syndrome was found to be most significant, yet not limited to, the more severe COVID-19 patients [30].

Many local health authorities worldwide are currently struggling to cope with the active COVID-19 patients, yet there is no doubt that future healthcare systems will be obligated to address the post-COVID19 syndrome.

3. Clinical presentation and epidemiology of post-COVID-19 syndrome

Considerable time has passed since the pandemic’s outbreak, and many recovered patients were found to suffer from vague symptoms connected to the SARS-CoV-2 infection. The understanding of such symptoms is crucial for identifying the underlying mechanism.

Numerous patients are experiencing chronic symptoms following the eradication of SARS-CoV-2, while only a minority of the infected completely recover with no symptoms left [31–33]. A wide variety of chronic/late symptoms were described following the coronavirus infection, including pulmonary (dyspnea, cough, pulmonary fibrosis), cardiovascular (chest pain, palpitations, stress cardiomyopathy, myocarditis), hematologic (thromboembolic events), neuropsychiatric (headache, chronic fatigue, depression, anxiety, sleep disturbances, cognitive imbalance anosmia, paraneesthesia), renal (renal failure), dermatologic and endocrine (diabetic ketoacidosis, Hashimoto’s thyroiditis, Graves’ disease) sequelae [29] (Table 1).

Recent studies indicate that manifestations related to post-COVID-19 syndrome exist among approximately 50–80% of symptomatic COVID-19 patients who recovered (Fig. 1) [31–33]. A German study, focusing on cardiovascular injury, had found that 78% of recently recovered COVID-19 patients had a persisting symptom; the most prevalent abnormality was myocardial inflammation (60%) [31]. An Italian study that assessed patients after a mean of 60.3 days from the onset of the first COVID-19 symptom found only 18 (12.6%) completely free of any COVID-19-related symptom, while 32% had 1 or 2 symptoms and 55% had three or more; worsened quality of life was observed among 44% of patients [32]. Finally, a study that analyzed 355 symptomatic COVID-19 patients had found that 46% developed post-COVID-19 symptoms, with post-viral fatigue being the most prevalent symptom in 70% of cases

Table 1
Description of the common symptoms of the post COVID19 syndrome, classified by systems involved.

Clinical Presentation of Post-Covid19
General symptoms
<i>Chronic Fatigue</i>
<i>General pain</i>
<i>Intermittent fever</i>
<i>Red eyes</i>
<i>Hair loss</i>
<i>Excessive sweat</i>
Neuropsychiatric
<i>Headache</i>
<i>Depression</i>
<i>Anxiety</i>
<i>Sleep disturbances</i>
<i>Cognitive imbalance</i>
<i>Dysautonomia</i>
<i>Anosmia/hyposmia</i>
<i>PTSD</i>
<i>Stroke</i>
Cardiovascular
<i>Chest pain</i>
<i>Palpitations</i>
<i>Stress cardiomyopathy</i>
<i>Myocarditis</i>
<i>Post-Orthostatic tachycardia syndrome</i>
Respiratory
<i>Cough</i>
<i>Dyspnoea</i>
<i>Pulmonary fibrosis</i>
<i>Pulmonary dysfunction</i>
Renal
<i>Renal failure</i>
Gastrointestinal
<i>Anorexia</i>
<i>Digestive disorders</i>
Endocrinal
<i>Hashimoto's thyroiditis</i>
<i>Grave's disease</i>
<i>Diabetic Ketoacidosis</i>
Rheumatologic
<i>Joint pain</i>
<i>Fibromyalgia</i>
Dermatologic
<i>Non-specific rash</i>
Hematologic
<i>Thromboembolic events</i>

[33]. The frequency of post-COVID19 syndrome in asymptomatic patients is still uncertain.

An extensive systematic review and meta-analysis performed by a Mexican group compiled more than 50 possible long-term effects of the coronavirus infection [29]. A “long-term symptom” was considered as any symptom that persisted for more than two weeks after the disease onset and was not completely reversible. From all the COVID-19 patients included in their data, 80% presented with at least one long-term complaint. The most common symptom was fatigue (58%) [29]. Regarding persistent respiratory-related consequences of SARS-CoV-2 infection, the most were dyspnea and cough, yet more severe sequelae such as pulmonary fibrosis and dysfunction were also reported [29]. Similar results were found in cohort studies from Italy, England, and China [32,34,35].

Headache is also on the top of the chronic COVID-19 complaints list [36,37], and according to the mentioned meta-analysis, it is present in near 45% of those patients [29]. Other neuropsychiatric symptoms are also frequently reported, such as anosmia (21%), attention disorder (27%) and depression (12%), and anxiety (13%). Depression, anxiety, foggy, and post-trauma-stress-disorder (PTSD) were previously correlated with COVID-19 in other studies [35,38].

At least one of the following non-specific symptoms was described in

10–20% of the patients: hair loss, joint pain, excessive sweat, digestive disorders, general pain, intermittent fever, and red eyes [29,39]. Less common but more severe manifestations, such as strokes and renal failure, were also described [29,39].

Regarding epidemiology, some studies have shown a predominance of post-COVID19 syndrome among Black, Asian and other minority ethnicities when compared with Caucasians [40,41]. In contrast, others found conflicting results, showing a higher prevalence among white people [42,43]. The syndrome is more common among patients older than 40, and there is no consensus whether females or males are more susceptible [42,43].

4. Post-COVID19 syndrome and autonomic nervous system dysfunction

In the past two decades, researching the role of autoimmunity in developing some autonomic nervous system dysfunction disorders has been considerably accelerating. Autoantibodies against G-coupled receptors (GPCRs) are in the spotlight, including adrenergic, muscarinic, endothelin, and angiotensin receptors [44,45]. These autoantibodies are suspected of playing an active role in many autoimmune disorders [44,45]. For example, the involvement of β 2-AR signaling was reported in the pathogenesis and progress of autoimmune diseases such as rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis, and myasthenia gravis. In addition, autoantibodies against muscarinic acetylcholine receptor M3 have been found to affect the progression of Sjogren's syndrome [45,46]. SARS-CoV-2 may contribute greatly to triggering autoimmune manifestations, including the development of various autoantibodies [8,25,27,47], thus some believe specific autoantibodies lead to autonomic nervous system dysfunction, which could explain many of the post-COVID19 syndrome manifestations.

Functionally active autoantibodies against different GPCRs, which are known to disturb the balance of neuronal and vascular activity, were found present in post-COVID-19 syndrome patients [48]. In that study, 31 post-COVID19 patients were included; all reported having between 2 and 7 different functionally active autoantibodies targeting GPCRs, that acted as receptor agonists, such as β 2-adrenoceptor, α 1-adrenoceptor, and Angiotensin II receptor type 1 receptor [48]. These autoantibodies, found before the infection, might explain the triggering of neurological and cardiovascular symptoms of the post-COVID-19 syndrome [48]. Further investigation of autoimmunity against the autonomic nervous system is carousal to extend the knowledge of this mechanism that may be the root of post-COVID19 syndrome's symptoms.

Our group had found close to 20 distinct autoantibodies which target GPCR of the nervous system and renin-angiotensin system-related molecules that are all significantly associated with the clinical severity of COVID-19 (Otavio Cabral-Marquesa, Gilad Halperdt, Lena F. Schimke, et al.; nature communications in press) (Fig. 1). Further investigation of autoimmunity against the autonomic nervous system is carousal to extend the knowledge of this mechanism that may be the root of post-COVID19 syndrome's symptoms.

The post-COVID19 syndrome phenomenon, which the WHO had well defined, seems to involve many different organs and leads to a broad spectrum of symptoms that are difficult to attribute to a single mechanism, virus- or immune-mediated. Nevertheless, the clinical presentation of the post-COVID19 syndrome can help us better evaluate the most reasonable underlying pathophysiologic process worth exploring. In this context, many had identified overwhelming similarities between manifestations related to post-COVID19 syndrome and other disorders, such as chronic fatigue syndrome (CFS) [49], fibromyalgia [50–52], postural orthostatic tachycardia syndrome (POTS), [53], and manifestations associated with silicone breast implants [45]. The typical presentation of the post-COVID19 syndrome, CFS, and fibromyalgia includes chronic fatigue and widespread pain accompanied by cognitive disorders that reduce the quality of life [29,54–58] (Table 2). Interestingly, these conditions are suspected of having an underlying autoimmune

Long-term manifestations of COVID-19




Fig. 1. In the center appears the SARS-CoV-2. Around it, at the upper part of the figure, appear some of the functionally active autoantibodies linked to COVID-19. At the bottom part of the figure appear some of the chronic symptoms associated with COVID-19.

AGTR1: Angiotensin II Receptor Type 1; AGTR2: Angiotensin II Receptor Type 2; BDKRB1: Bradykinin Receptor B1; MAS1: MAS1 Proto-Oncogene; CXCR3: C-X-C Motif Chemokine Receptor 3; CHRMs: Cholinergic Receptor Muscarinic 2; CHRMs: Cholinergic Receptor Muscarinic 3; CHRMs: Cholinergic Receptor Muscarinic 5; F2R: Coagulation Factor II Thrombin Receptor; NOR: nociception-like opioid receptor; ADRB1: Adrenergic receptor beta-1; ADRB2: Adrenergic receptor beta-2; ADRA1: Adrenoceptor Alpha 1A; NRP1: Neupilin 1; STAB1: Stabilin 1; ETA: Selective endothelin-A.

Table 2

The most common symptoms of Post-COVID19 syndrome, Fibromyalgia, and Chronic Fatigue Syndrome / Myalgic encephalomyelitis [29,54–58].

Post-COVID19 syndrome	Fibromyalgia	Chronic Fatigue Syndrome /Myalgic encephalomyelitis
Fatigue	Fatigue	Fatigue
Headache and Joint pain	Chronic widespread pain	Chronic widespread pain
Hair loss	Unrefreshing sleep	Unrefreshing sleep
Cognitive impairment	Fogginess (“fibrofog”)	Impaired memory
Dyspnea-related	Physical exhaustion	Impaired concentration
Mood related disorders	Cognitive impairment	Post-exertional malaise
Ageusia and anosmia		
Sweat		

dysautonomic-related disorder [59–61]. Furthermore, CFS and fibromyalgia are also associated with various viral infections [62–64].

Importantly, alongside the presence of functionally active autoantibodies against the autonomic nervous system components, dysfunction of the system was also found in patients with the post-COVID19 syndrome [48,65]. For example, a series of individuals with post-COVID-19 symptoms had resulted in orthostatic intolerance syndromes, possibly related to a virus- or immune-mediated disruption of the autonomic nervous system [65].

Many post-COVID-19 patients describe a clinical condition similar to the CFS: overwhelming and incapacitating fatigue, sleep disorders, cognition impairments, different kinds of autonomic dysfunction, which are exacerbated in exercises [29,35,66]. CFS causes disturbances of the central nervous system and immune regulation, leading to debilitating disorders with an autoimmune etiology [67]. The triggering of the disease, like many autoimmune diseases, it is believed to be activated by some specific viral pathogens, such as EBV [67,68]. In addition, CFS has been shown to be associated with higher serum levels of autoantibodies

against the autonomic receptors, such as beta-adrenergic and muscarinic receptors [69]. The same autoantibodies were previously related to other autonomic syndromes, such as POTS and small-fiber neuropathy [70–72]. Interestingly, those autonomic dysfunctions are some of the non-specific symptoms described in the post-COVID19 symptomatic patients [53,73,74].

The similarity between fibromyalgia and the post-COVID19 syndrome was demonstrated by a study that included a sample of 616 COVID-19 recovered patients. All the participants filled a web-based survey 6 ± 3 months after the COVID-19 diagnosis had found that 189 (30.7%) satisfied the American College of Rheumatology criteria for fibromyalgia (56.6% women). A further multivariate logistic regression model had shown that male gender and obesity were the strongest predictors of being classified as having post-COVID19 fibromyalgia (both $p < 0.0001$) [50]. A recent study demonstrated the pathogenicity of fibromyalgia patients’ IgG [75]. The researchers treated mice with IgG from fibromyalgia patients, which led to the increased sensitivity to noxious mechanical and cold stimulation and nociceptive fibers in skin-nerve preparations compared with control [75]. These findings support the belief that fibromyalgia has an autoimmune basis, and therapy that reduces patient IgG concentrations could benefit fibromyalgia [75]. It should be emphasized that there is substantial controversy relating to the etiology and pathophysiology of fibromyalgia. Many clinicians still label the signs and symptoms of fibromyalgia patients as merely psychological while not acknowledging this disorder as an ‘organic’ disease, despite strong evidence [76]. The post-COVID19 symptoms, similar to fibromyalgia, are not clearly understood thus might be disregarded by many healthcare professionals if not raised to their awareness.

5. Therapeutic options for post-COVID-19 patients

Numerous patients are at need for medical solution for new onset of symptoms following the COVID-19 recovery, in order to improve their quality of life [32]. In assessing the most beneficial therapy from post-COVID19 syndrome patients, the most probable underlying mechanism should be targeted. As SARS-CoV-2 contributes to

hyperstimulation of the immune system in the acute phase and may lead to the development of functional autoantibodies against the autonomic nervous system in the chronic phase, immunomodulatory therapy could be beneficial for post-COVID19 patients. Steroids could be considered the first line of treatment for the hyperstimulation phase of the disease [77], while other immunosuppressive treatments could be prescribed for persistent use [78]. In severe cases, plasmapheresis and intravenous immunoglobulin (IVIg) could be considered [79]. In addition, it was suggested that therapies that reduce autoantibodies levels, such as rituximab and extracorporeal apheresis, could help ease the post-COVID19 symptoms [49]. Those possibly beneficial therapeutic options should be carefully assessed in clinical in well-organized clinical trials due to the significant potential adverse effects and their high cost.

Physical exercise was found to be significantly beneficial in CFS and fibromyalgia patients [80–82]. Thus, exercise therapy might be a safer and more effective remedy in post-COVID19 syndrome in comparison to the potent immunomodulatory and immunosuppressive treatments. Exercise therapy, particularly graded aerobic and strength training, had been found to produce considerable improvements in pain, increase physical function and improve quality of life [80,83]. In controlled randomized trials, possible adverse effects of exercise, such as pain, fatigue, and musculoskeletal problems, had not been consistently reported. Therefore, a prescribed graded exercise is considered safe for these patients [80,82].

Although not much research had been done on the connection between the autonomic nervous system and autoimmune diseases, it is known that cardiovascular autonomic dysfunction is common in autoimmune disorders, specifically rheumatic diseases; systemic lupus erythematosus rheumatoid arthritis scleroderma and primary Sjögren's syndrome have an association with autonomic nervous system dysfunction, with or without cardiological [84].

The mechanism as which therapeutic physical exercise stabilizes the autonomic nervous system and restores its normal function remains unclear. Yet some findings suggest that exercise therapy can improve the outcome of patients with autonomic nervous dysfunction by increasing vagal modulation, mainly demonstrated on patients and animal models with chronic cardiac disease [85]. Further research is necessary in order to assess the benefit of physical exercise, as a therapy, in autoimmune disorders.

6. Conclusion

It is of great importance to recognize the autoimmune manifestations of COVID-19 and post-COVID19 syndrome in order to properly cope with their outcomes in the ongoing pandemic and the long-term post-pandemic period, giving the possible autoimmune-mediated underlying mechanisms. The key to managing post-COVID19 syndrome might be concealed in our knowledge of similar presentations, such as CFS and fibromyalgia. As SARS-CoV-2 contributes to hyperstimulation of the immune system in the acute phase and may lead to the development of functional autoantibodies against the autonomic nervous system in the chronic phase, some medication strategies may be beneficial. Therapeutic options such as immunomodulatory and immunosuppressive therapy could be favorable for some post-COVID19 patients, while plasmapheresis and IVIG could be considered in severe cases. Nevertheless, due to the potential adverse effects of the available therapeutic options, exercise therapy might be a safer and more effective therapeutic option in post-COVID19 syndrome. Yet, currently, further research is still necessary in order to assess the optimal treatment for the diverse symptoms and severity of the post-COVID19 syndrome.

Declaration of Competing Interest

None.

References

- [1] Huang YC, Lee PI, Hsueh PR. Evolving reporting criteria of COVID-19 in Taiwan during the epidemic. *J Microbiol Immunol Infect* 2020 Jun;53(3):413–8.
- [2] Lai CC, Ko WC, Lee PI, Jean SS, Hsueh PR. Extra-respiratory manifestations of COVID-19. *Int J Antimicrob Agents* 2020 Aug;56(2):106024.
- [3] Salzberger B, Buder F, Lampl B, et al. Epidemiology of SARS-CoV-2. *Infection*. 2021 Apr;49(2):233–9.
- [4] Gao Z, Xu Y, Sun C, et al. A systematic review of asymptomatic infections with COVID-19. *J Microbiol Immunol Infect* 2021 Feb;54(1):12–6.
- [5] Kim JS, Lee JY, Yang JW, et al. Immunopathogenesis and treatment of cytokine storm in COVID-19. *Theranostics*. 2021;11(1):316–29.
- [6] Watad A, Bragazzi NL, Amital H, Shoenfeld Y. Hyperstimulation of adaptive immunity as the common pathway for silicone breast implants, autoimmunity, and lymphoma of the breast. *Israel Med Assoc J* 2019 Aug 1;21(8):517–9.
- [7] Ferrara JL, Abhyankar S, Gilliland DG. Cytokine storm of graft-versus-host disease: a critical effector role for interleukin-1. *Transplant Proc* 1993 Feb 1;25(1 Pt 2): 1216–7.
- [8] Dotan A, Muller S, Kanduc D, David P, Halpert G, Shoenfeld Y. The SARS-CoV-2 as an instrumental trigger of autoimmunity. *Autoimmun Rev* 2021 Feb;19:102792.
- [9] Bastard P, Rosen LB, Zhang Q, et al. Autoantibodies against type I IFNs in patients with life-threatening COVID-19. *Science*. 2020 Oct 23;370(6515).
- [10] Cheng L, Li H, Li L, et al. Ferritin in the coronavirus disease 2019 (COVID-19): A systematic review and meta-analysis. *J Clin Lab Anal* 2020;34(10):e23618.
- [11] McGonagle D, Sharif K, O'Regan A, Bridgewood C. The role of cytokines including interleukin-6 in COVID-19 induced pneumonia and macrophage activation syndrome-like disease. *Autoimmun Rev* 2020 Jun;19(6):102537.
- [12] Hu B, Huang S, Yin L. The cytokine storm and COVID-19. *J Med Virol* 2021;93(1): 250–6.
- [13] Watad A, Rosenberg V, Tiosano S, et al. Silicone breast implants and the risk of autoimmune/rheumatic disorders: a real-world analysis. *Int J Epidemiol* 2018 Dec 1;47(6):1846–54.
- [14] Dotan A, Kanduc D, Muller S, Makatsariya A, Shoenfeld Y. Molecular mimicry between SARS-CoV-2 and the female reproductive system. *Am J Reprod Immunol* 2021 Dec;86(6):e13494.
- [15] Kanduc D, Shoenfeld Y. Molecular mimicry between SARS-CoV-2 spike glycoprotein and mammalian proteomes: implications for the vaccine. *Immunol Res* 2020 Sep 18:1–4.
- [16] Gao Z-W, Zhang H-Z, Liu C, Dong K. Autoantibodies in COVID-19: frequency and function. *Autoimmun Rev* 2021 Mar;20(3):102754.
- [17] Duarte-García A, Pham MM, Crowson CS, et al. The epidemiology of antiphospholipid syndrome: a population-based study. *Arthritis Rheumatol* 2019; 71(9):1545–52.
- [18] Lui DTW, Lee CH, Chow WS, et al. Long COVID in patients with mild to moderate disease: do thyroid function and autoimmunity play a role? *Endocr Pract* 2021 Sep 1;27(9):894–902.
- [19] Dotan A, Shoenfeld Y. Post-COVID syndrome: the aftershock of SARS-CoV-2. *Int J Infect Dis* 2022 Jan 1;(114):233–5.
- [20] Toscano G, Palmerini F, Ravaglia S, et al. Guillain-Barré syndrome associated with SARS-CoV-2. *N Engl J Med* 2020 Jun 25;382(26):2574–6.
- [21] Toscano G, Palmerini F, Ravaglia S, et al. Guillain-Barré syndrome associated with SARS-CoV-2. *N Engl J Med* 2020 Jun 25;382(26):2574–6.
- [22] Restivo DA, Centonze D, Alesina A, Marchese-Ragona R. Myasthenia gravis associated with SARS-CoV-2 infection. *Ann Intern Med* 2020 Dec 15;173(12): 1027–8.
- [23] Rahimi K. Guillain-Barre syndrome during COVID-19 pandemic: an overview of the reports. *Neurol Sci* 2020 Sep 2:1–8.
- [24] Akca UK, Kesici S, Ozsurekci Y, et al. Kawasaki-like disease in children with COVID-19. *Rheumatol Int* 2020 Dec;40(12):2105–15.
- [25] David P, Dotan A, Mahroum N, Shoenfeld Y. Immune thrombocytopenic purpura (ITP) triggered by COVID-19 infection and vaccination. *Isr Med Assoc J IMAJ* 2021 Jun;23(6):378–80.
- [26] Watad A, De Marco G, Mahajna H, et al. Immune-mediated disease flares or new-onset disease in 27 subjects following mRNA/DNA SARS-CoV-2 vaccination. *Vaccines*. 2021 May;9(5):435.
- [27] Dotan A, Shoenfeld Y. Perspectives on vaccine induced thrombotic thrombocytopenia. *J Autoimmun* 2021 Jul 1;(121):102663.
- [28] Flatby AV, Himmels JP, Brurberg KG, Gravningen KM. COVID-19: Post COVID-19 condition.
- [29] Lopez-Leon S, Wegman-Ostrosky T, et al. More than 50 long-term effects of COVID-19: a systematic review and meta-analysis. *Sci Rep* 2021 Aug 9;11(1):16144.
- [30] Taquet M, Geddes JR, Husain M, Luciano S, Harrison PJ. 6-month neurological and psychiatric outcomes in 236379 survivors of COVID-19: a retrospective cohort study using electronic health records. *Lancet Psychiatry* 2021 May;8(5):416–27.
- [31] Puntmann VO, Carerj ML, Wieters I, et al. Outcomes of cardiovascular magnetic resonance imaging in patients recently recovered from coronavirus disease 2019 (COVID-19). *JAMA Cardiol* 2020 Nov 1;5(11):1265–73.
- [32] Carfi A, Bernabei R, Landi F. Persistent symptoms in patients after acute COVID-19. *Jama*. 2020 Aug 11;324(6):603–5.
- [33] Mahmud R, Rahman MM, Rassel MA, et al. Post-COVID-19 syndrome among symptomatic COVID-19 patients: A prospective cohort study in a tertiary care center of Bangladesh. *PLoS One* 2021 Apr 8;16(4):e0249644.
- [34] Garrigues E, Janvier P, Kherabi Y, et al. Post-discharge persistent symptoms and health-related quality of life after hospitalization for COVID-19. *J Infect* 2020 Dec; 81(6):e4–6.

- [35] Huang C, Huang L, Wang Y, et al. 6-month consequences of COVID-19 in patients discharged from hospital: a cohort study. *Lancet* 2021 Jan 16;397(10270):220–32.
- [36] Al-Ramadan A, Rabab' h O, Shah J, Gharaibeh A. Acute and post-acute neurological complications of COVID-19. *Neurol Int* 2021 Mar 9;13(1):102–19.
- [37] Liu JWTW, de Luca RD, Mello Neto HO, Barcellos I. Post-COVID-19 syndrome? New daily persistent headache in the aftermath of COVID-19. *Arq Neuropsiquiatr* 2020 Nov;78(11):753–4.
- [38] Rogers JP, Chesney E, Oliver D, et al. Psychiatric and neuropsychiatric presentations associated with severe coronavirus infections: a systematic review and meta-analysis with comparison to the COVID-19 pandemic. *Lancet Psychiatry* 2020 Jul 1;7(7):611–27.
- [39] Kamal M, Abo Omirah M, Hussein A, Saeed H. Assessment and characterisation of post-COVID-19 manifestations. *Int J Clin Pract* 2021 Mar;75(3):e13746.
- [40] Halpin SJ, McIvor C, Whyatt G, et al. Postdischarge symptoms and rehabilitation needs in survivors of COVID-19 infection: A cross-sectional evaluation. *J Med Virol* 2021 Feb;93(2):1013–22.
- [41] Chopra V, Flanders SA, O'Malley M, Malani AN, Prescott HC. Sixty-day outcomes among patients hospitalized with COVID-19. *Ann Intern Med* 2021 Apr;174(4):576–8.
- [42] Carvalho-Schneider C, Laurent E, Lemaigen A, et al. Follow-up of adults with noncritical COVID-19 two months after symptom onset. *Clin Microbiol Infect* 2021 Feb 1;27(2):258–63.
- [43] Arnold DT, Hamilton FW, Milne A, et al. Patient outcomes after hospitalisation with COVID-19 and implications for follow-up: results from a prospective UK cohort. *Thorax*. 2021 Apr;76(4):399–401.
- [44] Halpert G, Amital H, Shoenfeld Y. Dysregulation of G protein-coupled receptors of the autonomic nervous system, adrenergic and muscarinic acetylcholine receptors, in patients with autoimmune dysautonomic-related disorders. *Brain Behav Immun - Health* 2020 Mar 3;(4):100056.
- [45] Halpert G, Watad A, Tsur AM, et al. Autoimmune dysautonomia in women with silicone breast implants. *J Autoimmun* 2021 Jun;120:102631.
- [46] Wu L, Tai Y, Hu S, et al. Bidirectional role of β 2-adrenergic receptor in autoimmune diseases. *Front Pharmacol* 2018;9:1313.
- [47] Dotan A, Mahroum N, Bogdanos DP, Shoenfeld Y. COVID-19 as an infectome paradigm of autoimmunity. *J Allergy Clin Immunol* 2022 Jan 1;149(1):63–4.
- [48] Wallukat G, Hohberger B, Wenzel K, et al. Functional autoantibodies against G-protein coupled receptors in patients with persistent long-COVID-19 symptoms. *J Transl Autoimmun* 2021 Jan 1;(4):100100.
- [49] Bornstein SR, Voit-Bak K, Donate T, et al. Chronic post-COVID-19 syndrome and chronic fatigue syndrome: is there a role for extracorporeal apheresis? *Mol Psychiatry* 2021 Jun 17:1–4.
- [50] Ursini F, Ciaffi J, Mancarella L, et al. Fibromyalgia: a new facet of the post-COVID-19 syndrome spectrum? Results from a web-based survey. *RMD Open* 2021 Aug 1;7(3):e001735.
- [51] Solano C, Martinez A, Becerril L, et al. Autonomic dysfunction in fibromyalgia assessed by the composite autonomic symptoms scale (COMPASS). *J Clin Rheumatol* 2009 Jun 1;15(4):172–6.
- [52] Martínez-Lavín M, Hermosillo AG. Autonomic nervous system dysfunction may explain the multisystem features of fibromyalgia. *Semin Arthritis Rheum* 2000 Feb;29(4):197–9.
- [53] Blitshteyn S, Whitelaw S. Postural orthostatic tachycardia syndrome (POTS) and other autonomic disorders after COVID-19 infection: a case series of 20 patients. *Immunol Res* 2021 Apr;69(2):205–11.
- [54] Häuser W, Ablin J, Fitzcharles A, et al. Fibromyalgia. *Nat Rev Dis Primers* 2015 Aug 13;1(1):1–16.
- [55] Rahman A, Underwood M, Carnes D. Fibromyalgia. *BMJ* 2014 Feb 24:348.
- [56] Yancey JR, Thomas SM. Chronic fatigue syndrome: diagnosis and treatment. *Am Fam Physician* 2012 Oct 15;86(8):741–6.
- [57] Brurberg KG, Fønhus MS, Larun L, Flottorp S, Malterud K. Case definitions for chronic fatigue syndrome/myalgic encephalomyelitis (CFS/ME): a systematic review. *BMJ Open* 2014 Feb 1;4(2):e003973.
- [58] Yong SJ. Long COVID or post-COVID-19 syndrome: putative pathophysiology, risk factors, and treatments. *Infect Dis Lond Engl* 2021 Oct;53(10):737–54.
- [59] Scheibenbogen C, Loebel M, Freitag H, Krueger A, Bauer S, Antelmann M, et al. Immunoabsorption to remove β 2 adrenergic receptor antibodies in chronic fatigue syndrome CFS/ME. *PLoS One* 2018;13(3):e0193672.
- [60] Hartwig J, Sotzny F, Bauer S, et al. IgG stimulated β 2 adrenergic receptor activation is attenuated in patients with ME/CFS. *Brain Behav Immun Health* 2020 Mar 1;3:100047.
- [61] Shoenfeld Y, Ryabkova VA, Scheibenbogen C, et al. Complex syndromes of chronic pain, fatigue and cognitive impairment linked to autoimmune dysautonomia and small fiber neuropathy. *Clin Immunol* 2020 May 1;214:108384.
- [62] Bansal AS, Bradley AS, Bishop KN, Kiani-Alikhan S, Ford B. Chronic fatigue syndrome, the immune system and viral infection. *Brain Behav Immun* 2012 Jan 1;26(1):24–31.
- [63] Low PA, Sandroni P. Postural tachycardia syndrome (POTS). *Prim Auton Nerv Syst* 2012 Jan 1:517–9.
- [64] Giacomelli C, Talarico R, Bombardieri S, Bazzichi L. The interaction between autoimmune diseases and fibromyalgia: risk, disease course and management. *Expert Rev Clin Immunol* 2013 Nov 1;9(11):1069–76.
- [65] Dani M, Dirksen A, Taraborrelli P, et al. Autonomic dysfunction in 'long COVID': rationale, physiology and management strategies. *Clin Med* 2021 Jan;21(1):e63–7.
- [66] Wostyn P. COVID-19 and chronic fatigue syndrome: is the worst yet to come? *Med Hypotheses* 2021 Jan;146:110469.
- [67] Rasa S, Nora-Krukke Z, Henning N, et al. Chronic viral infections in myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS). *J Transl Med* 2018 Oct 1;16(1):268.
- [68] Straus SE, Tosato G, Armstrong G, et al. Persisting illness and fatigue in adults with evidence of Epstein-Barr virus infection. *Ann Intern Med* 1985 Jan;102(1):7–16.
- [69] Loebel M, Grabowski P, Heidecke H, et al. Antibodies to β adrenergic and muscarinic cholinergic receptors in patients with chronic fatigue syndrome. *Brain Behav Immun* 2016 Feb 1;52:32–9.
- [70] Borba V, Malkova A, Basantsova N, et al. Classical examples of the concept of the ASIA syndrome. *Biomolecules*. 2020 Oct;10(10):1436.
- [71] Gunning WT, Kvale H, Kramer PM, Karabin BL, Grubb BP. Postural orthostatic tachycardia syndrome is associated with elevated G-protein coupled receptor autoantibodies. *J Am Heart Assoc* 2019 Sep 17;8(18):e013602.
- [72] Ruzieh M, Batizy L, Dasa O, Oostra C, Grubb B. The role of autoantibodies in the syndromes of orthostatic intolerance: a systematic review. *Scand Cardiovasc J SCJ* 2017 Oct;51(5):243–7.
- [73] Goldstein DS. The possible association between COVID-19 and postural tachycardia syndrome. *Heart Rhythm* 2021 Apr 1;18(4):508–9.
- [74] Goodman BP, Khoury JA, Blair JE, Grill MF. COVID-19 Dysautonomia. *Front Neurol* 2021;12:624968.
- [75] Goebel A, Krock E, Gentry C, et al. Passive transfer of fibromyalgia symptoms from patients to mice. *J Clin Invest* 2021 Jul 7;131(13).
- [76] Cohen H. Controversies and challenges in fibromyalgia: a review and a proposal. *Adv Musculoskelet Dis* 2017 May 1;9(5):115–27.
- [77] Chatterjee K, Wu CP, Bhardwaj A, Siuba M. Steroids in COVID-19: An overview. *Cleve Clin J Med* 2020 Aug 20;(8):1–4.
- [78] Schoot TS, Kerckhoffs AP, Hilbrands LB, Van Marum RJ. Immunosuppressive drugs and COVID-19: a review. *Front Pharmacol* 2020 Aug 28;11:1333.
- [79] Pourahmad R, Moazzami B, Rezaei N. Efficacy of plasmapheresis and immunoglobulin replacement therapy (IVIG) on patients with COVID-19. *SN Compr Clin Med* 2020 Sep 1;2(9):1407–11.
- [80] Busch AJ, Webber SC, Brachaniec M, et al. Exercise therapy for fibromyalgia. *Curr Pain Headache Rep* 2011 Oct;15(5):358–67.
- [81] Larun L, Brurberg KG, Odgaard-Jensen J, Price JR. Exercise therapy for chronic fatigue syndrome. *Cochrane Database Syst Rev* 2019;10.
- [82] Fulcher KY, White PD. Randomised controlled trial of graded exercise in patients with the chronic fatigue syndrome. *BMJ*. 1997 Jun 7;314(7095):1647.
- [83] Andrade A, Dominski FH. Infographic. Effects of exercise in patients with fibromyalgia: an umbrella review. *Br J Sports Med* 2021 Mar 1;55(5):279–80.
- [84] Stojanovich L. Autonomic dysfunction in autoimmune rheumatic disease. *Autoimmun Rev* 2009 Jun 1;8(7):569–72.
- [85] Goldsmith RL, Bloomfield DM, Rosenwinkel ET. Exercise and autonomic function. *Coron Artery Dis* 2000 Mar;11(2):129–35.