


Article

Aggregation Pheromone for an Invasive Mussel Consists of a Precise Combination of Three Common Purines



Jian He, Qi Dai,
Yuxuan Qi, ..., J.
Grant Burgess,
Caihuan Ke,
Danqing Feng

chke@xmu.edu.cn (C.K.)
dqfeng@xmu.edu.cn (D.F.)

HIGHLIGHTS

M. sallei uses a blend of three common simple purines as an aggregation pheromone

The three purines synergistically induce *M. sallei* larvae to settle

Larvae are highly sensitive to the ratio of purines released by conspecific adults

Common metabolites in precise combinations can act as species-specific pheromones

He et al., iScience 19, 691–702
September 27, 2019 © 2019
The Author(s).
<https://doi.org/10.1016/j.isci.2019.08.022>

Article

Aggregation Pheromone for an Invasive Mussel Consists of a Precise Combination of Three Common Purines

Jian He,¹ Qi Dai,¹ Yuxuan Qi,¹ Zhiwen Wu,¹ Qianyun Fang,¹ Pei Su,^{1,2} Miaoqin Huang,^{1,2} J. Grant Burgess,³ Caihuan Ke,^{1,2,*} and Danqing Feng^{1,4,*}

SUMMARY

Most marine benthic invertebrates have a pelagic larval phase, after which they settle preferentially on or near conspecific adults, forming aggregations. Although settlement pheromones from conspecific adults have been implicated as critical drivers of aggregation for more than 30 years, surprisingly few have been unambiguously identified. Here we show that in the invasive dreissenid mussel *Mytilopsis sallei* (an ecological and economic pest), three common purines (adenosine, inosine, and hypoxanthine) released from adults in a synergistic and precise ratio (1:1.125:3.25) serve as an aggregation pheromone by inducing conspecific larval settlement and metamorphosis. Our results demonstrate that simple common metabolites can function as species-specific pheromones when present in precise combinations. This study provides important insights into our understanding of the ecology and communication processes of invasive organisms and indicates that the combination and ratio of purines might be critical for purine-based signaling systems that are fundamental and widespread in nature.

INTRODUCTION

Animal aggregation is one of the most striking behaviors in biology that affects many spatial and temporal processes in ecological systems (Toonen and Pawlik, 1994; Parrish and Edelman-Keshet, 1999). Despite the cost of increased intraspecific competition, for example, for space, food, and oxygen, aggregation has often been viewed as an evolutionarily advantageous state, in which individuals derive the benefits of protection and reproduction (Danchin and Wagner, 2000; Dzierżyńska-Białończyk et al., 2018). Gregarious settlement is a common phenomenon among marine benthic invertebrates, including mussels, barnacles, oysters, and polychaetes. Most benthic marine invertebrates have a pelagic larval phase, after which they settle preferentially on or near conspecific adults, forming aggregations (Toonen and Pawlik, 1994). The transition from a planktonic to a benthic mode of life is generally accepted as a critical point in their life cycle and is fundamental to understanding population and community dynamics (Shikuma et al., 2014). Although many studies have repeatedly implicated a critical role for pheromones from conspecific adults in the induction of larval settlement forming dense aggregates for more than 30 years, surprisingly few settlement pheromones have been isolated and structurally identified (Burke, 1986; Dreanno et al., 2006). The present understanding of aggregation mechanisms and the evolution of aggregation pheromones is limited.

Invasive dreissenid mussels commonly foul submerged structures with typical high-density aggregations and are well-known ecological and economic pests in aquatic ecosystems (Pimentel et al., 2005; Michalak, 2017). These include the zebra mussel *Dreissena polymorpha*, the quagga mussel *Dreissena rostriformis bugensis* in North America and Europe (Michalak, 2017; Stokstad, 2007), *Mytilopsis leucophaeata* in Europe (Kennedy, 2011), and *Mytilopsis trautwineana* in South America (Aldridge et al., 2008). Dreissenids are dioecious with gametes released directly into the water and fertilized externally (Ram et al., 1996). After a brief free-swimming veliger stage, the pediveliger larvae settle and metamorphose to benthic juveniles, which attach to most substrates with secreted byssal threads leading to fouling. The gregarious settlement of dreissenid mussels causes adverse impacts on aquatic systems and serious cost to industries. The introduction of dreissenid mussels into water pipelines in power plants and water treatment plants causes damage worth billions of dollars in the Great Lakes area (Aldridge et al., 2006). Much research has focused on antifouling compounds in preventing invertebrate settlement (Yebra et al., 2004; Almeida and Vasconcelos, 2015; Qian et al., 2015; Martins et al., 2018). For the control of dreissenid mussels, chlorine has been commonly used in pipelines, but there are environmental concerns about this approach (Meehan et al., 2014). Furthermore, at present no practical

¹State-Province Joint Engineering Laboratory of Marine Bioproducts and Technology, College of Ocean & Earth Sciences, Xiamen University, Xiamen 361102, China

²State Key Laboratory of Marine Environmental Science, Xiamen University, Xiamen 361102, China

³School of Natural and Environmental Sciences, Newcastle University, Newcastle NE1 7RU, UK

⁴Lead Contact

*Correspondence: chke@xmu.edu.cn (C.K.), dqfeng@xmu.edu.cn (D.F.)

<https://doi.org/10.1016/j.isci.2019.08.022>






Figure 1. *M. sallei* Adults Release Chemical Cue(s) into Seawater to Induce Conspecific Larval Settlement and Metamorphosis

(A and B) Percentage settlement and metamorphosis of *M. sallei* larvae after 48-h exposure to conspecific adult-conditioned seawater (A) and conspecific adult shell-conditioned seawater (B). ACS, adult-conditioned seawater; SCS, shell-conditioned seawater. The preparation of 2-, 4-, and 8-h ACS, and that of 2-, 4-, and 8-h SCS, is described in the [Methods](#).

(C) Percentage settlement and metamorphosis of *M. sallei* larvae after 48-h exposure to three fractions of mantle cavity fluid from conspecific adults. Preparation of the three fractions is described in the [Methods](#). EtOAc-fr, the ethyl acetate fraction; butanol-fr, the n-butanol fraction; Aq-fr, the aqueous fraction; Control, filtered (0.22 µm) seawater. Results are shown as mean ± SD (n = 3). Asterisk denotes significant difference compared with the control (*p < 0.05, **p < 0.01, ***p < 0.001, Dunnett's test).

technology is available for inhibiting the gregarious settlement and invasion of dreissenid mussels in open waters ([Molloy et al., 2013](#)). Research on aggregation mechanisms of dreissenid mussels may shed light on developing environment-friendly and effective methods for their control. Although the formation of aggregates by adults of dreissenid mussels and factors that affect this process have been previously studied ([Dzierżyńska-Białończyk et al., 2018](#); [Tošenovský and Kobak, 2016](#)), there is still very little understanding of the biochemical mechanisms of gregarious settlement of dreissenid mussels.

The Caribbean false mussel *Mytilopsis sallei* (Recluz, 1849) ([Figure S1](#)), a close relative of *D. polymorpha* and *D. rostriformis bugensis*, was introduced into the Pacific via the Panama Canal ([Morton, 1981](#)). This dreissenid mussel has wide environmental tolerance, grows rapidly, matures fast, and has a high fecundity, contributing to its success as an invasive species ([Morton, 1981, 1989](#)). *M. sallei* has been found in Australasia, East Asia, and India ([Willan et al., 2000](#); [Wong et al., 2011](#); [Cai et al., 2014](#)). Large aggregations of *M. sallei* cause serious fouling problems on monsoon drains, concrete walls, floating rafts, aquaculture facilities, and other submerged artificial structures. Here, we hypothesized that *M. sallei* adults could release aggregation pheromone to induce settlement of conspecific larvae. The aim of the present study was to investigate whether such a pheromone exists in *M. sallei* adults and to attempt to isolate such an aggregation pheromone by bioassay-guided fractionation to determine its chemical structure. This work may provide insights into our understanding of the population dynamics and ecology of invasive dreissenid mussels and allow the development of methods for their control.

RESULTS AND DISCUSSION

An Aggregation Pheromone Exists in *M. sallei*

To confirm the existence of the conspecific cue(s), we examined larval settlement and metamorphosis of *M. sallei* in response to conspecific adult-conditioned seawater (ACS). Larval settlement and metamorphosis

Bioassay	Cue	Response	df between Groups	df within Groups	F Value	p Value
Effect of conditioned seawater	ACS	Settlement	3	8	55.035	0.000
		Metamorphosis	3	8	22.839	0.000
	SCS	Settlement	3	8	1.724	0.239
		Metamorphosis	3	8	2.303	0.154
Effect of MCF	EtOAc fraction	Settlement	3	8	8.514	0.003
		Metamorphosis	3	8	0.385	0.814
	Butanol fraction	Settlement	3	8	3.047	0.070
		Metamorphosis	3	8	0.771	0.568
	Aqueous fraction	Settlement	3	8	8.500	0.003
		Metamorphosis	3	8	12.458	0.001
Effect of subfractions obtained during the bioassay-guided fractionation of MCF	F1	Settlement	4	10	8.859	0.003
		Metamorphosis	4	10	4.400	0.026
	F2	Settlement	4	10	6.100	0.009
		Metamorphosis	4	10	1.150	0.388
	F3	Settlement	4	10	7.597	0.004
		Metamorphosis	4	10	10.750	0.001
	F3-1	Settlement	4	10	0.849	0.526
		Metamorphosis	4	10	0.885	0.507
	F3-2	Settlement	4	10	0.827	0.537
		Metamorphosis	4	10	0.512	0.729
	F3-3	Settlement	4	10	2.226	0.139
		Metamorphosis	4	10	1.726	0.220
	F3-4	Settlement	4	10	7.264	0.005
		Metamorphosis	4	10	16.469	0.000
	F3-4-1	Settlement	4	10	0.394	0.809
		Metamorphosis	4	10	0.310	0.865
	F3-4-2	Settlement	4	10	13.676	0.000
		Metamorphosis	4	10	9.342	0.002
	F3-4-3	Settlement	4	10	18.015	0.000
		Metamorphosis	4	10	12.375	0.001
	F3-4-4	Settlement	4	10	3.438	0.052
		Metamorphosis	4	10	1.314	0.329
	F3-4-3-1	Settlement	4	10	15.886	0.000
		Metamorphosis	4	10	11.452	0.001
	F3-4-3-2	Settlement	4	10	32.947	0.000
		Metamorphosis	4	10	17.717	0.000
	F3-4-3-3	Settlement	4	10	5.857	0.011
		Metamorphosis	4	10	1.521	0.269

Table 1. ANOVA Results for the Effect of Chemical Cues on Larval Settlement and Metamorphosis of *M. sallei*

(Continued on next page)

Bioassay	Cue	Response	df between Groups	df within Groups	F Value	p Value
Effect of purine compounds	Ado	Settlement	6	14	13.991	0.000
		Metamorphosis	6	14	16.123	0.000
	Ino	Settlement	6	14	26.119	0.000
		Metamorphosis	6	14	18.028	0.000
	Hyp	Settlement	6	14	13.604	0.000
		Metamorphosis	6	14	18.028	0.000
	Ade	Settlement	6	14	1.127	0.396
		Metamorphosis	6	14	0.673	0.673
Xan	Settlement	6	14	2.497	0.074	
	Metamorphosis	6	14	0.220	0.964	
Effect of dilution of ACS	Different dilutions of ACS	Settlement	4	10	10	0.000
		Metamorphosis	4	10	10	0.000
Synergistic effect of Ado, Ino, and Hyp	Mixture of Ado, Ino and Hyp, and individual components	Settlement	5	12	12	0.000
		Metamorphosis	5	12	12	0.000
Effects of different ratios of Ado, Ino, and Hyp	Different ratios of Ado, Ino and Hyp	Settlement	34	70	70	0.000
		Metamorphosis	34	70	70	0.000

Table 1. Continued

ACS, adult-conditioned seawater; SCS, shell-conditioned seawater; MCF, mantle cavity fluid; Ado, adenosine; Ino, inosine; Hyp, hypoxanthine; Ade, adenine; Xan, xanthine; EtOAc, ethyl acetate; df, degree of freedom.

were both significantly induced by ACS (Figure 1A, Table 1). The three treatments using ACS prepared by placing *M. sallei* adults in seawater for 2, 4, and 8 h, respectively, all showed inducing activity. However, treatments using shell-conditioned seawater prepared by placing empty shells of *M. sallei* adults in seawater for 2, 4, and 8 h, respectively, had no significant impact on larval settlement and metamorphosis compared with the control (Figure 1B, Table 1). These findings demonstrated that *M. sallei* adults could release chemical cue(s) to induce conspecific larval settlement and metamorphosis, and that the cue(s) were not derived from the shell of the adult *M. sallei*. We hypothesized that the inducing cue(s) in ACS were derived from mantle cavity fluid (MCF), as MCF is released with exhalant current from adults (Zimmer and Butman, 2000). MCF was successively partitioned with ethyl acetate and n-butanol, and the resultant three fractions, including the residual aqueous fractions, were examined for their effect on settlement and metamorphosis of *M. sallei* larvae. Only the aqueous fraction was active in inducing larval settlement and metamorphosis (Figure 1C, Table 1), further confirming the existence of waterborne inducing cues.

M. sallei Uses Three Common Simple Purines as Aggregation Pheromone

The active aqueous fraction was then subjected to bioassay-guided fractionation by ultrafiltration and column chromatography, which gave three pure active compounds (1, 2, and 3) (Figure 2). They all significantly induced settlement and metamorphosis of *M. sallei* larvae (Table 1). Based on analysis of their electrospray ionization mass spectrometry and nuclear magnetic resonance spectral data (Supplemental Information), compounds 1, 2, and 3 were identified as the purines, hypoxanthine (Hyp), inosine (Ino), and adenosine (Ado), respectively (Chenon et al., 1975; Saladino et al., 2006; Abou-Hussein et al., 2007; Ghose, 2009). This study identifies purines as pheromones for larval settlement or metamorphosis of a marine invertebrate. To further confirm the activity of compounds for larval settlement of *M. sallei* under hydrodynamic conditions, as in the natural environment, we performed larval bioassays using Ado (as an example of inductive purines) in a racetrack flume with seawater at a flow rate of 10.8–16.2 L min⁻¹ (Figure 3A). More than twice the number of *M. sallei* larvae were found to settle on substrates, which slowly released




Figure 2. Bioassay-Guided Fractionation of Compounds that Can Induce *M. sallei* Larval Settlement and Metamorphosis from Mantle Cavity Fluid of Conspecific Adults

Results are shown as mean \pm SD (n = 3). Asterisk denotes significant difference compared with the control (*p < 0.05, **p < 0.01, ***p < 0.001, Dunnett's test).




Figure 3. Adenosine Induced Larval Settlement of *M. sallei* under Flow Condition

(A) Experimental apparatus used in this study, with seawater in the racetrack flume at a flow rate of 10.8–16.2 L min⁻¹. (B–D) Number of settled larvae on the treated substrates that slowly released adenosine (Ado) and control substrates after 12 h (B), 24 h (C), and 48 h (D). Results are shown as mean ± SD (n = 3). Asterisk denotes significant difference compared with the control (*p < 0.05, Student's t test).

Ado compared with the controls (Figures 3B–3D), demonstrating the inducing activity of Ado for larval settlement under flow conditions.

Purines as signal molecules have been reported in fish and polychaetes. In fish, purines such as ATP, ADP, AMP, and Ado evoke attraction or feeding (appetitive) responses, whereas hypoxanthine 3-N-oxide can evoke avoidance or alarm (Wakisaka et al., 2017; Shamchuk et al., 2018). In polychaetes, the purine metabolites uric acid and Ino serve as the sperm- and egg-release pheromones, respectively, and were the first identified gamete-release pheromones in marine invertebrates (Zeeck et al., 1998a, 1998b). Here we demonstrate the use of purines for intraspecific communication in the phylum Mollusca. Purines are common metabolites in all organisms. Utilizing already existing purine metabolites as pheromones might be more energetically favorable than producing entirely new molecules. However, as suggested in a recent study (Shamchuk et al., 2018), the role of purines as chemical communication molecules in animals is substantially underestimated.

As Hyp, Ino, and Ado are purine metabolites, we wondered whether there were other active purine metabolites in *M. sallei* MCF. By using liquid chromatography-mass spectrometry, five purine metabolites, namely, Hyp, Ino, Ado, xanthine (Xan), and adenine (Ade), were detected in MCF of *M. sallei* (Figure 4A). However, unlike the significant inducing activity exhibited by Hyp, Ino, and Ado for *M. sallei* larval settlement and metamorphosis, Xan and Ade showed no significant effects (Figure 4B, Table 1). This finding further confirmed the efficiency of the bioassay-guided fractionation procedure for identification of active compounds used here.




Figure 4. Adenosine, Inosine, and Hypoxanthine Are Inducing Compounds from *M. sallei* Adults for Settlement and Metamorphosis of Conspecific Larvae

(A) Detection by liquid chromatography-mass spectrometry of purines in the mantle cavity fluid (MCF) of adults.

(B) Chemical structures of the purines detected in MCF, and percentages of settlement and metamorphosis of *M. sallei* larvae after 48-h exposure to each purine compound. Xan, xanthine; Hyp, hypoxanthine; Ino, inosine; Ado, adenosine; Ade, adenine. Results are shown as mean \pm SD ($n = 3$). Asterisk denotes significant difference compared with the control (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, Dunnett's test).

Pheromone Components Synergistically Induce Larval Settlement and Metamorphosis

To serve as effective pheromone signals in the natural environment, Hyp, Ino, and Ado should be released into seawater by *M. sallei* adults at effective concentrations. As ACS was active for conspecific larval settlement and metamorphosis, whether the three purines were released into ACS and responsible for its inducing activity was determined as follows. Figure 5A showed that when ACS was diluted with seawater, the inducing activity of ACS decreased with increasing dilution. A 5-fold dilution of ACS (5-d ACS) was the highest dilution to show inducing activity. Ado, Ino, and Hyp were found to be present in the 5-d ACS with concentrations of 8, 9, and 26 nM, respectively (Figures 5B and S2B), which were much lower than the lowest effective concentration of each compound when tested individually (0.5 μ M for Ado, 0.5 μ M for Ino, and 1.0 μ M for Hyp, Figure 4B), indicating that there might be a synergistic effect of these three purines on *M. sallei*. To determine whether there was a synergistic effect, the response of larval settlement and metamorphosis to a mixture of the three purines (8 nM for Ado, 9 nM for Ino, and 26 nM for Hyp) was investigated, and compared with the response to individual compounds under the same concentrations of each compound separately in the mixture. The 5-d ACS was used as positive control. Interestingly, the




Figure 5. Phormone Components Synergistically Induce Larval Settlement and Metamorphosis

(A) Percentage of larval settlement and metamorphosis in response to different dilutions of adult-conditioned seawater (ACS). Filtered seawater was used as control.

(B) Concentrations of adenosine (Ado), inosine (Ino), and hypoxanthine (Hyp) in the 5-fold dilution of ACS determined by high-performance liquid chromatography.

(C) Percentage of larval settlement and metamorphosis of *M. sallei* in response to 8 nM Ado, 9 nM Ino, 26 nM Hyp, and mixture of the three purines. Filtered seawater was used as a negative control. The 5-fold dilution of ACS (5-d ACS) was used as a positive control. Results are shown as mean \pm SD (n = 3). Asterisk denotes significant difference compared with the negative control (**p < 0.01, ***p < 0.001, Dunnett's test).

blend of the purines produced a significant response, whereas each individual component showed no inducing activity under the concentration tested in this bioassay (Figure 5C, Table 1), which strongly suggests a synergistic effect of Ado, Ino, and Hyp in ACS. By comparing the effective concentration of the mixture and that of the individual components, the mixture was 38.5- to 62.5-fold more potent than the individual compounds. Furthermore, the synthetic mixture and the 5-d ACS were equally effective at inducing settlement and metamorphosis.

Phormone Components Function in a Precise Ratio

As Ado, Ino, and Hyp are common metabolites found in aquatic environments (Cunliffe, 2015), we wondered how *M. sallei* larvae can distinguish conspecific adult-derived purine signals from background levels. The use of a blend of phormone components that act synergistically has also been reported in insects (Meier et al., 2016), although there are few published examples for marine organisms (Li et al., 2018). Usually the ratio of the phormone components is highly species specific in insects, allowing effective communication among related sympatric species (Symonds and Elgar, 2008). For example, two *Helicoverpa* (moth) species use the same sex phormone components (Z)-11-hexadecenal and (Z)-9-hexadecenal in nearly reverse ratios, 100:2 is used by *Helicoverpa armigera* and 6:100 is used by *Helicoverpa assulta*, thus ensuring segregation in nature (Wang et al., 2005). To determine whether *M. sallei* larvae had specific preference for particular ratios of Ado, Ino, and Hyp, thirty-three treatments with different ratios but with the same total concentration of compounds (50 nM) were prepared using simplex lattice mixture design (Figure 6A) (50 nM was chosen because the sum of concentrations of Ado, Ino, and Hyp in the effective 5-d ACS was close to 50 nM). These mixtures were then tested for larval response to settlement and metamorphosis. Results showed that different ratios produced quite different larval responses (Figures 6B–6E, Table 1). Twenty-five mixtures had no significant effect on settlement and metamorphosis. Six treatments with ratios of Ado, Ino, and Hyp of 0:4:1, 1:3:1, 0:1:4, 5:1:19, 1:5:19, and 1:1:3, gave a significant inducing activity (Figures 6B and 6C). The highest rates of settlement and metamorphosis were observed with a ratio of 1:1:3, which, interestingly was similar to the ratio of 8:9:26 (1:1.125:3.25) for Ado, Ino, and Hyp in natural ACS (Figures 5B, 6D, and 6E). The finding that *M. sallei* larvae are highly sensitive only to a particular ratio of Ado, Ino, and Hyp suggested that *M. sallei* larvae may have evolved species-specific responses to this ratio.

We further measured the ratios of these three purines in the conditioned seawater of the mussel *Perna viridis* and the oyster *Crassostrea angulata* (Figures S2B and S2C), two molluscan species with ecological niches similar to *M. sallei*, and found that the ratio of Ado, Ino, and Hyp in ACS was 6:6:1 for *P. viridis* and 1:10:5 for *C. angulata*. The clear difference of these ratios and that of *M. sallei* (1:1.125:3.25) supports the hypothesis that *M. sallei* larvae recognize conspecific adult-derived phormones from background levels of purines, by only responding to a particular ratio of phormone components. The other noteworthy finding was that two ratios of Ado, Ino, and Hyp (20:4:1 and 20:1:4) significantly inhibited settlement of *M. sallei* (Figure 6B), suggesting that even under the same total concentration of compounds, the larval




Figure 6. Specific Ratios of Adenosine, Inosine, and Hypoxanthine Induce Conspecific Larval Settlement and Metamorphosis

(A) Thirty three different ratios of Ado, Ino, and Hyp were designed based on simplex lattice mixture design. The test levels used for each purine compound were designed as 0, 1/25, 1/5, and 1. Fractions of each level were used, and the sum of contributory fractions for each ratio treatment was one. The positions of the coded levels are shown in the triangle, and the corresponding ratios are shown in the table on the right. All treatments were at the same total concentration of compounds (50 nM). Scale bar, 2 nM (i.e., 1/25 of total concentration).

(B and C) Percentage of larval settlement (B) and metamorphosis (C) of *M. sallei* in response to different ratios of Ado, Ino, and Hyp. Filtered seawater was used as a negative control (NC). The 5-fold dilution of adult-conditioned seawater was

Figure 6. Continued

used as a positive control (PC). Results are shown as mean \pm SD (n = 3). Asterisk denotes significant difference compared with the negative control (*p < 0.05, **p < 0.01, ***p < 0.001, Dunnett's test).

(D and E) Mixture contour plots show the effect of different ratios of Ado, Ino, and Hyp on settlement (D) and metamorphosis (E) of *M. sallei* larvae. Star shows the measured ratio of Ado:Ino:Hyp in ACS, i.e., 1:1.125:3.25.

response to certain mixtures could change from induction of settlement to its prevention. The discovery that changing the ratio of the pheromone components can result in the opposite observed effect is reported here. It suggests a potential method for interfering with the gregarious settlement of *M. sallei* and other dreissenid mussels by simply changing the ratio of components in an aggregation pheromone blend.

Purines have received increasing attention as intracellular and intercellular signaling messengers (Massé et al., 2007; Idzko et al., 2014; Verkhatsky and Burnstock, 2014). Sensitivity to purines is widespread across prokaryotes, plants, and animals. The purinergic signaling system is not only ancient in evolution but also omnipresent across species and tissues, involved in highly diverse functions (Verkhatsky and Burnstock, 2014). This signaling system is essential in living organisms because it mediates numerous cellular processes, including neurotransmission, neuromodulation, immune responses, cell proliferation, differentiation and death in development, regeneration, wound healing, cancer, and aging (Burnstock, 2012). Purines as signaling molecules in internal tissues of organisms are now widely accepted. Purinergic communication between individuals has also been found in a few organisms such as fish and polychaetes as mentioned above. As purines are most likely to occur as mixtures both in internal tissues and external environments (Shamchuk et al., 2018), our finding that the combination and ratio of purines is critical for communication in aquatic mussels has significant implications for the study of purinergic signaling in general.

We have shown that *M. sallei* MCF contains three purines, Ado, Ino, and Hyp, which can induce conspecific larval settlement and metamorphosis. Our discovery that the aggregation pheromone of *M. sallei* consists of a synergistic blend of these three purines, most active in a specific ratio allows us to begin to understand how different species of marine bivalve can communicate effectively in the same environment. We further suggest that a blend of Hyp, Ino, and Ado at the optimum ratio induces *M. sallei* larvae to settle and is a critical driver for the development of dense aggregations of this bivalve in the natural environment. Similar pheromone-driven aggregation mechanisms are likely to exist in other invasive dreissenid mussels. Throughout the chemical ecology literature, pheromone molecules are commonly reported to be unique compounds (Wyatt, 2014). However, this work indicates that further study of simple common metabolites that can act as species-specific pheromones when present in specific combinations, like a chemical combination lock, is warranted. In addition, the discovery of settlement inhibition by specific mixtures of purines suggests approaches to the prevention of marine biofouling.

Limitations of the Study

Our results suggested that *M. sallei* adults release a blend of three purines as the aggregation pheromone to induce settlement and metamorphosis of conspecific larvae. However, the mechanism of the synergetic effect of the three purines, Ado, Ino, and Hyp, on settlement of *M. sallei* larvae is unknown. Moreover, whether this pheromone plays an important role in *M. sallei* aggregation in the complex and dynamic natural environment remains to be confirmed, although we have demonstrated that Ado could induce larval settlement of *M. sallei* under flow conditions in laboratory. Further study will be needed to address these issues.

METHODS

All methods can be found in the accompanying [Transparent Methods supplemental file](#).

SUPPLEMENTAL INFORMATION

Supplemental Information can be found online at <https://doi.org/10.1016/j.isci.2019.08.022>.

ACKNOWLEDGMENTS

This research was supported by the National Natural Science Foundation of China under grant 41276127, the National Key Research and Development Program of China under grant 2018YFC1407505, the National

Marine Economic Development Demonstration Project in Xiamen under grant 16CZB023SF12, the Science and Technology Project of Fujian Province under grant 2018Y0076, and the Project of Subsidy Funds for Marine Economic Development in Fujian Province under grant FJHJF-L-2018-2. We thank S. Tang for help in structural identification of compounds.

AUTHOR CONTRIBUTIONS

J.H., D.F., P.S., M.H., and C.K. designed research. J.H., D.F., Q.D., Y.Q., Z.W., and Q.F. performed research. J.G.B. provided constructive suggestions. All authors wrote the paper.

DECLARATION OF INTERESTS

The authors declare no competing interests.

Received: March 13, 2019

Revised: June 25, 2019

Accepted: August 13, 2019

Published: September 27, 2019

REFERENCES

- Abou-Hussein, D.R., Badr, J.M., and Youssef, D.T.A. (2007). Nucleoside constituents of the Egyptian tunicate *Eudistoma laysani*. *Nat. Prod. Sci.* 13, 229–233.
- Aldridge, D.C., Salazar, M., Serna, A., and Cock, J. (2008). Density-dependent effects of a new invasive false mussel, *Mytilopsis trautwineana* (Tryon 1866), on shrimp, *Litopenaeus vannamei* (Boone 1931), aquaculture in Colombia. *Aquaculture* 281, 34–42.
- Aldridge, D.C., Elliott, P., and Moggridge, G.D. (2006). Microencapsulated biobullets for the control of biofouling zebra mussels. *Environ. Sci. Technol.* 40, 975–979.
- Almeida, J.R., and Vasconcelos, V. (2015). Natural antifouling compounds: effectiveness in preventing invertebrate settlement and adhesion. *Biotechnol. Adv.* 33, 343–357.
- Burke, R.D. (1986). Pheromones and gregarious settlement of marine invertebrate larvae. *Bull. Mar. Sci.* 39, 323–331.
- Burnstock, G. (2012). Purinergic signalling: its unpopular beginning, its acceptance and its exciting future. *Bioessays* 34, 218–225.
- Cai, L.Z., Hwang, J.S., Dahms, H.U., Fu, S.J., Zhuo, Y., and Guo, T. (2014). Effect of the invasive bivalve *Mytilopsis sallei* on the macrofaunal fouling community and the environment of Yundang Lagoon, Xiamen, China. *Hydrobiologia* 741, 101–111.
- Chenon, M.T., Pugmire, R.J., Grant, D.M., Panzica, R.P., and Townsend, L.B. (1975). Carbon-13 magnetic resonance. XXVI. A quantitative determination of the tautomeric populations of certain purines. *J. Am. Chem. Soc.* 97, 4636–4642.
- Cunliffe, M. (2015). Purine catabolic pathway revealed by transcriptomics in the model marine bacterium *Ruegeria pomeroyi* DSS-3. *FEMS Microbiol. Ecol.* 92, 1–6.
- Danchin, E., and Wagner, R.H. (2000). Benefits of membership. *Science* 287, 803.
- Dreanno, C., Matsumura, K., Dohmae, N., Takio, K., Hirota, H., Kirby, R.R., and Clare, A.S. (2006). An α_2 -macroglobulin-like protein is the cue to gregarious settlement of the barnacle *Balanus amphitrite*. *Proc. Natl. Acad. Sci. U S A* 103, 14396–14401.
- Dzierżyńska-Białończyk, A., Skrzypczak, A., and Kobak, J. (2018). Happy together? Avoidance of conspecifics by gregarious mussels. *Curr. Zool.* 64, 53–61.
- Ghose, R. (2009). Interaction of adenosine, guanosine and inosine with ruthenium hydride complexes. *J. Chem. Res.* 2009, 52–55.
- Idzko, M., Ferrari, D., and Eltzschig, H.K. (2014). Nucleotide signalling during inflammation. *Nature* 509, 310–317.
- Kennedy, V.S. (2011). The invasive dark false mussel *Mytilopsis leucophaeata* (Bivalvia: Dreissenidae): a literature review. *Aquat. Ecol.* 45, 163–183.
- Li, K., Buchinger, T.J., and Li, W. (2018). Discovery and characterization of natural products that act as pheromones in fish. *Nat. Prod. Rep.* 35, 501–513.
- Martins, S.E., Fillmann, G., Lillicrap, A., and Thomas, K. (2018). Reviews: ecotoxicity of organic and organo-metallic antifouling co-biocides and implications for environmental hazard and risk assessments in aquatic ecosystems. *Biofouling* 34, 34–52.
- Massé, K., Bhamra, S., Eason, R., Dale, N., and Jones, E.A. (2007). Purine-mediated signalling triggers eye development. *Nature* 449, 1058–1062.
- Meehan, S., Gruber, B., and Lucy, F.E. (2014). Zebra mussel control using Zequanox® in an Irish waterway. *Manag. Biol. Invasion* 5, 279–286.
- Meier, L.R., Zou, Y., Millar, J.G., Mongold-Diers, J.A., and Hanks, L.M. (2016). Synergism between enantiomers creates species-specific pheromone blends and minimizes cross-attraction for two species of cerambycid beetles. *J. Chem. Ecol.* 42, 1181–1192.
- Michalak, A.M. (2017). Environmental sciences: troubled waters on the Great Lakes. *Nature* 543, 488.
- Molloy, D.P., Mayer, D.A., Giamberini, L., and Gaylo, M.J. (2013). Non-target trials with *Pseudomonas fluorescens* strain CL145A, a lethal control agent of dreissenid mussels (Bivalvia: Dreissenidae). *Manag. Biol. Invasion* 4, 71–79.
- Morton, B. (1981). The biology and functional morphology of *Mytilopsis sallei* (Recluz) (Bivalvia: Dreissenacea) fouling Visakhapatnam Harbour, Andhra Pradesh, India. *J. Molluscan Stud.* 47, 25–42.
- Morton, B. (1989). Life-history characteristics and sexual strategy of *Mytilopsis sallei* (Bivalvia: Dreissenacea), introduced into Hong Kong. *J. Zool.* 219, 469–485.
- Parrish, J.K., and Edelman-Keshet, L. (1999). Complexity, pattern, and evolutionary trade-offs in animal aggregation. *Science* 284, 99–101.
- Pimentel, D., Zuniga, R., and Morrison, D. (2005). Update on the environmental and economic costs associated with alien-invasive species in the United States. *Ecol. Econ.* 52, 273–288.
- Qian, P.Y., Li, Z., Xu, Y., Li, Y., and Fusetani, N. (2015). Mini-review: marine natural products and their synthetic analogs as antifouling compounds: 2009–2014. *Biofouling* 31, 101–122.
- Ram, J.L., Fong, P.P., and Garton, D.W. (1996). Physiological aspects of zebra mussel reproduction: maturation, spawning, and fertilization. *Am. Zool.* 36, 326–338.
- Saladino, R., Crestini, C., Neri, V., Ciciiriello, F., Costanzo, G., and Di Mauro, E. (2006). Origin of informational polymers: the concurrent roles of formamide and phosphates. *ChemBiochem* 7, 1707–1714.
- Shamchuk, A.L., Blunt, B.J., Lyons, D.D., Wang, M.Q., Gasheva, A., Lewis, C.R., Tomlin, K., Hazard, E.S., Hardiman, G., and Tierney, K.B. (2018). Nucleobase-containing compounds evoke behavioural, olfactory, and transcriptional responses in model fishes. *FACETS* 3, 79–102.

- Shikuma, N.J., Pilhofer, M., Weiss, G.L., Hadfield, M.G., Jensen, G.J., and Newman, D.K. (2014). Marine tubeworm metamorphosis induced by arrays of bacterial phage tail-like structures. *Science* 343, 529–533.
- Stokstad, E. (2007). Feared quagga mussel turns up in western United States. *Science* 315, 453.
- Symonds, M.R., and Elgar, M.A. (2008). The evolution of pheromone diversity. *Trends. Ecol. Evol.* 23, 220–228.
- Toonen, R.J., and Pawlik, J.R. (1994). Foundations of gregariousness. *Nature* 370, 511–512.
- Tošenovský, E., and Kobak, J. (2016). Impact of abiotic factors on aggregation behaviour of the zebra mussel *Dreissena polymorpha*. *J. Molluscan Stud.* 82, 55–66.
- Verkhatsky, A., and Burnstock, G. (2014). Biology of purinergic signalling: its ancient evolutionary roots, its omnipresence and its multiple functional significance. *Bioessays* 36, 697–705.
- Willan, R.C., Russell, B.C., Murfet, N.B., Moore, K.L., McEnnulty, F.R., Horner, S.K., Hewitt, C.L., Dally, G.M., Campbell, M.L., and Bourke, S.T. (2000). Out-break of *Mytilopsis sallei* (Recluz, 1849)(Bivalvia: Dreissenidae) in Australia. *Molluscan Res.* 20, 25–30.
- Wakisaka, N., Miyasaka, N., Koide, T., Masuda, M., Hiraki-Kajiyama, T., and Yoshihara, Y. (2017). An adenosine receptor for olfaction in fish. *Curr. Biol.* 27, 1437–1447.
- Wang, H.L., Zhao, C.H., and Wang, C.Z. (2005). Comparative study of sex pheromone composition and biosynthesis in *Helicoverpa armigera*, *H. assulta* and their hybrid. *Insect Biochem. Mol. Biol.* 35, 575–583.
- Wong, Y.T., Meier, R., and Tan, K.S. (2011). High haplotype variability in established Asian populations of the invasive Caribbean bivalve *Mytilopsis sallei* (Dreissenidae). *Biol. Invasions* 13, 341–348.
- Wyatt, T.D. (2014). *Pheromones and Animal Behavior: Chemical Signals and Signatures* (Cambridge University Press).
- Yebra, D.M., Kiil, S., and Dam-Johansen, K. (2004). Antifouling technology—past, present and future steps towards efficient and environmentally friendly antifouling coatings. *Progr. Org. Coating* 50, 75–104.
- Zeeck, E., Harder, T., and Beckmann, M. (1998a). Uric acid: the sperm-release pheromone of the marine polychaete *Platynereis dumerilii*. *J. Chem. Ecol.* 24, 13–22.
- Zeeck, E., Harder, T., and Beckmann, M. (1998b). Inosine, L-glutamic acid and L-glutamine as components of a sex pheromone complex of the marine polychaete *Nereis succinea* (Annelida: Polychaeta). *Chemoecology* 8, 77–84.
- Zimmer, R.K., and Butman, C.A. (2000). Chemical signaling processes in the marine environment. *Biol. Bull.* 198, 168–187.

ISCI, Volume 19




Supplemental Information

**Aggregation Pheromone for an Invasive
Mussel Consists of a Precise Combination
of Three Common Purines**

Jian He, Qi Dai, Yuxuan Qi, Zhiwen Wu, Qianyun Fang, Pei Su, Miaoqin Huang, J. Grant Burgess, Caihuan Ke, and Danqing Feng

Supplementary Information

Table S1. ESIMS and NMR data, and chemical structures of compounds 1-3, related to Figure 2 and 4.

Compound	Compound 1		Compound 2		Compound 3	
ESI-MS (<i>m/z</i>)	135.03 [M-H] ⁻		267.07 [M-H] ⁻		268.10 [M+H] ⁺	
Position	δ_{H}	δ_{C}	δ_{H}	δ_{C}	δ_{H}	δ_{C}
2	8.20 (1H, br s)	145.7 (CH)	8.08 (1H, d, <i>J</i> = 1.2 Hz)	146.4 (CH)	8.35 (1H, s)	152.9 (CH)
3	13.46 (1H, br s, N-H)		12.40 (1H, s, N-H)			
4		154.6 (C)		148.7 (C)	7.35 (2H, s, -NH ₂)	149.5 (C)
5		115.6 (C)		124.9 (C)		119.8 (C)
6		157.5 (C)		157.0 (C)		156.6 (C)
7	12.24 (1H, br s, N-H)					
8	7.98 (1H, s)	142.3 (CH)	8.35 (1H, s)	139.2 (CH)	8.14 (1H, s)	140.4 (CH)
1'			5.88 (1H, d, <i>J</i> = 5.4 Hz)	87.9 (CH)	5.88 (1H, d, <i>J</i> = 6.6 Hz)	88.4 (CH)
2'			4.49 (1H, t, <i>J</i> = 5.4 Hz)	74.6 (CH)	4.62 (1H, m)	73.5 (CH)
2'-OH			5.43 (1H, m)		5.43 (1H, m)	
3'			4.14 (1H, m)	70.8 (CH)	4.15 (1H, m)	71.1 (CH)
3'-OH			5.23 (1H, s)		5.18 (1H, d, <i>J</i> = 4.8 Hz)	
4'			3.94 (1H, m)	86.1 (CH)	3.97 (1H, m)	86.4 (CH)
5'a			3.65 (1H, dd, <i>J</i> = 12.0, 3.6 Hz)	61.7 (CH ₂)	3.68 (1H, m)	62.2 (CH ₂)
5'b			3.55 (1H, <i>J</i> = 12.0, 3.6 Hz)		3.56 (1H, m)	
5'-OH			5.05 (1H, s)		5.43 (1H, m)	
Structure						

δ in ppm, ¹H-NMR (600MHz, DMSO-*d*₆), ¹³C-NMR (150 MHz, DMSO-*d*₆).

Supplemental Figure Legends

Figure S1. *M. sallei* and its high-density aggregation and fouling, related to Figure 1, 4 and 5.

(A) *M. sallei* adult with byssus.

(B) Pediveliger larva of *M. sallei*.

(C) Natural aggregation of *M. sallei* on the submerged facilities of an oyster farm, Dongshan, China.

(D) Enlarged view of (C).

Figure S2. Detection of adenosine, inosine and hypoxanthine in ACS of three bivalve species by

HPLC, related to Figure 5 and 6. HPLC spectra of ACS of *M. sallei* (A), *P. viridis* (B) and *C. angulata* (C). (D) HPLC spectrum of a mixture of adenosine, inosine and hypoxanthine standards.

Figure S1





Figure S2



Transparent Methods

Larval culture of *Mytilopsis sallei*

M. sallei adults were collected from the submerged ropes of a fish farm in Maluan Bay, Xiamen, China (24°33'N, 118°01'E). Spawning induction and larval culture were carried out in the laboratory as previously described (He et al., 2015). Veligers were fed with *Dicrateria zhanjiangensis* (Chrysophyta) at a concentration of $1.0\text{-}5.0\times 10^4$ cells mL⁻¹. Pediveligers competent to settle and metamorphose were harvested after 6-8 d of incubation and used for settlement assays.

Preparation of conditioned seawater

Conditioned seawater was prepared as follows. *M. sallei* adults were cleaned by scrubbing and rinsing their shells to remove epifaunal organisms, and their byssus was gently removed with scissors. Adult-conditioned seawater (ACS) was prepared by placing 50 healthy individuals (2-3 cm shell length) in 4 L of 0.22- μm -filtered artificial seawater (FSW) for 2 h, 4 h or 8 h. The artificial seawater was prepared according to Berges et al. (2001). The shell-conditioned seawater (SCS) of *M. sallei* was prepared by placing the empty shells from 50 freshly killed individuals in 4 L of FSW for 2 h, 4 h or 8 h. The conditioned seawater was 0.22- μm filtered before bioassays.

Preparation of fractions of mantle cavity fluid (MCF)

MCF of *M. sallei* adults was collected as follows. *M. sallei* adults were placed on a petri dish and broken open with scissors, then the fluid was collected with an Eppendorf pipette. All operations were performed on ice. After centrifugation of the MCF (12000 rpm at 4 °C for 20 min), 50 mL of the supernatant was successively partitioned with equal volumes of ethyl acetate (EtOAc) and n-butanol, and these fractions were evaporated to dryness under reduced pressure. The residual aqueous fraction was freeze-dried.

Isolation of the conspecific inducing cues

One liter aqueous fraction of MCF collected from 3000 *M. sallei* adults was prepared as described above. The aqueous fraction was sequentially separated by ultrafiltration with centrifugal filters (Amicon[®] Ultra-15, Millipore Corporation, USA) with cutoffs of 100 and 10 KDa. Three fractions were obtained: >100, 10-100 and <10 KDa. They were freeze-dried and tested for activity on larval settlement and metamorphosis following the procedures described below. Inducing activity was recognized in the <10 KDa (F3) fraction which was further subjected to Sephadex LH-20 chromatography (25×1500 mm; Amersham, Piscataway, NJ) eluted with 70% methanol at a flow rate of 0.1 ml min⁻¹. Four sub-fractions (F3-1, F3-2, F3-3 and F3-4) were obtained. F3-4 showed significant settlement-inducing activity. Accordingly, F3-4 was further purified via HPLC (HPLC-pump 420, gradient former 425, detector Uvikon 720 LC, 254 nm) with a preparative column Xbridge[®] BEH Prep Amide (250×20 mm, 5 µm, Waters), eluted by a mobile phase of 85% acetonitrile at 7 mL min⁻¹, to yield four fractions (F3-4-1, F3-4-2, F3-4-3 and F3-4-4). It was found that F3-4-2 and F3-4-3 were settlement-inducing. F3-4-2 was a pure compound (**1**). F3-4-3 was further purified by HPLC using a preparative column Xbridge[®] BEH Prep Amide (250×10 mm, 5 µm, Waters), eluted by a mobile phase of 85% acetonitrile at 4 mL min⁻¹, to yield active compounds **2** and **3**.

Settlement and metamorphosis assays

To confirm that *M. sallei* adults can release conspecific cue(s), ACS and SCS, prepared as described above, were tested for effects on larval settlement and metamorphosis. There were six treatments, including 2 h- ACS, 4 h- ACS, 8 h- ACS, 2 h- SCS, 4 h- SCS and 8 h- SCS. Assays of larval settlement and metamorphosis were performed as previously described (He et al., 2017). Briefly, assays were conducted in sterile 6-well polystyrene plates (each well with 3.5 cm in diameter and 2.0 cm in height). A volume of 50 µL of each treatment was added to each well containing 9.95 mL FSW. Approximately 30-40 pediveligers of *M. sallei* were randomly added to each well containing

10 mL of a test solution. Petri plates were then maintained at 28 °C in the dark. After a 48 h incubation, the total number of settled larvae and metamorphosed larvae were counted through a Leica inverted microscope (DM IL LED). Throughout this work, experiments were carried out in triplicate. FSW was used as a control.

To determine the existence of conspecific cue(s) in MCF, bioassays were carried out to examine the larval response to the three fractions of MCF following the procedures as described above. Each fraction of MCF was tested under the concentrations of 10, 20, 50 and 100 $\mu\text{g mL}^{-1}$.

To examine the activity of the sub-fractions obtained during the bioassay-guided fractionation of conspecific cue(s), bioassays were performed to test the larval response to each sub-fraction following the procedures as described above. The tested concentrations for each sub-fraction were shown in Figure 2.

To confirm that the conspecific cues (identified as three purines, see the results) are not only active under still-water condition, but also active under hydrodynamic conditions, as in the natural environment, we performed the flume experiment with compound **2**, which was identified as adenosine (Ado, see the results). The reason for running this test with Ado was that we aimed to choose one active purine as an example to confirm the inducing activity under flow, and Ado exhibited the highest inducing activity amongst the three active purines in the still-water tests (Figure 4B). The flow experiment was conducted in a plastic racetrack flume (Figure 3A), consisting of two semicircular ends (10-cm radius) and two straight sections (40-cm long). FSW (10 L) were added into the flume, with a seawater depth of 9 cm. Water flow was generated through the use of a controlled motor-driven paddle wheel installed in one straight section. The direction of flow is shown in Figure 3A, indicated by the arrows. In the other straight section, six substrates (three treatments and three controls) were fixed 1 cm depth below the seawater surface. The substrate was prepared with a 30- μm mesh net fixed to the bottom of a PVC pipe (inner diameter 3.6 cm, height 2.5 cm). The pipes were filled with plastic sponge and stuck to the flume wall with double-sided tape,

leaving the substrates (i.e. the mesh nets at the bottom of the pipe) 1 cm below the seawater surface. In the flume, the flow speed was controlled to 2-3 cm/s using a flow controller. About 20,000 larvae were added into the flume. In the treatment pipe, 100 mL test solution of 10 μ M Ado in FSW was added using a separating funnel and slowly released into the plastic sponge at rate of 2 mL/h. Ado was thus absorbed by the sponge and slowly released into the seawater surrounding the substrate. In the control pipe, there was no compound released into the sponge. After 12, 24 and 48 h, the numbers of larvae settled onto the mesh nets of the treatments and controls were counted.

Identification of the conspecific inducing cues

Nuclear magnetic resonance (NMR) spectra of the purified compounds (**1**, **2**, and **3**) dissolved in deuterium oxide were recorded with a Bruker Advance III spectrometer operating at 600 MHz for ^1H and 150 MHz for ^{13}C , respectively. Chemical shifts (δ) are reported in parts per million (ppm), and coupling constants (J) are in Hz. ESIMS spectral data were measured in the positive- and negative-ion mode on the Thermo Fisher Q-Exactive mass spectrometer. Mass spectra were recorded across the range m/z 100-1000. Structural elucidation of the compounds was based on their spectral data (NMR, ESIMS) and comparison with published values.

Identification of purines in MCF by LC-MS and their effect on larval settlement and metamorphosis

A Thermo Fisher Q-Exactive mass spectrometer was used with electrospray ionization in positive or negative ion mode in the 100–1000 AMU range. Solvents were water with 10 mM ammonium acetate (A) and methanol (B). The column temperature was maintained at 30 $^{\circ}\text{C}$ and the solvent flow was 1.0 mL min^{-1} . A 10- μ L MCF was separated on a reversed-phase column C18 SB-AQ column (250 mm \times 4.6 mm, 5 μ m), which was eluted with a solvent composition starting with 98:2 (A:B) for 10 min followed by a gradient to 20:80 in 20 min and an additional 10 min at 20:80. UV absorption was monitored at 254 nm.

Five purine metabolites, namely hypoxanthine (Hyp), inosine (Ino), Ado, xanthine (Xan), and adenine (Ade), were detected in MCF. They were tested for activity on larval settlement and metamorphosis. The standard compounds Ado, Ino, Hyp, Ade and Xan (purity \geq 99%), purchased from Sigma-Aldrich Co. Ltd (St Louis, MO, USA), were used. Test solutions of each compound in FSW at the concentrations of 0.1, 0.5, 1, 5, 10 and 50 μ M were assayed with *M. sallei* larvae according to the procedures described above.

Determination of the synergistic effect of Ado, Ino and Hyp from *M. sallei* ACS

To determine the lowest effective concentrations of Ado, Ino and Hyp needed for ACS to exhibit inducing activity, we performed the following assays. Firstly, 4 h-ACS was further diluted with FSW into 2, 5 and 10-fold dilutions. These were then tested for activity on larval settlement and metamorphosis, to determine the highest active dilution. Secondly, the concentrations of Ado, Ino and Hyp in *M. sallei* ACS were determined by HPLC. The quantification was performed by external calibration with standards. Each compound (3.0 mg) was dissolved in 20 mL Milli-Q water to a concentration of 150 μ g mL⁻¹. The three different purine solutions were mixed, then the mixed purine solution was diluted to 2, 5, 10, 20, and 50 μ g mL⁻¹. A standard curve was obtained by plotting the peak area vs. concentration by HPLC using a C18 SB-Aq column (250 mm \times 4.6 mm, 5 μ m). The solvents were 7.5 mM potassium phosphate buffer (A) and methanol (B). The column temperature was maintained at 30 $^{\circ}$ C and the solvent flow was at 1.0 mL min⁻¹. Elution with a solvent composition starting with 98:2 (A:B) for 10 min followed by a gradient to 20:80 in 20 min and an additional 10 min at 20:80 was performed. The eluant was monitored by UV detection at 254 nm. Preparation of ACS samples was performed as above and subject to HPLC. Purine quantification was achieved based on regression analysis of peak area against concentration.

Since the concentrations of Ado, Ino and Hyp in the highest effective dilution of ACS, the 5-d ACS, were found to be much lower than the lowest effective concentration of each compound when tested individually, the synergistic effect of these three compounds was tested. The response of

larvae to a mixture of the three purines (8 nM for Ado, 9 nM for Ino, and 26 nM for Hyp, i.e., the same concentrations as determined for the 5-d ACS) was investigated, and compared to the response to individual compounds under the same concentrations of each compound separately in the mixture, with the 5-d ACS used as positive control.

Effects of different ratios of Ado, Ino and Hyp on larval settlement and metamorphosis

A simplex-lattice mixture design (Das et al., 2016), which was programmed using Design-Expert (version 11.0) software (Stat-Ease Inc., Minneapolis, USA), was used to evaluate the effects of different ratios of the three purines on larval settlement and metamorphosis. Here, the test levels used for each purine compound were designed as 0, 1/25, 1/5 and 1. Fractions of each level were used, and the sum of contributory fractions for each ratio treatment was one. For example, for the treatment of 5:5:15, the sum of its contributory fractions was $5/25 + 5/25 + 15/25 = 1$. A total of 33 treatments with different ratios of Ado, Ino and Hyp with the same total concentration of compounds (50 nM) were designed. The coded level of simplex lattice mixture design is shown in Figure 6A. The total concentration of compounds was 50 nM and was chosen because the sum concentration of the three purines in the effective 5-d ACS was close to 50 nM. Each mixture was tested for its effect on larval settlement and metamorphosis.

Determination of concentrations of Ado, Ino and Hyp in ACS of two species of bivalve molluscs

Two other bivalve species, the mussel *Perna viridis* and the oyster *Crassostrea angulata*, with habitats similar to *M. sallei*, were studied. The concentrations of Ado, Ino and Hyp in their ACS measured as follows. ACS of *P. viridis* and *C. angulata* were prepared by placing five healthy individuals (with total weight 200-250 g, similar to the weight of *M. sallei* used in preparing *M. sallei* ACS) in 4 L FSW for 4 h. Their ACS was subjected to HPLC as described above. Purine quantification was carried out based on regression analysis of compound peak area against concentration.

Statistical analysis

Results were analyzed with SPSS 17.0 software. All data expressed in percentages of larval settlement and metamorphosis were arcsine-transformed prior to analysis. For the bioassays listed in Table 1, one-way analysis of variance (ANOVA) was performed with a Dunnett's post-hoc test for multiple comparisons of treatment means with a control. For the flume experiment, Student's t-test was used to evaluate the statistical significance between two data sets.

Supplemental References

- Berges, J. A., Franklin, D. J., and Harrison, P. J. (2001). Evolution of an artificial seawater medium: improvements in enriched seawater, artificial water over the last two decades. *J. Phycol.* 37, 1138–1145.
- Das, S. K., Khanam, J., and Nanda, A. (2016). Optimization of preparation method for ketoprofen-loaded microspheres consisting polymeric blends using simplex lattice mixture design. *Mat. Sci. Eng. C-Mater.* 69, 598–608.
- He, J., Qi, J. F., Feng, D. Q., and Ke, C. H. (2015). Embryonic and larval development of the invasive biofouler *Mytilopsis sallei* (Récluz, 1849)(Bivalvia: Dreissenidae). *J. Mollus. Stud.* 82, 23–30.
- He, J. Qi, J. F., Huang Y. Q., Sheng, Y. Q., Su, P., Feng, D. Q., and Ke, C. H. (2017). Larval settlement and metamorphosis of the invasive biofouler, *Mytilopsis sallei*, in response to ions and neuroactive compounds. *Mar. Biol. Res.* 13, 645–658.