

Research article

Open Access

Antisense RNA decreases AP33 gene expression and cytoadherence by *T. vaginalis*

V Mundodi, AS Kucknoor and JF Alderete*

Address: Department of Microbiology, University of Texas Health Science Center, 7703 Floyd Curl Drive, San Antonio, TX 78229-3900, USA

Email: V Mundodi - mundodi@uthscsa.edu; AS Kucknoor - kucknoor@uthscsa.edu; JF Alderete* - alderete@uthscsa.edu

* Corresponding author

Published: 3 July 2007

Received: 31 October 2006

BMC Microbiology 2007, **7**:64 doi:10.1186/1471-2180-7-64

Accepted: 3 July 2007

This article is available from: <http://www.biomedcentral.com/1471-2180/7/64>

© 2007 Mundodi et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Host parasitism by *Trichomonas vaginalis* is complex. Adherence to vaginal epithelial cells (VECs) is mediated by surface proteins. We showed before that antisense down-regulation of expression of adhesin AP65 decreased amounts of protein, which lowered levels of *T. vaginalis* adherence to VECs. We now perform antisense down-regulation of expression of the *ap33* gene to evaluate and confirm a role for AP33 in adherence by *T. vaginalis*. We also used an established transfection system for heterologous expression of AP33 in *T. foetus* as an additional confirmatory approach.

Results: We successfully select stable trichomonads with sense (S) and antisense (AS) plasmids. RT-PCR confirmed decreased amounts of *ap33* mRNA in AS-transfected parasites, and decreased amounts of AP33 had no effect on growth and viability when compared to wild-type (wt) trichomonads. Immunoblots of proteins from AS-transfectants gave significant decreased amounts of functional AP33 capable of binding to host cells compared to wt- and S-transfected trichomonads. As expected, AS-transfectants had lower levels of adherence to VECs, which was related to reduction in surface expression of AP33. Stable expression of *T. vaginalis* AP33::HA fusion in *T. foetus* was confirmed by immunoblots and fluorescence. The episomally-expressed surface AP33::HA fusion increased adherence of trichomonads to human VECs, which was abrogated with anti-AP33 serum.

Conclusion: These results using both antisense inhibition of gene expression and AP33 synthesis and the heterologous expression of AP33 in *T. foetus* confirms a role for this protein as an adhesin in *T. vaginalis*.

Background

The protozoan *Trichomonas vaginalis* is responsible for the number one, non-viral sexually transmitted disease (STD) worldwide [1]. There are ~9 million new cases of vaginitis in the US alone [2-4]. Trichomonosis causes serious health consequences for women, including preterm delivery, low birth weight infants, infertility cervical cancer, pelvic inflammatory disease and infection by other STD

agents [5-9]. Trichomonosis also predisposes humans to HIV by increasing the portal of entry and exit of virus [10]. A recent study showed a relationship between trichomonosis and prostate cancer [11]. Unlike other STDs, the prevalence of *T. vaginalis* does not decrease with age [12,13]. Given the significant human morbidity caused by *T. vaginalis*, there is an urgency towards identifying virulence fac-

tors, elucidating the mechanisms of pathogenesis, and developing interference strategies.

Adherence by *T. vaginalis* to vaginal epithelial cells (VECs) is preparatory for colonization and infection. Identification and characterization of five surface proteins (AP120, AP65, AP51, AP33 and AP23) involved in attachment to VECs has provided an understanding, in part, of the molecular basis of host cell adherence by this parasite [14-17]. The adhesins interact with host cells via ligand-receptor interactions [16-18]. Not surprisingly and as expected, there is a direct relationship between surface expression of adhesins and levels of cytoadherence [16,19]. All members of the adhesin gene families are coordinately up-regulated by iron [16,17,20], and iron appears important for compartmentalization and surface placement of the proteins [17]. Interestingly, the adhesins have sequence identity to metabolic enzymes located in the double membrane bound hydrogenosome organelle [18,21-24].

Adding to the complexity of these proteins is the fact that the genes encoding the adhesins are members of multi-gene families [14,21,22], which makes individual gene knockout approaches impractical for genetic-molecular studies. Therefore, a recent study by us used antisense technology as a genetic approach to confirm the importance of the prominent AP65 adhesin in adherence to VECs [17,25]. In addition, as an alternative approach to confirm AP65 function we established a transfection system for heterologous expression of the *T. vaginalis* AP65 in *T. foetus* and showed surface placement of AP65 that led to higher levels of *T. foetus* attachment to VECs [26]. In this report, we demonstrate a role of AP33 in parasite adherence to the host cells by antisense inhibition of *ap33* expression. Furthermore, we show heterologous expression of AP33 on the surface of *T. foetus*, which elevated *T. foetus* cytoadherence.

Results

Plasmid construction and isolation of stable transfectants

We wanted to use the antisense approach to silence expression of *ap33*, as before [25]. Plasmid constructs containing the DNA fragment representing the coding region of the *ap33-1* gene [24] in the sense (S) or antisense (AS) orientations were generated. Transfected *T. vaginalis* isolate T016 parasites were selected using 200 µg per ml G418, and resistant S- and AS-transfected trichomonads were cloned in soft agar. To confirm the presence of the plasmids in drug-resistant trichomonads, PCR was performed to amplify a 795-bp coding region of *neo* using template DNA from wild type (wt) organisms and S- and AS-transfectants. As expected, Figure 1A shows the *neo* PCR products in S- and AS-transfected trichomonads harboring the plasmids. No PCR product was obtained from wt organisms.




Figure 1
Transfection and RT-PCR showing reduced *ap33* mRNA levels in *T. vaginalis* trichomonads transfected with the antisense plasmid. Part A shows PCR amplification of the *neo* coding region in transfected parasites. The ethidium bromide (EtBr)-stained band after electrophoresis in 1% agarose is the PCR product of the *neo* gene that was amplified using DNA from transfected *T. vaginalis* parasites. As expected, no PCR product was expected from wt organisms, and a predicted product was obtained from the plasmid used directly during PCR as a control (part A, lane labeled C). Part B gives bands after electrophoresis as in part A showing RT-PCR products for the *ap33* S (panel a) and AS transcript (panel b). The RT-PCR product for *ap65* and α -tubulin are presented in panels c and d, respectively. Part C illustrates the quantitation of the transcript bands in part B (panel a) for the *ap33* transcript in AS-transfected trichomonads compared to S-transfected and wt parasites. The bar graph shows the relative amounts of RT-PCR products for *ap33*. The amount of wild type *ap33* transcript was normalized 100%. Quantification was done following Scion image β program.

Antisense mRNA modulates amounts of *ap33*

We next performed RT-PCR on total RNA isolated from S- and AS-transfectants and wt *T. vaginalis* as shown in Figure 1B. Results show the decreased intensity of the 580-bp PCR product of the *ap33* coding region in the AS-trichomonads compared to S- and wt organisms (panel a). In contrast, the primers to the control *ap65* gene gave equal intensities of 500-bp RT-PCR product (panel c), and this confirms specificity in the antisense inhibition of *ap33* and the use of equivalent amounts of total RNA in the reactions. Only the AS-transfectants yielded a product from RT-PCR when specific primers were used to amplify a 525-bp product of the antisense transcript (panel b). As an additional control to show equal amounts of RNA, RT-PCR was performed to obtain a 650-bp product with primers specific to the α -tubulin gene (panel d), which is constitutively expressed in trichomonads. Finally, we performed the Scion image β program to quantitate RT-PCR products. Figure 1C shows an ~70% decrease in amounts

of *ap33* transcript (Figure 1B, panel a) when compared to the wt organisms and S-transfectants. As additional controls to show that transfection alone had no effect on *ap33* mRNA levels, trichomonads were transfected with plasmid without insert and with plasmid carrying the *ap65* antisense [25]. No effect on *ap33* transcript levels by RT-PCR were seen for these controls. It is noteworthy that the primers used for RT-PCR of *ap33* cross-hybridize with all three *ap33* genes of the multigene family. Therefore, as all three *ap33* genes are expressed under these growth conditions and given the decrease in mRNA by $\geq 70\%$, the RT-PCR data indicate that antisense RNA expression decreases the amounts of endogenous transcripts of all three *ap33* genes.

Antisense decreased amounts of AP33

An immunoblot was performed using as probe monoclonal antibody (mAb) F5.2 to AP33 [24]. As shown in panel 1 of Figure 2A, the lanes with total proteins derived from equal numbers of wt (lane 1) and S-transfected parasites (lane 2) had readily detectable AP33 compared to AS-transfected parasites (lane 3). In contrast, the mAb 12G4 to AP65 shown in panel 2 had equal amount of protein for each sample in duplicate blots of panel 1, indicating that the effect of *ap33* antisense RNA is specific. In panel 3 the mAb to α -tubulin reaffirms that equal amounts of protein was added to each lane and served as another internal control. Figure 2B presents an immunoblot using mAb F5.2 to AP33 after a ligand assay to measure amounts of functional AP33. There was decreased amounts of AP33 bound to VECs from extracts of AS-transfectants (lane 3) compared to wt (lane 1) and S-transfected parasites (lane 2). The amount of AP33 seen in lane 3 as determined by the Scion image β program was decreased by $\sim 70\%$. This extent of inhibition of both mRNA seen above (Figure 1B, panel a) and protein synthesis is evidence that all three *ap33* genes were down-regulated in expression.

Reduction in amount of AP33 does not affect parasite growth

Because AP33 is β -succinyl coenzyme synthetase also localized to the hydrogenosome where ATP is generated through the oxidation of pyruvate [27], we felt it important to determine whether inhibition of expression of this gene affected overall parasite growth and multiplication. As illustrated in Figure 3, there was no difference in the growth rates between wt and AS-transfected parasites nor was there any adverse effect on the viability and motility of trichomonads, as before [25].

Decreased surface expression of AP33 and lower levels of adherence by AS-transfectants

Immunofluorescence experiments were then performed to examine surface expression of AP33. Figure 4 demon-




Figure 2
Immunoblot analysis of S- and AS-transfected trichomonads and wt *T. vaginalis* organisms. Part A presents triplicate blots of total protein from 10^7 trichomonads after SDS-PAGE on 10% acrylamide before blotting onto Hybond-P membranes. The blots were probed with mAb F5.2 to AP33 (panel 1), mAb 12G4 to AP65 (panel 2) [17], and mAb to α -tubulin (panel 3). Part B presents immunoblot results from a ligand assay, which was performed using extracts from equal numbers of organisms.

strates the decreased intensity of fluorescence by polyclonal rabbit anti-AP33 serum antibody characterized previously [28] in the non-permeabilized AS-transfected organisms (panel C) compared with wt (panel A) and S-transfected organisms (panel B) handled identically. These results are in agreement with data presented above on the decreased amounts of total AP33 in AS-transfectants. Under no conditions was any fluorescence detected using normal rabbit serum antibody as a control (not shown). Furthermore, the level of adherence by AS trichomonads was measured and found to be $\sim 30\%$ lower than wt and S-transfected parasites (Figure 5), and this decrease in extent of adherence is not unexpected given the availability of alternative surface adhesins, such as the prominent AP65 adhesin [17], for cytoadherence. In separate control experiments performed simultaneously, trichomonads transfected with plasmid without insert and plasmid carrying the *ap65* antisense had no effect on amounts of AP33. As with the S transfectants, no decrease




Figure 3
Representative growth kinetics of wt organisms and S- and AS-transfected trichomonads cultivated in batch culture. Medium was inoculated with 10^5 parasites, and cell densities at different time points were enumerated using a hemocytometer. Similar results were obtained from three growth experiments performed independently at different days.

in growth and adherence was evident by trichomonads transfected with plasmid without insert. Thus, these data suggest strongly that the effect was due to absence of surface AP33 and not to nonspecific events such as those that might result from transfection alone.

Episomal expression of *T. vaginalis* AP33 elevates adherence levels in *T. foetus*

As before [26], heterologous expression in *T. foetus* parasites is an additional approach for studying the function of *T. vaginalis* virulence genes. Therefore, *T. foetus* were transfected with the S plasmid to generate a fusion protein containing hemagglutinin (AP33::HA). Stable transfectants were obtained and characterized, also as before [26]. Episomal expression of AP33::HA was verified by RT-PCR (data not shown), and synthesis of AP33::HA was demonstrated by immunoblot analysis using mAb F5.2. Figure 6A (lane 2) shows the higher molecular weight AP33::HA fusion protein band in total protein blots of transfected *T. foetus*, but not wt *T. foetus* (lane 1). Bands common to lanes 1 and 2 illustrate the crossreactivity by mAb 5.2 with the *T. foetus* equivalent protein(s). Not unexpectedly based on our previous work [28], the lower-sized bands in lane 2 detected by mAb F5.2 are degraded AP33. Duplicate blots were also probed with mAb to HA (Figure 6B),




Figure 4
Immunofluorescence showing decreased surface expression of AP33 in non-permeabilized, AS-transfected (panel C) compared to S-transfected (panel B) and wt trichomonads (panel A). Rabbit polyclonal anti-AP33 serum characterized previously [24] was used as described in Methods.

and as expected, this mAb detected the fusion AP33::HA protein in transfectants (lane 2) but not in wt *T. foetus* (lane 1). The bands common to lanes 1 and 2 are nonspecific reactions by mAb with *T. foetus* proteins. The absence of detection of lower-sized, degraded AP33 in lane 2 is because the mAb is to HA, not AP33. We then performed fluorescence on non-permeabilized wt and AP33::HA transfected organisms. As can be seen in Figure 6C, the HA mAb only detected *T. foetus* trichomonads expressing the fusion protein, again providing evidence for the surface placement of AP33::HA. Finally and importantly and as before [26], *T. foetus* transfected with plasmid without insert gave results in all assay identical to those seen for wt *T. foetus*.




Figure 5
Relative adherence of AS-transfectants (bar AS) compared with wt-(bar wt) and S-transfected (bar S) parasites. *T. vaginalis* parasites expressing antisense display decreased adherence to MS-74 VECs. The extent of adherence by wt organisms was normalized to 100% for comparative purposes. The results are the average from four different experiments. Each experiment was performed in quadruplicate samples. The asterisk illustrates that this extent of decrease was statistically significant ($p < 0.05$).

Lastly, we performed an adherence assay for *T. foetus* expressing AP33::HA. As can be seen in Figure 7, episomal expression of AP33::HA (bar Tf-AP33) had elevated levels of adherence to VECs when compared to background levels seen for wt *T. foetus* (bar Tf). Levels of adherence was compared with those of *T. vaginalis* (bar Tv), which was normalized to 100%. Importantly, the increased level of adherence was attributed to surface AP33, as anti-AP33 antibody pretreatment of the transfectants (bar Tf-AP33 + antibody) gave levels of adherence equal to wt organisms (bar Tf). These experiments reinforce the idea that AP33 indeed is surface expressed and involved in the property of trichomonal adherence to VECs.

Discussion

Our recent work with antisense RNA-mediated inhibition of expression of AP65 [25] coupled with the alternative approach of heterologous expression in the *T. foetus* trichomonad of AP65 [26] provided additional confirmatory evidence for a role of this surface protein in *T. vaginalis* adherence to VECs. These approaches are important to obtaining experimental evidence on a functional role for trichomonad virulence factors, such as adhesins.




Figure 6
Immunoblot and fluorescence analyses showing episomal expression of AP33::HA in *T. foetus*. A and B: Total protein from 10^7 parasites were separated on 10% SDS-PAGE and blotted on to Hybond-P membrane. The blots were probed with mAb F 5.2 to AP33 (A) and mAb to HA (B). C: Immunofluorescence was performed using anti-HA mAb on non-permeabilized wt (Tf) and transfected trichomonads (Tf-AP33::HA). Brightfield microscopy shows the same trichomonads in the field used for fluorescence microscopy.

This is especially the case when virulence genes are members of multigene families [21-24], all of which are coordinately expressed [20-24] and in which individual gene knockouts may be difficult or impossible to achieve.




Figure 7

***T. foetus* with surface AP33::HA displays enhanced level of adherence to immortalized MS-74 VECs.**

Transfected trichomonads with AP33::HA (bar labeled Tf-AP33) had higher levels of adherence compared to wt *T. foetus* (bar Tf), as before for heterologously-expressed AP65 [26]. The percent level of adherence was adjusted to that seen for *T. vaginalis* (Tv). Trichomonads expressing AP33::HA were pretreated with anti-AP33 antiserum (bar Tf-AP33 + antibody). The results are the average from four different experiments, and each experiment was carried out using quadruplicate samples.

We now undertook similar characterization of AP33. AS-transfectants decreased in *ap33* mRNA expression (Figure 1) had correspondingly less amounts of total and surface AP33 (Figures 2 and 4), which reflected lower levels of adherence to host cells (Figure 5). The fact that there was at best a 30% reduction in adherence to VECs in AS-transfectants is not problematic. These trichomonads have four additional surface proteins (AP120, AP65, AP51 and AP23) that provide redundant and alternative functions for adherence [14,17,23]. In fact, this extent of lowered levels of adherence is in agreement with previous data from inhibition experiments using anti-AP33 antibody and recombinant AP33 [24]. In addition, heterologous expression and surface placement of *T. vaginalis* AP33 (Figure 6) gave enhanced binding by *T. foetus* to human VECs (Figure 7), consistent with a recent report on heterologous expression of AP65 in *T. foetus* [26]. Altogether, this data affirms a role for AP33 in trichomonal adherence to host cells.

It is important to point out that the mechanism is unknown by which antisense inhibits gene expression in trichomonads. In our earlier work [25] and in this study we see decreased amounts of *ap33* mRNA and AP33. As in

other systems [29,30], translation arrest or degradation of mRNA following antisense-RNA-mRNA interactions may be occurring to account for the lower levels of *ap33* transcript.

It was not surprising that decreased amounts of AP33 (α -SCS) did not affect overall trichomonal growth and multiplication (Figure 3) despite the fact that AP33 has sequence identity to α -succinyl CoA synthetase (α -SCS). No effect on parasite viability and growth parameters was also evident upon down-regulation of *ap65* expression [25]. It is known that the parasites are capable of surviving without the hydrogenosome enzymes. For instance, MR100 is a drug-resistant isolate [31] lacking the adhesins-enzymes [17], and these parasites are unable to cytoadhere to VECs [17]. Indeed, parasites are capable of generating energy (ATP) through alternative metabolic pathways [32].

Recent reports [33,34] have suggested that trichomonad LPG may also be a mediator of adherence to host cells. LPG mutants showed a decreased ability to cytoadhere. These findings may suggest additional mechanisms by which the parasite successfully colonizes the host. Such redundancy is not surprising given the existence of multiple mechanisms by which pathogens target host cells and tissues [35]. In this report and an earlier paper [25] we decrease synthesis of two protein adhesins by antisense and, as predicted, show concomitant lower levels of adherence. These latter findings are also consistent with the numerous reports supporting a role for surface proteins as adhesins [15-18,20-26,28,36]. In addition, recent work on the relationship between polyamine metabolism and putrescine secretion on *T. vaginalis* adherence and cytotoxicity [36], while reinforcing the role of surface protein adhesins, illustrates the complexity of this adherence phenotype, which also involves host protein acquisition [37] and cysteine proteinases [38,39]. This work [36] establishes a testable hypothesis that has the potential to integrate the various adherence models. Therefore, further work will be needed to fully understand how these seemingly distinct mechanisms of adherence are integrated.

Conclusion

These results using both antisense inhibition of gene expression and AP33 synthesis and the heterologous expression of AP33 in *T. foetus* confirms a role for this protein as an adhesin in *T. vaginalis*. Further, it is clear that antisense technology and heterologous expression in a different *Trichomonas* species are enabling experimental approaches to investigate and dissect the complex process of parasite adherence to host cells. This is especially crucial when other genetic approaches are untenable. Notwithstanding the possible existence of additional adherence mechanisms, the available data continue to

support a role for surface AP33, and the other protein adhesins, in parasite recognition and binding to host cells.

Methods

Parasite culture and host cells

Trichomonas vaginalis isolate T016 and *Tritrichomonas foetus* (02–97) were grown in trypticase-yeast extract-maltose (TYM) medium with 10% heat-inactivated horse serum [40]. Immortalized MS-74 VECs used before [17,25] were grown in D-MEM (Invitrogen-Life Technologies, Carlsbad, CA) supplemented with 10 % fetal bovine serum at 37°C in presence of 5% CO₂.

Generation of sense (S) and antisense (AS) plasmids with ap33 coding region

The S and AS plasmids, designated pBS-*neo-ap33*-S and pBS-*neo-ap33*-AS, respectively, were constructed by cloning the coding region of *ap33* gene in forward (S primer, 5'-CATACGCATATGCTCGCAGGCGACTTCTC-3' and AS primer, 5'-GATCTTGGTACCATTCTCTTCATCTCCTCG-3') and reverse orientation (S primer, 5'-CATACGGGTAC-CATGCTCGCAGGCGACTTCTC-3' and AS primer, 5'-GATCTTCATATGCCATTCTCTTCATCTCCTCG-3'). The original plasmid [41] was used to generate the plasmid used by us recently [25] to study the effect of antisense down-regulation of *ap65* expression on adherence. This plasmid was modified as described [25] to replace the S and AS *ap65* with the S and AS *ap33*. Briefly, after removal of the *ap65* gene in the plasmid by partial digestion, the *ap33-1* gene [24] of 0.9-kb was cloned into the *Nde*I and *Asp*718 sites. The authenticity of S- and AS-*ap33* plasmids were confirmed by sequencing. Plasmid DNA for transfection was purified using maxi prep columns (Qiagen, Inc., Valencia, CA).

Stable transfection and selection for G418 resistance

Transfection of *T. vaginalis* and *T. foetus* cells was carried out by electroporation [42]. Parasites at early logarithmic phase of growth were used for transfection. Briefly, 4 × 10⁷ parasites were centrifuged at 1800 rpm at 4°C, and the pellet resuspended in 400 µl fresh TYM before transferring into a 4-mm gap cuvette (BTX®, Genetronics, Inc., San Diego, CA) with 25 µg of plasmid DNA. Electroporation was performed at 320 V, 1000 microfarads and 725 ohms using ECM 630 Electro cell manipulator (BTX®). Following the pulse, cells were placed on ice for 10 min and transferred into two T25 flasks with 50 ml of fresh TYM-serum medium. The cells were grown free of drug for 24 h followed by the addition of Geneticin (G418) (Invitrogen) at 200 µg ml⁻¹. Single cells were cloned using soft-agar plates containing 25 µg ml⁻¹ G418. Four different clones were further analyzed, and representative data from one of the clones is presented in this report. The DNA was isolated from single cell cultures using DNAzol (Invitrogen) and further purified by phenol-chloroform

extraction. The presence of plasmid in numerous single cell clones was confirmed by PCR amplification of the *neo* gene using the *neo*-sense primer 5'GATCGGTACCAT-GATTGATTGAACAAGATGGATTG-3' and *neo*-antisense primer 5'CTTTAGACCAAGTTCGTGTCAGAAGAAGCTCGT-CAA-G-3, as shown in Figure 1A.

RNA isolation and RT-PCR analysis

Total RNA was isolated from both wt and transfected parasites using the Trizol reagent (Invitrogen). For RT-PCR, 1 µg of total RNA was reverse transcribed using SuperScript II RNase H-Reverse Transcriptase (Invitrogen). Then, 10% of the reverse transcribed cDNA was used as template for the PCR reactions. The primers used for PCR amplifications of the *ap33* transcript were as follows: *ap33*-sense primer, 5'CTCATTTTCGTCCCAGCTCC-3' and *ap33*-antisense primer, 5'AAACAATACCGATCTTACCG-3'. For α -*tubulin* the sense primer was 5'-ACTCTGCTGCCCTCGAG-CACGGTATC-3', and antisense primer was 5'-GAAAT-GACTGGTGCATAAAGAGC-3'. To demonstrate the synthesis of AP33 antisense transcript, total RNA was reverse transcribed using a gene-specific primer (5'-CAT-ACGCATATGCTCGCAGGCGACTTCTC-3'). The forward and reverse primers for PCR of the antisense transcript were 5'CTCATTTTCGTCCCAGCTCC-3' and 5'AAACAATACCGATCTTACCG-3', respectively.

Immunoblot detection of AP65, AP33 and AP33::HA

Total proteins of 10⁷ *T. vaginalis* and *T. foetus* organisms were obtained as before using trichloroacetic acid (TCA) [43] for sodium dodecylsulfate-polyacrylamide gel electrophoresis (SDS-PAGE) [44] prior to blotting onto Hybond-P membranes (Amersham) for immunoblot detection with mAb 12G4 to AP65, mAb F5.2 to AP33, and mAb to hemagglutinin (HA) (Sigma) for detection of the fusion AP33-HA. TCA-precipitated proteins were solubilized using electrophoresis dissolving buffer [44]. SDS-PAGE was carried out using 10% acrylamide gels. The mAbs and epitope reactivity have been described before [17,24]. Following reactivity with the mAb probes, the bands were visualized by the chemiluminescence assay with horseradish peroxidase as the color developer (Bio-Rad Laboratories, Hercules, CA).

Ligand assay to assay for functional AP33

The ligand assay to detect adhesins that bind the host cells was carried out as before [16]. Briefly, after fixation and processing of HeLa cells with glutaraldehyde, 10⁶ *T. vaginalis* cells were incubated with a trichomonad detergent extract derived from 2 × 10⁷ solubilized parasites. After incubation, cells were vigorously washed to remove unbound and loosely-associated trichomonad proteins. Cells were boiled in electrophoresis dissolving buffer to elute the HeLa cell-binding proteins followed by SDS-PAGE in 10% acrylamide. The gels were further stained

with Coomassie brilliant blue for visualization, and duplicate gel was blotted onto Hybond-P membrane for immunoblot analysis using the mAb F5.2 to AP33.

Surface AP33 detected by fluorescence

Immunofluorescence of AP33 on the surface was carried out using a modification of a recently-described procedure [17]. Briefly, 1×10^6 logarithmic phase organisms were washed twice with cold PBS and fixed with 4% paraformaldehyde for 30 min at RT. Fixed non-permeabilized organisms were washed in PBS, then blocked with 5% BSA for 1 h at RT prior to incubation for 1 h at RT of *T. vaginalis* with rabbit polyclonal anti-AP33 serum diluted 1:100 (v/v). Parasites were washed with PBS and incubated for 1 h at 37°C with fluoresceine isothiocyanate-conjugated anti-rabbit IgG (Sigma) diluted 1:100. Finally, parasites were washed twice with PBS and observed under 100× magnification using the Olympus BX41 microscope. For both wt *T. foetus* and *T. foetus* transfected with the plasmid encoding the AP33::HA fusion protein, fluorescence was performed similarly except that the mAb to hemagglutinin (HA) diluted 1:100 (v/v) was used.

VEC adherence assay

Immortalized MS-74 human VECs [17,45] were used for adherence experiments. Confluent monolayers of MS-74 VECs on individual 96-well microtiter plates were stabilized with 3% glutaraldehyde, as before [16], prior to addition of labeled parasites. Late logarithmic growth phase parasites (5×10^6) of *T. vaginalis* and *T. foetus* were washed in PBS and suspended in 1 ml of TYM without serum for labeling with calcein for 30 min at 37°C. Labeling was followed washing with PBS, and washed parasites (2.5×10^5) were added to individual wells of confluent, fixed VECs. After plates were incubated at 37°C for 30 min, the wells were washed three times with PBS. The adherent parasites were lysed with 200 µl of 0.1% Triton X-100. The level of adherence to VECs was determined by the intensity of fluorescence of the lysate at excitation absorbance of 485 nm and emission wavelength of 528 nm with Synergy™ HT Multi-Detection Microplate Reader (Bio-Tek Instruments, Inc., Winooski, VT). For adherence assays using wt and *T. foetus* transfected with the plasmid encoding the AP33::HA fusion protein, trichomonads were also pretreated with anti-AP33 serum (1:100; v/v) prior to the adherence assay. This was done to insure that the increase in adherence by *T. foetus* was due to surface AP33::HA, as before [26].

Reproducibility of experiments

Unless otherwise stated in the text, all experiments were performed numerous times and no less than on three different occasions.

Abbreviations

AP65, adhesin protein of molecular weight 65-kDa; AP33, adhesin protein of molecular weight 33-kDa; AS, anti-sense; AP33::HA, fusion protein of AP33 and hemagglutinin (HA), BSA, bovine serum albumin; mAb, monoclonal antibody; PBS, phosphate buffered saline; SDS-PAGE, sodium dodecylsulfate polyacrylamide gel electrophoresis; RT, room temperature; S, sense; Tf, *Trichomonas foetus*; Tv, *Trichomonas vaginalis* VEC, vaginal epithelial cell

Authors' contributions

VM carried out the design of the study and performed plasmid constructions, RT-PCR, immunoblots, ligand assay, growth curve, adherence assay and drafted the manuscript. ASK performed transfection and immunofluorescence. JFA participated in the design of the experiments, offered suggestions during the experiments, and helped to write the manuscript. All the authors read and approved the final manuscript.

Acknowledgements

This work was supported by Public Health Service grant AI43940 from the National Institutes of Health. Members of the laboratory are also acknowledged for their suggestions and discussion of our work.

References

1. Health Organization: **three hundred, thirty-three million new, STD curable cases in 1995.** *AIDS Wkly* 1995:15-16.
2. Cates W Jr: **Estimates of the incidence and prevalence of sexually transmitted diseases in the United States.** *American Social Health Association Panel.* *Sex Transm Dis* 1999, **26**:S2-7.
3. Weinstock H, Berman S, Cates W Jr: **Sexually transmitted diseases among American youth: incidence and prevalence estimates, 2000.** *Perspect Sex Reprod Health* 2004, **36**:6-10.
4. World Health Organization: **Global Prevalence and Incidence of Selected Curable Sexually Transmitted Infections. Overview and Estimates.** WHO, Geneva, Switzerland; 2001.
5. Cotch MF, Pastorek JG 2nd, Nugent RP, Hillier SL, Gibbs RS, Martin DH, Eschenbach DA, Edelman R, Carey JC, Regan JA: **Trichomonas vaginalis associated with low birth weight and preterm delivery. The Vaginal Infections and Prematurity Study Group.** *Sex Transm Dis* 1997, **24**:353-360.
6. Moodley P, Wilkinson D, Connolly C, Moodley J, Sturm AW: **Trichomonas vaginalis is associated with pelvic inflammatory disease in women infected with human immunodeficiency virus.** *Clin Infect Dis* 2002, **34**:519-522.
7. Viikki M, Pukkala E, Nieminen P, Hakama M: **Gynaecological infections as risk determinants of subsequent cervical neoplasia.** *Acta Oncol* 2000, **39**:71-75.
8. Zhang ZF, Begg CB: **Is Trichomonas vaginalis a cause of cervical neoplasia? Results from a combined analysis of 24 studies.** *Int J Epidemiol* 1994, **23**:682-690.
9. Gottlieb SL, Douglas JM Jr, Schmid DS, Bolan G, Iatesta M, Malotte CK, Zenilman J, Foster M, Baron AE, Steiner JF: **Seroprevalence and correlates of herpes simplex virus type 2 infection in five sexually transmitted-disease clinics.** *J Infect Dis* 2002, **186**:1381-1389.
10. Sorvillo F, Smith L, Kerndt P, Ash L: **Trichomonas vaginalis, HIV, and African-Americans.** *Emerg Infect Dis* 2001, **7**:927-932.
11. Sutcliffe S, Giovannucci E, Alderete JF, Chang TH, Gaydos CA, Zenilman JM, De Marzo AM, Willette WC, Platz EA: **Plasma antibodies against Trichomonas vaginalis and subsequent risk of prostate cancer.** *Cancer Epidemiol Biomarkers Prev* 2006, **15**:939-945.
12. Huppert JS, Batteiger BE, Braslins P, Feldman JA, Hobbs MM, Sankey HZ, Sena AC, Wendel KA: **Use of an immunochromatographic assay for rapid detection of Trichomonas vaginalis in vaginal specimens.** *J Clin Microbiol* 2005, **43**:684-687.

13. Ohlemeyer CL, Hornberger LL, Lynch DA, Swierkosz EM: **Diagnosis of *Trichomonas vaginalis* in adolescent females: InPouch TV culture versus wet-mount microscopy.** *J Adolesc Health* 1998, **22**:205-208.
14. Moreno-Brito V, Yanez-Gomez C, Meza-Cervantes P, Avila-Gonzales L, Rodriguez MA, Ortega-Lopez J, Gonzales-Robles A, Arroyo R: **A *Trichomonas vaginalis* 120 kDa protein with identity to hydrogenosome pyruvate:ferredoxin oxidoreductase is a surface adhesin induced by iron.** *Cell Microbiol* 2005, **7**:245-258.
15. Alderete JF, Garza GE: **Specific nature of *Trichomonas vaginalis* parasitism of host cell surfaces.** *Infect Immun* 1985, **50**:701-708.
16. Arroyo R, Engbring J, Alderete JF: **Molecular basis of host epithelial cell recognition by *Trichomonas vaginalis*.** *Mol Microbiol* 1992, **6**:853-862.
17. Garcia AF, Chang TH, Benchimol M, Klumpp DJ, Lehker MW, Alderete JF: **Iron and contact with host cells induce expression of adhesins on surface of *Trichomonas vaginalis*.** *Mol Microbiol* 2003, **47**:1207-1224.
18. Alderete JF, Garza GE: **Identification and properties of *Trichomonas vaginalis* proteins involved in cytoadherence.** *Infect Immun* 1988, **56**:28-33.
19. Kucknoor AS, Mundodi V, Alderete JF: **Adherence to human vaginal epithelial cells signals for increased expression of *Trichomonas vaginalis* genes.** *Infect Immun* 2005, **73**:6472-6478.
20. Lehker MW, Arroyo R, Alderete JF: **The regulation by iron of the synthesis of adhesins and cytoadherence levels in the protozoan *Trichomonas vaginalis*.** *J Exp Med* 1991, **174**:311-318.
21. Alderete JF, O'Brien JL, Arroyo R, Engbring JA, Musatovova O, Lopez O, Lauriano C, Nguyen J: **Cloning and molecular characterization of two genes encoding adhesion proteins involved in *Trichomonas vaginalis* cytoadherence.** *Mol Microbiol* 1995, **17**:69-83.
22. O'Brien JL, Lauriano CM, Alderete JF: **Molecular characterization of a third malic enzyme-like ap65 adhesin gene of *Trichomonas vaginalis*.** *Microb Pathogen* 1996, **20**:335-349.
23. Alderete JF, Engbring J, Lauriano CM, O'Brien JL: **Only two of the *Trichomonas vaginalis* triplet AP51 adhesins are regulated by iron.** *Microb Pathogen* 1998, **24**:1-16.
24. Engbring JA, Alderete JF: **Characterization of *Trichomonas vaginalis* AP33 adhesin and cell surface interactive domains.** *Microbiol* 1998, **144**:3011-3018.
25. Mundodi V, Kucknoor AS, Klumpp DJ, Chang TH, Alderete JF: **Silencing the ap65 gene reduces adherence to vaginal epithelial cells by *Trichomonas vaginalis*.** *Mol Microbiol* 2004, **53**:1099-1108.
26. Kucknoor AS, Mundodi V, Alderete JF: **Heterologous expression in *Tritrichomonas foetus* of functional *Trichomonas vaginalis* AP65 adhesin.** *BMC Mol Biol* 2005, **6**:5.
27. Müller M: **The hydrogenosome.** *J Gen Microbiol* 1993, **139**:2879-2889.
28. Engbring JA, Alderete JF: **Three genes encode distinct AP33 proteins involved in *Trichomonas vaginalis* cytoadherence.** *Mol Microbiol* 1998, **28**:305-313.
29. Lee LK, Roth CM: **Antisense technology in molecular and cellular bioengineering.** *Curr Opin Biotech* 2003, **14**:505-511.
30. Wagner EG, Simons RW: **Antisense RNA control in bacteria, phages, and plasmids.** 1994, **48**:713-742.
31. Kulda J: ***Trichomonads*, hydrogenosomes and drug resistance.** *Int J Parasitol* 1999, **29**:199-212.
32. Brown DM, Upcroft JA, Dodd HN, Chen N, Upcroft P: **Alternative 2-keto acid oxidoreductase activities in *Trichomonas vaginalis*.** *Mol Biochem Parasitol* 1999, **98**:203-214.
33. Fichorova RN, Trifonova RT, Gilbert RO, Costello CE, Hayes GR, Lucas JJ, Singh BN: ***Trichomonas vaginalis* lipophosphoglycan triggers a selective upregulation of cytokines by human female reproductive tract epithelial cells.** *Infect Immun* 2006, **74**:5773-5779.
34. Bastida-Courcuera FD, Okumura CY, Colocoussi A, Johnson PJ: ***Trichomonas vaginalis* lipophosphoglycan mutants have reduced adherence and cytotoxicity to human ectocervical cells.** *Eukaryot Cell* 2005, **4**:1951-1958.
35. Stabbins GE, Galan JE: **Structural mimicry in bacterial virulence.** *Nature* 2001, **412**:701-705.
36. Garcia AF, Benchimol M, Alderete JF: ***Trichomonas vaginalis* polyamine metabolism is linked to host cell adherence and cytotoxicity.** *Infect Immun* 2005, **73**:2602-2610.
37. Peterson KM, Alderete JF: **Host plasma proteins on the surface of pathogenic *Trichomonas vaginalis*.** *Infect Immun* 1982, **37**:755-762.
38. Arroyo R, Alderete JF: ***Trichomonas vaginalis* surface proteinase activity is necessary for parasite adherence to epithelial cells.** *Infect Immun* 1989, **57**:2991-2997.
39. Mendoza-Lopez MR, Becerril-Garcia C, Fattel-Facenda LV, Avila-Gonzales L, Ruiz-Tachiquin ME, Ortega-Lopez J, Arroyo R: **CP30, a cysteine proteinase involved in *Trichomonas vaginalis* cytoadherence.** *Infect Immun* 2000, **68**:4907-4912.
40. Diamond LS: **The establishment of various trichomonads of animals and man in axenic cultures.** *J Parasitol* 1957, **43**:488-490.
41. Land KM, Delgadillo MG, Johnson PJ: **In vivo expression of ferredoxin in a drug resistant trichomonad increases metronidazole susceptibility.** *Mol Biochem Parasitol* 2002, **121**:153-157.
42. Tsai C-D, Liu H-W, Tai J-H: **Characterization of an iron-responsive promoter in the protozoan pathogen *Trichomonas vaginalis*.** *J Biol Chem* 2002, **277**:5153-5162.
43. Alderete JF: **Antigen analysis of several pathogenic strains of *Trichomonas vaginalis*.** *Infect Immun* 1983, **39**:1041-1047.
44. Laemmli UK: **Cleavage of structural proteins during the assembly of the head of bacteriophage T4.** *Nature* 1970, **227**:680-685.
45. Klumpp DJ, Forrestal SG, Karr JE, Mudge CS, Anderson BE, Schaeffer AJ: **Epithelial differentiation promotes the adherence of type I-piliated *Escherichia coli* to human vaginal cells.** *J Infect Dis* 2002, **186**:1631-1638.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

