

## RESEARCH ARTICLE

# Arbovirus risk perception as a predictor of mosquito-bite preventive behaviors in Ponce, Puerto Rico

Josée M. Dussault<sup>1\*</sup>, Gabriela Paz-Bailey<sup>2</sup>, Liliana Sánchez-González<sup>2</sup>, Laura E. Adams<sup>2</sup>, Dania M. Rodríguez<sup>2</sup>, Kyle R. Ryff<sup>2</sup>, Chelsea G. Major<sup>2</sup>, Olga Lorenzi<sup>2</sup>, Vanessa Rivera-Amill<sup>3</sup>

**1** Department of Epidemiology, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, United States of America, **2** Division of Vector-borne Diseases, Centers for Disease Control and Prevention, San Juan, Puerto Rico, **3** Ponce Health Sciences University/Ponce Research Institute, Ponce, Puerto Rico

\* [josee.dussault@unc.edu](mailto:josee.dussault@unc.edu)



## OPEN ACCESS

**Citation:** Dussault JM, Paz-Bailey G, Sánchez-González L, Adams LE, Rodríguez DM, Ryff KR, et al. (2022) Arbovirus risk perception as a predictor of mosquito-bite preventive behaviors in Ponce, Puerto Rico. *PLoS Negl Trop Dis* 16(7): e0010653. <https://doi.org/10.1371/journal.pntd.0010653>

**Editor:** Arunasalam Pathmeswaran, Faculty of Medicine, University of Kelaniya, SRI LANKA

**Received:** August 20, 2021

**Accepted:** July 11, 2022

**Published:** July 26, 2022

**Copyright:** This is an open access article, free of all copyright, and may be freely reproduced, distributed, transmitted, modified, built upon, or otherwise used by anyone for any lawful purpose. The work is made available under the [Creative Commons CC0](https://creativecommons.org/licenses/by/4.0/) public domain dedication.

**Data Availability Statement:** Data cannot be shared publicly because of U.S. CDC data security and confidentiality guidelines. Data are available from CDC and PHSU for researchers that send in a formal request. Any external researchers can request access to a restricted use dataset by submitting a concept proposal. Data requests can be sent to [dengue@cdc.gov](mailto:dengue@cdc.gov).

**Funding:** All authors who are federal employees have nothing to disclose. VRA received funding from the National Center for Emerging and

## Abstract

Mosquito-borne arboviruses are an important cause of morbidity and mortality in the Caribbean. In Puerto Rico, chikungunya, dengue, and Zika viruses have each caused large outbreaks during 2010–2022. To date, the majority of control measures to prevent these diseases focus on mosquito control and many require community participation. In 2018, the U.S. Centers for Disease Control and Prevention launched the COPA project, a community-based cohort study in Ponce, Puerto Rico, to measure the impact of novel vector control interventions in reducing arboviral infections. Randomly selected households from 38 designated cluster areas were offered participation, and baseline data were collected from 2,353 households between May 2018 and May 2019. Household-level responses were provided by one representative per home. Cross-sectional analyses of baseline data were conducted to estimate 1) the association between arboviral risk perception and annual household expenditure on mosquito control, and 2) the association between arboviral risk perception and engagement in  $\geq 3$  household-level risk reduction behaviors. In this study, 27% of household representatives believed their household was at high risk of arboviruses and 36% of households engaged in at least three of the six household-level preventive behaviors. Households where the representative perceived their household at high risk spent an average of \$35.9 (95% confidence interval: \$23.7, \$48.1) more annually on mosquito bite prevention compared to households where the representative perceived no risk. The probability of engaging in  $\geq 3$  household-level mosquito-preventive behaviors was 10.2 percentage points greater (7.2, 13.0) in households where the representatives perceived high risk compared to those in which the representatives perceived no risk. Paired with other research, these results support investment in community-based participatory approaches to mosquito control and providing accessible information for communities to accurately interpret their risk.

Zoonotic Infectious Diseases (U01CK000437 & U01CK000580; <https://www.cdc.gov/ncezid/index.html>). JMD received funding from the National Institute of Mental Health (3U19MH113202-03S1; <https://www.nimh.nih.gov/>) and the National Institute of Allergy and Infectious Diseases (T32 AI 070114; <https://www.niaid.nih.gov/>). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing interests:** The authors have declared that no competing interests exist.

## Author summary

Mosquito-borne disease is an important cause of illness and death in the Caribbean, including Puerto Rico. Most tactics to prevent these diseases rely on stopping mosquito bites, either by reducing the mosquito population or creating barriers between mosquitoes and humans. These methods vary in the degree of community involvement required. This study used data collected from 2,353 households in Ponce, Puerto Rico from May 2018 to May 2019 to understand how household perception of risk of contracting these diseases related to 1) the amount of money households spent annually to prevent mosquito bites, and 2) the number of activities their household engaged in to prevent mosquito bites. We found that 27% of households perceived themselves at high risk of contracting these diseases, and 36% of households engaged in at least three activities to reduce their risk. On average, households that perceived themselves at high risk spent more money on mosquito bite prevention and engaged in more activities to prevent mosquito bites, compared to households that perceived no risk. Paired with other research in this area, these results support investment in community-based approaches to mosquito control and ensuring that communities have accessible information to understand their risk of mosquito-borne disease.

## Introduction

Mosquito-borne illnesses are an important cause of morbidity and mortality in the Caribbean [1]. In Puerto Rico, chikungunya, dengue, and Zika viruses have each caused large outbreaks during 2010–2022. Although chikungunya and Zika viruses were recently introduced to Puerto Rico, dengue is endemic and seasonal, with outbreaks occurring every 3–5 years [1,2]. All three viruses are transmitted via *Aedes* mosquito vectors and can cause acute febrile illness, although each of these diseases has the potential for severe or long-term outcomes [2–4]. Patients with chikungunya can have persistent joint pain for months or years. Perinatal Zika infections are known to cause birth defects [5,6]. Evidence suggests that individuals infected with chikungunya or Zika virus are conferred immunity against the respective virus [5,7]. However, dengue virus has four serotypes, and individuals who have been infected with one serotype have a higher risk of severe dengue if later infected with a different serotype. Severe dengue can be lethal and is characterized by severe plasma leakage, severe hemorrhage, and/or organ impairment [8]. There are no specific treatments for any of these diseases and management is primarily supportive.

The lack of treatment options for arboviruses is compounded by the scarcity of pharmaceutical prevention methods. There is currently one vaccine for dengue, CYD-TDV, approved by the U.S. Food and Drug Administration (FDA) and recommended by the Advisory Committee on Immunization Practices (ACIP) for use in children 9–16 years old who live in an endemic area and who have a lab-confirmed history of a previous dengue infection [9,10]. However, CYD-TDV is only available to a small proportion of the population at risk for dengue [10], and there are no other FDA-approved vaccines to protect people from these arboviruses. Thus, the public health community relies on vector control tools to prevent or reduce arboviral infections in endemic populations. Some of these vector control tools focus on killing adult mosquitoes or reducing population growth through methods like disposal of stagnant water to decrease *Aedes* mosquito breeding sites, fumigation, and mosquito traps. Other methods focus on creating barriers between humans and mosquitoes through the use of screens in

the doors and windows of buildings, centralized air conditioning, topical mosquito repellents applied to skin or clothing, and bed nets.

There are several barriers to mosquito-borne disease prevention, both at the individual and household level. In Puerto Rico, where the median household income was \$20,296 in 2018 [11], the cost of supplies and services to reduce mosquito populations and mosquito bites can represent a considerable financial burden. Moreover, individuals may perceive themselves at low risk of contracting arboviral disease, either because they do not see illnesses in their communities or do not feel that mosquitos are a problem [12]. People who perceive themselves at low risk may be less motivated to protect themselves and other members of their household from contracting an arboviral infection. Whether due to financial capacity or motivation, the underutilization of mosquito prevention techniques and services puts populations at higher risk of arboviral infection.

Most models of health behavior generally emphasize the role of risk perception in decision making [13,14], but empirical studies on arboviral risk perception have produced conflicting results [12,15], and there have been few studies investigating arboviral risk perception and health behaviors in Puerto Rico [16]. Understanding the association between risk perception and risk reduction behaviors among people living in Puerto Rico can help guide selection of appropriate public health messaging and interventions to reduce the incidence of arboviral diseases. If risk perception correlates to increased protective behaviors, increasing campaigns to raise community awareness about arboviral disease risks could be beneficial. Alternatively, in the presence of a null relationship between risk perception and engagement in protective behaviors, then public health agencies may find it more effective to focus efforts on disease prevention methods that do not require active community engagement.

Accordingly, the objective of this study is to better understand the relationship between risk perception of arboviral diseases and adoption of behaviors to prevent mosquito bites among people living in southern Puerto Rico communities with high historical arboviral disease incidence rates. The central hypothesis is that individuals who perceive themselves at high risk for arboviral infection will be more likely to engage in mosquito-preventive behaviors than those who perceive no risk. Specifically, this study aims to (1) estimate the association between perceived household risk of arboviral infection and annual household expenditure on mosquito control, and (2) estimate the association between perceived household risk of arboviral infection and reported engagement in mosquito-controlling behavior (i.e., spraying insecticide, burning citronella candles/coils, etc.) among households in Ponce, Puerto Rico.

## Methods

### Ethics statement

All participants provided written, informed consent. The COPA study was approved by the Institutional Review Boards (IRBs) of Ponce Medical School Foundation, Inc., and the U.S. Centers for Disease Control and Prevention (CDC). An exemption for this secondary statistical analysis was approved by the IRB at the University of North Carolina at Chapel Hill.

### Parent study

COPA (Communities Organized to Prevent Arboviruses) is a prospective community-based cohort study in Ponce, Puerto Rico initiated in 2018 to evaluate the impact of novel vector control interventions in reducing arboviral infections. Households were selected randomly from 38 study-defined cluster areas and visited up to three times to invite them to participate in COPA; field-level strategies are detailed elsewhere [17]. From May 2018 to May 2019, 2,353 households were enrolled in COPA. Household residents were eligible if they slept an average

of four or more nights per week in the home, were 1–50 years old, and did not intend to move in the next six months. Study staff administered a standardized questionnaire to participants in Spanish regarding their knowledge, attitudes, and practices relating to arboviral infections and prevention methods. For each household, we asked one resident (an adult or emancipated minor) present during data collection to act as the household representative and answer an additional series of questions specific to their household. The following is an analysis of cross-sectional interview data from the COPA household representatives.

### Exposure

The exposure of interest was the household representative's perception of their household's risk of contracting arboviral illnesses in the next 12 months, which was assessed separately for Zika, dengue, and chikungunya. However, as the responses to these questions were highly correlated, dengue was used as a surrogate for all three arboviruses to measure the household representatives' perceived risk of arboviral infection with three levels: high risk, low risk, and no risk. Models were tested in which the exposure was coded as a single 3-level ordinal variable versus as a set of two disjoint indicator (dummy) variables. Disjoint indicator variables allowed the model to relax the assumption of linearity between the three levels of the exposure variable by creating two binary variables that represented the low risk and high risk groups, with no risk as the referent level. The Bayesian information criterion (BIC) and Akaike information criterion (AIC) were used to compare the fit of the two models. Generally, models with lower AIC and BIC values are preferred. Where the values produced different conclusions, the BIC was preferred rather than the AIC because it usually selects the correct model more frequently than the AIC [18].

### Outcomes

There were two outcomes of interest. The first outcome of interest was annual household expenditure on mosquito prevention products and services, reported in United States dollars (USD) by the household representative. The second outcome was regular engagement in three or more mosquito preventive behaviors. In total, there were six possible household-level behaviors that the household representative could report during the baseline interview: (1) Eliminating stagnant water around the house (ex: cleaning containers that collect water like flowerpots and/or gutters; (2) Covering containers that collect water; (3) Emptying trash cans, drums, or in-ground trash cans if they had standing water; (4) Cleaning or removing tires and/or debris from the yard; (5) Spraying insecticide or fumigating indoors or outdoors; and (6) Burning citronella candles and/or mosquito coils. These behaviors were reported as the frequency with which any members of the household engaged in these activities in the previous 12 months, with responses including "daily", "weekly", "monthly", "rarely", or "never". To dichotomize each of the six activities, households that reported daily or weekly engagement in these behaviors were counted as engaging in these behaviors regularly, and those who reported performing the activities monthly, rarely, or never were categorized as not conducting these activities regularly. This outcome was then dichotomized as regular engagement in three or more of these behaviors. The cutoff of three behaviors was selected because some activities (e.g., behaviors 1 and 3) appeared to be correlated, likely because it would be easy to accomplish one task while completing the other.

### Covariates

Covariates included in this study were from the household representatives' responses to questionnaires from the same study visit. These variables included: mosquito bite frequency,

household income, exposure to community educational campaigns, education level, sex, and the community cluster to which the household belonged. Reported mosquito bites was a binary variable produced from a COPA question asking about where the participants most often were bitten. If the participant responded saying that they rarely/never were bitten, this was encoded into the bite frequency variable as “never bitten”. Otherwise, participants were categorized as “bitten”. Household income was reported as one of eight categories and remained an ordinal categorical variable in statistical analyses. Education level was collapsed into five categories due to small counts in some of the original categories. Sex and educational campaign exposure were binary variables, and age was maintained as a continuous variable. Education level and community cluster were modeled as disjoint indicator variables. Rooted in Health Belief Model theory and additional readings [12–16,18–28], we developed directed acyclic graphs (DAGs) that assisted in the identification of mediating paths and confounders [29–32]. Based on these DAGs, the aforementioned covariates included in this study represented a sufficient set of confounders. Because participating households were selected from community clusters, the intra-cluster correlation (ICC) coefficient was calculated to assess the need to use multilevel modeling techniques; the ICC was determined to be small enough (0.02) to include community clusters as disjoint indicator variables rather than using a 2-level model [33].

### Missing data

Approximately 11% of participants were missing data on at least one of the covariates, and most missing data came from the household income variable. A complete case analysis of these data could result in loss of precision and validity by not including the full possible sample of households [34–39]. Thus, this analysis used multiple stochastic conditional-mean imputation with coefficient resampling to assign values onto all missing covariates based on each participant’s observed characteristics, assuming that data were missing at random, conditional on the additional covariates in the imputation model [37,38,40]. The covariates included as predictive variables in the imputation models were employment status, whether the house had screens on the windows, and the six risk reduction behaviors described in the outcomes section. All exposure, outcome, and confounding variables that would be included in the analysis model were also included in the multiple imputation model. Multiple imputation methods were applied using Markov chain Monte Carlo (MCMC) method in SAS software, version 9.4.9 [41], with 50 imputed datasets, 500 burn-in iterations, and 500 additional iterations between imputations; the method used a single chain. The MCMC method was necessary because the missing data structure was arbitrary, and the number of burn-in iterations, additional iterations, and imputed datasets were selected based on recommendations to reduce Monte Carlo error [42–44]. Because two functional forms for the exposure variable were compared using model fit statistics that are not available in multiple imputation models, complete case models were used to assess the proper functional form of the exposure variable, and the multiple imputation model with that exposure coding was preferred over the complete case model.

### Analytical approach

**Household expenditure.** In this analysis, we were interested in understanding the association between arboviral disease risk perception and annual household expenditure on risk reduction products and services as the primary outcome. To estimate the association, we fit a normal linear regression model. To meet the assumption of homoscedasticity in the model residuals [45], five outlier households whose expenditures ranged from \$1800 to \$3600 were

set to missing values. For comparison, the sixth-largest expenditure value in the data was \$1440, and the mean expenditure amount was \$185. In the regression model, we included the following potential confounders: reported mosquito bites, household income, exposure to community educational campaigns, sex and education level of the household representative, and the community cluster to which the household belonged.

**Household-level risk reduction.** The objective of this analysis was to understand the association between household perception of arboviral disease risk and the prevalence of engaging in household-level risk reduction behaviors. To estimate the prevalence difference, we fit a linear binomial model including the same set of potential confounders as described previously. Missing data on all variables were addressed using multiple imputation as described above. Data were analyzed using SAS software, version 9.4.9 [41].

## Results

Among 2,353 household representatives who provided responses, 635 (27%) perceived their household at high risk of contracting dengue in the next 12 months. Comparatively, 1,249 (53%) believed their households were at low risk, and 425 (18%) perceived no risk (Table 1). The mean age of household representatives was 37 years (standard deviation [SD]: 8.8 years), 1,596 (68%) were female, and 1,429 (61%) reported having completed some postsecondary schooling. Most household representatives reported that they were bitten by mosquitoes ( $n = 2,304$ ; 98%); few reported that they were rarely or not bitten ( $n = 45$ ; 2%). The most commonly reported household annual income category was  $< \$10,000$  ( $n = 915$ ; 39%). Mean annual expenditure on mosquito products was \$185 (SD: \$225). A majority ( $n = 1,884$ ; 80%) of household representatives perceived at least some risk of contracting dengue in the next year (Tables 1 & 2).

Household representatives who perceived themselves at no risk of arboviral diseases generally had lower levels of education compared to the low-risk and high-risk perception groups. Contrary to expectation, the no-risk perception group was also more likely to report having seen an educational campaign on mosquito prevention and control in the last 12 months compared to the other two risk perception groups. In total, 1,318 (56%) household representatives reported using mosquito repellent in the last 30 days (Table 2).

There was generally low uptake of the six household-level preventive behaviors assessed. About half of household representatives (51%) reported emptying trash cans with standing water, 44% reported eliminating other sources of stagnant water, and 44% reported spraying insecticide around the house or fumigating indoors or outdoors. A small number of households reported covering containers that could collect water (25%), cleaning debris from around the outside of the house (25%) or burning citronella candles or coils (17%) (Table 2). In total, 840 (36%) households engaged in at least three of the six household-level preventive behaviors. Households that were engaged in three or more preventive behaviors generally had household representatives with higher levels of education, a larger household income, and

**Table 1. Distribution of COPA Household Representatives' Perceived Risk by Arbovirus (N = 2,353), 2018–2019.**

	Dengue, N (%)	Zika, N (%)	Chikungunya, N (%)
Perceived high risk	635 (28)	606 (26)	613 (27)
Perceived low risk	1249 (54)	1264 (55)	1221 (53)
Perceived no risk	425 (18)	435 (19)	473 (21)
Missing	44	48	46
Total	2309	2305	2307

<https://doi.org/10.1371/journal.pntd.0010653.t001>



Table 2. Characteristics of COPA household representatives by perceived risk of dengue (N = 2,353), 2018–2019.

	Perceived high risk (n = 635) <sup>a</sup>		Perceived low risk (n = 1,249) <sup>a</sup>		Perceived no risk (n = 425) <sup>a</sup>		Total (N = 2,353) <sup>a</sup>
	N	%	N	%	N	%	N (%)
<b>Education Level</b>							
< High school GED	41	6	76	6	67	16	184 (8)
Finished high school/GED	171	27	359	29	166	39	704 (30)
Technical/Associate degree	188	30	334	27	90	21	620 (27)
Bachelor's degree	163	26	344	28	77	18	592 (26)
Professional/Post-graduate	69	11	123	10	23	5	217 (9)
<b>Income level</b>							
< \$10,000	253	42	412	36	239	61	915 (39)
\$10,000 - \$19,999	123	21	248	21	58	15	435 (18)
\$20,000 - \$29,999	69	12	202	17	41	11	319 (14)
\$30,000 - \$39,999	60	10	111	10	23	6	199 (8)
\$40,000 - \$49,999	34	6	71	6	8	2	113 (5)
\$50,000 - \$59,999	18	3	41	4	7	2	68 (3)
\$60,000 - \$69,999	15	3	27	2	7	2	50 (2)
≥ \$70,000	25	4	45	4	7	2	78 (3)
<b>Bitten by mosquitos</b>	627	99	1224	98	410	96	2304 (98)
<b>Female</b>	439	69	819	66	307	72	1596 (68)
<b>Exposed to educational campaigns, last 12 months</b>	144	23	300	24	131	31	587 (25)
<b>Engaged in protective behavior:</b>	302	48	547	44	153	36	1018 (44)
<b>#1: Eliminating stagnant water</b>							
<b>#2: Covering containers that collect water</b>	158	25	320	26	95	23	584 (25)
<b>#3: Emptying trash cans if they have standing water</b>	328	52	654	53	175	41	1181 (51)
<b>#4: Cleaning or removing tires and/or debris from yard</b>	151	24	305	25	98	24	565 (25)
<b>#5: Spraying insecticide or fumigate indoors or outdoors</b>	336	54	500	41	155	37	1006 (44)
<b>#6: Burning citronella candles and/or mosquito coils</b>	160	25	182	15	51	12	399 (17)
<b>Household representative wore repellent in the last 30 days</b>	443	70	661	53	206	48	1318 (56)
<b>Household engaged in &gt; = 3 protective behaviors<sup>b</sup></b>	265	42	445	36	114	27	840 (36)
	<b>Mean</b>	<b>SD</b>	<b>Mean</b>	<b>SD</b>	<b>Mean</b>	<b>SD</b>	<b>Mean (SD)</b>
<b>Age</b>	38	8.4	36	8.9	37	8.8	37 (8.8)
<b>Household Expenditure on Mosquito Products</b>	\$211	\$242	\$176	\$206	\$173	\$253	\$185 (\$225)
<b>Number of household mosquito-preventive behaviors reported<sup>b</sup></b>	2.3	1.5	2.0	1.5	1.7	1.5	2.0 (1.5)

<sup>a</sup> Percentages presented as proportion of non-missing observations. Total number of missing observations on each variable are as follows: a) risk perception (exposure variable), 44; b) education level, 36; c) mosquito bite frequency, 4; d) sex, 0; e) income level, 176; f) exposure to educational campaigns, 51; g) protective behavior #1, 26; h) protective behavior #2, 46; i) protective behavior #3, 26; j) protective behavior #4, 58; k) protective behavior #5, 44; l) protective behavior #6, 29; m) repellent use in the last 30 days, 0; n) age, 0; o) household expenditure on products, 20; p) number of household mosquito-preventive behaviors, 17.

<sup>b</sup> Household-level mosquito-preventive behaviors: (1) Eliminating stagnant water around the house (ex: cleaning containers that collect water like flowerpots and/or gutters); (2) Covering containers that collect water; (3) Emptying trash cans, drums, or in-ground trash cans if they had standing water; (4) Cleaning or removing tires and/or debris from yard; (5) Spraying insecticide or fumigate indoors or outdoors (ex: Blackjack, Real Kill, Raid); (6) Burning citronella candles and/or mosquito coils.

<https://doi.org/10.1371/journal.pntd.0010653.t002>

spent more money on mosquito products compared to households that engaged in fewer than three preventive behaviors (Table 3).

The regression models that encoded risk perception as an ordinal exposure variable (Models 1, 3, 5, and 7) produced similar outputs to those in which the exposure was coded as disjoint indicator variables (Models 2, 4, 6, and 8). Nonetheless, the Bayesian information criterion (BIC) and Akaike information criterion (AIC) values (see S1 Table) preferred the

Table 3. Characteristics of COPA household representatives by protective behavior level (N = 2,353), 2018-2019<sup>a</sup>.

	Engaged in < 3 protective behaviors (n = 1,496) <sup>b</sup>		Engaged in ≥ 3 protective behaviors (n = 840) <sup>b</sup>		Total (N = 2,353) <sup>b</sup>
	N	%	N	%	N (%)
<b>Education Level</b>					
< High school GED	134	9	49	6	184 (8)
Finished high school/GED	485	33	215	26	704 (30)
Technical/Associate degree	370	25	245	30	620 (27)
Bachelor's degree	353	24	235	28	592 (26)
Professional/Post-graduate	134	9	83	10	217 (9)
<b>Income level</b>					
< \$10,000	634	42	281	33	915 (39)
\$10,000 - \$19,999	251	17	184	22	435 (18)
\$20,000 - \$29,999	198	13	121	14	319 (14)
\$30,000 - \$39,999	112	7	87	10	199 (8)
\$40,000 - \$49,999	71	5	42	5	113 (5)
\$50,000 - \$59,999	44	3	24	3	68 (3)
\$60,000 - \$69,999	30	2	20	2	50 (2)
≥ \$70,000	53	4	25	3	78 (3)
<b>Regularly bitten by mosquitos</b>	1463	98	824	98	2304 (98)
<b>Female</b>	1039	69	547	65	1596 (68)
<b>Exposure to educational campaigns, last 12 months</b>	368	25	219	26	587 (25)
	<b>Mean</b>	<b>SD</b>	<b>Mean</b>	<b>SD</b>	<b>Mean (SD)</b>
<b>Age</b>	37	8.8	37	8.8	37 (8.8)
<b>Household Expenditure on Mosquito Products</b>	\$151	\$195	\$244	\$259	\$185 (\$225)

<sup>a</sup> There were 6 household-level mosquito-preventive behaviors total: (1) Eliminating stagnant water around the house (ex: cleaning containers that collect water like flowerpots and/or gutters); (2) Covering containers that collect water; (3) Emptying trash cans, drums, or in-ground trash cans if they had standing water; (4) Cleaning or removing tires and/or debris from yard; (5) Spraying insecticide or fumigate indoors or outdoors (ex: Blackjack, Real Kill, Raid); (6) Burning citronella candles and/or mosquito coils.


<sup>b</sup> Percentages presented as proportion of non-missing observations. Total number of missing observations on each variable are as follows: a) Number of preventive behaviors engaged in (outcome variable), 17; b) education level, 33; c) mosquito bite frequency, 4; d) sex, 0; e) income level, 159; f) exposure to educational campaigns, 35; g) age, 0; h) household expenditure on products, 3.

<https://doi.org/10.1371/journal.pntd.0010653.t003>

models in which the exposure is coded as an ordinal variable (Models 1, 3, 5, and 7). These models also produced estimates with greater precision (Figs 1 & 2, S1 Table). Therefore, the subsequent description of results will focus on reporting model outputs from models that used ordinal coding of the exposure variable and multiple imputation (models 3 and 7).

Fig 1 shows the output from the regression analysis for the first outcome of interest, household expenditure on mosquito-preventive products and services. The adjusted linear regression model with ordinal exposure coding (Model 3) demonstrated that a 1-unit increase in dengue risk perception (from no risk perception to low, or from low to high risk perception) was associated with a \$17.9 (95% CI: \$5.8, \$30.1) increase in annual household expenditure to prevent mosquito bites. On average, households where the household representative perceived their household at high risk of contracting dengue in the next year spent \$35.9 (\$23.7, \$48.1) more annually on mosquito bite prevention compared to a household where the household representative considered their household at no risk. Fig 1 provides a graphical representation of the model estimates and 95% confidence intervals. For a detailed comparison of Model 1 – Model 4 outputs, see S1 Table.






**Fig 1. Estimates and 95% confidence intervals from four models of the difference in household expenditure (USD) on mosquito-preventive products per risk perception group, COPA, 2018-2019<sup>a</sup>.** <sup>a</sup> All models included the full set of covariates as described in the methods section. Point estimates (dots) represent the average difference in household expenditure (USD) on mosquito preventive products when the household representative perceived their household at either low risk or high risk (based on the x-axis) compared to when the representative perceived their household at no risk. The bands around each dot represents the 95% confidence intervals for each point estimate. Positive values indicate increased spending; negative values indicate decreased spending. Model 1 used an ordinal exposure variable and complete case analysis; Model 2 used disjoint indicator coding for the exposure variable and complete case analysis; Model 3 used ordinal exposure variable and multiple imputation to address missing observations; Model 4 used disjoint indicator coding for the exposure variable and multiple imputation. Model fit statistics can be found in Table A in [S1 Table](#). Based on superior model fit statistics, Model 3 results are reported in the text.

<https://doi.org/10.1371/journal.pntd.0010653.g001>

The regression model results for the second outcome of interest—the probability of engaging in at least three household-level protective behaviors—are shown in [Fig 2](#). This second set of regression models demonstrated that the probability of engaging in at least three household-level mosquito-preventive behaviors also increased with increasing arboviral disease risk perception. Model 7, which used ordinal exposure coding and multiple imputation, estimated that each 1-unit increase of risk perception was associated with an absolute increase of 5.1 percentage points (95% CI: 0.02, 0.08) in the probability that the household would report engagement in three or more household-level mosquito-preventive behaviors. Thus, households with a household representative who reported a high risk perception had a probability of engaging in three or more household-level mosquito-preventive behaviors that was 10.2 percentage points greater (95% CI: 7.2, 13.0), compared to households in which the representatives



**Fig 2. Prevalence difference estimates and 95% confidence intervals from four models representing the probability of household engagement in at least three protective behaviors per risk perception group, COPA, 2018-2019<sup>a</sup>.** <sup>a</sup> All models included the full set of covariates as described in the methods section. Point estimates (dots) represent the average difference in probability that the household engaged in 3 or more protective behaviors when the household representative perceived their household at either low risk or high risk (based on the x-axis) compared to when the representative perceived their household at no risk. The bands around each dot represents the 95% confidence intervals for each point estimate. Positive values indicate increased spending; negative values indicate decreased spending. Model 5 used an ordinal exposure variable and complete case analysis; Model 6 used disjoint indicator coding for the exposure variable and complete case analysis; Model 7 used ordinal exposure variable and multiple imputation to address missing observations; Model 8 used disjoint indicator coding for the exposure variable and multiple imputation. Model fit statistics can be found in Table B in S1 Table. Based on superior model fit statistics, Model 7 results are reported in the text.

<https://doi.org/10.1371/journal.pntd.0010653.g002>

perceived their household at no risk of arboviral disease. Fig 2 provides a graphical representation of the model estimates and 95% confidence intervals. For a detailed comparison of Model 5 –Model 8 outputs, see S1 Table.

## Discussion

This study analyzed a large (N = 2,353) sample of households in Ponce, Puerto Rico, with a median annual household income of \$10,000-\$19,999. This analysis found that after adjusting for income and other confounders, households with high perceived risk of contracting arboviruses reported spending \$35.9 more per year (95% CI: \$23.7, \$48.1) on preventive products and services compared to households with no perceived risk. In a population where the median annual household income was less than \$20,000, these results demonstrate that households are willing to invest in mosquito-bite prevention despite potentially limited resources if there is a high perceived risk of arboviral disease. Study data also showed a positive association between the household representative's perception of risk and the household's probability of engaging in at least three risk-reducing behaviors. These findings highlight the role of arboviral risk perception in influencing risk-reducing behaviors and support the continued

development of community-based programs to increase awareness of arboviral disease risk across Puerto Rico.

The association between heightened arbovirus risk perception and increased prevalence of behaviors to prevent arboviral infections in this study fits well with previous research related to community engagement in mosquito vector control in Puerto Rico. For example, in a CDC case study of dengue control measures during an epidemic in the 1990s, visible public spraying efforts, while largely ineffective, reduced residents' engagement in mosquito bite preventive activities [21]. As residents shifted the responsibility of mosquito control to the government, they perceived themselves at low risk of arboviral infection and were less likely to engage in further protective behaviors. Moreover, a pilot study conducted in Manuel A. Pérez Public Housing in San Juan, Puerto Rico in 2018 demonstrated the potency of community-based participatory campaigns to increase employment of mosquito-control tactics [22]. In that study, community members were directly engaged in the process of planning and implementing a risk communication initiative for their community. After the initiative, there was a significant increase in community members' recognition of personal and community responsibility for the prevention of mosquito-borne disease, and their engagement in preventive behaviors for mosquito control increased [22].

Key strengths of this study include its sample size and the detailed information captured for each household, which allowed for comprehensive analyses. Another strength of this analysis is the use of multiple imputation, which allowed us to use all 2,353 households in our analysis and mitigate the risk of selection bias introduced by complete case analyses [46,47].

As in most studies on attitudes and practices that utilize in-person interviews, there may be some desirability bias in exposure and outcome measurement. Specifically, because the data collection teams visited participants at home after enrolling in a study focused on arboviruses, participants may have overreported protective behaviors, particularly if they felt that these were actions expected or supported by study staff. Likewise, participants who truly perceived themselves at no risk may have reported a low perceived risk of dengue to satisfy perceived interviewer expectations.

Another limitation of our study is the cross-sectional nature of the data, which does not allow us to assess the temporal sequence between the exposure and outcomes. While this study assessed whether households perceiving themselves at high risk would participate in preventive behaviors, it is possible that some participants perceived themselves at low or no risk of arbovirus in the upcoming 12 months because of their high engagement in prevention in the previous 12 months. Nevertheless, if this were the overarching trend, we would have expected to see a negative correlation between the exposure and outcome variables named in this analysis [27]. One final limitation in this study is residual confounding bias in the mosquito bite variable. Due to data unavailability, we dichotomized this variable, but there is plausible heterogeneity within the "bitten" category, particularly when compared to the "never bitten" level.

Our findings add to the growing body of evidence in support of community education interventions for populations at risk of contracting arboviral diseases. Paired with other findings [19,21,22], our results suggest that key stakeholders should invest in community-based participatory approaches to mosquito control, with particular focus on providing information in an accessible manner for community members to accurately interpret their risk and make more informed choices to reduce their risk of arboviral disease.

## Supporting information

**S1 Table. Model 1–8 Outputs.** Model outputs presented in this supporting document provide the statistics that support Figs 1 and 2 in this manuscript. Table A in S1 Table. Model outputs

describing the relationship between perceived risk and household expenditure (USD) on mosquito-preventive products, COPA, 2018–2019a. Table B in S1 Table. Model outputs describing the relationship between perceived arboviral risk and the probability of engaging in at least three household-level protective behaviors, COPA, 2018–2019a. (DOCX)

## Author Contributions

**Conceptualization:** Gabriela Paz-Bailey, Laura E. Adams, Vanessa Rivera-Amill.

**Data curation:** Gabriela Paz-Bailey, Liliana Sánchez-González, Dania M. Rodríguez, Kyle R. Ryff.

**Formal analysis:** Josée M. Dussault.

**Funding acquisition:** Gabriela Paz-Bailey.

**Project administration:** Gabriela Paz-Bailey, Liliana Sánchez-González, Laura E. Adams.

**Supervision:** Gabriela Paz-Bailey.

**Writing – original draft:** Josée M. Dussault.

**Writing – review & editing:** Josée M. Dussault, Gabriela Paz-Bailey, Liliana Sánchez-González, Laura E. Adams, Dania M. Rodríguez, Kyle R. Ryff, Chelsea G. Major, Olga Lorenzi, Vanessa Rivera-Amill.

## References

1. Lowe R, Gasparrini A, Meerbeeck CJV, Lippi CA, Mahon R, Trotman AR, et al. Nonlinear and delayed impacts of climate on dengue risk in Barbados: A modelling study. *PLOS Medicine*. 2018; 15: e1002613. <https://doi.org/10.1371/journal.pmed.1002613> PMID: 30016319
2. Adams LE, Martin SW, Lindsey NP, Lehman JA, Rivera A, Kolsin J, et al. Epidemiology of Dengue, Chikungunya, and Zika Virus Disease in the U.S. States and Territories, 2017. *Am J Trop Med Hyg*. 2019. <https://doi.org/10.4269/ajtmh.19-0309> PMID: 31436154
3. Dengue Virus Infections | 2015 Case Definition. [cited 28 Aug 2019]. Available from: /nndss/conditions/dengue-virus-infections/case-definition/2015/.
4. Petersen LR, Jamieson DJ, Powers AM, Honein MA. Zika Virus. *N Engl J Med*. 2016; 374: 1552–1563. <https://doi.org/10.1056/NEJMr1602113> PMID: 27028561
5. Zika Virus. In: CDC [Internet]. 5 Nov 2014 [cited 29 Apr 2019]. Available from: <https://www.cdc.gov/zika/vector/range.html>.
6. Marinho PS, Cunha AJ, Amim Junior J, Prata-Barbosa A. A review of selected Arboviruses during pregnancy. *Maternal Health, Neonatology and Perinatology*. 2017; 3: 17. <https://doi.org/10.1186/s40748-017-0054-0> PMID: 29026637
7. Chikungunya. [cited 28 Aug 2019]. Available from: <https://www.who.int/news-room/fact-sheets/detail/chikungunya>.
8. Soo K-M, Khalid B, Ching S-M, Chee H-Y. Meta-Analysis of Dengue Severity during Infection by Different Dengue Virus Serotypes in Primary and Secondary Infections. *PLoS One*. 2016;11. <https://doi.org/10.1371/journal.pone.0154760> PMID: 27213782
9. First FDA-approved vaccine for the prevention of dengue disease in endemic regions. In: FDA [Internet]. 11 Sep 2019 [cited 14 Feb 2020]. Available from: <http://www.fda.gov/news-events/press-announcements/first-fda-approved-vaccine-prevention-dengue-disease-endemic-regions>.
10. Dengue Vaccine. In: CDC [Internet]. 25 Jun 2021 [cited 9 Jul 2021]. Available from: <https://www.cdc.gov/dengue/prevention/dengue-vaccine.html>.
11. Puerto Rico | Data USA. [cited 13 Feb 2020]. Available from: <https://datausa.io/profile/geo/puerto-rico/>.
12. Raude J, Chinfatt K, Huang P, Betansedi CO, Katumba K, Vernazza N, et al. Public perceptions and behaviours related to the risk of infection with Aedes mosquito-borne diseases: a cross-sectional study

- in Southeastern France. *BMJ Open*. 2012; 2: e002094. <https://doi.org/10.1136/bmjopen-2012-002094> PMID: 23194955
13. Ferrer R, Klein WM. Risk perceptions and health behavior. *Curr Opin Psychol*. 2015; 5: 85–89. <https://doi.org/10.1016/j.copsyc.2015.03.012> PMID: 26258160
  14. Brewer NT, Chapman GB, Gibbons FX, Gerrard M, McCaul KD, Weinstein ND. Meta-analysis of the relationship between risk perception and health behavior: The example of vaccination. *Health Psychology*. <https://doi.org/10.1037/0278-6133.26.2.136> PMID: 17385964; 26: 136.
  15. Banneheke H, Paronavitane S, Jayasuriya V, Banneheka S. Perceived Risk of Dengue in Ones' Living Environment as a Determinant of Behavior Change through Social Mobilization and Communication: Evidence from a High Risk Area in Sri Lanka. *J Arthropod Borne Dis*. 2016; 10: 392–402. PMID: 27308297
  16. Pérez-Guerra CL, Zielinski-Gutierrez E, Vargas-Torres D, Clark GG. Community beliefs and practices about dengue in Puerto Rico. *Rev Panam Salud Publica*. 2009; 25: 218–226. <https://doi.org/10.1590/s1020-49892009000300005> PMID: 19454149
  17. Rodríguez DM, Ryff K, Sánchez-Gonzalez L, Rivera-Amill V, Paz-Bailey G, Adams L. HTrack: A new tool to facilitate public health field visits and electronic data capture. *Angkurawaranon C, editor. PLoS ONE*. 2020; 15: e0244028. <https://doi.org/10.1371/journal.pone.0244028> PMID: 33320896
  18. Neath AA, Cavanaugh JE. The Bayesian information criterion: background, derivation, and applications: The Bayesian information criterion. *WIREs Comp Stat*. 2012; 4: 199–203. <https://doi.org/10.1002/wics.199>
  19. Winch PJ, Leontsini E, Rigau-Pérez JG, Ruiz-Pérez M, Clark GG, Gubler DJ. Community-based dengue prevention programs in Puerto Rico: impact on knowledge, behavior, and residential mosquito infestation. *Am J Trop Med Hyg*. 2002; 67: 363–370. <https://doi.org/10.4269/ajtmh.2002.67.363> PMID: 12452490
  20. Fritzell C, Raude J, Adde A, Dusfour I, Quenel P, Flamand C. Knowledge, Attitude and Practices of Vector-Borne Disease Prevention during the Emergence of a New Arbovirus: Implications for the Control of Chikungunya Virus in French Guiana. *PLOS Neglected Tropical Diseases*. 2016; 10: e0005081. <https://doi.org/10.1371/journal.pntd.0005081> PMID: 27802275
  21. Gubler DJ, Clark GG. Community involvement in the control of *Aedes aegypti*. *Acta Tropica*. 1996; 61: 169–179. [https://doi.org/10.1016/0001-706x\(95\)00103-1](https://doi.org/10.1016/0001-706x(95)00103-1) PMID: 8740894
  22. Juarbe-Rey D, Pérez AO, Santoni RPCP, Ramírez MR, Vera M. Using Risk Communication Strategies for Zika Virus Prevention and Control Driven by Community-Based Participatory Research. *Int J Environ Res Public Health*. 2018;15. <https://doi.org/10.3390/ijerph15112505> PMID: 30423904
  23. Piltch-Loeb R, Merdjanoff AA, Bhanja A, Abramson DM. Support for vector control strategies in the United States during the Zika outbreak in 2016: The role of risk perception, knowledge, and confidence in government. *Preventive Medicine*. 2019; 119: 52–57. <https://doi.org/10.1016/j.ypmed.2018.12.019> PMID: 30594531
  24. Miranda E, Santos-Pinto C, Antunes-de-Lima C, Osorio-de-Castro C. Risk Perception of Zika Virus Infection Among Vulnerable Women in Rio de Janeiro. *Prehospital and Disaster Medicine*. 2019; 34: s36–s37. <https://doi.org/10.1017/S1049023X19000906>
  25. Rosenstock IM. The Health Belief Model and Preventive Health Behavior. *Health Education Monographs*. 1974; 2: 354–386. <https://doi.org/10.1177/109019817400200405>
  26. Rosenstock IM, Strecher VJ, Becker MH. Social Learning Theory and the Health Belief Model. *Health Education Quarterly*. 1988; 15: 175–183. <https://doi.org/10.1177/109019818801500203> PMID: 3378902
  27. Glanz K. *Health Behavior and Health Education: Theory, Research, and Practice*. Jossey-Bass; 2002.
  28. Jones CL, Jensen JD, Scherr CL, Brown NR, Christy K, Weaver J. The Health Belief Model as an Explanatory Framework in Communication Research: Exploring Parallel, Serial, and Moderated Mediation. *Health Commun*. 2015; 30: 566–576. <https://doi.org/10.1080/10410236.2013.873363> PMID: 25010519
  29. PEARL J. Causal diagrams for empirical research. *Biometrika*. 1995; 82: 669–688. <https://doi.org/10.1093/biomet/82.4.669>
  30. Hernán MA, Hernández-Díaz S, Werler MM, Mitchell AA. Causal Knowledge as a Prerequisite for Confounding Evaluation: An Application to Birth Defects Epidemiology. *American Journal of Epidemiology*. 2002; 155: 176–184. <https://doi.org/10.1093/aje/k155.2.176> PMID: 11790682
  31. Greenland S, Pearl J, Robins JM. Causal diagrams for epidemiologic research. *Epidemiology*. 1999; 10: 37–48. PMID: 9888278

32. VanderWeele TJ, Robins JM. Directed Acyclic Graphs, Sufficient Causes, and the Properties of Conditioning on a Common Effect. *American Journal of Epidemiology*. 2007; 166: 1096–1104. <https://doi.org/10.1093/aje/kwm179> PMID: 17702973
33. Killip S, Mahfoud Z, Pearce K. What Is an Intracluster Correlation Coefficient? Crucial Concepts for Primary Care Researchers. *Ann Fam Med*. 2004; 2: 204–208. <https://doi.org/10.1370/afm.141> PMID: 15209195
34. Stuart EA, Azur M, Frangakis C, Leaf P. Multiple Imputation With Large Data Sets: A Case Study of the Children's Mental Health Initiative. *American Journal of Epidemiology*. 2009; 169: 1133–1139. <https://doi.org/10.1093/aje/kwp026> PMID: 19318618
35. Madley-Dowd P, Hughes R, Tilling K, Heron J. The proportion of missing data should not be used to guide decisions on multiple imputation. *Journal of Clinical Epidemiology*. 2019; 110: 63–73. <https://doi.org/10.1016/j.jclinepi.2019.02.016> PMID: 30878639
36. Dong Y, Peng C-YJ. Principled missing data methods for researchers. Springerplus. 2013; 2: 222. <https://doi.org/10.1186/2193-1801-2-222> PMID: 23853744
37. Little RJA, Rubin DB. *Statistical Analysis with Missing Data*. John Wiley & Sons; 2019.
38. Sterne JAC, White IR, Carlin JB, Spratt M, Royston P, Kenward MG, et al. Multiple imputation for missing data in epidemiological and clinical research: potential and pitfalls. *BMJ*. 2009; 338: b2393. <https://doi.org/10.1136/bmj.b2393> PMID: 19564179
39. Greenland S, Finkle WD. A Critical Look at Methods for Handling Missing Covariates in Epidemiologic Regression Analyses. *American Journal of Epidemiology*. 1995; 142: 1255–1264. <https://doi.org/10.1093/oxfordjournals.aje.a117592> PMID: 7503045
40. Zhao Y, Long Q. Variable Selection in the Presence of Missing Data: Imputation-based Methods. *Wiley Interdiscip Rev Comput Stat*. 2017;9. <https://doi.org/10.1002/wics.1402> PMID: 29085552
41. Copyright 2020 SAS Institute Inc. SAS and all other SAS Institute Inc. product or service names are registered trademarks or trademarks of SAS Institute Inc., Cary, NC, USA.
42. Yuan Y. Multiple Imputation Using SAS Software. *J Stat Soft*. 2011;45. <https://doi.org/10.18637/jss.v045.i06>
43. Harel O, Mitchell EM, Perkins NJ, Cole SR, Tchetgen Tchetgen EJ, Sun B, et al. Multiple Imputation for Incomplete Data in Epidemiologic Studies. *American Journal of Epidemiology*. 2018; 187: 576–584. <https://doi.org/10.1093/aje/kwx349> PMID: 29165547
44. Schafer JL. *Analysis of Incomplete Multivariate Data*. CRC Press; 1997.
45. Schützenmeister A, Jensen U, Piepho H-P. Checking Normality and Homoscedasticity in the General Linear Model Using Diagnostic Plots. *Communications in Statistics—Simulation and Computation*. 2012; 41: 141–154. <https://doi.org/10.1080/03610918.2011.582560>
46. Westreich D. Berkson's bias, selection bias, and missing data. *Epidemiology*. 2012; 23: 159–164. <https://doi.org/10.1097/EDE.0b013e31823b6296> PMID: 22081062
47. Perkins NJ, Cole SR, Harel O, Tchetgen Tchetgen EJ, Sun B, Mitchell EM, et al. Principled Approaches to Missing Data in Epidemiologic Studies. *Am J Epidemiol*. 2018; 187: 568–575. <https://doi.org/10.1093/aje/kwx348> PMID: 29165572