



<https://doi.org/10.1038/s41467-021-24833-y>

OPEN

Publisher Correction: Neurovascular coupling and oxygenation are decreased in hippocampus compared to neocortex because of microvascular differences

K. Shaw , L. Bell, K. Boyd, D. M. Grijseels , D. Clarke, O. Bonnar , H. S. Crombag  & C. N. Hall 

Correction to: *Nature Communications* <https://doi.org/10.1038/s41467-021-23508-y>, published online 27 May 2021

The original version of this Article contained errors in Eq. (1), in which $Hbt(t)$ was used as a numerator and denominator instead of $Hbr(t)$.

Equation 1 incorrectly read:

$$CMRO_2(t) = CBF(t) \times \frac{Hbt(t)}{Hbt(t)}$$

The correct form of Eq. (1) is:

$$CMRO_2(t) = CBF(t) \times \frac{Hbr(t)}{Hbt(t)} \quad (1)$$

This has been corrected in the PDF and HTML versions of the Article.

Published online: 19 July 2021



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2021