

# Identification of Putative Potassium Channel Homologues in Pathogenic Protozoa

David L. Prole<sup>1\*</sup>, Neil V. Marrion<sup>2</sup>

**1** Department of Pharmacology, University of Cambridge, Cambridge, United Kingdom, **2** School of Physiology and Pharmacology, University of Bristol, Bristol, United Kingdom

## Abstract

K<sup>+</sup> channels play a vital homeostatic role in cells and abnormal activity of these channels can dramatically alter cell function and survival, suggesting that they might be attractive drug targets in pathogenic organisms. Pathogenic protozoa lead to diseases such as malaria, leishmaniasis, trypanosomiasis and dysentery that are responsible for millions of deaths each year worldwide. The genomes of many protozoan parasites have recently been sequenced, allowing rational design of targeted therapies. We analyzed the genomes of pathogenic protozoa and show the existence within them of genes encoding putative homologues of K<sup>+</sup> channels. These protozoan K<sup>+</sup> channel homologues represent novel targets for anti-parasitic drugs. Differences in the sequences and diversity of human and parasite proteins may allow pathogen-specific targeting of these K<sup>+</sup> channel homologues.

**Citation:** Prole DL, Marrion NV (2012) Identification of Putative Potassium Channel Homologues in Pathogenic Protozoa. PLoS ONE 7(2): e32264. doi:10.1371/journal.pone.0032264

**Editor:** Gordon Langsley, Institut national de la santé et de la recherche médicale - Institut Cochin, France

**Received:** November 20, 2011; **Accepted:** January 24, 2012; **Published:** February 21, 2012

**Copyright:** © 2012 Prole, Marrion. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** This work was funded by a Meres Senior Research Associateship from St. John's College, Cambridge (to DLP). The funder had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

\* E-mail: dp350@cam.ac.uk


## Introduction

Protozoan parasites are major contributors to worldwide disease [1]. They include apicomplexan parasites such as *Plasmodium* spp. (malaria), *Toxoplasma gondii* (toxoplasmosis), *Cryptosporidium* spp. (cryptosporidiosis, diarrhoea) and *Babesia bovis* (babesiosis), as well as the kinetoplastid parasites *Trypanosoma* spp. (sleeping sickness, Chagas' disease) and *Leishmania* spp. (leishmaniasis). These parasites are together responsible for billions of infections and hundreds of thousands of deaths each year [1,2]. Other protozoan parasites causing widespread disease include *Giardia intestinalis* (giardiasis), *Entamoeba histolytica* (dysentery) and *Trichomonas vaginalis* (trichomoniasis). Current treatments for diseases caused by protozoa are often ineffective or poorly tolerated, and emergence of drug resistance is an imminent threat to their efficacy [3–5]. New therapeutic targets and drugs are therefore needed.

K<sup>+</sup> channels are a diverse family of transmembrane proteins, which form K<sup>+</sup>-selective pores and mediate K<sup>+</sup> flux across membranes [6,7]. K<sup>+</sup> channels are essential components in a multitude of homeostatic and signalling pathways and are present in animal cells [6], plants [8,9], fungi [10,11] and many bacteria [7,12]. Only a handful of organisms appear to lack K<sup>+</sup> channels completely, and most of these are bacteria that are obligate parasites [7,12]. Many K<sup>+</sup> channels are present in free-living protozoa such as *Paramecium* [13], but little is known about the existence and physiological role of K<sup>+</sup> channels in pathogenic protozoa, many of which spend part of their life cycles as intracellular parasites. K<sup>+</sup> channels are known to exist in *Plasmodium* spp. [14–16] and K<sup>+</sup>-conductive pathways have also been observed in *Trypanosoma cruzi* [17], but the molecular identity of the channels underlying these latter K<sup>+</sup> fluxes is unknown.

Several subtypes of K<sup>+</sup> channel exist, including voltage-gated (K<sub>v</sub>), inward rectifier (K<sub>ir</sub>), two-pore (K<sub>2P</sub>), calcium-gated (K<sub>Ca</sub>) and cyclic nucleotide-gated (K<sub>CNG</sub>) channels [6] (**Figure 1A**). These channels are all formed by a tetrameric arrangement of pore-forming domains, contributed to by each of four monomeric subunits in most channels, except K<sub>2P</sub> channels which exist as a dimer of subunits, with each subunit containing two domains that contribute to the pore (**Figure 1A**) [6,18]. The K<sup>+</sup>-conducting pore region is comprised of a tetrameric arrangement of re-entrant pore loops (P-loops), part of which forms the selectivity filter, together with the following pore-lining transmembrane domains (TMDs) from each subunit, which form the inner pore (**Figures 1A and 1B**). Diversity of K<sup>+</sup> channels is increased by subunit heteromerization and by the association of auxiliary subunits, such as K<sub>β</sub>, KCNE, KChIP, BKβ, and sulfonylurea (SUR) subunits, which alter the functional properties, trafficking, modulation and pharmacology of K<sup>+</sup> channels [19–22].

In many cell types K<sup>+</sup> channels are found mainly in the plasma membrane, but they are also found in the membranes of intracellular organelles such as mitochondria [23,24], nuclei [25–28], endosomes [29], endoplasmic reticulum [30], secretory vesicles [31,32] and intracellular vacuoles [33–35]. Physiological roles of K<sup>+</sup> flux include setting or altering membrane potentials, effecting osmolyte homeostasis, altering enzyme activity, promoting mitogenesis or apoptosis, and facilitating transmembrane transport processes [6–9,12,36]. Pharmacological or genetic perturbation of K<sup>+</sup> channel activity has profound effects on cell function in many organisms, suggesting that parasite homologues of these channels might represent novel drug targets. Consistent with this, disruption of K<sup>+</sup> channel function in *Plasmodium falciparum* and *Plasmodium berghei* is lethal to these parasites [16,37].



**Figure 1. K<sup>+</sup> channel families.** (A) Topology diagrams of K<sup>+</sup> channel subunits, showing locations of transmembrane domains (TMDs), functional domains and termini. Plus signs denote charged basic residues within the voltage sensor (S4) region of K<sub>v</sub> and K<sub>Ca1</sub> channels. In contrast to other K<sup>+</sup> channel subunits, K<sub>Ca1</sub> channel subunits have extracellular N-termini [145,146]. RCK denotes a Ca<sup>2+</sup>-binding regulator of conductance of K<sup>+</sup> channels domain, which also binds a variety of other ionic ligands in different channels [60–63]. CaM denotes calmodulin (CaM) bound to a CaM-binding site within the channel subunit [60,64]. CNBD denotes a cyclic nucleotide monophosphate (cNMP) binding site [65]; (B) A crystal structure of the KcsA pore domain is shown (PDB accession number 1K4C) [147], with only the TMDs and pore loops of two subunits depicted for clarity. Red circles represent a number of the K<sup>+</sup> ions in the selectivity filter.  
doi:10.1371/journal.pone.0032264.g001

Recent advances in genomics have resulted in whole-genome sequencing of many pathogenic protozoa [1,38–56]. In this study we examine the genomes of pathogenic protozoa comprehensively, using diverse K<sup>+</sup> channel sequences from mammals, plants, fungi, bacteria and archaea, to search for the presence of predicted proteins that may fulfil roles as K<sup>+</sup> channels. We show that genes encoding homologues of K<sup>+</sup> channels exist in all pathogenic protozoa examined. Sequence divergence of putative protozoan channels from their human counterparts in regions that are known to be important for channel activation, ion conduction or drug binding may result in distinct pharmacological profiles. These parasite channels may therefore represent novel targets for anti-parasitic therapy.

## Results

### Identification and classification of K<sup>+</sup> channel homologues

The defining feature of K<sup>+</sup> channels is their selectivity for K<sup>+</sup> ions, which is conferred by residues within the selectivity filter region of the pore [57] (Figure 1). Diverse mammalian K<sup>+</sup> channels show sequence similarity in the selectivity filter region,

with a core selectivity filter motif of XXGXXGX, most commonly TXGYGD [58]. K<sup>+</sup> selectivity is known to be tolerant of some sequence variation in this selectivity filter motif [59] as well as in the outer and inner pore regions, and such variation exists between channel subtypes [58]. For example, selectivity filter sequences of K<sup>+</sup>-selective channels include TIGYGF (eg. K<sub>ir</sub>2.1, K<sub>ir</sub>2.3), TIGYGL (eg. K<sub>ir</sub>2.2), XXGFGX (eg. K<sub>ir</sub>6.2, ERG, EAG, mouse K<sub>Ca1</sub>1.1), and XXGLGD (eg. some K<sub>2P</sub>) [58]. We therefore searched parasite genomes using diverse K<sup>+</sup> channel sequences from humans, plants, fungi, bacteria and archaea (see **Methods**), which together cover most known K<sup>+</sup>-selective pore sequences. We identified predicted protein products in the genomes of pathogenic protozoa, which display significant sequence similarity to K<sup>+</sup> channels in the pore region, including the selectivity filter (Table 1 and Figure 2). These proteins also satisfy other criteria for defining them as putative K<sup>+</sup> channel homologues, such as the presence of multiple TMDs (see **Methods**). These homologues may therefore function as K<sup>+</sup>-selective channels in protozoan parasites. Homologues were classified according to the family of human K<sup>+</sup> channel to which they showed greatest sequence similarity, and according to the presence of conserved functional domains (Figure 1A) such as putative voltage sensors, Ca<sup>2+</sup>-

**Table 1.** Identity of K<sup>+</sup> channel homologues in pathogenic protozoa.

Parasite	K <sub>v</sub> or K <sub>CNG</sub> *	K <sub>Ca</sub>	K <sub>ir</sub>
<i>Plasmodium falciparum</i>	XP_001350669 (12) (K <sub>v</sub> )	XP_001348796 (10)	NF
		XP_001350669 (12)	
<i>Plasmodium knowlesi</i>	XP_002262343 (13) (K <sub>v</sub> )	XP_002260211 (8)	NF
		XP_002262343 (13)	
<i>Plasmodium vivax</i>	XP_001617360 (10) (K <sub>v</sub> )	XP_001615733 (8)	NF
		XP_001617360 (10)	
<i>Toxoplasma gondii</i>	XP_002365940 (8) (K <sub>v</sub> )	XP_002366551 (10)	XP_002369151 (4)
		XP_002365940 (8)	
<i>Cryptosporidium hominis</i>	NF	XP_668687 (8)	XP_666498 (2)
<i>Cryptosporidium muris</i>	XP_002140632(10)(K <sub>v</sub> )	XP_002140632 (10)	XP_002140833 (2)
		XP_002141624 (11)	
<i>Cryptosporidium parvum</i>	NF	XP_626777 (8)	XP_626299 (2)
		XP_001388321 (6) <sup>b</sup> (K <sub>Ca</sub> 4)	
<i>Babesia bovis</i>	NF	XP_001609692 (2)	NF
		XP_001610013 (5)	
<i>Giardia intestinalis</i>	NF	EFO63588 (9)	NF
<i>Entamoeba histolytica</i>	NF	XP_655083 (6)	NF
<i>Leishmania major</i>	NF	XP_001687475 (8)	NF
		XP_001687474 (6)	
		XP_001682763 (7)	
		XP_001687653 (6) <sup>b</sup> (K <sub>Ca</sub> 2/K <sub>Ca</sub> 3)	
		XP_001687652 <sup>a</sup>	
<i>Leishmania infantum</i>	NF	XP_001462697 (7)	NF
		XP_001462696 (6)	
		XP_001465142 (9)	
		XP_001464237 (6) <sup>b</sup> (K <sub>Ca</sub> 3/K <sub>Ca</sub> 2)	
		XP_001464236 <sup>a</sup>	
<i>Leishmania braziliensis</i>	NF	XP_001561516 (6)	NF
		XP_001564698 (6)	
		XP_001563345 (7) <sup>b</sup> (K <sub>Ca</sub> 3/K <sub>Ca</sub> 2)	
		XP_001563344 <sup>a</sup>	
<i>Trypanosoma brucei</i>	NF	EAN76555 (7)	NF
		XP_001219138 (6)	
<i>Trypanosoma cruzi</i>	NF	XP_821941 (8)	NF
		XP_816151 (8)	
		XP_820381 (6)	
		XP_820126 (6)	
		XP_818052 (8) <sup>b</sup> (K <sub>Ca</sub> 3/K <sub>Ca</sub> 2)	
		XP_813982 (8) <sup>b</sup> (K <sub>Ca</sub> 3/K <sub>Ca</sub> 2)	
		XP_813983 <sup>a</sup>	
		XP_818073 <sup>a</sup>	
<i>Trichomonas vaginalis</i>	XP_001324404 (6)*	NF	NF
		XP_001319471 (6)*	
		XP_001326965 (6)*	
		XP_001318022 (6)*	
		XP_001325751 (2) <sup>a</sup>	

Protein accession numbers are shown and NF denotes that no homologues were found. Number of predicted TMDs is indicated in parentheses. K<sup>+</sup> channel homologues are classified on the basis of closest similarity to a particular subtype of human K<sup>+</sup> channel subunit and according to the presence of characteristic functional domains, such as a charged TMD4 (K<sub>v</sub>), the presence of a cyclic nucleotide-binding domain CNBD (K<sub>CNG</sub>), or the presence of RCK domains (K<sub>Ca</sub>). Where a protein showed similarity to more than one class of K<sup>+</sup> channel, its accession number is shown in both relevant columns (eg. all K<sub>v</sub> homologues are also K<sub>Ca</sub> homologues).

\*K<sub>CNG</sub> homologues which contain a CNBD, but also a charged TMD4;

**Table 1.** Cont.

<sup>a</sup>likely non-selective or non-functional due to GYRD, GYSD or GYSE selectivity filter motifs;

<sup>b</sup>no putative RCK domains or calmodulin-binding domains (CaMBDs) were detected in these proteins by searching the Conserved Domains Database, but in BLASTP searches of the human genome these proteins showed greatest sequence similarity to the  $K_{Ca}$  channel subtypes indicated in parentheses. The *P. falciparum* proteins XP\_001609692 and XP\_001350669 are identical to the previously described PfKch2 and PfKch1 proteins respectively [14,16]. The *B. bovis* (XP\_001610013) and *C. hominis* (XP\_668687) proteins have been identified previously as orthologues of PfKch1 in *P. berghei* [16]. In addition to the  $K^+$  channel homologues shown, homologues of putative adenylyl cyclase/ $K^+$  channel fusion proteins [123,124] that contain GXG motifs after their TMD6 domains were also identified in *P. falciparum* (XP\_001348216), *P. knowlesi* (XP\_002260946), *P. vivax* (XP\_001616904), *T. gondii* (XP\_002368352, XP\_002370938 and XP\_002367966), *C. muris* (XP\_002140763), *C. hominis* (XP\_666311) and *C. parvum* (XP\_626352). Homologues of these proteins were absent in all other parasites examined.  
doi:10.1371/journal.pone.0032264.t001

sensing regulator of conductance (RCK) domains of  $K_{Ca}$  channels [60–63], calmodulin (CaM)-binding domains (CaMBDs) [60,64], or cyclic nucleotide-binding domains (CNBDs) [65] (**Table 1** and **Figure 2**). The *P. falciparum* proteins XP\_001609692 and XP\_001350669 are identical to the previously described PfKch2 and PfKch1 proteins respectively (also known as PfK2 and PfK1 respectively) [14,16]. The homologues in *B. bovis* (XP\_001610013) and *Cryptosporidium hominis* (XP\_668687) have been previously identified by similarity to a  $K^+$  channel homologue in *P. berghei* [16].

### Putative pore regions of $K^+$ channel homologues

Parasite  $K^+$  channel homologues have a variety of putative selectivity filter sequences, including XXGYGD, XXGFGD and XXGYGS (henceforth referred to as GYGD, GFGD and GYGS) (**Figure 2**). The sequence GYGD is common in mammalian  $K^+$  channels [58], while GFGD sequences also occur in several channels, including human  $K_{ir}$ , ERG and EAG, as well as mouse  $K_{Ca1.1}$  [58]. In contrast, GYGS is very rare in mammals [58], but occurs in several of the parasite  $K^+$  channel homologues examined, which are most closely related to human  $K_{ir}$  channels (**Figure 2**). The identity of the residue immediately following the second glycine of the selectivity filter motif affects conductance and rectification as well as gating, binding of  $K^+$  and potency of inhibitors [66,67]. An aspartate residue at this locus reduces block by  $Cd^{2+}$  [68] and tetraethylammonium (TEA) [66], suggesting that sensitivity to these and other blockers may differ between GYGD-containing human  $K^+$  channels and the GYGS-containing protozoan  $K^+$  channel homologues. These GYGS-containing parasite channels also contain a glutamate residue at an analogous position to the conserved P-loop glutamate residue of  $K_{ir}$  channels and KcsA (E138 in  $K_{ir2.1}$ , **Figure 2**) that is involved in gating at the selectivity filter [69,70]. In contrast, these homologues lack an aspartate residue (D172 in  $K_{ir2.1}$ ) within the inner helix that confers strong inward rectification on some mammalian  $K_{ir}$  channels [71]. These parasite GYGS-containing  $K^+$  channel homologues may therefore show rectification and sensitivity to extracellular blockers that differs from human  $K_{ir}$  channels, but may exhibit similarities in gating.

A separate group of proteins (henceforth termed GYX-containing proteins) was also defined in kinetoplastid *Leishmania* and *Trypanosoma* parasites as well as *T. vaginalis*. These proteins have GYSD/E or TIGYRD sequences respectively at the putative selectivity filter locus, but otherwise show pronounced similarity to various mammalian  $K^+$  channel pores (**Figure 2** and **Table 1**). Each of the genes encoding these GYX proteins in *Leishmania* spp. and *T. cruzi* is present in a chromosomal region that is closely adjacent to genes encoding one of the other  $K^+$  channel homologues identified, which have canonical GYG- or GFG-containing selectivity filters. This suggests that these GYX proteins may be paralogues that arose via gene duplication. Mutations of the second glycine in the GXG motif have been shown to

eliminate  $K^+$  selectivity and function in  $K^+$  channels [72], suggesting that these GYX proteins may be non-selective cation channels, or may be non-functional.


The selectivity filter and adjoining P-loop form the binding site for many drugs and toxins that block  $K^+$  channels [73–78]. Likewise, the inner pore region of  $K^+$  channels often binds a variety of drugs [77,79–84]. Sequence differences between human and parasite homologues in these regions (**Figure 2**) suggest that it may be possible to isolate drugs that bind specifically to the inner pore of parasite proteins. For example, many protozoan  $K_{Ca}$  homologues differ from human  $K_{Ca1.1}$  at a locus (equivalent to G376 of h $K_{Ca1.1}$  in **Figure 2**) that determines sensitivity to the potent  $K_{Ca1.1}$ -specific fungal toxin paxilline [85]. This suggests that human  $K_{Ca}$  channels and protozoan  $K_{Ca}$  channel homologues might exhibit different sensitivity to paxilline and related drugs.

### $K_v$ channel homologues

The genomes of *Plasmodium* spp., *T. gondii*, *Cryptosporidium muris* and *T. vaginalis* contain genes encoding homologues of  $K_v$  channels, with TMD4 regions that contain a number of regularly spaced basic residues, similar to the S4 voltage sensor sequences of  $K_v$  channels [60] (**Table 1**, **Figure 1** and **Figure 3**). The apicomplexan  $K_v$  homologues also show sequence similarity to the C-terminal tails of  $K_{Ca1}$  channels, including the RCK domains that bind  $Ca^{2+}$  or other ions [61,62,86] (*data not shown*). This suggests that these parasite homologues may be dually modulated by voltage and ions such as  $Ca^{2+}$ ,  $Mg^{2+}$  or  $H^+$ . In contrast,  $K_v$  homologues in *T. vaginalis* all contain conserved CNBDs in their C-terminal tails (*see later*), suggesting that they, like mammalian hyperpolarization-activated cyclic nucleotide-gated non-selective (HCN) channels [65], may be dually modulated by voltage and cyclic nucleotides such as cyclic adenosine monophosphate (cAMP) or cyclic guanosine monophosphate (cGMP).


### $K_{Ca}$ channel homologues

The most common genes encoding  $K^+$  channel homologues in protozoan parasites are those encoding  $K_{Ca}$  channel homologues (**Table 1** and **Figure 2**). This is consistent with the presence of complex  $Ca^{2+}$ -signalling machinery in protozoa [2,87,88]. Known  $K_{Ca}$  channels form two main families. Proteins in one family are activated by  $Ca^{2+}$  (and in some cases by other ions such as  $Mg^{2+}$ ,  $H^+$  and  $Na^+$ ) via direct binding to domains within the channel, including the C-terminal RCK domains of  $K_{Ca1}$  channels [60–63] (**Figure 1A**). Proteins in the second family, which includes  $K_{Ca2}$  and  $K_{Ca3}$  channels, are activated by  $Ca^{2+}$  via binding of their C-terminal tails to the accessory  $Ca^{2+}$ -binding protein CaM [60,64] (**Figure 1A**). Many of the protozoan  $K_{Ca}$  channel homologues show sequence similarity to the C-terminal tails of  $K_{Ca1}$  or  $K_{Ca2/3}$  channels, including the RCK domains and CaMBDs (*data not shown*), suggesting that their activity may be regulated by  $Ca^{2+}$ , or by other ions such as  $Mg^{2+}$ ,  $H^+$  or  $Na^+$ .



**Figure 2. Multiple sequence alignment of protozoan K<sup>+</sup> channel homologues with the pores of mammalian K<sup>+</sup> channels.** Predicted pore-lining TMD regions are underlined. The GXG motif of human K<sup>+</sup> channels is shaded in grey. Total number of residues in each protein is indicated in parentheses to the right of each sequence. *L. braz.* denotes *L. braziliensis*, and *G.intest.* denotes *G. intestinalis*. The proteins XP\_001609692 and XP\_001350669 encoded by the *P. falciparum* genome are identical to the previously described PfKch2 and PfKch1 proteins respectively [14,16]. The proteins XP\_001610013 and XP\_668687 are identical to previously identified K<sup>+</sup> channel homologues in *B. bovis* and *Cryptosporidium hominis* respectively [16]. The proteins labelled GYX have GYRD, GYSD or GYSE-containing selectivity filter regions, suggesting a lack of K<sup>+</sup> selectivity or function.

doi:10.1371/journal.pone.0032264.g002



**Figure 3. Protozoan K<sup>+</sup> channels containing charged TMD4 regions.** (A) Topology diagram of K<sub>v</sub>1.2, with the positively charged TMD4 shown in red; (B) Multiple sequence alignment of the TMD4 regions of human voltage-gated K<sub>v</sub>1.2 and K<sub>Ca</sub>1.1, plant voltage-gated KAT1 and the predicted TMD4 regions of those protozoan K<sup>+</sup> channel homologues containing at least three basic residues within this region. Asterisks above the alignment indicate basic residues involved in voltage sensing in K<sub>v</sub>1.2 channels. doi:10.1371/journal.pone.0032264.g003


We were most interested in the multiple K<sub>Ca</sub> channel homologues present in *Leishmania* parasites and chose to interrogate these homologues further (Figure 4). Based on full-length alignments and phylogenetic relationships (Figure 4A), K<sub>Ca</sub> homologues in *Leishmania* parasites could be grouped into K<sub>Ca</sub>1-like proteins (which showed similarity to the discontinuous RCK domains of hK<sub>Ca</sub>1.1, *data not shown*) and K<sub>Ca</sub>2/3-like proteins which had greater similarity to K<sub>Ca</sub>2/3 channels. The latter group contain homologues with canonical GYG-containing putative selectivity filter regions, as well as homologues with GYX-containing selectivity filters (Figure 4A and Figure 2). These K<sub>Ca</sub>2/3-like proteins have consensus IQ-motifs in their C-terminal tails, shortly after the final TMD, which are predicted to form CaMBDs (shown in Figure 4B for the GYG-containing homologues). These predicted CaMBDs overlap closely with the well-characterized CaMBD of K<sub>Ca</sub>2.2 (Figure 4B) and share sequence identity at several loci that are critical for binding of CaM to human K<sub>Ca</sub>2.2 (Figure 4B, open triangles) [64]. This suggests that these parasite homologues may form K<sub>Ca</sub> channels whose activity is controlled by Ca<sup>2+</sup> acting via bound CaM, similar to human K<sub>Ca</sub>2 channels. The K<sub>Ca</sub> homologues in *Leishmania* (as well as the other protozoa examined) show considerable sequence divergence from human K<sub>Ca</sub>2 channels in the P-loop and selectivity filter, including at positions implicated in the binding of isoform-specific drugs and toxins to K<sub>Ca</sub>2 channels (filled triangles in Figure 4B) [78]. These homologues also differ markedly from human K<sub>Ca</sub>2 channels in the predicted TMD3-TMD4 linker region (*data not shown*), which is also an important determinant of drug binding in K<sub>Ca</sub>2 channels [89]. This suggests that these parasite K<sub>Ca</sub> homologues may exhibit unique pharmacological profiles.

**K<sub>CNG</sub> homologues.** Several genes encoding K<sup>+</sup> channel homologues with conserved CNBDs (identified by searches of the Conserved Domains Database, NCBI) are present in *T. vaginalis*, but are absent from all other protozoa examined (Table 1 and Figure 5). These putative K<sup>+</sup> channel homologues exhibit sequence similarity to the CNBD of the well-characterized HCN2 channel that is activated by cAMP and cGMP, including some conserved residues involved in binding of cyclic nucleotides [65] (Figures 5A, 5B and 5C). Predicted secondary structure of the CNBD domain in HCN2 (Figure 5A) closely matched secondary

structure in the crystal structure of HCN2 [65] (Figure 5B), suggesting that secondary structure of the parasite proteins could also be predicted accurately. Predicted secondary structure of the putative CNBDs of protozoan K<sub>CNG</sub> channel homologues in most cases closely matched that of HCN2 (Figures 5A and 5B), suggesting that the structure of these domains is also conserved in human HCN2 and *T. vaginalis* K<sub>CNG</sub> homologues. One exception was the C-terminal end of this domain in the GYX protein XP\_001325751 (which as discussed earlier may not form functional or K<sup>+</sup>-selective channels), which lacked entirely the final alpha-helical stretch of HCN2 termed the C-helix, which is involved in binding and efficacy of cyclic nucleotides in HCN2 [90] (Figure 5A and Figure 5B). Hence most of the protozoan K<sub>CNG</sub> homologues identified here may contain conserved CNBDs that bind cyclic nucleotides. The cyclic nucleotide selectivity of binding and efficacy in these putative channels cannot at present be predicted on the basis of sequence alone, and will require experimental testing. Unlike the non-selective mammalian HCN and CNG channels, but similar to prokaryotic K<sub>CNG</sub> channels [7], these *T. vaginalis* proteins contain canonical K<sup>+</sup> channel selectivity filter motifs (T/AXGYGD), suggesting that they may be K<sup>+</sup>-selective. We also searched parasite genomes with sequences of the K<sup>+</sup>-permeable (but non-selective) mammalian cyclic nucleotide-gated non-selective cation (CNG) channels and hyperpolarization-activated cyclic nucleotide-gated non-selective cation (HCN) channels, but no additional homologues were found.

### K<sub>ir</sub> channel homologues

Among the protozoan genomes examined, genes encoding K<sub>ir</sub> channel homologues are found only in the genomes of *Cryptosporidium* spp. and *T. gondii* (Table 1 and Figure 2). K<sub>ir</sub> channels are widespread among many organisms, and many subtypes exist that are differentially regulated by diverse stimuli including adenosine triphosphate (ATP), G-protein activation, phospholipids, and divalent cations [71]. As discussed earlier, the predicted pore region of the protozoan K<sub>ir</sub> channel homologues shows some differences to human K<sub>ir</sub> channels, which may confer unique characteristics on these homologues. The K<sub>ir</sub> homologue in *T. gondii* shows most similarity to human ATP-modulated K<sub>ir</sub>6.2 and G-protein activated K<sub>ir</sub>3 channels (*data not shown*), suggesting that cytosolic ATP or G-proteins within parasites may regulate this



**B**

		TMD5	P-loop	selectivity filter	TMD6	
hK <sub>Ca</sub> 2.2	(NP_067627)	WTVRACERYHDQDVTSN	FLGAMWLISITFLS	IGYGMVNPNTYCGKGVCL		376
<i>L. major</i>	(XP_001687653)	WLTVGCLFARTQN---	LSLGDALWFSFQSL	STLGYGDITPSTLSGRAVAF		262
<i>L. infantum</i>	(XP_001464237)	WLTVGCLFARTQN---	LSLGDALWFSFQSL	STLGYGDITPSTLSGRVVAE		262
<i>L. braz.</i>	(XP_001563345)	WLTVGCLFARAQS---	LSLGDALWFSFQSL	ATLGYGDITPSTLSGRTVAE		319
		* * : *	: *::	:: : *::	* * * :	
hK <sub>Ca</sub> 2.2	(NP_067627)	L T G I M G A G C T A L V V A V V A R K L E L T K A E K H V H N F M M D T Q L T K R V K N A A A N V				426
<i>L. major</i>	(XP_001687653)	I A W I V S Y V I M A F L I I T M Y S L L Q A S D R S H N M Q T L M E C H E L A H S L R G R S A R V				312
<i>L. infantum</i>	(XP_001464237)	I A W I L S Y V I M A F L I I T M Y S L L Q A S D R S H N M Q T L M E C H D L A H S L R G R S A R V				312
<i>L. braz.</i>	(XP_001563345)	I A W I A S Y V I M A F L I T A M Y S L L Q A S D R L H N M Q A L M E C H E L I Q S L R G R S A R V				369
		:: * * * :	* : :	:: : * :	* : : :	
		CaM-binding domain				
hK <sub>Ca</sub> 2.2	(NP_067627)	L R E T W L I Y K N T K L V K K I D H A K V R K H Q R K F - L Q A I H Q L R S V K M E Q R K L N - -				473
<i>L. major</i>	(XP_001687653)	I Q L A W R L H K A R K V G P H E P S V K Q K V Y T L A L S W L L T H M I A A V R R T R Q Q L N G A				362
<i>L. infantum</i>	(XP_001464237)	I Q L A W R L R K A R K V G P H A P S L K Q K V Y T L A L S W L L T D M I S A V R R T R Q Q L N G A				362
<i>L. braz.</i>	(XP_001563345)	I Q L A W R L H K A R K V R L H A P S L K Q R M Y T L V L S W L V T H T I I A R R R T R Q Q L N S V				419
		:: : * * :	* : :	: : :	:: : * :	
hK <sub>Ca</sub> 2.2	(NP_067627)	-----DQANTLVLDLAKTQINIMYDMISDLNE-----RSEDFEKRIVTTLETK				513
<i>L. major</i>	(XP_001687653)	VRSLNETHNIHPLTGLSAYQYNEYLQLTEKAARQLQRAKDVDVRYLALRDR				412
<i>L. infantum</i>	(XP_001464237)	VRSLSETHNIHPLTGLSAYQYNAYLQLTEKAARQLQRAKDVDVRYLALRDR				412
<i>L. braz.</i>	(XP_001563345)	VRSLQETKTHPLTGLSEYQYNAYLQVTHKAARQLQRVKDVDRVYALALRDR				469
		: : * * :	* * :	: :	* * : :	
hK <sub>Ca</sub> 2.2	(NP_067627)	--LETLLIG-----SIHALPLGISQTIRQQQDFIEAQMESYDKHVTYNAE				556
<i>L. major</i>	(XP_001687653)	DVLASSLSRRDIESIIVLPGDSLEPIDTALPAALTSSSSGTAAASVAEMAEW				462
<i>L. infantum</i>	(XP_001464237)	DVLASSLSRRDIEIIVLPGDSLEPIGTALPVALTSSSSGTAAASVAEMAEW				462
<i>L. braz.</i>	(XP_001563345)	DVLASSLSRRDIESIIVLPGDSLEPIGAAPPTTFTGIGSTAVKMAEMAQW				519
		* : :	* * * :	* :	: :	
hK <sub>Ca</sub> 2.2	(NP_067627)	RSRSSRRRRSS-----STAPPTSSESS-				579 (579)
<i>L. major</i>	(XP_001687653)	RDRVAHLEQKCARLAEVL E A I S A V A A T H P P S M A V S Q M				499 (499)
<i>L. infantum</i>	(XP_001464237)	RDRVTQLEQKCARL TE V L E A I S A V A E T H P P S M A V S Q M				499 (499)
<i>L. braz.</i>	(XP_001563345)	RNQLTQLERKCARL TE V L E A M S A V A E T H A P S M S V S Q V				556 (556)
		* : :	: : :	* * :	* * :	

**Figure 4.  $K_{Ca}$  channel homologues in *Leishmania* parasites.** (A) Phylogram showing the relationship between the sequences of human  $K_{Ca}$  channels and  $K^+$  channel homologues in *Leishmania* spp. (see Methods). Branch length scale bar and branch support values are shown (see Methods). Two main groups of *Leishmania* proteins ( $K_{Ca1}$ -like and  $K_{Ca2/3}$ -like) are indicated. Selectivity filter GYG-containing  $K_{Ca2/3}$ -like channels and their GYG-containing putative paralogues are also indicated; (B) Multiple sequence alignment of human  $K_{Ca2.2}$  (small-conductance  $Ca^{2+}$ -activated SK2 channels) with the GYG-containing  $K_{Ca2/3}$ -like homologues in *Leishmania* spp. Selectivity filter, TMD and P-loop regions are indicated above the alignment. Filled triangles above the alignment indicate  $K_{Ca2.2}$  residues implicated in binding of inhibitory toxins. Those previously shown experimentally to alter toxin effects are indicated by red triangles, while additional residues implicated via molecular modelling are indicated by black triangles [78]. The yellow shaded region denotes the fragment of  $K_{Ca2.2}$  that binds CaM [64] and open triangles indicate specific  $K_{Ca2.2}$  residues known to be involved in binding CaM.  
doi:10.1371/journal.pone.0032264.g004

homologue.  $K_{ir}$  homologues in *Cryptosporidium* spp. were also most similar to human ATP-modulated  $K_{ir6}$ , as well as  $K_{ir2}$  channels (data not shown), suggesting that these homologues might also be regulated by cytosolic ATP within parasites.

### $K_{2P}$ channel homologues

We used the sequences of human and yeast  $K_{2P}$  channels to search for predicted proteins with both sequence similarity to  $K_{2P}$  channels [18,91] and at least four predicted TMDs in two distinct regions, each with credible potential as selectivity filter regions. Using these criteria, we found no evidence for genes encoding homologues of  $K_{2P}$  channels in the parasite genomes examined. Although the  $K_{ir}$  homologue in *T. gondii* contained four predicted TMDs in two regions, only the second pair of TMDs had an intervening sequence with similarity to canonical selectivity filters (data not shown).

### Protozoan parasites lack homologues of $K^+$ channel auxiliary subunits

Many  $K^+$  channels are associated with auxiliary proteins that can change their biophysical properties, localization or regulation by cellular signalling pathways [19,20]. These include the KCNE, KChIP and  $K_v\beta$  subunits that alter  $K_v$  channel activity and localization [20,21]. Other auxiliary subunits include the BK $\beta$  family of subunits that alter BK channel activity and pharmacology [20], and the sulfonylurea (SUR) subunits that are responsible for the nucleotide-diphosphate sensitivity and pharmacological profile of  $K_{ir6}$  channel complexes [92,93]. We searched the genomes of protozoan pathogens for genes encoding predicted proteins with similarity to these auxiliary subunits, but none were found. This suggests that the protozoan parasites examined here lack conventional auxiliary subunits of  $K^+$  channels.

### Discussion

All protozoan genomes examined contain genes encoding  $K^+$  channel homologues (Table 1), suggesting that these putative channels have widespread and conserved physiological functions in these organisms. Many of these putative  $K^+$  channels are not yet annotated in available pathogen databases (<http://eupathdb.org/eupathdb>) [94]. Genes encoding homologues of most of the major families of  $K^+$  channel ( $K_v$ ,  $K_{Ca}$ ,  $K_{ir}$  and  $K_{CNG}$  channels) are present in protozoan genomes, although genes encoding homologues of  $K_{2P}$  channels appear to be absent. Experimental studies will be required to confirm the expression and function in parasites of these putative  $K^+$  channel homologues.

In mammalian cells, most plasma membrane  $K^+$  channels mediate  $K^+$  efflux, due to the common occurrence of large transmembrane  $K^+$  concentration gradients and relatively depolarized membrane potentials. Outward rectification or depolarization-induced activation of the channels themselves also contributes to selective efflux of  $K^+$  in many cases. In contrast, some plasma membrane  $K^+$  channels are capable of mediating  $K^+$  influx [95,96], due to unusual  $K^+$  concentration gradients or

membrane potentials. Inward rectification [71] and hyperpolarization-induced activation [97–99] also facilitate selective  $K^+$  influx in some cases. Protozoan parasites in many cases spend part of their lifecycle within cells and part in an extracellular environment. The  $K^+$  concentrations of these environments in a mammalian host differ by ~40-fold (4 mM extracellular  $K^+$  versus 155 mM intracellular  $K^+$ ) [100]. Depending on the lifecycle stage, the concentration gradient for  $K^+$  flux across the parasite plasma membrane may therefore differ considerably. In this context it is interesting that *Plasmodium* parasites induce alterations of the  $Na^+/K^+$  ratio in host cells and thereby reduce intracellular  $K^+$  concentration [101–104]. Whether other parasites also exert similar effects on host cells is unknown. Whether influx of  $K^+$  occurs through  $K^+$  channel homologues in the plasma membrane of intracellularly located parasites, and whether this might be exploited therapeutically (see later) remains to be explored.

Protozoan  $K_v$  channel homologues may be regulated by transmembrane voltage. Most  $K_v$  channels are activated by depolarization [60], while a few are activated by hyperpolarization [98,99]. However, known depolarization-activated and hyperpolarization-activated channels show similar voltage sensor sequences [105], making it difficult to determine the polarity of voltage dependence on the basis of sequence alone. Further experimental studies will therefore be required to define the properties of these homologues. The vast majority of mammalian  $K_v$  channels are present and functional in the plasma membrane, which experiences the most substantial changes in transmembrane potential. Parasite  $K_v$  channel homologues therefore seem likely to reside within the plasma membrane of these organisms, although this will require experimental testing. Parasite  $K^+$  channels may also be expressed in the membranes of their host cells. For example, in the case of *P. falciparum*, the PfKch1 channel is located in the plasma membrane of the host erythrocyte, while PfKch2 is mainly located in the parasite [15]. The presence of putative  $K_v$  channel homologues in protozoa suggests that these organisms may experience dynamic physiological changes in membrane potential. The plasma membrane potentials of some protozoa have been estimated. For example, the intraerythrocytic form of *P. falciparum* has an estimated membrane potential of  $-95$  mV, which is contributed to by  $K^+$  flux [106]. Similarly, the plasma membrane potential of the bloodstream form of *T. brucei* has been measured as  $-82$  mV and is mainly due to  $K^+$  flux [107]. In addition, the membrane potential of *Leishmania donovani* amastigotes has been measured as between  $-90$  and  $-113$  mV and is also contributed to by  $K^+$  flux [108]. However, whether the membrane potentials of protozoan parasites change between stages of the lifecycle or in response to environmental stimuli, and whether the  $K_v$  channel homologues identified here respond to such changes is unknown.

Genes encoding  $K_{Ca}$  channel homologues are present in all protozoa except *T. vaginalis*. The  $K_{Ca2/3}$  channel homologues in many cases contain consensus CaMBDs [65], while the  $K_{Ca1}$  homologues show similarity to the discontinuous  $Ca^{2+}$ -binding RCK domains of  $K_{Ca1}$  channels [60–63]. Searches of parasite genomes using the sequence of human CaM showed that genes





encoding homologues of CaM are present in the genomes of all protozoa examined in this study, consistent with a previous report [109] except that we additionally identified a CaM homologue in *C. parvum* (*data not shown*). In addition, effects attributed to functionally expressed CaM have been reported in many pathogenic protozoa [110–117]. This suggests that CaM is a likely  $\text{Ca}^{2+}$ -binding modulator of protozoan  $\text{K}_{\text{Ca}}$  channel homologues with consensus CaMBDs. RCK domains of  $\text{K}_{\text{Ca}}$  channels can directly confer regulation by  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{Na}^{+}$  or  $\text{H}^{+}$  ions [61]. The discontinuous nature of RCK domain structure [118] and the occurrence of ion binding sites at interfaces between RCK subunits [62,63,119], as well as between RCK subunits and other regions of the channel [120], makes prediction of the ionic specificity of RCK domains difficult on the basis of sequence alone. Experimental testing will be required to determine which of these potential modulatory factors acts at the various parasite  $\text{K}_{\text{Ca}1}$  homologues identified.

CNBD-containing  $\text{K}_{\text{CNG}}$  channels show great variability in their occurrence within genomes. They are very rare in prokaryotes [7], but are abundant in *Paramecium* [12,13]. The relative abundance of CNBD-containing  $\text{K}_{\text{CNG}}$  homologues in *T. vaginalis* is consistent with a large number of genes encoding adenyl cyclases in this organism [121]. This suggests that these proteins may play roles in diverse cyclic nucleotide signalling pathways in this parasite and that these pathways may be possible therapeutic targets.

$\text{K}_{\text{ir}}$  channels are widespread and have diverse functions in many organisms [71]. Many different stimuli affect the activity of  $\text{K}_{\text{ir}}$  channels, including G-proteins, phospholipids, divalent and monovalent cations, as well as ATP [71]. Which stimuli affect the protozoan  $\text{K}_{\text{ir}}$  homologues identified here is difficult based on sequence alone, and will require future experimental testing.

Some parasite  $\text{K}^{+}$  channel subunit homologues had GYX-containing putative selectivity filters, suggesting that they may be non-selective or non-functional. Some of these proteins may be paralogues of  $\text{K}^{+}$  channels with canonical GYG-containing selectivity filters. Interestingly, GYX-containing  $\text{K}^{+}$  channel homologues exist in some prokaryotes, but their ionic selectivity and function are unknown [7]. In contrast, genes encoding GYX-containing  $\text{K}^{+}$  channel subunit homologues are not found in the human genome (*data not shown*). This suggests that if functional, the parasite GYX proteins (whether  $\text{K}^{+}$ -selective channels or not) may constitute unique drug targets in protozoan parasites.

In addition to the  $\text{K}^{+}$  channel homologues identified, it is possible that other  $\text{K}^{+}$ -permeable channels with novel architectures may exist in parasites. For example, a novel family of adenyl cyclases found in members of the Alveolata subkingdom of protozoa, such as *Paramecium* and *P. falciparum*, may also possess  $\text{K}^{+}$  channel activity [122–124]. These proteins are involved in parasite exocytosis and infectivity [125]. These putative adenyl cyclase/ $\text{K}^{+}$  channel fusion proteins have an architecture that is radically different from that of canonical  $\text{K}^{+}$  channels, with a GXG motif situated *after* TMD6 [124]. Only a single study has thus far indicated their possible intrinsic function as  $\text{K}^{+}$ -conductive (but relatively non-selective) ion channels, using a member of this protein family from *Paramecium* [122]. Hence this family of proteins cannot at present be definitively categorized as functional  $\text{K}^{+}$  channel homologues in parasites, but may in future be shown to contribute to  $\text{K}^{+}$  flux in these organisms. Consistent with previous reports [123–125] our analyses identified genes encoding homologues of these fusion proteins in *Plasmodium* spp., *T. gondii* and *Cryptosporidium* spp. (legend to **Table 1**). Genes encoding homologues of these proteins were absent in the genomes of all other parasites examined.

Genes encoding homologues of  $\text{K}^{+}$  channel auxiliary subunits are absent in the genomes of the protozoa examined. This suggests that functional  $\text{K}^{+}$  channel complexes in these organisms may lack the diversity of human  $\text{K}^{+}$  channel complexes. In addition, since auxiliary subunits can dictate the pharmacology of native  $\text{K}^{+}$  channel complexes in mammalian cells [20,92,93], the pharmacology of native  $\text{K}^{+}$  channel complexes in parasites may differ substantially from those of humans. It is also possible that unique and as yet unidentified auxiliary subunits exist in protozoa, which are unrelated to currently known auxiliary subunits of  $\text{K}^{+}$  channels in other organisms.

### $\text{K}^{+}$ channels in parasitic protozoa and their free-living relatives

In contrast to the relatively few genes encoding  $\text{K}^{+}$  channel subunit homologues in parasitic protozoa, the genome of their free-living ciliate relative *Paramecium tetraurelia* has several hundred genes encoding  $\text{K}^{+}$  channel subunits [13]. Whether this dramatic difference arose due to the acquisition of a parasitic existence in some protozoa is unclear.

BLAST searches using the sequences of  $\text{KcsA}$  as well as human  $\text{K}_{\text{v}1.2}$ ,  $\text{K}_{\text{ir}2.1}$  and  $\text{K}_{\text{Ca}1.1}$  proteins suggest that the genome of the free-living flagellate *Monosiga brevicollis* encodes at least nine distinct  $\text{K}^{+}$  channel subunit homologues (*data not shown*), compared with the smaller number in kinetoplastid parasites (*eg.* only two in *T. brucei*). Whether the fewer  $\text{K}^{+}$  channel homologues in the flagellate parasites examined here is due to a parasitic existence is unclear.

### Protozoan $\text{K}^{+}$ channels as therapeutic targets

$\text{K}^{+}$  channels are critical for cellular homeostasis and signal transduction, and pharmacological modulation of these channels can lead to marked changes in cell growth and viability [36,77,126,127]. Disrupting  $\text{K}^{+}$  channel function in *P. falciparum* and *P. berghei* severely compromises survival of these parasites [16,37], suggesting that parasite  $\text{K}^{+}$  channels may be attractive drug targets for treatment of parasitic disease. Interestingly, genetic disruption of  $\text{K}^{+}$  channel function affects different life cycle stages in *P. falciparum* and *P. berghei*. In *P. falciparum* the asexual blood stage is affected [15], while in *P. berghei* the sexual mosquito stage is severely affected but the asexual stage is relatively unaffected [16]. While the human genome encodes more than 70  $\text{K}^{+}$  channel subunits [13], the genomes of pathogenic protozoa each contain only a small number of genes encoding homologues of  $\text{K}^{+}$  channel subunits (**Table 1**). This striking lack of redundancy in the  $\text{K}^{+}$  channel complement of protozoan parasites further suggests that these channels might be effective therapeutic drug targets.

Several drugs that inhibit  $\text{K}^{+}$  channel activity are known to be toxic to protozoan parasites. In many cases the primary anti-parasitic mode of action of these drugs is likely to be on processes other than  $\text{K}^{+}$  flux. However,  $\text{K}^{+}$  channel-blocking drugs may alter the activity of the protozoan parasite  $\text{K}^{+}$  channel homologues described in this study, perturb cellular  $\text{K}^{+}$  homeostasis and contribute to the decreased survival of parasites. For example, chloroquine blocks  $\text{K}_{\text{ir}}$  channels [128] and is also toxic to various species of *Plasmodium* [129]. The anti-trypanosomal drug pentamidine also blocks  $\text{K}_{\text{ir}}$  channels [130]. The  $\text{K}_{\text{ir}6}$  channel antagonist glibenclamide [131] and the  $\text{K}^{+}$  channel blocker amiodarone [132] are both known to be lethal to *Leishmania mexicana*. The  $\text{K}^{+}$  channel blocker amantadine is toxic to *P. falciparum* [133]. In addition, bicuculline and tubocurarine, which block some  $\text{K}^{+}$  channels in addition to other targets [134,135] are also toxic to *P. falciparum* [37]. Functional evidence will be required before any

causal links between inhibition of parasite  $K^+$  channel activity and anti-parasitic effects can be established.

Drugs that *activate*  $K^+$  channels may also be toxic to protozoan parasites. This has not been experimentally tested in protozoa, although it has been reported to play a role in the action of some drugs against parasitic nematodes [136]. Yeast also exhibit a loss of viability in response to a  $K^+$  channel-activating toxin which leads to excessive  $K^+$  flux [11]. Protozoan parasites within mammalian cells could be uniquely susceptible to drug-induced activation of parasite plasma membrane  $K^+$  channels, as the relatively high cytosolic  $K^+$  concentration of the host cell might allow  $K^+$  overload of the parasite, a scenario rarely encountered by mammalian cells due to the relatively low concentration of extracellular  $K^+$ . Well-defined  $K^+$  channel openers exist which have minimal negative effects in humans, and indeed some of these are used as anti-nematode drugs [136], muscle relaxants, antiepileptics and analgesics [137,138]. It is possible that the effectiveness of this strategy would be reduced in parasites that lower the cytosolic  $K^+$  concentration of the host cell, such as *Plasmodium* spp. [101–104].

The well-characterized pharmacology of  $K^+$  channels, together with sequence differences between human and parasite proteins, suggests that specific block of parasite  $K^+$  channel homologues might be an achievable target. For example, determinants of isoform-specific block of  $K_{Ca}2$  channels by drugs and toxins have been described in the outer pore region [78,139,140], where sequence differences exist between human and parasite homologues. This suggests avenues for development of drugs that specifically block parasite  $K_{Ca}$  channel homologues rather than human isoforms. Pharmacological blockers of other  $K^+$  channels also bind within the pore region and in some cases their binding affinity can be predicted with confidence from primary protein sequence alone. For example, TEA binds to the outer pore and a residue following the selectivity filter (GXGXXX) is a critical determinant of its binding affinity [75]. Channels with an aromatic residue at this position almost universally display a high-affinity interaction with TEA. Hence it is likely that the  $K^+$  channel homologues in *Plasmodium* spp. (XP\_001348796, XP\_001615733 and XP\_001348796) as well as the homologue in *E. histolytica* (XP\_655083), which all possess a phenylalanine or tyrosine residue at this locus, will display high affinity block by TEA. As discussed earlier, the lack of homologues of mammalian  $K^+$  channel auxiliary subunits in the protozoan parasites examined here also suggests a lack of diversity in native parasite  $K^+$  channel complexes and an opportunity for selective targeting of these parasite  $K^+$  channels by drugs.

This study presents the opportunity for cloning and functional characterization of  $K^+$  channels in pathogenic protozoa, and suggests that rational design of therapeutic strategies targeted against parasite  $K^+$  channels may be an attractive prospect. Future studies of parasite genomes and cellular signalling will lead to a deeper understanding of the presence and function of these channels in pathogenic parasites.

## Materials and Methods

### Genome analysis, sequence alignments and topology analysis

Analysis of genomes, sequence alignments and topology analysis were conducted as reported previously [2]. BLASTP and TBLASTN searches of protozoan genomes were carried out against the National Center for Biotechnology (NCBI) genomic protein databases. In multiple sequence alignments (ClustalW2) asterisks below the alignment indicate positions that have a single

fully conserved residue, while colons below the alignment indicates positions that have residues with highly similar properties (scoring  $>0.5$  in the Gonnet PAM 250 matrix, ClustalW2).

BLASTP analysis was carried out using the sequences of the following diverse human  $K^+$  channels (protein accession number in parentheses):  $K_v1.2$  (NP\_004965.1),  $K_v7.1$  (NP\_000209.2) and  $K_v11.1$  (hERG1) (Q12809.1);  $K_{ir}1.1$  (ROMK1) (NP\_000211.1),  $K_{ir}2.1$  (IRK1) (NP\_000882.1),  $K_{ir}3.1$  (GIRK1) (NP\_002230.1),  $K_{ir}4.1$  (P78508.1),  $K_{ir}5.1$  (Q9NPI9.1),  $K_{ir}6.1$  ( $K_{ATP1}$ ) (Q15842.1),  $K_{ir}6.2$  (NP\_000516.3) and  $K_{ir}7.1$  (CAA06878.1);  $K_{2p}1.1$  (TWIK1) (NP\_002236.1),  $K_{2p}2.1$  (TREK1) (NP\_001017425.2),  $K_{2p}3.1$  (TASK1) (NP\_002237.1),  $K_{2p}13.1$  (THIK1) (NP\_071337.2),  $K_{2p}16.1$  (TALK1) (NP\_001128577.1) and  $K_{2p}18.1$  (TRESK2) (NP\_862823.1);  $K_{Ca}1.1$  (BK) (NP\_001154824.1),  $K_{Ca}2.1$  (SK1) (NP\_002239.2),  $K_{Ca}2.2$  (SK2) (NP\_067627),  $K_{Ca}3.1$  (IK/SK4) (NP\_002241.1) and  $K_{Ca}4.1$  (SLACK/ $K_{Na}$ ) (NP\_065873.2). Other  $K^+$  channel sequences were also used to search for parasite homologues, including: *Plasmodium falciparum* PfKch1 (XP\_001350669.2) and PfKch2 (XP\_001348796.2) [15], bacterial  $KcsA$  (P0A334), bacterial cyclic nucleotide-gated MlotiK1 (Q98GN8.1), archaeal depolarization-activated  $K_v$ AP (Q9YDF8.1), archaeal hyperpolarization-activated MVP (Q57603.1), archaeal  $Ca^{2+}$ -activated MthK (O27564.1), and fungal TOK1 (CAA89386.1). Plant  $K^+$  channel sequences were also used, including: the vacuolar outwardly rectifying, calcium-regulated vacuolar two-pore TPK1 channel (NP\_200374.1); vacuolar KCO3 (NP\_001190480.1); the pollen plasma membrane TPK4 (NP\_171752.1), the inward rectifier KAT1 (NP\_199436.1), the outward rectifier SKOR (pore region of NP\_186934.1, residues 271–340 to avoid ankyrin hits), and AKT1 (NP\_180233.1). Sequences of human CNGA1 (EAW93049; full-length, and TMD residues 200–420), and CNGB1 (NP\_00121288), as well as human HCN2 (NP\_001185.3; full-length, and TMD residues 200–470) were also used to search for parasite homologues. In addition, the sequence of a novel putative adenylyl cyclase/ $K^+$  channel fusion protein in *P. falciparum* (PfAC1 or PfAC $\alpha$ ; XP\_001348216) was used to search for homologues in other parasites. Sequences of  $K^+$  channel auxiliary subunits that were used to search for parasite homologues include: human KCNE1 (NP\_001121142.1), human  $K_v\beta 1$  (NP\_751892.1), human KChIP1 (NP\_001030009), human  $BK\beta$  (NP\_004128.1), human SUR1 (NP\_000343) and human SUR2A (NP\_005682). The sequences of both the isolated nucleotide-binding domains of SUR1 and SUR2A and the sequences outside these nucleotide-binding regions were also used to search for parasite homologues. Results of BLASTP analysis were confirmed using TBLASTN analysis in all cases. Default BLAST parameters for assessing statistical significance and for filtering were used in all cases (*ie.* an Expect threshold of 10, and SEG filtering).

Several procedures ensured that hits were probable  $K^+$  channel homologues. Firstly, the occurrence of multiple putative TMDs was confirmed using TOPCONS [141]. Secondly, reciprocal BLASTP searches (non-redundant protein database at NCBI) were undertaken, using identified parasite hits as bait, and only proteins that gave the original mammalian protein family as hits were analyzed further. Thirdly, conserved domains were identified using the Conserved Domains Database (NCBI). Lastly, only hits with regions of sequence similarity that encompassed the selectivity filter sequence of the  $K^+$  channel subunit used as bait were acknowledged.  $K_{CNG}$  channel parasite homologues were identified as proteins showing sequence similarity in both the pore and the CNBD regions. Where a hit showed similarity to more than one human  $K^+$  channel, the parasite protein was designated as a homologue of the human channel to which it showed greatest sequence similarity (ClustalW2) and which contained similar putative functional domains.

For phylogenetic analysis, multiple sequence alignments were constructed with MUSCLE v3.7 using default parameters. After use of GBLOCKS at low stringency to remove regions of low confidence, and removal of gaps, Maximum Likelihood analysis was undertaken using PhyML v3.0 (WAG substitution model; 4 substitution rate categories; default estimated gamma distribution parameters; default estimated proportions of invariable sites; 100 bootstrapped data sets). The phylogenetic tree is shown using TreeDyn (v1.98.3). MUSCLE, GBLOCKS, PhyML and TreeDyn are all functions of Phylogeny.fr (<http://www.phylogeny.fr/>) [142].

## References

- Wiser MF (2011) Protozoa and Human disease. New York: Garland Science. 218 p.
- Prole DL, Taylor CW (2011) Identification of intracellular and plasma membrane calcium channel homologues in pathogenic parasites. *PLoS One* 6(10): e26218.
- Castillo E, Dea-Ayuela MA, Bolás-Fernández F, Rangel M, González-Rosende ME (2010) The kinetoplastid chemotherapy revisited: current drugs, recent advances and future perspectives. *Curr Med Chem* 17: 4027–4051.
- Monzote L, Siddiq A (2011) Drug development to protozoan diseases. *Open Med Chem J* 5: 1–3.
- Petersen I, Eastman R, Lanzer M (2011) Drug-resistant malaria: molecular mechanisms and implications for public health. *FEBS Lett* 585: 1551–1562.
- Miller C (2000) An overview of the potassium channel family. *Genome Biol* 1(4): REVIEWS0004.
- Kuo MM, Haynes WJ, Loukin SH, Kung C, Saimi Y (2005) Prokaryotic K<sup>+</sup> channels: from crystal structures to diversity. *FEMS Microbiol Rev* 29: 961–985.
- Ward JM, Maser P, Schroeder JI (2009) Plant ion channels: gene families, physiology, and functional genomics analyses. *Annu Rev Physiol* 71: 59–82.
- Gajdanowicz P, Michard E, Sandmann M, Rocha M, Corrêa LG, et al. (2011) Potassium (K<sup>+</sup>) gradients serve as a mobile energy source in plant vascular tissues. *Proc Natl Acad Sci U S A* 108: 864–869.
- Roberts SK (2003) TOK homologue in *Neurospora crassa*: first cloning and functional characterization of an ion channel in a filamentous fungus. *Eukaryot Cell* 2: 181–190.
- Ahmed A, Sesti F, Ilan N, Shih TM, Sturley SL, et al. (1999) A molecular target for viral killer toxin: TOK1 potassium channels. *Cell* 99: 283–291.
- Loukin SH, Kuo MM, Zhou XL, Haynes WJ, Kung C, et al. (2005) Microbial K<sup>+</sup> channels. *J Gen Physiol* 125: 521–527.
- Haynes WJ, Ling KY, Saimi Y, Kung C (2003) PAK paradox: *Paramecium* appears to have more K<sup>+</sup>-channel genes than humans. *Eukaryot Cell* 2: 737–745.
- Ellekvist P, Ricke CH, Litman T, Salanti A, Colding H, et al. (2004) Molecular cloning of a K<sup>+</sup> channel from the malaria parasite *Plasmodium falciparum*. *Biochem Biophys Res Commun* 318: 477–484.
- Waller KL, McBride SM, Kim K, McDonald TV (2008) Characterization of two putative potassium channels in *Plasmodium falciparum*. *Malar J* 7: 19.
- Ellekvist P, Maciel J, Mlambo G, Ricke CH, Colding H, et al. (2008) Critical role of a K<sup>+</sup> channel in *Plasmodium berghei* transmission revealed by targeted gene disruption. *Proc Natl Acad Sci U S A* 105: 6398–6402.
- Jimenez V, Henriquez M, Galanti N, Riquelme G (2011) Electrophysiological characterization of potassium conductive pathways in *Trypanosoma cruzi*. *J Cell Biochem* 112: 1093–1102.
- Goldstein SA, Wang KW, Ilan N, Pausch MH (1998) Sequence and function of the two P domain potassium channels: implications of an emerging superfamily. *J Mol Med (Berl)* 76: 13–20.
- Mangubat EZ, Tseng TT, Jakobsson E (2003) Phylogenetic analyses of potassium channel auxiliary subunits. *J Mol Microbiol Biotechnol* 5: 216–224.
- Pongs O, Schwarz JR (2010) Ancillary subunits associated with voltage-dependent K<sup>+</sup> channels. *Physiol Rev* 90: 755–796.
- Campomanes CR, Carroll KI, Manganas LN, Hershberger ME, Gong B, et al. (2002) K<sub>v</sub>β subunit oxidoreductase activity and K<sub>v</sub>1 potassium channel trafficking. *J Biol Chem* 277: 8298–8305.
- Rodrigo GC, Standen NB (2005) ATP-sensitive potassium channels. *Curr Pharm Des* 11: 1915–1940.
- Jarmuszkiewicz W, Matkovic K, Koszela-Piotrowska I (2010) Potassium channels in the mitochondria of unicellular eukaryotes and plants. *FEBS Lett* 584: 2057–2062.
- Szewczyk A, Jarmuszkiewicz W, Kunz WS (2009) Mitochondrial potassium channels. *IUBMB Life* 61: 134–143.
- Mazzanti M, DeFelice LJ, Cohn J, Malter H (1990) Ion channels in the nuclear envelope. *Nature* 343: 764–767.
- Quesada I, Rovira JM, Martín F, Roche E, Nadal A, et al. (2002) Nuclear K<sub>ATP</sub> channels trigger nuclear Ca<sup>2+</sup> transients that modulate nuclear function. *Proc Natl Acad Sci U S A* 99: 9544–9549.
- Yamashita M, Sugioka M, Ogawa Y (2006) Voltage- and Ca<sup>2+</sup>-activated potassium channels in Ca<sup>2+</sup> store control Ca<sup>2+</sup> release. *FEBS J* 273: 3585–3597.
- Chen Y, Sánchez A, Rubio ME, Kohl T, Pardo LA, et al. (2011) Functional K<sub>v</sub>10.1 channels localize to the inner nuclear membrane. *PLoS One* 6(5): e19257.
- Bao L, Hadjiolova K, Coetzee WA, Rindler MJ (2011) Endosomal K<sub>ATP</sub> channels as a reservoir after myocardial ischemia: a role for SUR2 subunits. *Am J Physiol Heart Circ Physiol* 300: H262–70.
- Ng KE, Schwarzer S, Duchon MR, Tinker A (2010) The intracellular localization and function of the ATP-sensitive K<sup>+</sup> channel subunit K<sub>v</sub>6.1. *J Membr Biol* 234: 137–147.
- Geng X, Li L, Watkins S, Robbins PD, Drain P (2003) The insulin secretory granule is the major site of K<sub>ATP</sub> channels of the endocrine pancreas. *Diabetes* 52: 767–776.
- Kelly ML, Abu-Hamdah R, Jeremic A, Cho SJ, Ilie AE, et al. (2005) Patch clamped single pancreatic zymogen granules: direct measurements of ion channel activities at the granule membrane. *Pancreatology* 5: 443–449.
- Ward JM, Schroeder JI (1994) Calcium-Activated K<sup>+</sup> Channels and Calcium-Induced Calcium Release by Slow Vacuolar Ion Channels in Guard Cell Vacuoles Implicated in the Control of Stomatal Closure. *Plant Cell* 6: 669–683.
- Isayenkov S, Isner JC, Maathuis FJ (2011) Rice two-pore K<sup>+</sup> channels are expressed in different types of vacuoles. *Plant Cell* 23: 756–768.
- Gobert A, Isayenkov S, Voelker C, Czempinski K, Maathuis FJ (2007) The two-pore channel TPK1 gene encodes the vacuolar K<sup>+</sup> conductance and plays a role in K<sup>+</sup> homeostasis. *Proc Natl Acad Sci U S A* 104: 10726–10731.
- Remillard CV, Yuan JX (2004) Activation of K<sup>+</sup> channels: an essential pathway in programmed cell death. *Am J Physiol Lung Cell Mol Physiol* 286: L49–67.
- Waller KL, Kim K, McDonald TV (2008) *Plasmodium falciparum*: growth response to potassium channel blocking compounds. *Exp Parasitol* 120: 280–285.
- Gardner MJ, Hall N, Fung E, White O, Berriman M, et al. (2002) Genome sequence of the human malaria parasite *Plasmodium falciparum*. *Nature* 419: 498–511.
- Gardner MJ, Shallom SJ, Carlton JM, Salzberg SL, Nene V (2002) Sequence of *Plasmodium falciparum* chromosomes 2, 10, 11 and 14. *Nature* 419: 531–534.
- Carlton JM, Adams JH, Silva JC, Bidwell SL, Lorenzi H, et al. (2008) Comparative genomics of the neglected human malaria parasite *Plasmodium vivax*. *Nature* 455: 757–763.
- Lorenzi H, Caler E, Galinsky K, Schobel S, Brunk BP, et al. (2008) Annotation of *Toxoplasma gondii* ME49. EMBL/GenBank/DBJ databases.
- Xu P, Widmer G, Wang Y, Ozaki LS, Alves JM, et al. (2004) The genome of *Cryptosporidium hominis*. *Nature* 431: 1107–1112.
- Bankier AT, Spriggs HF, Fartmann B, Konfortov BA, Madera M, et al. (2003) Integrated mapping, chromosomal sequencing and sequence analysis of *Cryptosporidium parvum*. *Genome Res* 13: 1787–1799.
- Abrahamsen MS, Templeton TJ, Enomoto S, Abrahante JE, Zhu G, et al. (2004) Complete genome sequence of the apicomplexan, *Cryptosporidium parvum*. *Science* 304: 441–445.
- Brayton KA, Lau AO, Herndon DR, Hannick L, Kappmeyer LS, et al. (2007) Genome sequence of *Babesia bovis* and comparative analysis of apicomplexan hemoprotozoa. *PLoS Pathog* 3(10): 1401–1413.
- Laurentino EC, Ruiz JC, Fazelinia G, Myler PJ, Degraeve W, et al. (2004) A survey of *Leishmania braziliensis* genome by shotgun sequencing. *Mol Biochem Parasitol* 137: 81–86.
- Ivens AC, Peacock CS, Worthey EA, Murphy L, Aggarwal G, et al. (2005) The genome of the kinetoplastid parasite, *Leishmania major*. *Science* 309: 436–442.
- Peacock CS, Seeger K, Harris D, Murphy L, Ruiz JC, et al. (2007) Comparative genomic analysis of three *Leishmania* species that cause diverse human disease. *Nat Genet* 39: 839–847.
- Berriman M, Ghedin E, Hertz-Fowler C, Blandin G, Renauld H, et al. (2005) The genome of the African trypanosome *Trypanosoma brucei*. *Science* 309: 416–422.
- El-Sayed NM, Myler PJ, Blandin G, Berriman M, Crabtree J, et al. (2005) Comparative genomics of trypanosomatid parasitic protozoa. *Science* 309: 404–409.

51. Loftus B, Anderson I, Davies R, Alsmark UC, Samuelson J, et al. (2005) The genome of the protist parasite *Entamoeba histolytica*. *Nature* 433: 865–868.
52. Franzén O, Jernström-Hultqvist J, Castro E, Sherwood E, Ankarklev J, et al. (2009) Draft genome sequencing of *Giardia intestinalis* assemblage B isolate GS: is human giardiasis caused by two different species? *PLoS Pathog* 5(8): e1000560.
53. Morrison HG, McArthur AG, Gillin FD, Aley SB, Adam RD, et al. (2007) Genomic minimalism in the early diverging intestinal parasite *Giardia lamblia*. *Science* 317: 1921–1926.
54. McArthur AG, Morrison HG, Nixon JE, Passamaneck NQ, Kim U, et al. (2000) The *Giardia* genome project database. *FEMS Microbiol Lett* 189: 271–273.
55. Jernström-Hultqvist J, Franzén O, Ankarklev J, Xu F, Nohýnková E, et al. (2010) Genome analysis and comparative genomics of a *Giardia intestinalis* assemblage E isolate. *BMC Genomics* 11: 543.
56. Carlton JM, Hirt RP, Silva JC, Delcher AL, Schatz M, et al. (2007) Draft genome sequence of the sexually transmitted pathogen *Trichomonas vaginalis*. *Science* 315: 207–212.
57. Heginbotham L, Lu Z, Abramson T, MacKinnon R (1994) Mutations in the K<sup>+</sup> channel signature sequence. *Biophys J* 66: 1061–1067.
58. Shealy RT, Murphy AD, Ramarathnam R, Jakobsson E, Subramaniam S (2003) Sequence-function analysis of the K<sup>+</sup>-selective family of ion channels using a comprehensive alignment and the KcsA channel structure. *Biophys J* 84: 2929–2942.
59. So I, Ashmole I, Davies NW, Sutcliffe MJ, Stanfield PR (2001) The K<sup>+</sup> channel signature sequence of murine K<sub>v</sub>2.1: mutations that affect microscopic gating but not ionic selectivity. *J Physiol* 531: 37–50.
60. Yellen G (2002) The voltage-gated potassium channels and their relatives. *Nature* 419: 35–42.
61. Salkoff L, Butler A, Ferreira G, Santi C, Wei A (2006) High-conductance potassium channels of the SLO family. *Nat Rev Neurosci* 7: 921–931.
62. Wu Y, Yang Y, Ye S, Jiang Y (2010) Structure of the gating ring from the human large-conductance Ca<sup>2+</sup>-gated K<sup>+</sup> channel. *Nature* 466: 393–397.
63. Yuan P, Leonetti MD, Pico AR, Hsiung Y, MacKinnon R (2010) Structure of the human BK channel Ca<sup>2+</sup>-activation apparatus at 3.0 Å resolution. *Science* 329: 182–186.
64. Schumacher MA, Crum M, Miller MC (2004) Crystal structures of apocalmodulin and an apocalmodulin/SK potassium channel gating domain complex. *Structure* 12: 849–860.
65. Zagotta WN, Olivier NB, Black KD, Young EC, Olson R, et al. (2003) Structural basis for modulation and agonist specificity of HCN pacemaker channels. *Nature* 425: 200–205.
66. Kirsch GE, Pascual JM, Shieh CC (1995) Functional role of a conserved aspartate in the external mouth of voltage-gated potassium channels. *Biophys J* 68: 1804–1813.
67. Haug T, Olcese R, Toro L, Stefani E (2004) Regulation of K<sup>+</sup> flow by a ring of negative charges in the outer pore of BK<sub>Ca</sub> channels. Part I: Aspartate 292 modulates K<sup>+</sup> conduction by external surface charge effect. *J Gen Physiol* 124: 173–184.
68. Chapman ML, Blanke ML, Krovetz HS, VanDongen AM (2006) Allosteric effects of external K<sup>+</sup> ions mediated by the aspartate of the GYGD signature sequence in the K<sub>v</sub>2.1 K<sup>+</sup> channel. *Pflugers Arch* 451: 776–792.
69. Cordero-Morales JF, Cuello LG, Perozo E (2006) Voltage-dependent gating at the KcsA selectivity filter. *Nat Struct Mol Biol* 13: 319–322.
70. Cordero-Morales JF, Cuello LG, Zhao Y, Jogini V, Cortes DM, et al. (2006) Molecular determinants of gating at the potassium-channel selectivity filter. *Nat Struct Mol Biol* 13: 311–318.
71. Hibino H, Inanobe A, Furutani K, Murakami S, Findlay I, et al. (2010) Inwardly rectifying potassium channels: their structure, function, and physiological roles. *Physiol Rev* 90: 291–366.
72. Kuo MM, Saimi Y, Kung C (2003) Gain-of-function mutations indicate that *Escherichia coli* Kch forms a functional K<sup>+</sup> conduit *in vivo*. *EMBO J* 22: 4049–4058.
73. MacKinnon R, Yellen G (1990) Mutations affecting TEA blockage and ion permeation in voltage-activated K<sup>+</sup> channels. *Science* 250: 276–279.
74. MacKinnon R, Heginbotham L, Abramson T (1990) Mapping the receptor site for charybdotoxin, a pore-blocking potassium channel inhibitor. *Neuron* 5: 767–771.
75. Heginbotham L, MacKinnon R (1992) The aromatic binding site for tetraethylammonium ion on potassium channels. *Neuron* 8: 483–491.
76. Chatelain FC, Gazzarrini S, Fujiwara Y, Arrigoni C, Domigan C, et al. (2009) Selection of inhibitor-resistant viral potassium channels identifies a selectivity filter site that affects barium and amantadine block. *PLoS One* 4(10): e7496.
77. Wulff H, Castle NA, Pardo LA (2009) Voltage-gated potassium channels as therapeutic targets. *Nat Rev Drug Discov* 8: 982–1001.
78. Weatherall KL, Goodchild SJ, Jane DE, Marrion NV (2010) Small conductance calcium-activated potassium channels: from structure to function. *Prog Neurobiol* 91: 242–55.
79. Yellen G, Jurman ME, Abramson T, MacKinnon R (1991) Mutations affecting internal TEA blockage identify the probable pore-forming region of a K<sup>+</sup> channel. *Science* 251: 939–42.
80. Kirsch GE, Shieh CC, Drewe JA, Vener DF, Brown AM (1993) Segmental exchanges define 4-aminopyridine binding and the inner mouth of K<sup>+</sup> pores. *Neuron* 11: 503–512.
81. Sanguinetti MC, Chen J, Fernandez D, Kamiya K, Mitcheson J, et al. (2005) Physicochemical basis for binding and voltage-dependent block of hERG channels by structurally diverse drugs. *Novartis Found Symp* 266: 159–166.
82. Mitcheson J, Perry M, Stansfeld P, Sanguinetti MC, Witchel H, et al. (2005) Structural determinants for high-affinity block of hERG potassium channels. *Novartis Found Symp* 266: 136–150.
83. Decher N, Streit AK, Rapedius M, Netter MF, Marzian S, et al. (2010) RNA editing modulates the binding of drugs and highly unsaturated fatty acids to the open pore of K<sub>v</sub> potassium channels. *EMBO J* 29: 2101–2113.
84. Madeja M, Steffen W, Mesic I, Garic B, Zhorov BS (2010) Overlapping binding sites of structurally different antiarrhythmic flecainide and propafenone in the subunit interface of potassium channel K<sub>v</sub>2.1. *J Biol Chem* 285: 33898–33905.
85. Zhou Y, Tang QY, Xia XM, Lingle CJ (2010) Glycine311, a determinant of paxilline block in BK channels: a novel bend in the BK S6 helix. *J Gen Physiol* 135: 481–494.
86. Lee US, Cui J (2010) BK channel activation: structural and functional insights. *Trends Neurosci* 33: 415–423.
87. Nagamune K, Moreno SN, Chini EN, Sibley LD (2008) Calcium regulation and signaling in apicomplexan parasites. *Subcell Biochem* 47: 70–81.
88. Billker O, Lourido S, Sibley LD (2009) Calcium-dependent signaling and kinases in apicomplexan parasites. *Cell Host Microbe* 5: 612–22.
89. Weatherall KL, Seutin V, Liégeois JF, Marrion NV (2011) Crucial role of a shared extracellular loop in apamin sensitivity and maintenance of pore shape of small-conductance calcium-activated potassium (SK) channels. *Proc Natl Acad Sci U S A* 108: 18494–18499.
90. Zhou L, Siegelbaum SA (2007) Gating of HCN channels by cyclic nucleotides: residue contacts that underlie ligand binding, selectivity, and efficacy. *Structure* 15: 655–70.
91. Ketchum KA, Joiner WJ, Sellers AJ, Kaczmarek LK, Goldstein SA (1995) A new family of outwardly rectifying potassium channel proteins with two pore domains in tandem. *Nature* 376: 690–5.
92. Aittoniemi J, Fotinou C, Craig TJ, de Wet H, Proks P, et al. (2009) SUR1: a unique ATP-binding cassette protein that functions as an ion channel regulator. *Philos Trans R Soc Lond B Biol Sci* 364: 257–67.
93. Akrouh A, Halcomb SE, Nichols CG, Sala-Rabanal M (2009) Molecular biology of K<sub>ATP</sub> channels and implications for health and disease. *IUBMB Life* 61: 971–8.
94. Aurrecochea C, Heiges M, Wang H, Wang Z, Fischer S, et al. (2007) ApiDB: integrated resources for the apicomplexan bioinformatics resource center. *Nucleic Acids Res* 35: D427–30.
95. Hirsch RE, Lewis BD, Spalding EP, Sussman MR (1998) A role for the AKT1 potassium channel in plant nutrition. *Science* 280: 918–21.
96. Fairman C, Zhou X, Kung C (1999) Potassium uptake through the TOK1 K<sup>+</sup> channel in the budding yeast. *J Membr Biol* 168: 149–57.
97. Latorre R, Olcese R, Basso C, Gonzalez C, Munoz F, et al. (2003) Molecular coupling between voltage sensor and pore opening in the *Arabidopsis* inward rectifier K<sup>+</sup> channel KAT1. *J Gen Physiol* 122: 459–69.
98. Schachtman DP, Schroeder JI, Lucas WJ, Anderson JA, Gaber RF (1992) Expression of an inward-rectifying potassium channel by the *Arabidopsis* KAT1 cDNA. *Science* 258: 1634–8.
99. Sesti F, Rajan S, Gonzalez-Coloso R, Nikolaeva N, Goldstein SA (2003) Hyperpolarization moves S4 sensors inward to open MVP, a methanococcal voltage-gated potassium channel. *Nat Neurosci* 6: 353–61.
100. Hille B (1992) *Ionic Channels of Excitable Membranes*. Second Edition. Sunderland/Massachusetts: Sinauer Associates Inc. 15 p.
101. Dunn MJ (1969) Alterations of red blood cell sodium transport during malarial infection. *J Clin Invest* 48: 674–84.
102. Ginsburg H, Handeli S, Friedman S, Gorodetsky R, Krugliak M (1986) Effects of red blood cell potassium and hypertonicity on the growth of *Plasmodium falciparum* in culture. *Z Parasitenkd* 72: 185–99.
103. Lee P, Ye Z, Van Dyke K, Kirk RG (1988) X-ray microanalysis of *Plasmodium falciparum* and infected red blood cells: effects of quinghaosu and chloroquine on potassium, sodium, and phosphorus composition. *Am J Trop Med Hyg* 39: 157–65.
104. Overman RR (1948) Reversible cellular permeability alterations in disease. *In vivo* studies on sodium, potassium and chloride concentrations in erythrocytes of the malarious monkey. *Am J Physiol* 152: 113–21.
105. Mannikko R, Elinder F, Larsson HP (2002) Voltage-sensing mechanism is conserved among ion channels gated by opposite voltages. *Nature* 419: 837–41.
106. Allen RJ, Kirk K (2004) The membrane potential of the intraerythrocytic malaria parasite *Plasmodium falciparum*. *J Biol Chem* 279: 11264–72.
107. Nolan DP, Voorheis HP (2000) Factors that determine the plasma-membrane potential in bloodstream forms of *Trypanosoma brucei*. *Eur J Biochem* 267: 4615–23.
108. Glaser TA, Utz GL, Mukkada AJ (1992) The plasma membrane electrical gradient (membrane potential) in *Leishmania donovani* promastigotes and amastigotes. *Mol Biochem Parasitol* 51: 9–15.
109. Nagamune K, Sibley LD (2006) Comparative genomic and phylogenetic analyses of calcium ATPases and calcium-regulated proteins in the apicomplexa. *Mol Biol Evol* 23: 1613–27.
110. Matsumoto Y, Perry G, Scheibel LW, Aikawa M (1987) Role of calmodulin in *Plasmodium falciparum*: implications for erythrocyte invasion by the merozoite. *Eur J Cell Biol* 45: 36–43.

111. Mazumder S, Mukherjee T, Ghosh J, Ray M, Bhaduri A (1992) Allosteric modulation of *Leishmania donovani* plasma membrane  $\text{Ca}^{2+}$ -ATPase by endogenous calmodulin. *J Biol Chem* 267: 18440–6.
112. Orfa Rojas M, Wasserman M (1995) Stage-specific expression of the calmodulin gene in *Plasmodium falciparum*. *J Biochem* 118: 1118–23.
113. Bernal RM, Tovar R, Santos JI, Muñoz ML (1998) Possible role of calmodulin in excystation of *Giardia lamblia*. *Parasitol Res* 84: 687–93.
114. Yakubu MA, Majumder S, Kierszenbaum F (1994) Changes in *Trypanosoma cruzi* infectivity by treatments that affect calcium ion levels. *Mol Biochem Parasitol* 66: 119–25.
115. Makioka A, Kumagai M, Kobayashi S, Takeuchi T (2002) Possible role of calcium ions, calcium channels and calmodulin in excystation and metacystic development of *Entamoeba invadens*. *Parasitol Res* 88: 837–43.
116. Song HO, Ahn MH, Ryu JS, Min DY, Joo KH, et al. (2004) Influence of calcium ion on host cell invasion and intracellular replication by *Toxoplasma gondii*. *Korean J Parasitol* 42: 185–93.
117. Pezzella N, Bouchot A, Bonhomme A, Pingret L, Klein C, et al. (1997) Involvement of calcium and calmodulin in *Toxoplasma gondii* tachyzoite invasion. *Eur J Cell Biol* 74: 92–101.
118. Yusifov T, Savalli N, Gandhi CS, Otolia M, Olcese R (2008) The RCK2 domain of the human  $\text{BK}_{\text{Ca}}$  channel is a calcium sensor. *Proc Natl Acad Sci U S A* 105: 376–81.
119. Pau VP, Smith FJ, Taylor AB, Parfenova LV, Samakai E, et al. (2011) Structure and function of multiple  $\text{Ca}^{2+}$ -binding sites in a  $\text{K}^+$  channel regulator of  $\text{K}^+$  conductance (RCK) domain. *Proc Natl Acad Sci U S A* 108: 17684–9.
120. Yang H, Hu L, Shi J, Delaloye K, Horrigan FT, et al. (2007)  $\text{Mg}^{2+}$  mediates interaction between the voltage sensor and cytosolic domain to activate BK channels. *Proc Natl Acad Sci U S A* 104: 18270–5.
121. Cui J, Das S, Smith TF, Samuelson J (2010) *Trichomonas* transmembrane cyclases result from massive gene duplication and concomitant development of pseudogenes. *PLoS Negl Trop Dis* 4: e782.
122. Schultz JE, Klumpp S, Benz R, Schürhoff-Goeters WJ, Schmid A (1992) Regulation of adenylyl cyclase from *Paramecium* by an intrinsic potassium conductance. *Science* 255: 600–3.
123. Muhia DK, Swales CA, Eckstein-Ludwig U, Saran S, Polley SD, et al. (2003) Multiple splice variants encode a novel adenylyl cyclase of possible plastid origin expressed in the sexual stage of the malaria parasite *Plasmodium falciparum*. *J Biol Chem* 278: 22014–22.
124. Weber JH, Vishnyakov A, Hambach K, Schultz A, Schultz JE, et al. (2004) Adenylyl cyclases from *Plasmodium*, *Paramecium* and *Tetrahymena* are novel ion channel/enzyme fusion proteins. *Cell Signal* 16: 115–25.
125. Ono T, Cabrita-Santos L, Leitao R, Bettiol E, Purcell LA, et al. (2008) Adenylyl cyclase alpha and cAMP signaling mediate *Plasmodium* sporozoite apical regulated exocytosis and hepatocyte infection. *PLoS Pathog* 4: e1000008.
126. Burg ED, Remillard CV, Yuan JX (2006)  $\text{K}^+$  channels in apoptosis. *J Membr Biol* 209: 3–20.
127. Leung YM (2010) Voltage-gated  $\text{K}^+$  channel modulators as neuroprotective agents. *Life Sci* 86: 775–80.
128. Rodriguez-Menchaca AA, Navarro-Polanco RA, Ferrer-Villada T, Rupp J, Sachse FB, et al. (2008) The molecular basis of chloroquine block of the inward rectifier  $\text{K}_{\text{ir}}2.1$  channel. *Proc Natl Acad Sci U S A* 105: 1364–8.
129. Foley M, Tilley L (1998) Quinoline antimalarials: mechanisms of action and resistance and prospects for new agents. *Pharmacol Ther* 79: 55–87.
130. de Boer TP, Nalos L, Stary A, Kok B, Houtman MJ, et al. (2010) The antiprotozoal drug pentamidine blocks  $\text{K}_{\text{IR}}2.x$ -mediated inward rectifier current by entering the cytoplasmic pore region of the channel. *Br J Pharmacol* 159: 1532–41.
131. Serrano-Martin X, Payares G, Mendoza-Leon A (2006) Glibenclamide, a blocker of  $\text{K}^+$  channels, shows antileishmanial activity in experimental murine cutaneous leishmaniasis. *Antimicrob Agents Chemother* 50: 4214–6.
132. Serrano-Martin X, García-Marchan Y, Fernandez A, Rodriguez N, Rojas H, et al. (2009) Amiodarone destabilizes intracellular  $\text{Ca}^{2+}$  homeostasis and biosynthesis of sterols in *Leishmania mexicana*. *Antimicrob Agents Chemother* 53: 1403–10.
133. Evans SG, Havlik I (1993) *Plasmodium falciparum*: effects of amantadine, an antiviral, on chloroquine-resistant and -sensitive parasites *in vitro* and its influence on chloroquine activity. *Biochem Pharmacol* 45: 1168–70.
134. Khawaled R, Bruening-Wright A, Adelman JP, Maylie J (1999) Bicuculline block of small-conductance calcium-activated potassium channels. *Pflügers Arch* 438: 314–21.
135. Rossokhin A, Teodorescu G, Grissmer S, Zhorov BS (2006) Interaction of *d*-tubocurarine with potassium channels: molecular modeling and ligand binding. *Mol Pharmacol* 69: 1356–65.
136. Martin RJ, Buxton SK, Neveu C, Charvet CL, Robertson AP (2011) Emodesipide and SL0-1 potassium channels: A review. *Exp Parasitol*. In press.
137. Devulder J (2010) Flupirtine in pain management: pharmacological properties and clinical use. *CNS Drugs* 24: 867–81.
138. Czuczwar P, Wojtak A, Ciozek-Czuczwar A, Parada-Turska J, Maciejewski R, et al. (2010) Retigabine: the newer potential antiepileptic drug. *Pharmacol Rep* 62: 211–9.
139. Ishii TM, Maylie J, Adelman JP (1997) Determinants of apamin and *d*-tubocurarine block in SK potassium channels. *J Biol Chem* 272: 23195–200.
140. Lamy C, Goodchild SJ, Weatherall KL, Jane DE, Liégeois JF, et al. (2010) Allosteric block of  $\text{K}_{\text{Ca}}2$  channels by apamin. *J Biol Chem* 285: 27067–77.
141. Bernsel A, Viklund H, Hennerdal A, Elofsson A (2009) TOPCONS: consensus prediction of membrane protein topology. *Nucleic Acids Res* 37: W465–8.
142. Dereeper A, Guignon V, Blanc G, Audic S, Buffet S, et al. (2008) Phylogeny.fr: robust phylogenetic analysis for the non-specialist. *Nucleic Acids Res* 36: W465–9.
143. Yap KL, Kim J, Truong K, Sherman M, Yuan T, et al. (2000) Calmodulin target database. *J Struct Funct Genomics* 1: 8–14.
144. Adamczak R, Porollo A, Meller J (2005) Combining prediction of secondary structure and solvent accessibility in proteins. *Proteins* 59: 467–75.
145. Wallner M, Meera P, Toro L (1996) Determinant for  $\beta$ -subunit regulation in high-conductance voltage-activated and  $\text{Ca}^{2+}$ -sensitive  $\text{K}^+$  channels: an additional transmembrane region at the N terminus. *Proc Natl Acad Sci U S A* 93: 14922–7.
146. Meera P, Wallner M, Song M, Toro L (1997) Large conductance voltage- and calcium-dependent  $\text{K}^+$  channel, a distinct member of voltage-dependent ion channels with seven N-terminal transmembrane segments (S0–S6), an extracellular N terminus, and an intracellular (S9–S10) C terminus. *Proc Natl Acad Sci U S A* 94: 14066–71.
147. Zhou Y, Morais-Cabral JH, Kaufman A, MacKinnon R (2001) Chemistry of ion coordination and hydration revealed by a  $\text{K}^+$  channel-Fab complex at 2.0 Å resolution. *Nature* 414: 43–8.
148. Flynn GE, Black KD, Islas LD, Sankaran B, Zagotta WN (2007) Structure and rearrangements in the carboxy-terminal region of SpIH channels. *Structure* 15: 671–82.
149. Wallace AC, Laskowski RA, Thornton JM (1995) LIGPLOT: a program to generate schematic diagrams of protein-ligand interactions. *Protein Eng* 8: 127–34.