

Original Article
Occupation &
Environmental Medicine



Job Stress-attributable Burden of Disease in Korea

Kyungjoon Lee ¹ and Inah Kim ^{2,3}

¹Department of Occupational and Environmental Medicine, Hanyang University Hospital, Seoul, Korea

²Department of Occupational and Environmental Medicine, College of Medicine, Hanyang University, Seoul, Korea

³Department of Occupational and Environmental Health, Graduate School of Public Health, Hanyang University, Seoul, Korea



Received: Apr 18, 2017

Accepted: Jan 16, 2018

Address for Correspondence:

Inah Kim, MD, MPH, PhD

Department of Occupational and Environmental Medicine, College of Medicine, Hanyang University, 222-1 Wangsimni-ro, Seongdong-gu, Seoul 04763, Korea.
E-mail: inahkim@hanyang.ac.kr

© 2018 The Korean Academy of Medical Sciences.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

Kyungjoon Lee

<https://orcid.org/0000-0001-8436-4212>

Inah Kim

<https://orcid.org/0000-0003-3568-4484>

Funding

This study was supported by Ministry of Environment as "The Environmental Health Action program (2014001350001)."

Disclosure

The authors have no potential conflicts of interest to disclose.

ABSTRACT

Background: As job stress is associated with various diseases and psychiatric conditions, we aimed to estimate the job stress-attributable burden of disease in Korea based on the concept of disability-adjusted life years (DALY).

Methods: We selected ischemic heart disease (IHD), stroke, major depressive disorder (MDD), and suicide as health outcomes from job stress, because of the ease of access to data estimating burdens and of important meaning of them in Korean occupational background.

Results: Our findings demonstrated that approximately 21% of Korean workers were exposed to high job strain, which was attributable for approximately 6.7% of IHDs, 6.9% of strokes, 13.6% of MDDs, and 4% of suicides. In terms of job stress-attributable DALY, the burdens of disease per 100,000 people were 38 for IHD, 72 for stroke, 168 for MDDs, and 44 for suicides.

Conclusion: The present findings suggested that one-fifth of Korean workers were suffering from high job strain. Although the figures may not be accurate due to several assumptions, job stress is an important risk factor for health in working environment in Korea.

Keywords: Job Stress; Ischemic Heart Disease; Stroke; Major Depressive Disorder; Burden of Disease; DALY

INTRODUCTION

Workers must adapt to rapidly changing working conditions and psychosocial factors in the labor market. Thus, appropriate and comprehensive approaches are needed to address job stress-related issues in the workplace. In addition, psychosocial factors (including job stress and mental health) have been important issues in the field of occupational health, especially in developed countries. In this context, researchers, management, health providers, trade unions, and policy makers were interested in productivity, sustainable employment, and the burden of social change. Furthermore, there was increasing evidence that job stress was significantly associated with mental disorders,¹⁻³ coronary heart disease,^{4,5} stroke,⁶ diabetes mellitus,⁷ hypertension,^{8,9} and musculoskeletal disorders.¹⁰ Therefore, it is important to measure accurately the job stress-related burden of disease (BOD), in order to efficiently allocate limited resources and facilitate rational decision-making to promote workers' health.

Author Contributions

Conceptualization: Kim I. Data curation: Lee KJ, Kim I. Formal analysis: Lee KJ. Investigation: Kim I. Methodology: Lee KJ, Kim I. Writing - original draft: Lee KJ. Writing - review & editing: Kim I.

The World Health Organization (WHO) and the Global Burden of Disease (GBD) study suggested that death and disability-adjusted life years (DALY) was a useful metric for quantifying BOD. DALY is defined as the sum of years of life lost (YLL: burden due to death) and years lived with a disability (YLD: burden due to disability). Another metric for quantifying the effect of job stress on BOD is the population attributable fraction (PAF). PAF or BOD for specific causes were rare in Korea but it is worthwhile to estimate the social impact or to decide policy priority. Thus, the job stress-attributable BOD can be calculated by multiplying DALY and PAF.^{11,12}

Psychiatric disorders account for a large part of the global BOD, and the WHO has predicted that unipolar depressive disorder will become the leading cause of global BOD by 2030 (6.2% of the total DALY, up from 4.3% in 2004).¹³ In addition, for several years during the 21st century, Korea has ranked highest in suicide rate among the members of the Organization for Economic Co-operation and Development.¹⁴ Thus, Korean companies became increasingly interested in the mental health of their workers. Furthermore, the Samsung Economic Research Institute reported that the number of mental illnesses patients increased by 50% from 1.58 million people in 2004 to 2.31 million people in 2010. Moreover, the estimated socioeconomic cost of mental illness in 2010 was approximately 23 trillion KRW (approximately 19 billion USD), which corresponds to approximately 2.01% of the Korean gross domestic product (GDP) in 2010. The indirect cost of lost productivity was estimated to be approximately 20 trillion KRW. This significant economic burden highlights the importance of job stress-related mental health issues (e.g., psychiatric disorders and suicide).¹⁵

Previous studies have revealed relatively consistent associations between job stress and several cardiovascular diseases, which were listed as occupational diseases in the Enforcement Decree of the Korean Industrial Accident Compensation Insurance Act. These cardiovascular diseases include intracerebral hemorrhage, subarachnoid hemorrhage, cerebral infarction, myocardial infarction, and dissecting aortic aneurysm with work-related causes (including psychological strain), rather than natural disease progression.¹⁶ Cardiovascular diseases and cerebrovascular diseases were the second and third leading causes of death in Korea, which suggests that these diseases are important targets when estimating the BOD from job stress in Korea. Interestingly, a French study from 2003 revealed that the attributable fractions of job stress were 25% for cardiovascular disease, 34% for anxiety and depression, and 27% for musculoskeletal disorders.^{17,18}

The present study aimed to evaluate the role of job stress in disease development by estimating PAF for job stress in selected health outcomes and job stress related DALY in Korea.

METHODS

Health outcomes

We selected health outcomes as target diseases, which were defined by the following conditions: 1) the relative risk (RR) available (for calculating PAF) by reviewing literature, 2) disease prevalences available or calculable based on representative data, and 3) health outcomes with high priority with respect to the Korean occupational backgrounds and policy making; diseases that were the leading cause of death or were increasing in incidence would be eligible for this condition. Finally, the present study considered ischemic heart disease (IHD), stroke, major depressive disorder (MDD), and suicide as the health outcomes from job stress.

PAF

PAF was calculated using the following formula, as suggested by Levin and used in many previous studies^{19,20}:

$$\text{PAF} = P \times (\text{RR} - 1) / (1 + P \times [\text{RR} - 1])$$

In this equation, P denotes the proportion of workers who were exposed to job stress and RR denotes the RR of job stress related diseases.

The PAFs for ischemic stroke and hemorrhagic stroke were calculated using their respective RRs, and the total PAF for stroke was calculated by adding the PAF values for ischemic and hemorrhagic strokes.

The Korean National Police Agency has reported that approximately 4% of suicides were related to work- or workplace-related issues.²¹ Thus, we used this value to estimate PAF because no other studies have reported RRs or PAFs for job stress-related suicide in Korea.

Proportions of workers exposed to job stress

Based on Karasek's job strain model, high job strain occurs "when job demands are high and job decision latitude is low."²² Approximately 21.2% of Korean workers were estimated to have high job strain exposure, based on the 2006 Korean Working Conditions Survey.²³ The survey population represented active workers aged 15–65 years as the minimum age for employment in Korea was 15 years. The survey was conducted through face-to-face interviews using questionnaires, which was almost identical to the Fourth European Working Conditions Survey conducted in 2005.²⁴ In that study, job strain was categorized into four levels (low, active, passive, and high strain groups) using the combination of job demands exposure and job control. Prior to 2011, the year for which we estimated the prevalence of the target diseases, the Korean Working Conditions Survey had been conducted twice (in 2006 and 2010). Thus, we used data from 2006, which included the results of the job strain exposure proportion.

RR

The RRs in the present study were obtained from studies of meta-analyses that we selected after reviewing the literature. We considered these studies appropriate for our purposes, as they reported recent advances in this field and used well-designed strategies. Kivimäki and Kawachi²⁵ reported job strain-related RRs for incident coronary heart disease (1.33–1.44) in their review of 17 cohort studies since 2003. In addition, an RR of 1.34 (95% confidence interval [CI]: 1.18–1.51) was reported based on the IPD-Work Consortium data (10 unpublished studies, with previously published studies excluded).⁴ Moreover, job strain was associated with hazard ratios of 1.24 (95% CI, 1.05–1.46) for ischemic stroke and 1.01 (95% CI, 0.75–1.36) for hemorrhagic stroke.²⁶

In their meta-analysis, Theorell et al.²⁷ reported that job strain exposure was associated with a weighted odds ratio (OR) of 1.74 (95% CI, 1.53–1.96) for depression. However, Theorell et al.²⁷ evaluated both diagnosed depression and depressive symptoms (a much milder state), and Jauregui and Schnall²⁸ assumed that these depression outcomes could increase the duration of sickness or reduce workplace productivity and quality. As we agree with this approach, we believe that it was acceptable to use their OR which approximates RR based on the assumption that the disease prevalence was low in the population.²⁹

DALY

The following formula was used to estimate YLD based on the burden of disability:

$$YLD = P \times DW$$

In this equation, P denotes the prevalence of disease and DW denotes the disability weight, which quantifies health levels that are associated with non-fatal outcomes. We initially considered the DW values that were suggested by the GBD study, although they need to be modified to be properly used in the present study. Thus, we used the average values if there were multiple DW choices available from the GBD study.³⁰ For example, the GBD study classified the DW of MDD into 1) mild episodes = 0.145, 2) moderate episodes = 0.396, and 3) severe episodes = 0.658; thus, the final DW of MDD was defined as the average value (0.400). We used this method because we could not classify our prevalence data according to the disease classification approach used in the GBD study.

Disease prevalence was the preferred measure for calculating DALY, although it was difficult to directly calculate the prevalences in Korea because of the lack of available and representative data. Thus, we estimated the numbers of IHD and stroke cases based on 2011 claims data from the Health Insurance Review and Assessment Service.³¹ This estimation used the following parameters: 1) the health outcomes were matched only based on the code of the main diagnosis (excluding sub-diagnoses), 2) the health outcomes were defined as IHD (International Classification of Diseases-10 codes: I20–I25) and stroke (I60–I64), and 3) the claims were only submitted for inpatients. Based on these parameters, we calculated that there were 203,648 cases of IHD and 215,630 cases of stroke.

The MDD prevalence was estimated using data from the 2011 Korean Epidemiological Survey of Mental Disorders,³² which reported a 1-year prevalence of 3.1% for MDD among 18–74-year-old adults. Based on the mid-registration population during 2013 (the consensus standard), this prevalence corresponded to 1,271,312 MDD cases. As we assumed that the diseases' minimum latency periods were 0 years, we targeted individuals who were ≥ 15 years old (i.e., the minimum age of the economically active population). However, adults and children who were < 18 years old have different classification systems for mental illnesses and there were no data, that was why we only considered individuals who were 18–74 years old for the MDD calculations.

The following formula was used to estimate YLL based on the burden of death:

$$YLL = N \times L$$

In this equation, N denotes the number of deaths and L denotes the life expectancy. For example, if a woman dies of stroke at the age of 50 years, the YLL for stroke would be 36, based on a life expectancy of 86 years for Korean women. N and L were estimated from the national cause-of-death statistics (N) and life table (L), which were published by Statistics Korea in 2013 (IHD: I20–I25, stroke: I60–I64, I67, and I69, suicide: X60–X84).³³

Ethics statement

This study was approved by the Institutional Review Board of Sungkyunkwan University (#2014-06-009).

RESULTS

Approximately 21.2% of Korean workers were exposed to high job strain. When we evaluated the effects of job strain, we found that it was attributable for approximately 6.7% of IHD cases, 6.9% of stroke cases (4.8% for ischemic stroke and 2.1% for hemorrhagic stroke), 13.6% of MDD cases, and 4% of suicides (**Table 1**). The estimated prevalences per 100,000 people were 473 for IHD, 501 for stroke, and 3,100 for MDD. The deaths per 100,000 people were estimated to be 32 for IHD, 59 for stroke, and 67 for suicide. The total DALY values per 100,000 people were estimated to be 563 for IHD, 1,048 for stroke, 1,240 for MDD, and 1,102 for suicide. The total BOD per 100,000 people in terms of job stress-attributable DALY were estimated to be 38 for IHD, 72 for stroke, 168 for MDD, and 44 for suicide (**Table 2**). The detailed data that were used to calculate these findings are shown in **Table 3**.

DISCUSSION

As a result of the present study, the PAFs (%) for IHD, ischemic stroke, hemorrhagic stroke, MDD, and suicide were estimated to be 6.72, 4.84, 2.08, 13.6, and 4.00, respectively. Also, job stress-attributable BODs were estimated to be 38 years per 100,000 people for IHD, 72 years per 100,000 people for stroke, 168 years per 100,000 people for MDD, and 44 years per 100,000 people for suicide.

Sultan-Taïeb et al.^{17,18} reported that 8.8%–10.2% (men) and 0.2%–1.1% (women) of coronary heart disease morbidity and 15.2%–19.8% (men) and 14.3%–27.1% (women) of mental disorders (defined as clinical depression) were attributable to job strain in France. This French study was the only one that estimated the PAF of job strain. Similar to the present study, the French job stress exposure data were calculated using the Karasek model,²² which

Table 1. Estimates of RR and PAF for selected stress-related health outcomes

Health outcomes	RR (95% CI) ^a	PAF (%) ^b
IHD	1.34 (1.18–1.52) ^{4,25}	6.72
Stroke (ischemic)	1.24 (1.05–1.46) ^{25,29}	4.84
Stroke (hemorrhagic)	1.01 (0.75–1.36) ^{25,29}	2.08
Stroke (all)	-	6.92
MDD	1.74 (1.53–1.96) ¹	13.60
Suicide	-	4.00 ²⁵

PAF was calculated using the following formula: $PAF = P \times (RR - 1) / (1 + P \times [RR - 1])$, where P denotes the proportion of workers exposed to job stress (21.2% was used in the present study).

RR = relative risk, PAF = population attributable fraction, CI = confidence interval, IHD = ischemic heart disease, MDD = major depressive disorder.

^aThis pooled RR value from prospective studies was obtained by comparing to employees free of job strain;

^bApproximately 21.2% of Korean workers were thought to have high job strain exposure, based on the 2006 Korean Working Conditions Survey.²³

Table 2. Prevalences, deaths, total DALYs, and JS-DALYs for selected health outcomes in 2013 (unit: per 100,000 people)

Health outcomes	Prevalence (people)	Deaths rate ³³ (people)	YLD, yr	YLL, yr	Total DALYs, yr	JS-DALYs, yr
IHD	473 ³¹	32	74	489	563	38
Stroke	501 ³¹	59	155	893	1,048	72
MDD	3,100 ³²	-	1,240	-	1,240	168
Suicide	-	33	-	1,102	1,102	44

The mid-year registration population was used in 2013 (persons who were ≥ 15 years old [43,037,646 people] or were ≥ 18 years old and had major depressive disorders [41,010,081 people]). All values were reported per 100,000 people.

DALYs = disability-adjusted life years, JS-DALYs = job stress-attributable DALYs, IHD = ischemic heart disease, MDD = major depressive disorder, YLD = years lived with disability, prevalence × disability weight³⁰, YLL = years of life lost, number of deaths × life expectancy³³, Total DALYs = YLD + YLL, JS-DALYs = Total DALY × population attributable fraction (PAF).

Table 3. Age-specific data needed to calculate total DALYs for selected health outcomes

Age, yr	Prevalence (people) ³¹		Death (people) ³²		Life expectancy, yr ³²		DW ³⁰	YLD, yr		YLL, yr		Total DALYs, yr	
	Male	Female	Male	Female	Male	Female		Male	Female	Male	Female	Male	Female
IHD													
15-19	131	46	0	0	63.90	70.41	0.157	20.5	7.2	0.0	0.0	20.5	7.2
20-24	238	138	2	0	58.99	65.46	0.157	37.4	21.7	118.0	0.0	155.4	21.7
25-29	492	292	16	1	54.13	60.54	0.157	77.3	45.9	866.1	60.5	943.4	106.4
30-34	946	377	28	4	49.29	55.64	0.157	148.5	59.2	1,380.1	222.6	1,528.7	281.7
35-39	2,415	749	90	10	44.48	50.77	0.157	379.2	117.5	4,003.2	507.7	4,382.4	625.2
40-44	4,354	1,923	175	21	39.73	45.92	0.157	683.5	301.9	6,952.8	964.3	7,636.3	1,266.2
45-49	7,954	3,423	295	38	35.07	41.11	0.157	1,248.8	537.4	10,345.7	1,562.2	11,594.4	2,099.6
50-54	12,939	7,308	528	65	30.57	36.34	0.157	2,031.3	1,147.3	16,141.0	2,362.1	18,172.3	3,509.4
55-59	15,046	8,962	583	98	26.22	31.63	0.157	2,362.2	1,407.0	15,286.3	3,099.7	17,648.5	4,506.7
60-64	16,246	10,869	590	159	22.03	26.96	0.157	2,550.7	1,706.4	12,997.7	4,286.6	15,548.4	5,993.1
65-69	17,062	14,315	645	249	18.00	22.39	0.157	2,678.7	2,247.5	11,610.0	5,575.1	14,288.7	7,822.6
70-74	15,393	17,561	1,029	677	14.19	17.98	0.157	2,416.6	2,757.1	14,601.5	12,172.5	17,018.1	14,929.6
75-79	10,738	14,308	1,238	1,123	10.83	13.89	0.157	1,685.9	2,246.3	13,407.5	15,598.5	15,093.5	17,844.8
80+	7,522.96	11,900	1,934	3,969	8.00	10.26	0.157	1,181.1	1,868.3	15,472.0	40,721.9	16,653.1	42,590.2
Total	111,477	92,171	7,153	6,414	-	-	-	17,502	14,471	123,182	87,134	140,684	101,605
Stroke													
15-19	115	38	9	1	63.90	70.41	0.309	35.7	11.9	575.1	70.4	610.8	82.3
20-24	169	154	13	7	58.99	65.46	0.309	52.3	47.5	766.9	458.2	819.2	505.8
25-29	423	292	20	10	54.13	60.54	0.309	130.7	90.3	1,082.6	605.4	1,213.3	695.7
30-34	746	423	42	34	49.29	55.64	0.309	230.6	130.7	2,070.2	1,891.8	2,300.7	2,022.5
35-39	1,508	785	102	43	44.48	50.77	0.309	465.9	242.4	4,537.0	2,183.1	5,002.8	2,425.6
40-44	3,492	1,969	209	116	39.73	45.92	0.309	1,079.1	608.5	8,303.6	5,326.7	9,382.7	5,935.2
45-49	5,908	3,408	386	170	35.07	41.11	0.309	1,825.5	1,053.0	13,537.0	6,988.7	15,362.5	8,041.7
50-54	10,046	5,777	590	300	30.57	36.34	0.309	3,104.3	1,785.1	18,036.3	10,902.0	21,140.6	12,687.1
55-59	11,562	7,177	690	279	26.22	31.63	0.309	3,572.5	2,217.7	18,091.8	8,824.8	21,664.3	11,042.4
60-64	13,315	8,669	801	320	22.03	26.96	0.309	4,114.5	2,678.8	17,646.0	8,627.2	21,760.5	11,306.0
65-69	15,862	13,000	1,072	547	18.00	22.39	0.309	4,901.3	4,016.9	19,296.0	12,247.3	24,197.3	16,264.3
70-74	17,308	19,977	1,963	1,398	14.19	17.98	0.309	5,348.1	6,172.8	27,855.0	25,136.0	33,203.1	31,308.8
75-79	14,415	20,931	2,486	2,459	10.83	13.89	0.309	4,454.3	6,467.6	26,923.4	34,155.5	31,377.7	40,623.1
80+	12,708	25,454	3,706	7,658	8.00	10.26	0.309	3,926.6	7,865.2	29,648.0	78,571.1	33,574.6	86,436.3
Total	107,577	108,053	12,089	13,342	-	-	-	33,241	33,388	188,369	195,988	221,610	229,377
MDD													
Total	1,271,312		-		-		0.400	508,525		-		508,525	
Suicide													
15-19	-	-	165	106	63.90	70.41	-	-	-	10,543.5	7,463.5	10,543.5	7,463.5
20-24	-	-	308	177	58.99	65.46	-	-	-	18,168.9	11,586.4	18,168.9	11,586.4
25-29	-	-	412	287	54.13	60.54	-	-	-	22,301.6	17,375.0	22,301.6	17,375.0
30-34	-	-	714	411	49.29	55.64	-	-	-	35,193.1	22,868.0	35,193.1	22,868.0
35-39	-	-	775	375	44.48	50.77	-	-	-	34,472.0	19,038.8	34,472.0	19,038.8
40-44	-	-	1,025	434	39.73	45.92	-	-	-	40,723.3	19,929.3	40,723.3	19,929.3
45-49	-	-	1,076	332	35.07	41.11	-	-	-	37,735.3	13,648.5	37,735.3	13,648.5
50-54	-	-	1,236	409	30.57	36.34	-	-	-	37,784.5	14,863.1	37,784.5	14,863.1
55-59	-	-	1,030	288	26.22	31.63	-	-	-	27,006.6	9,109.4	27,006.6	9,109.4
60-64	-	-	746	212	22.03	26.96	-	-	-	16,434.4	5,715.5	16,434.4	5,715.5
65-69	-	-	608	202	18.00	22.39	-	-	-	10,944.0	4,522.8	10,944.0	4,522.8
70-74	-	-	752	302	14.19	17.98	-	-	-	10,670.9	5,430.0	10,670.9	5,430.0
75-79	-	-	638	312	10.83	13.89	-	-	-	6,909.5	4,333.7	6,909.5	4,333.7
80+	-	-	553	504	8.00	10.26	-	-	-	4,424.0	5,171.0	4,424.0	5,171.0
Total	-	-	10,038	4,351	-	-	-	-	-	313,312	161,055	313,312	161,055

DALYs = disability-adjusted life years, IHD = ischemic heart disease, MDD = major depressive disorder, DW = disability weight, YLD = years lived with disability, prevalence × disability weight³⁰, YLL = years of life lost, number of deaths × life expectancy³³, Total DALYs = YLD + YLL.

defines job strain as a combination of high psychological demands at work (i.e., task quantity and complexity, and time pressure) and low decision-making latitude (i.e., decision authority and skill discretion). However, direct comparison with the French results was difficult for some reasons: 1) Unlike the French study, we did not consider gender differences, 2) The

prevalence of job strain exposure was different (19.6% for men and 28.2% for women came from the national survey conducted in France in 2003), and 3) We used the RR values given in the studies of a meta-analysis, while they derived RRs from their own systematic review.

The exposure prevalence of job strain among Korean workers was estimated to be 21.2%, based on data from a cross-sectional questionnaire survey of 7,856 Korean workers.²³ As the PAF is very sensitive to changes in this value, care should be taken when selecting the data. We tried to reflect the Korean working environment and find the latest value at the same time; however, we had no choice but to select the result. This value would be representative, but there might be a gap with the current situation because it was from 2006. Several other studies have used different epidemiological data and reported similar findings: 19% among Finnish men and 23% among Finnish women,^{19,34} 18.6% among Australian men and 25.5% among Australian women,³⁵ and 23.2% among the general French population (19.6% for men and 28.2% for women).³⁶ Although we were unable to provide sex-specific Korean estimates, our data provide useful perspective regarding the Korean working environment.

In the present study, we estimated the prevalence of IHD and stroke using 2011 claims data from the Health Insurance Review and Assessment Service, and also estimated the MDD prevalence based on data from the Korean suicide prevention center. According to the 2011 Korea National Health and Nutrition Examination Survey report, the prevalence of physician-diagnosed disease were 2.4% for angina or myocardial infarction among > 30-year-old individuals, 3.8% for stroke among > 50-year-old individuals, and 4.5% for depression among > 30-year-old individuals.³⁷ However, these values were noticeably different from our estimates: 0.6% for IHD among > 30-year-old individuals, 1.3% for stroke among > 50-year-old individuals, and 3.9% for MDD among > 30-year-old individuals. This result was dependent on the respondents' reports and did not cover detailed diagnosis that we would like to estimate. On the other hand, our results would be more relevant in that they contained detailed and accurate diagnoses, and especially for stroke, we analyzed the claims data entered by doctors, not survey data. However, since the health insurance data was for the general population rather than workers, strictly speaking, the estimates do not reflect the prevalence among workers. Thus, it is possible that the prevalence was overestimated. We could not estimate the accurate prevalence among Korean workers by analyzing the claims data from the Health Insurance Review and Assessment Service. Self-employed workers, who accounted for 23.1% of the economically active population in 2011³⁸ and laborers are classified as district subscribers, which means they could not be distinguished from the general population.

In the GBD report, Korea was included in a group of countries with DALY values of 600–1,200 per 100,000 person-years for stroke and 0–300 per 100,000 person-years for IHD.³⁹ The present study estimated a relatively high DALY value for IHD, although the DALY value was similar for stroke. However, it was difficult to guess why the DALY for IHD in this study was high, because we could not figure out which data was used to estimate the DALYs. Similarly, Hong et al.⁴⁰ have reported that the Korean DALY for ischemic stroke was 483 per 100,000 person-years. This result was obtained from using the incidence of ischemic stroke by analyzing the health insurance data of 2004. Unlike these studies, we used the prevalence, not the incidence that was commonly used in estimating BOD. It was also impossible to measure the portion of job stress-attributable BOD among the total DALY.

The assessment of job stress-related health effects should be performed based on an understanding of the specific working environments and situations in each country.

However, there were limited Korean studies regarding this topic, which made it difficult to estimate job stress-related BOD using domestic data. Inevitably, the present study has several limitations that should be considered. First, we did not consider other diseases that are relevant to job strain, such as musculoskeletal disorders, because it was difficult to access prevalence data and reasonable RR values of many types of musculoskeletal disorders to calculate DALY. In addition, musculoskeletal disorders were excluded from the analysis because of possible reverse causation. Second, the data were not analyzed according to sex. However, it is reasonable to estimate the BOD with DALY as it is obtained by summation and there were no available study results to measure job stress exposure proportion by sex. Third, our result may be overestimated, and not accurately reflect the Korean working environment, as it was impossible to access data on prevalence or death rate targeting the Korean working population. Fourth, we replaced the RR with an OR for depression and with a hazard ratio for stroke. However, the prevalences of diseases were so low that the results obtained by using OR were not expected to make a large difference. Fifth, the proportion of workers exposed to job stress may not reflect the current working environment as there is a gap between the time of data collection and the present. Despite these limitations, the present study provides a useful perspective regarding the Korean working environment, and was designed to estimate the outcomes in a way that closely reflected this working environment. In particular, we tried to increase the validity of the calculated value by using representative data such as claim data from the Health Insurance Review and Assessment Service.

Despite several limitations due to data, the present study would be meaningful in that studies on PAF or BOD for specific causes were rare in Korea, and especially since studies on the calculation of BOD from job stress were also rare globally. However, in order to estimate the BOD more accurately, further studies would be necessary to: 1) Estimate the disease prevalence according to the employment type by using nationally representative data such as data of the National Health Insurance Service, 2) Calculate accurate RR values through a large-scale cohort study of workers, and 3) Evaluate the proportions of workers exposed to other job stressors including job strain.

BOD could be used to quantitatively estimate major health impacts and could be applied as the main outcome of public health policy, unlike other estimates of health impact such as mortality or morbidity. This is the first study to attempt to estimate the Korean BOD from job stress using the latest and best available data. As society becomes more concerned about suicides or deaths due to cerebrovascular disease in the working-age population in relation to burden at the workplace, estimating the BOD from job stress is essential in determining the priority of the national policy on prevention.

REFERENCES

1. Theorell T, Hammarström A, Aronsson G, Träskman Bendz L, Grape T, Hogstedt C, et al. A systematic review including meta-analysis of work environment and depressive symptoms. *BMC Public Health* 2015;15(1):738.
[PUBMED](#) | [CROSSREF](#)
2. Bonde JP. Psychosocial factors at work and risk of depression: a systematic review of the epidemiological evidence. *Occup Environ Med* 2008;65(7):438-45.
[PUBMED](#) | [CROSSREF](#)
3. Stansfeld S, Candy B. Psychosocial work environment and mental health--a meta-analytic review. *Scand J Work Environ Health* 2006;32(6):443-62.
[PUBMED](#) | [CROSSREF](#)

4. Kivimäki M, Nyberg ST, Batty GD, Fransson EI, Heikkilä K, Alfredsson L, et al. Job strain as a risk factor for coronary heart disease: a collaborative meta-analysis of individual participant data. *Lancet* 2012;380(9852):1491-7.
[PUBMED](#) | [CROSSREF](#)
5. Nyberg ST, Fransson EI, Heikkilä K, Alfredsson L, Casini A, Clays E, et al. Job strain and cardiovascular disease risk factors: meta-analysis of individual-participant data from 47,000 men and women. *PLoS One* 2013;8(6):e67323.
[PUBMED](#) | [CROSSREF](#)
6. Fransson EI, Nyberg ST, Heikkilä K, Alfredsson L, Bjorner JB, Borritz M, et al. Job strain and the risk of stroke: an individual-participant data meta-analysis. *Stroke* 2015;46(2):557-9.
[PUBMED](#) | [CROSSREF](#)
7. Nyberg ST, Fransson EI, Heikkilä K, Ahola K, Alfredsson L, Bjorner JB, et al. Job strain as a risk factor for type 2 diabetes: a pooled analysis of 124,808 men and women. *Diabetes Care* 2014;37(8):2268-75.
[PUBMED](#) | [CROSSREF](#)
8. Babu GR, Jotheeswaran AT, Mahapatra T, Mahapatra S, Kumar A Sr, Detels R, et al. Is hypertension associated with job strain? A meta-analysis of observational studies. *Occup Environ Med* 2014;71(3):220-7.
[PUBMED](#) | [CROSSREF](#)
9. Gilbert-Ouimet M, Trudel X, Brisson C, Milot A, Vézina M. Adverse effects of psychosocial work factors on blood pressure: systematic review of studies on demand-control-support and effort-reward imbalance models. *Scand J Work Environ Health* 2014;40(2):109-32.
[PUBMED](#) | [CROSSREF](#)
10. Lang J, Ochsmann E, Kraus T, Lang JW. Psychosocial work stressors as antecedents of musculoskeletal problems: a systematic review and meta-analysis of stability-adjusted longitudinal studies. *Soc Sci Med* 2012;75(7):1163-74.
[PUBMED](#) | [CROSSREF](#)
11. Forouzanfar MH, Alexander L, Anderson HR, Bachman VF, Biryukov S, Brauer M, et al. Global, regional, and national comparative risk assessment of 79 behavioural, environmental and occupational, and metabolic risks or clusters of risks in 188 countries, 1990-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 2015;386(10010):2287-323.
[PUBMED](#) | [CROSSREF](#)
12. World Health Organization (WHO). Global burden of disease (GBD). http://www.who.int/healthinfo/global_burden_disease/gbd/en. Accessed October 14, 2016.
13. World Health Organization (WHO). Global burden of disease (GBD): 2004 update. http://www.who.int/healthinfo/global_burden_disease/GBD_report_2004update_part.pdf. Accessed October 14, 2016.
14. Organization for Economic Co-operation and Development (OECD). Suicide rates (indicator). <https://data.oecd.org/healthstat/suicide-rates.htm>. Accessed October 14, 2016.
15. Samsung Economic Research Institute. Conditions for healthy company: mental health of workers. In: Lee KJ, translator. *CEO Information*. Seoul, Korea: Samsung Economic Research Institute; 2012, 1-3.
16. Enforcement decree of the industrial accident compensation insurance act: presidential decree No. 27751 (2016.12.30). <http://www.law.go.kr/lsSc.do?menuId=0&p1=&subMenu=1&nwYn=1&query=22804&x=0&y=0#AJAX>. Accessed January 9, 2017.
17. Sultan-Taïeb H, Lejeune C, Drummond A, Niedhammer I. Fractions of cardiovascular diseases, mental disorders, and musculoskeletal disorders attributable to job strain. *Int Arch Occup Environ Health* 2011;84(8):911-25.
[PUBMED](#) | [CROSSREF](#)
18. Sultan-Taïeb H, Niedhammer I. Evaluating the burden of job stress from the public-health and economic viewpoints: perspectives and methodological pitfalls. *Travail et Emploi* 2013;5(1):45-54.
[CROSSREF](#)
19. Levin ML. The occurrence of lung cancer in man. *Acta Unio Int Contra Cancrum* 1953;9(3):531-41.
[PUBMED](#)
20. Nurminen M, Karjalainen A. Epidemiologic estimate of the proportion of fatalities related to occupational factors in Finland. *Scand J Work Environ Health* 2001;27(3):161-213.
[PUBMED](#) | [CROSSREF](#)
21. Ministry of Health & Welfare. *Suicide Prevention White Book*. Seoul, Korea: Ministry of Health & Welfare; 2015.
22. Karasek RA. Job demands, job decision latitude, and mental strain: implications for job redesign. *Adm Sci Q* 1979;24(2):285-308.
[CROSSREF](#)
23. Lu ML, Nakata A, Park JB, Swanson NG. Workplace psychosocial factors associated with work-related injury absence: a study from a nationally representative sample of Korean workers. *Int J Behav Med* 2014;21(1):42-52.
[PUBMED](#) | [CROSSREF](#)

24. Parent-Thirion A, Fernández Macías E, Hurley J, Vermeylen G. *Fourth European Working Conditions Survey*. Dublin, Ireland: European Foundation for the Improvement of Living and Working Conditions; 2006.
25. Kivimäki M, Kawachi I. Work stress as a risk factor for cardiovascular disease. *Curr Cardiol Rep* 2015;17(9):630.
[PUBMED](#) | [CROSSREF](#)
26. Fransson EI, Nyberg ST, Heikkilä K, Alfredsson L, Bjorner JB, Borritz M, et al. Job strain and the risk of stroke: an individual-participant data meta-analysis. *Stroke* 2015;46(2):557-9.
[PUBMED](#) | [CROSSREF](#)
27. Theorell T, Hammarström A, Aronsson G, Träskman Bendz L, Grape T, Hogstedt C, et al. A systematic review including meta-analysis of work environment and depressive symptoms. *BMC Public Health* 2015;15(1):738.
[PUBMED](#) | [CROSSREF](#)
28. Jauregui M, Schnall PL. Work, psychosocial stressors and the bottom line, unhealthy work. In: Schnall PL, Dobson M, Rosskam E, editors. New York, NY: Baywood Publishers; 2009, 153-67.
29. Walter SD. The distribution of Levin's measure of attributable risk. *Biometrika* 1975;62(2):371-2.
[CROSSREF](#)
30. Salomon JA, Haagsma JA, Davis A, de Noordhout CM, Polinder S, Havelaar AH, et al. Disability weights for the global burden of disease 2013 study. *Lancet Glob Health* 2015;3(11):e712-23.
[PUBMED](#) | [CROSSREF](#)
31. Health Insurance Review and Assessment Services. <http://www.hira.or.kr/eng/index.html>. Accessed December 26, 2017.
32. Seoul National University College of Medicine. *The Epidemiological Survey of Mental Disorders in Korea 2011*. Seoul, Korea: Ministry of Health and Welfare; 2012.
33. Statistics Korea. <http://kostat.go.kr/portal/korea/index.action>. Accessed December 26, 2017.
34. Kauppinen T, Toikkanen J, Pukkala E. From cross-tabulations to multipurpose exposure information systems: a new job-exposure matrix. *Am J Ind Med* 1998;33(4):409-17.
[PUBMED](#) | [CROSSREF](#)
35. LaMontagne AD, Keegel T, Vallance D, Ostry A, Wolfe R. Job strain - attributable depression in a sample of working Australians: assessing the contribution to health inequalities. *BMC Public Health* 2008;8(1):181.
[PUBMED](#) | [CROSSREF](#)
36. Sultan-Taïeb H, Chastang JF, Mansouri M, Niedhammer I. The annual costs of cardiovascular diseases and mental disorders attributable to job strain in France. *BMC Public Health* 2013;13(1):748.
[PUBMED](#) | [CROSSREF](#)
37. Ministry of Health & Welfare, Korea Centers for Disease Control and Prevention. Korea Health Statistics 2011: Korea National Health and Nutrition Examination Survey (KNHANES). Seoul, Korea: Ministry of Health & Welfare; 2012.
38. Statistics Korea. Economically active population survey. http://www.index.go.kr/potal/main/EachDtlPageDetail.do?idx_cd=2779. Accessed December 26, 2017.
39. Kim AS, Johnston SC. Global variation in the relative burden of stroke and ischemic heart disease. *Circulation* 2011;124(3):314-23.
[PUBMED](#) | [CROSSREF](#)
40. Hong KS, Kim J, Cho YJ, Seo SY, Hwang SI, Kim SC, et al. Burden of ischemic stroke in Korea: analysis of disability-adjusted life years lost. *J Clin Neurol* 2011;7(2):77-84.
[PUBMED](#) | [CROSSREF](#)