

RESEARCH ARTICLE

Pain and Function Recovery Trajectories following Revision Hip Arthroplasty: Short-Term Changes and Comparison with Primary Hip Arthroplasty in the ADAPT Cohort Study

Erik Lenguerrand*, Michael R. Whitehouse, Vikki Wylde, Rachael Gooberman-Hill, Ashley W. Blom

Musculoskeletal Research Unit, School of Clinical Sciences, University of Bristol, Level 1 Learning and Research Building, Southmead Hospital, Westbury-on-Trym, Bristol, BS10 5NB, United Kingdom

* erik.lenguerrand@bristol.ac.uk



OPEN ACCESS

Citation: Lenguerrand E, Whitehouse MR, Wylde V, Gooberman-Hill R, Blom AW (2016) Pain and Function Recovery Trajectories following Revision Hip Arthroplasty: Short-Term Changes and Comparison with Primary Hip Arthroplasty in the ADAPT Cohort Study. PLoS ONE 11(10): e0164839. doi:10.1371/journal.pone.0164839

Editor: Luigi F. Rodella, University of Brescia, ITALY

Received: March 24, 2016

Accepted: September 30, 2016

Published: October 14, 2016

Copyright: © 2016 Lenguerrand et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: Ethical restrictions prevent public sharing of data. The restrictions prohibiting the authors from making this dataset publicly available are related to the ethical permissions for the study (NHS Research Ethics committee South West, REC reference 09/H0102/72, IRAS project ID 36293). Patients were asked to provide their written informed consent to participate and for members of the University of Bristol to have access to the data for research purposes. They did not provide consent to the

Abstract

Background and Purpose

Patients report similar or better pain and function before revision hip arthroplasty than before primary arthroplasty but worse results are reported after revision surgery than after primary surgery. The trajectory of post-operative recovery during the first months and any differences by type of surgery have received little attention. We explored the trajectories of change in pain and function after revision hip arthroplasty to 12-months post-operatively and compare them with those observed after primary hip arthroplasty.

Methods

This study is a prospective cohort study of patients undergoing primary (n = 80 with 92% for an indication of osteoarthritis) and revision (n = 43) hip arthroplasties. WOMAC pain and function scores and walking speed were collected pre-operatively, at 3 and 12-months post-operatively. Multilevel regression models were used to chart and compare the trajectories of change (0–3 months and 3–12 months) between types of surgery.

Results

The improvements in pain and function following revision arthroplasty occurred within the first 3-months with no evidence of further change beyond this initial period. While the pattern of recovery was similar to the one observed after primary arthroplasty, improvements in the first 3-months were smaller after revision compared to primary arthroplasty. Patients listed for revision surgery reported lower pre-operative pain levels but similar post-operative levels compared to those undergoing primary surgery. At 12-months post-operation patients who underwent a revision arthroplasty had not reached the same level of function achieved by those who underwent primary arthroplasty.

sharing of the data beyond University of Bristol staff. However, external researchers wishing to view or re-use the minimal dataset may submit their request to the study sponsor, North Bristol NHS Trust, to obtain information about how to apply for appropriate NHS Research Ethics and Governance approvals: Research & Innovation North Bristol NHS Trust Floor 3 Learning & Research building Southmead Hospital, Westbury-on-Trym Bristol BS10 5NB United Kingdom E-mail: research@nbt.nhs.uk. With this in place, the principal investigator of the ADAPT Study (Professor Ashley Blom, University of Bristol, Musculoskeletal Research Unit) will facilitate access to the data under the terms defined by the NHS Research Ethics and Governance approvals and the study sponsor.

Funding: This article presents independent research funded by the National Institute for Health Research (NIHR) (<http://www.nihr.ac.uk/>) under its Programme Grants for Applied Research programme (RP-PG-0407-10070). The funder had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. The views expressed in this article are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health.

Competing Interests: The authors have declared that no competing interests exist.

Conclusion

The post-operative improvements in pain and function are larger following primary hip arthroplasty than following revision hip arthroplasty. Irrespective of surgery type, most of the improvements occur in the first three post-operative months. More research is required to identify whether the recovery following revision surgery could be improved with specific post-operative interventions.

Introduction

The volume of primary hip arthroplasties rose by approximately 26% between 2010 and 2015 [1, 2]. Over 88,000 primary total hip arthroplasties are performed per year in England and Wales [2]. These figures will continue to rise due to increases in obesity and an aging community [3]. The revision burden is approximately 11% and over 9,500 revision hip arthroplasties were performed in England and Wales in 2015 [1, 2].

Both primary and revision hip arthroplasty have been shown to improve patient-reported pain and function for the majority of patients [4–8]. While patients tend to have similar [5, 6, 8] or better [7] pain and function prior to revision arthroplasty than prior to primary arthroplasty, patients who undergo primary surgery report better post-operative outcomes than those who undergo revision surgery [5–7].

Typically, the outcome of both primary and revision is reported at 12-months or more after surgery [6–8]. The pattern of recovery trajectories within the first 12-months after surgery, and differences between primary and revision surgery in this period have received little attention. Pain and function have previously been reported not to improve further after 6-months following revision arthroplasty [9, 10]. In the absence of assessment prior to 6-months in these studies the pattern of improvement in the first months post-operation requires further investigation.

Evidence from the ADAPT cohort study suggests that in primary hip arthroplasty, the improvement in patient-reported pain and function post-operation plateaus at 3-months [11]. It is not currently clear if the pattern of recovery following revision hip arthroplasty is similar or if the complexity, extent of surgery and surgical trauma leads to a different pattern.

To describe and explore potential disparities in the degree and pattern of post-operative recovery following revision hip arthroplasty, we analysed data collected pre-operatively, at 3- and 12-months post-operation from the ADAPT prospective cohort study. Specifically our research aims were 1. to describe the early trajectories of pain and function after revision hip arthroplasty, 2. compare these trajectories with those observed after primary hip arthroplasty and 3. compare the post-operative outcomes achieved after these two types of surgery. We also investigated whether the pattern of recovery was similar when function is objectively assessed with standardised performance tests compared to patient-reported outcome measures.”

Materials and Methods

This study followed the STROBE (STrengthening the Reporting of OBservational studies in Epidemiology) guidelines for reporting observational studies in epidemiology (Appendix A in [S1 File](#)).

Study design

ADAPT is a single-centre UK prospective cohort study including patients undergoing hip or knee arthroplasty (UKCRN ID 8311). National Health Service Research Ethics Committee

approval was granted for the study (09/H0102/72) and all patients provided informed, written consent.

Detailed information on study design, patient recruitment, inclusion-exclusion criteria, and assessment methods are provided in the published study protocol [12]. Briefly, between February 2010 and November 2011, patients waiting for hip or knee arthroplasty at a high-volume elective orthopaedic centre were invited to participate in the study. Approximately 250 patients were recruited to ensure a sufficient number of patients to perform meaningful data analysis. Patients were due to undergo a range of primary and revision arthroplasty procedures (primary total knee arthroplasty, unicompartmental knee arthroplasty, patellofemoral arthroplasty, revision total knee arthroplasty, primary total hip arthroplasty, primary hip resurfacing or revision total hip arthroplasty) so that functional measures could be investigated across a range of patients with diverse indications for surgery and degrees of functional impairment. The majority of patients listed for primary arthroplasty had an indication of osteoarthritis. Exclusion criteria included an inability to provide written informed consent, to complete English language questionnaires (not all the questionnaires we used have been translated or validated for use in other languages), participation to another study, and severe functional limitations which would prevent completion of a performance test. In particular, patients using wheelchairs were excluded.

This analysis was restricted to patients who underwent primary total, resurfacing or revision hip arthroplasty.

Data collection

Assessments were conducted before surgery (median 19 days) and then at 3 and 12-months after surgery. At each post-operative assessment time, participants completed a postal questionnaire.

Participant and surgical characteristics

Data on gender, age, living arrangement, level of education, working status and number of joints affected by arthritis were collected in the pre-operative questionnaire. The indication for surgery, type of surgery, surgical approach, height and weight were extracted from participants' medical records.

Patients undergoing primary arthroplasty had a total hip replacement ($n = 74$) or hip resurfacing ($n = 6$). Osteoarthritis was the indication for surgery in 92% of primary cases. Patients undergoing revision arthroplasty had revision of a total hip arthroplasty (86%, $n = 37$), hip resurfacing (9%, $n = 4$), or hemiarthroplasty (5%, $n = 2$). The most common indication for revision arthroplasty was aseptic loosening (67%, $n = 29$); the remaining indications were pain (9%, $n = 4$), aseptic lymphocyte-dominated vasculitis-associated lesion (9%, $n = 4$) and other reasons (11%, $n = 6$). Primary (87%, $n = 70$) and revision arthroplasties (98%, $n = 42$) were mostly commonly performed via a posterior surgical approach.

Patient-reported measures

Self-reported pain and function were assessed using the Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC) function and pain sub-scales [13]. The WOMAC-function measure consists of 17 questions assessing the extent of functional limitation when performing a range of daily activities. WOMAC-pain consists of five questions assessing pain during walking, using stairs, in bed, sitting or lying. Each sub-score ranges from 0–100 (worst to best). The WOMAC score has good psychometric properties with test-retest reliability above 0.8 for the physical function subscale and above 0.7 for the pain subscale [14].

Performance test

An objective measure of function was obtained using a timed walk test [15]. Participants were timed and supervised by a research nurse as they walked a 20 metres straight distance on level ground at their normal, comfortable speed. Speed (metres per second) was derived by dividing the distance walked by the time required to complete the task. The test-retest reliability of the 20 metres has been showed to be high (> 0.9) [16, 17].

Statistical analysis

Three random intercept and slope linear regression models, one for each studied outcome (WOMAC-pain, WOMAC-function and walking speed), were used to investigate the pattern of post-operative changes following revision hip arthroplasty (aim 1) and compare the changes with those following primary hip arthroplasty (aim 2). This approach accounts for repeated and unequal numbers of measurements per participant while producing estimations valid under the missing at random assumption [18]. In this modelling framework, all available pre- and post-operative assessments of the outcome of interest were modelled. The outcomes were standardised (using the pre-operative mean and standard deviation of the score of interest) to produce estimates comparable across models. Those outcomes were regressed on an intercept (mean of standardised outcome on day of surgery at the sample mean age), age (centred at 65.2, the overall sample mean age) and two time splines (with random effect on their associated effects): one spline (a line between two points) for the “short-term change” occurring between the pre-operative assessment and the second assessment (3-months post-operative) and another spline for the “long-term change” occurring between the two post-operative assessments (3 and 12-months). Changes between assessment points were modelled rather than the actual scores achieved at 3- or 12-months. This was because the distributions of the scores were strongly skewed but the changes were normally distributed and could be analysed with the model framework presented above (as evidenced by the residuals plots). These models were stratified by primary/revision status to produce estimates specific to each type of surgery. Comparisons of the short- and long-term changes by surgery type were performed using their fixed effects and contrasts.

The equation structure of these models is described in more details in appendix B in [S1 File](#) with the code used to compute them.

The short- and long-term changes were also plotted by surgery type. For this purpose, the random intercept and slope linear model framework described above was re-run unadjusted for age on the unstandardised outcomes of pain and function. The fixed effects associated with the intercepts, and time splines of the primary and revision arthroplasty equations were used to produce the mean changes and their 95%CI.


Finally, the post-operative outcomes achieved at 3- and 12-months post-operatively were compared by surgery type (aim 3). As explained, the actual post-operative outcomes were strongly skewed and could not be investigated within the regression framework. Mann-Whitney tests were used for this purpose.

All models were fitted using Stata SE 13.1 (StataCorp LP, College Station, Texas, USA) and MLwiN v2.31 using Stata runmlwin command [19]. A p-value of <0.05 was considered as evidence of statistical significance.

Results

Sample description

Overall, 664 patients were identified on the waiting list for primary or revision hip arthroplasty (Fig 1). A total of 447 patients were not approached or refused to discuss the study. Forty-six



*Patients who underwent surgery, with at least one pre- or post-surgery assessment

Fig 1. Flow diagram of recruitment and participation.

doi:10.1371/journal.pone.0164839.g001

patients were also ineligible among which 15 were wheelchair users including two with severe balance issue. A total of 171 patients were eligible and 131 agreed to take part (77%). Eight patients did not subsequently undergo hip arthroplasty and therefore 123 patients were included in the final analysis. Of these patients 80 had a primary and 43 had a revision hip arthroplasty.

All these 123 participants had at least one assessment (pre-operative, 3 and/or 12-months) for any of the investigated measures (WOMAC-pain, WOMAC-function scores or walking speed) and were considered in the analyses. A description of the available number of assessments at each data collection points is provided in Table 1. The percentage of participants with complete information on WOMAC-pain, WOMAC-function scores and walking speed at 12-months post-operation was comparable between type of surgery (89% and 84% for primary and revision arthroplasty respectively).

The characteristics of the cohort are shown in Table 2. The mean age was 65 years (SD 11) and 66 years (SD 11) respectively for participants who underwent a primary and revision arthroplasty respectively. The median body mass index was 26 kg/m² (Interquartile range (IQR) 24–29) and 28 (24–28) respectively.

Pain and function trajectories after revision arthroplasty

Revision hip arthroplasty lead to a significant improvement in both pain and function (Fig 2). Changes in pain and function occurred within the first 3-months post-operation (WOMAC-pain, $p < 0.0001$; WOMAC-function, $p < 0.0001$; Walking speed, $p < 0.0001$; S1 Table). No evidence of further improvement in pain or function was found between 3 and 12-months

Table 1. Pain and function by assessment period and revision/primary profile.

N = 123	Pre-operative			3-months			12-months		
	Total	Primary	Revision	Total	Primary	Revision	Total	Primary	Revision
WOMAC Pain	121	78	43	112	76	36	108	71	37
missing, n =	2	2	0	11	4	7	15	9	6
%	98.4	64.5	35.5	91.1	67.9	32.1	87.8	65.7	34.3
Median	55	55	60	95	95	95	100	100	95
Interquartile range ^a	[35, 70]	[30, 70]	[50, 75]	[80, 100]	[85, 100]	[73, 100]	[85, 100]	[90, 100]	[80, 100]
p-value ^b	0.031			0.479			0.268		
WOMAC function	121	78	43	112	76	36	109	71	38
missing, n =	2	2	0	11	4	7	14	9	5
%	98.4	64.5	35.5	91.1	67.9	32.1	88.6	65.1	34.9
Median	56	54	62	90	90	89	94	96	93
Interquartile range ^a	[38, 71]	[38, 71]	[41, 75]	[81, 96]	[81, 96]	[79, 96]	[84, 99]	[87, 100]	[76, 97]
p-value ^b	0.165			0.678			0.015		
Walking-speed^c	118	77	41	107	74	33	108	72	36
missing, n =	5	3	2	16	6	10	15	8	7
%	95.9	65.3	34.8	87	69.2	30.8	87.8	66.7	33.3
Median	0.91	0.91	0.83	1.11	1.11	1.05	1.18	1.18	1.11
Interquartile range ^a	[0.71, 1.11]	[0.71, 1.11]	[0.67, 1.11]	[0.91, 1.25]	[0.95, 1.25]	[0.91, 1.18]	[0.95, 1.33]	[1.03, 1.38]	[0.87, 1.18]
p-value ^b	0.464			0.343			0.004		

a First and third quartiles: 25th and 75th percentiles

b Mann-Whitney test to compare median scores by primary/revision profile.

c Walking-speed expressed in metres per second: 20 metres / Completion-time

doi:10.1371/journal.pone.0164839.t001

Table 2. Participant characteristics.

		N = 123		Primary n = 80		Revision n = 43	
			%		%		%
Sex	Men	61	49.6	38	47.5	23	53.5
	Women	62	50.4	42	52.5	20	46.5
Number of other joints with OA	None	25	20.3	20	25.0	5	11.6
	One joint	29	23.6	22	27.4	7	16.3
	Two joints	22	17.9	13	16.3	9	20.9
	3 joints	18	14.6	8	10.0	10	23.3
	> = 4 joints	22	17.9	13	16.3	9	20.9
	Unknown	7	5.7	4	5.0	3	7.0
Living alone	Living with someone	90	73.2	59	73.7	31	72.1
	Living alone	30	24.4	18	22.5	12	27.9
	Unknown	3	2.4	3	3.8	0	0.0
Education	Normal school leaving age	66	53.7	41	51.2	25	58.1
	College	26	21.1	20	25.0	6	14.0
	University	28	22.8	16	20.0	12	27.9
	Unknown	3	2.4	3	3.8	0	0.0
Working status	Full time	55	44.7	34	42.5	21	48.8
	Retired	60	48.8	38	47.5	22	51.2
	Unemployed	7	5.7	7	8.7	0	0.0
	Unknown	1	0.8	1	1.3	0	0.0

doi:10.1371/journal.pone.0164839.t002

(S1 Table). Pain and function trajectories after primary arthroplasty are reported in Fig 2 and S1 Table and have been previously described [11]. Changes mainly occur within the first 3-months following surgery and there is no evidence of further changes after 3-months with the exception of the walking speed which continued to marginally improved between 3 and 12-months ($p = 0.005$).

Comparisons between revision and primary arthroplasty

Pre-operatively, the level of pain reported by participants listed for primary surgery was worse than for those listed for revision surgery (median 55 vs. 66, $p = 0.031$, Table 1). However greater short-term improvements in WOMAC-pain (Fig 2A) were assessed during the first 3-months following primary arthroplasty compare to the changes found after revision arthroplasty ($p < 0.0001$, S1 Table). No evidence of change was found between 3 and 12-months for either type of surgery. As a result there was no more significant difference at 3 ($p = 0.479$, Table 1) or 12-months ($p = 0.268$, Table 1). Pre-operatively, the median WOMAC-function scores were not different between those that underwent primary and revision hip arthroplasty (Table 1). The mean short-term change in the WOMAC-function score (Fig 2B) following revision arthroplasty was smaller than that following primary arthroplasty ($p < 0.001$, S1 Table). No evidence of long-term change (between 3 and 12-months) was observed for either type of surgery. At 12-months post-operation, the median WOMAC-function score was higher after primary surgery than after revision surgery (96 vs 93, $p = 0.015$, Table 1).

The walking speed was comparable pre-operatively for participants who subsequently underwent primary and revision arthroplasty ($p = 0.464$, Table 1). The trajectory of recovery exhibited by the walking speed differed between patients with revision and primary arthroplasty (Fig 2C). The speed improved to a similar extent during the first 3-months following both types of surgery. However, the improvement continued after 3-months for primary




Fig 2. Mean trajectories^a for WOMAC-pain, WOMAC-function and walking speed (Unstandardised outcomes) by revision/primary surgery. The mean trajectories are derived using the fixed effects terms of the linear mixed models stratified on primary-revision profile and regressing each outcome on the time of assessment parameterised as two linear splines.

doi:10.1371/journal.pone.0164839.g002

arthroplasty ($p = 0.005$, [S1 Table](#)) but not for revision arthroplasty ($p = 0.300$, [S1 Table](#)). Participants who had undergone revision arthroplasty reported a slower walking speed at 12-months post-operatively than those who had undergone primary ($p = 0.004$; [Table 1](#)).

Discussion

Investigation of the early outcome trajectories after revision hip arthroplasty has revealed that the improvements in pain and function were mainly observed within the first 3 post-operative months with no evidence of further change beyond this initial period. The comparisons of these trajectories with those observed after primary hip arthroplasty have shown that while pain or function recovery was plateauing before six-months for both surgery types, the extent of improvements was different with smaller short-term changes after revision arthroplasty than after primary arthroplasty. Comparisons of the achieved post-operative outcomes reveal that patients who underwent revision surgery reported less pre-operative pain than those who underwent primary surgery but due to the difference in the extent of post-operative changes this advantage was not sustained post-operatively. While function was comparable pre-operatively, at 12-months post-operation patients who underwent revision arthroplasty had not reached the level of function achieved by those who underwent primary arthroplasty. Finally, some difference in the pattern of recovery was observed when function was objectively assessed. Contrary to patient-reported function, minor but statistically significant improvements in walking-speed was observed between 3 and 12-months after primary arthroplasty and this long-term changes were not observed after revision surgery.

The observed effectiveness of revision hip arthroplasty to improve patient-reported pain and function is consistent with the existing evidence [[9](#), [20–25](#)]. The few studies measuring outcomes prior to 12 months post-operation report that changes in outcomes following revision hip arthroplasty plateau at 6-months post-operation [[9](#), [10](#)]. None of these studies has measured pain or function at 3-months post-operation and the current study filled this gap. This “plateau” was reached at least 3-months earlier than previously shown and was lower than the one observed after primary arthroplasty. This suggests that the higher complexity or degree of trauma related to revision surgery as compared to primary surgery limits the extent of the recovery but does not increase the time taken to recover: patients undergoing revision arthroplasty will improve but should not expect to achieve outcomes as high as those reached after their primary surgery.

The differential between the degree of improvement following revision and/or the outcome level reached post-operatively compared to primary surgery have also been shown but only in the post-operative period starting 12 months or more after surgery [[4–8](#), [25–28](#)]. In this respect, the current findings fill another gap in the literature.

There is limited evidence on the improvement in objective function following revision hip arthroplasty but the findings are in agreement with ours [[20](#)]. Aghayev et al. demonstrated the benefit of revision hip arthroplasty on the ability to walk, reporting an improvement in the percentage of their patients unable to walk for more than 30 minutes 12-months after surgery from 65% pre-operatively to 50% 12-months after surgery. Similar to our observations, the improvement was less than that observed following primary surgery.

The strengths of this study are the availability of patient reported outcomes measured at 3-months post-operation in addition to objective function assessment with a performance test. Using an objective measurement allows us to ascertain that the lack of functional improvement beyond 3-months among the patients who underwent a revision surgery is not due to the inherent ceiling effect associated with using a score-bounded patient-reported outcome such as the WOMAC score [29–32]: the lack of long-term improvement was also observed when function was measured with the walking speed, an objective tool that may be less subject to ceiling-effect. Moreover, among patients undergoing primary surgery, the mean long-term improvement of objective function was significant but small compared to the short-term mean improvement (0.02 vs 0.15, Table 2), confirming that most of the functional changes, whether objectively or subjectively measured, occur before the 3-month post-operative time point.

This study is not without limitations. The findings were obtained on patients from a single-centre orthopaedic unit limiting their external validity. The modest sample size restricted our ability to adjust for factors known to be associated with post-operative outcomes such as gender, mental health and co-morbidities [26, 33, 34]. However, our findings were adjusted for age. A larger sample would nevertheless have been required to adjust for additional confounding factors, in particular type or indication for surgery. As all patients undergoing revision surgery and 93% of those undergoing primary surgery received a total hip replacement, our comparisons between revision and primary arthroplasty are more generalizable to patients undergoing total joint replacement. The remaining 7% (n = 6) of patients undergoing primary arthroplasty were listed for resurfacing surgery. While they exhibit comparable pre- and post-operative functional outcomes their pain at 12 months post-operation was worse than for those listed for primary total arthroplasty (Medians: 75 vs 100; p-values = 0.03). The group of patients undergoing revision surgery was modest in size (n = 43) but relatively homogeneous with 86% being revision of a primary total arthroplasty rather than after a previous revision. A larger revision group would have allowed the stratification of the analysis by indication for surgery. The post-operative outcomes following revision arthroplasty have been shown to be influenced by the indication for surgery [22, 35] and therefore our findings are more reflective of those revised for aseptic loosening (>67% of the revised participants). No information on the pre- and post-operative treatment received by the participants was available. They were offered standard care as provided at the treating centre. This comprised a pre-operative educational class focusing on preparation for surgery and the hospital stay, and post-operative outpatient physiotherapy on a needs basis. Finally, the inclusion of additional assessment points prior to 3-months would have allowed more detailed investigation of the very early recovery trajectories. We considered that additional assessment points would have represented an excess burden for participants with a probability of increased levels of attrition in the cohort.

Conclusions

Patients undergoing revision hip arthroplasty should be informed that the expected improvement following such surgery will be less marked than that expected and experienced for primary surgery and the majority of their improvement will occur in the first 3 post-operative months.

More research is now required to identify whether specific in-patient and post-discharge rehabilitation tailored towards patients undergoing revision arthroplasty would improve or achieve equivalent outcomes to primary surgery and whether patients who are achieving limited improvements at 3-months post-operative would benefit from longer or more intensive rehabilitation. This will become all the more important with the increasing volume of

revision surgery and the high expectations of patients who aspire to a disease-free and active life [36–38].

Supporting Information

S1 File. Appendices.

(DOCX)

S1 Table. Univariate age-adjusted linear mixed regression models of function and pain stratified by revision/primary profiles.

(DOCX)

Acknowledgments

The research team acknowledges the support of the NIHR, through the Comprehensive Clinical Research Network. The authors would like to acknowledge and thank the participants of the ADAPT study, Professor Paul Dieppe, members of the Project Steering Committee and the following research team members: Cindy Mann, Debbie Delgado, Connie Jamera, Louise Hawkins, Sue Bowman, Leigh Morrison, Dave Rea and Jenny Tyler.

Author Contributions

Conceptualization: EL MRW VW RGH AWB.

Data curation: EL AWB.

Formal analysis: EL.

Funding acquisition: VW RGH AWB.

Investigation: EL MRW VW RGH AWB.

Methodology: EL MRW.

Project administration: VW RGH AWB.

Resources: VW RGH AWB.

Supervision: MRW AWB.

Visualization: EL MRW VW RGH AWB.

Writing – original draft: EL MRW VW RGH AWB.

Writing – review & editing: EL MRW VW RGH AWB.

References

1. National Joint Registry for England and Wales. 11th Annual Report 2014. Available from: <http://www.njrreports.org.uk/Portals/0/PDFdownloads/NJR%2011th%20Annual%20Report%202014.pdf>.
2. National Joint Registry. 2015 [29/02/2016]. Available from: http://www.njrreports.org.uk/hips-all-procedures-activity/H01v2NJR?reportid=06011166-BAC8-4202-8161-F42B58C34CD7&defaults=JYS_Filter_Calendar_Year_From_To=%22max-max%22,H_Filter_Joint=%22Hip%22.
3. Kurtz S, Ong K, Lau E, Mowat F, Halpern M. Projections of primary and revision hip and knee arthroplasty in the United States from 2005 to 2030. *J Bone Joint Surg Am.* 2007; 89(4):780–5. doi: [10.2106/JBJS.F.00222](https://doi.org/10.2106/JBJS.F.00222) PMID: [17403800](https://pubmed.ncbi.nlm.nih.gov/17403800/).
4. Adelani MA, Crook K, Barrack RL, Maloney WJ, Clohisey JC. What is the prognosis of revision total hip arthroplasty in patients 55 years and younger? *Clin Orthop Relat Res.* 2014; 472(5):1518–25. doi: [10.1007/s11999-013-3377-9](https://doi.org/10.1007/s11999-013-3377-9) PMID: [24249534](https://pubmed.ncbi.nlm.nih.gov/24249534/); PubMed Central PMCID: [PMC3971212](https://pubmed.ncbi.nlm.nih.gov/PMC3971212/).

5. Espehaug B, Havelin LI, Engesaeter LB, Langeland N, Vollset SE. Patient satisfaction and function after primary and revision total hip replacement. *Clin Orthop Relat Res*. 1998;(351):135–48. PMID: [9646756](#).
6. Field RE, Cronin MD, Singh PJ. The Oxford hip scores for primary and revision hip replacement. *J Bone Joint Surg Br*. 2005; 87(5):618–22. doi: [10.1302/0301-620X.87B5.15390](#) PMID: [15855360](#).
7. Patil S, Garbuz DS, Greidanus NV, Masri BA, Duncan CP. Quality of life outcomes in revision vs primary total hip arthroplasty: a prospective cohort study. *J Arthroplasty*. 2008; 23(4):550–3. doi: [10.1016/j.arth.2007.04.035](#) PMID: [18514873](#).
8. Robinson AH, Palmer CR, Villar RN. Is revision as good as primary hip replacement? A comparison of quality of life. *J Bone Joint Surg Br*. 1999; 81(1):42–5. PMID: [10068000](#).
9. Davis AM, Agnidis Z, Badley E, Kiss A, Waddell JP, Gross AE. Predictors of functional outcome two years following revision hip arthroplasty. *J Bone Joint Surg Am*. 2006; 88(4):685–91. doi: [10.2106/JBJS.E.00150](#) PMID: [16595456](#).
10. Atroshi I, Ornstein E, Franzen H, Johnsson R, Stefansdottir A, Sundberg M. Quality of life after hip revision with impaction bone grafting on a par with that 4 years after primary cemented arthroplasty. *Acta Orthop Scand*. 2004; 75(6):677–83. PMID: [15762256](#).
11. Lenguerrand E, Wylde V, Goberman-Hill R, Sayers A, Brunton L, Beswick AD, et al. Trajectories of Pain and Function after Primary Hip and Knee Arthroplasty: The ADAPT Cohort Study. *PLoS One*. 2016; 11(2):e0149306. doi: [10.1371/journal.pone.0149306](#) PMID: [26871909](#).
12. Wylde V, Blom AW, Bolink S, Brunton L, Dieppe P, Goberman-Hill R, et al. Assessing function in patients undergoing joint replacement: a study protocol for a cohort study. *BMC Musculoskelet Disord*. 2012; 13:220. doi: [10.1186/1471-2474-13-220](#) PMID: [23148591](#); PubMed Central PMCID: [PMC3520823](#).
13. Bellamy N, Buchanan WW, Goldsmith CH, Campbell J, Stitt LW. Validation study of WOMAC: a health status instrument for measuring clinically important patient relevant outcomes to antirheumatic drug therapy in patients with osteoarthritis of the hip or knee. *J Rheumatol*. 1988; 15:1833–40. PMID: [3068365](#)
14. McConnell S, Kolopack P, Davis AM. The Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC): a review of its utility and measurement properties. *Arthritis Rheum*. 2001; 45(5):453–61. PMID: [11642645](#).
15. Senden R, Grimm B, Heyligers IC, Savelberg HH, Meijer K. Acceleration-based gait test for healthy subjects: reliability and reference data. *Gait & posture*. 2009; 30(2):192–6. doi: [10.1016/j.gaitpost.2009.04.008](#) PMID: [19477649](#).
16. Gill S, McBurney H. Reliability of performance-based measures in people awaiting joint replacement surgery of the hip or knee. *Physiother Res Int*. 2008; 13(3):141–52. doi: [10.1002/pri.411](#) PMID: [18697226](#); PubMed Central PMCID: [PMC10.1002/pri.411](#).
17. Kennedy DM, Stratford PW, Wessel J, Gollish JD, Penney D. Assessing stability and change of four performance measures: a longitudinal study evaluating outcome following total hip and knee arthroplasty. *BMC Musculoskelet Disord*. 2005; 6:3. PMID: [15679884](#); PubMed Central PMCID: [PMCPMC549207](#) [10.1186/1471-2474-6-3](#). doi: [10.1186/1471-2474-6-3](#)
18. Little RJA. Modeling the Drop-Out Mechanism in Repeated-Measures Studies. *Journal of the American Statistical Association*. 1995; 90:1112–21.
19. Leckie G, Charlton C. Runmlwin: a program to run the MLwiN multilevel modelling software from within Stata. *Journal of Statistical Software*. 2013;(52):1–40.
20. Aghayev E, Teuscher R, Neukamp M, Lee EJ, Melloh M, Egli S, et al. The course of radiographic loosening, pain and functional outcome around the first revision of a total hip arthroplasty. *BMC Musculoskelet Disord*. 2013; 14:167. doi: [10.1186/1471-2474-14-167](#) PMID: [23672339](#); PubMed Central PMCID: [PMCPMC3658875](#).
21. Barlow BT, Boles JW, Lee YY, Ortiz PA, Westrich GH. Short-Term Outcomes and Complications After Rejuvenate Modular Total Hip Arthroplasty Revision. *J Arthroplasty*. 2015. doi: [10.1016/j.arth.2015.10.041](#) PMID: [26777570](#).
22. Biring GS, Masri BA, Greidanus NV, Duncan CP, Garbuz DS. Predictors of quality of life outcomes after revision total hip replacement. *J Bone Joint Surg Br*. 2007; 89(11):1446–51. doi: [10.1302/0301-620X.89B11.19617](#) PMID: [17998179](#).
23. Izquierdo RJ, Northmore-Ball MD. Long-term results of revision hip arthroplasty. Survival analysis with special reference to the femoral component. *J Bone Joint Surg Br*. 1994; 76(1):34–9. PMID: [8300678](#).
24. Romano CL, Romano D, Logoluso N, Meani E. Septic versus aseptic hip revision: how different? *J Orthop Traumatol*. 2010; 11(3):167–74. doi: [10.1007/s10195-010-0106-y](#) PMID: [20811922](#); PubMed Central PMCID: [PMCPMC2948127](#).

25. Saleh KJ, Celebrezze M, Kassim R, Dykes DC, Gioe TJ, Callaghan JJ, et al. Functional outcome after revision hip arthroplasty: a metaanalysis. *Clin Orthop Relat Res*. 2003;(416):254–64. doi: [10.1097/01.blo.0000093006.90435.43](https://doi.org/10.1097/01.blo.0000093006.90435.43) PMID: [14646768](https://pubmed.ncbi.nlm.nih.gov/14646768/).
26. Lubbeke A, Katz JN, Perneger TV, Hoffmeyer P. Primary and revision hip arthroplasty: 5-year outcomes and influence of age and comorbidity. *J Rheumatol*. 2007; 34(2):394–400. PMID: [17143967](https://pubmed.ncbi.nlm.nih.gov/17143967/).
27. Singh JA, Lewallen DG. Patient-level clinically meaningful improvements in activities of daily living and pain after total hip arthroplasty: data from a large US institutional registry. *Rheumatology (Oxford)*. 2013; 52(6):1109–18. doi: [10.1093/rheumatology/kes416](https://doi.org/10.1093/rheumatology/kes416) PMID: [23382362](https://pubmed.ncbi.nlm.nih.gov/23382362/); PubMed Central PMCID: PMC3651614.
28. Stevens M, Hoekstra T, Wagenmakers R, Bulstra SK, van den Akker-Scheek I. People who undergo revision arthroplasty report more limitations but no decrease in physical activity compared with primary total hip arthroplasty: an observational study. *Aust J Physiother*. 2009; 55(3):185–9. PMID: [19681740](https://pubmed.ncbi.nlm.nih.gov/19681740/).
29. Quintana JM, Escobar A, Bilbao A, Arostegui I, Lafuente I, Vidaurreta I. Responsiveness and clinically important differences for the WOMAC and SF-36 after hip joint replacement. *Osteoarthritis Cartilage*. 2005; 13(12):1076–83. doi: [10.1016/j.joca.2005.06.012](https://doi.org/10.1016/j.joca.2005.06.012) PMID: [16154777](https://pubmed.ncbi.nlm.nih.gov/16154777/).
30. Ramisetty N, Kwon Y, Mohtadi N. Patient-reported outcome measures for hip preservation surgery—a systematic review of the literature. *Journal of Hip Preservation Surgery Advance*. 2015; 0(0):1–13.
31. Garbuz DS, Xu M, Sayre EC. Patients' outcome after total hip arthroplasty: a comparison between the Western Ontario and McMaster Universities index and the Oxford 12-item hip score. *J Arthroplasty*. 2006; 21(7):998–1004. doi: [10.1016/j.arth.2006.01.014](https://doi.org/10.1016/j.arth.2006.01.014) PMID: [17027542](https://pubmed.ncbi.nlm.nih.gov/17027542/).
32. Nilsson A, Bremander A. Measures of hip function and symptoms: Harris Hip Score (HHS), Hip Disability and Osteoarthritis Outcome Score (HOOS), Oxford Hip Score (OHS), Lequesne Index of Severity for Osteoarthritis of the Hip (LISOH), and American Academy of Orthopedic Surgeons (AAOS) Hip and Knee Questionnaire. *Arthritis Care Res (Hoboken)*. 2011; 63 Suppl 11:S200–7. doi: [10.1002/acr.20549](https://doi.org/10.1002/acr.20549) PMID: [22588745](https://pubmed.ncbi.nlm.nih.gov/22588745/).
33. Judge A, Cooper C, Williams S, Dreinhoefer K, Dieppe P. Patient-reported outcomes one year after primary hip replacement in a European Collaborative Cohort. *Arthritis care & research*. 2010; 62(4):480–8. doi: [10.1002/acr.20038](https://doi.org/10.1002/acr.20038) PMID: [20391502](https://pubmed.ncbi.nlm.nih.gov/20391502/).
34. Singh JA, Lewallen D. Age, gender, obesity, and depression are associated with patient-related pain and function outcome after revision total hip arthroplasty. *Clin Rheumatol*. 2009; 28(12):1419–30. doi: [10.1007/s10067-009-1267-z](https://doi.org/10.1007/s10067-009-1267-z) PMID: [19727914](https://pubmed.ncbi.nlm.nih.gov/19727914/); PubMed Central PMCID: PMC2963019.
35. Singh JA, Lewallen DG. Operative diagnosis for revision total hip arthroplasty is associated with patient-reported outcomes (PROs). *BMC Musculoskelet Disord*. 2013; 14:210. doi: [10.1186/1471-2474-14-210](https://doi.org/10.1186/1471-2474-14-210) PMID: [23866848](https://pubmed.ncbi.nlm.nih.gov/23866848/); PubMed Central PMCID: PMC3722075.
36. Barrack RL, McClure JT, Burak CF, Clohisey JC, Parvizi J, Hozack W. Revision total hip arthroplasty: the patient's perspective. *Clin Orthop Relat Res*. 2006; 453:173–7. doi: [10.1097/01.blo.0000246537.67500.50](https://doi.org/10.1097/01.blo.0000246537.67500.50) PMID: [17016214](https://pubmed.ncbi.nlm.nih.gov/17016214/).
37. Haddad FS, Garbuz DS, Chambers GK, Jagpal TJ, Masri BA, Duncan CP. The expectations of patients undergoing revision hip arthroplasty. *J Arthroplasty*. 2001; 16(1):87–91. doi: [10.1054/arth.2001.17937](https://doi.org/10.1054/arth.2001.17937) PMID: [11172276](https://pubmed.ncbi.nlm.nih.gov/11172276/).
38. Eisler T, Svensson O, Tengstrom A, Elmstedt E. Patient expectation and satisfaction in revision total hip arthroplasty. *J Arthroplasty*. 2002; 17(4):457–62. PMID: [12066276](https://pubmed.ncbi.nlm.nih.gov/12066276/).