

ADOPTED: 21 November 2019

doi: 10.2903/j.efsa.2020.5930

List of non-EU phytoplasmas of *Cydonia* Mill., *Fragaria* L., *Malus* Mill., *Prunus* L., *Pyrus* L., *Ribes* L., *Rubus* L. and *Vitis* L.

EFSA Panel on Plant Health (PLH),

Claude Bragard, Katharina Dehnen-Schmutz, Paolo Gonthier, Josep Anton Jaques Miret, Annemarie Fejer Justesen, Alan MacLeod, Christer Sven Magnusson, Panagiotis Milonas, Juan A Navas-Cortes, Stephen Parnell, Roel Potting, Philippe Lucien Reignault, Hans-Hermann Thulke, Wopke Van der Werf, Antonio Vicent Civera, Jonathan Yuen, Lucia Zappalà, Domenico Bosco, Michela Chiumenti, Francesco Di Serio, Luciana Galetto, Cristina Marzachi, Marco Pautasso and Marie-Agnès Jacques

Abstract

Following a request from the European Commission, the EFSA Panel on Plant Health prepared a list of non-EU phytoplasmas of *Cydonia* Mill., *Fragaria* L., *Malus* Mill., *Prunus* L., *Pyrus* L., *Ribes* L., *Rubus* L. and *Vitis* L. A systematic literature review and search of databases identified 27 phytoplasmas infecting one or more of the host genera under consideration. These phytoplasmas were assigned to three categories. The first group (a) consists of 10 non-EU phytoplasmas, known to occur only outside the EU ('*Candidatus* Phytoplasma australiense', '*Ca. P. hispanicum*', '*Ca. P. pruni*'-related strain (NAGYIII), '*Ca. P. pyri*'-related strain (PYLR) and Buckland valley grapevine yellows phytoplasma) or having only limited presence in the EU ('*Ca. P. aurantifolia*'-related strains, '*Ca. P. fraxini*', '*Ca. P. phoenicium*', '*Ca. P. trifolii*' and '*Ca. P. ziziphi*'). The second group (b) consists of three non-EU phytoplasmas, whose presence in the target plant species is not fully supported by the available literature. The third group (c) consists of 14 phytoplasmas with substantial presence in the EU (i.e. they are originally described or reported from the EU or known to occur or be widespread in some EU Member States or frequently reported in the EU). Phytoplasmas of categories (b) and (c) were excluded at this stage from further categorisation efforts. One phytoplasma from category (a) ('*Ca. P. phoenicium*') was excluded from further categorisation, as a pest risk assessment has been performed by EPPO. Comments provided by the EU Member States were integrated in the opinion. The main uncertainties of this listing concern: the geographic distribution and prevalence, the taxonomy, biology and host range. The phytoplasmas considered as non-EU and whose presence in target plant species is fully supported by literature (category (a)) are categorised by the Panel in a separate opinion.

© 2020 European Food Safety Authority. *EFSA Journal* published by John Wiley and Sons Ltd on behalf of European Food Safety Authority.

Keywords: *Crotalaria* witches' broom phytoplasma, North American grapevine yellows, peach yellow leaf roll, pear decline, pest risk, plant health, plant pest, quarantine, sweet potato little leaf

Requestor: European Commission

Question numbers: EFSA-Q-2018-00272, EFSA-Q-2018-00633, EFSA-Q-2018-00634, EFSA-Q-2018-00635, EFSA-Q-2018-00636, EFSA-Q-2018-00637, EFSA-Q-2018-00638, EFSA-Q-2018-00639

Correspondence: alpha@efsa.europa.eu

Panel members: Claude Bragard, Katharina Dehnen-Schmutz, Francesco Di Serio, Paolo Gonthier, Marie-Agnès Jacques, Josep Anton Jaques Miret, Annemarie Fejer Justesen, Alan MacLeod, Christer Sven Magnusson, Panagiotis Milonas, Juan A Navas-Cortes, Stephen Parnell, Roel Potting, Philippe L Reignault, Hans-Hermann Thulke, Wopke Van der Werf, Antonio Vicent Civera, Jonathan Yuen and Lucia Zappalà.

Acknowledgments: This document was prepared in cooperation with the Istituto per la Protezione Sostenibile delle Piante, Consiglio Nazionale delle Ricerche (Italy) under the tasking grant (GP/EFSA/ALPHA/2017/02). The Panel thanks for the information provided to this scientific output: Franco Finelli (Phytosanitary Service, Italy). The Panel acknowledges all European competent institutions, Member State bodies and other organisations that provided data for this scientific output.

Competing interests: In line with EFSA's policy on declarations of interest, Panel member Francesco Di Serio did not participate in the adoption of this scientific output.

Suggested citation: EFSA PLH Panel (EFSA Panel on Plant Health), Bragard C, Dehnen-Schmutz K, Gonthier P, Jaques Miret JA, Justesen AF, MacLeod A, Magnusson CS, Milonas P, Navas-Cortes JA, Parnell S, Potting R, Reignault PL, Thulke H-H, Van der Werf W, Vicent Civera A, Yuen J, Zappalà L, Bosco D, Chiumenti M, Di Serio F, Galetto L, Marzachi C, Pautasso M and Jacques M-A, 2020. Scientific Opinion on the list of non-EU phytoplasmas of *Cydonia* Mill., *Fragaria* L., *Malus* Mill., *Prunus* L., *Pyrus* L., *Ribes* L., *Rubus* L. and *Vitis* L. EFSA Journal 2020;18(1):5930, 25 pp. <https://doi.org/10.2903/j.efsa.2020.5930>

ISSN: 1831-4732

© 2020 European Food Safety Authority. *EFSA Journal* published by John Wiley and Sons Ltd on behalf of European Food Safety Authority.

This is an open access article under the terms of the [Creative Commons Attribution-NoDerivs](https://creativecommons.org/licenses/by/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited and no modifications or adaptations are made.



The EFSA Journal is a publication of the European Food Safety Authority, an agency of the European Union.



Table of contents

Abstract.....	1
1. Introduction.....	4
1.1. Background and Terms of Reference as provided by the requestor.....	4
1.1.1. Background	4
1.1.2. Terms of Reference	4
1.1.2.1. Terms of Reference: Appendix 1.....	5
1.1.2.2. Terms of Reference: Appendix 2.....	6
1.1.2.3. Terms of Reference: Appendix 3.....	7
1.1.3. Interpretation of the Terms of Reference.....	8
2. Data and methodologies	9
2.1. Data.....	9
2.1.1. Literature search	9
2.1.2. Database search	9
2.2. Methodology	9
3. Listing of phytoplasmas	10
3.1. Phytoplasmas considered as non-EU	10
3.2. Phytoplasmas excluded from further categorisation in the frame of the present mandate	10
3.3. Uncertainties.....	10
4. Conclusions.....	10
References.....	11
Abbreviations	14
Appendix A – Non-EU phytoplasmas of <i>Cydonia</i> , <i>Fragaria</i> , <i>Malus</i> , <i>Prunus</i> , <i>Pyrus</i> , <i>Ribes</i> , <i>Rubus</i> and <i>Vitis</i>	16
Appendix B – Phytoplasmas of <i>Cydonia</i> , <i>Fragaria</i> , <i>Malus</i> , <i>Prunus</i> , <i>Pyrus</i> , <i>Ribes</i> , <i>Rubus</i> and <i>Vitis</i> excluded from further categorisation as their presence in the species is not fully supported by available literature	19
Appendix C – Phytoplasmas of <i>Cydonia</i> , <i>Fragaria</i> , <i>Malus</i> , <i>Prunus</i> , <i>Pyrus</i> , <i>Ribes</i> , <i>Rubus</i> and <i>Vitis</i> excluded from further categorisation	20
Annex A – List of phytoplasmas considered in the opinion	25

1. Introduction

1.1. Background and Terms of Reference as provided by the requestor

1.1.1. Background

Council Directive 2000/29/EC¹ on protective measures against the introduction into the Community of organisms harmful to plants or plant products and against their spread within the Community establishes the present European Union plant health regime. The Directive lays down the phytosanitary provisions and the control checks to be carried out at the place of origin on plants and plant products destined for the Union or to be moved within the Union. In the Directive's 2000/29/EC annexes, the list of harmful organisms (pests) whose introduction into or spread within the Union is prohibited, is detailed together with specific requirements for import or internal movement.

Following the evaluation of the plant health regime, the new basic plant health law, Regulation (EU) 2016/2031² on protective measures against pests of plants, was adopted on 26 October 2016 and will apply from 14 December 2019 onwards, repealing Directive 2000/29/EC. In line with the principles of the above mentioned legislation and the follow-up work of the secondary legislation for the listing of EU regulated pests, EFSA is requested to provide pest categorisations of the harmful organisms included in the annexes of Directive 2000/29/EC, in the cases where recent pest risk assessment/pest categorisation is not available.

1.1.2. Terms of Reference

EFSA is requested, pursuant to Article 22(5.b) and Article 29(1) of Regulation (EC) No 178/2002³, to provide scientific opinion in the field of plant health.

EFSA is requested to prepare and deliver a pest categorisation (step 1 analysis) for each of the regulated pests included in the appendices of the annex to this mandate. The methodology and template of pest categorisation have already been developed in past mandates for the organisms listed in Annex II Part A Section II of Directive 2000/29/EC. The same methodology and outcome is expected for this work as well.

The list of the harmful organisms included in the annex to this mandate comprises 133 harmful organisms or groups. A pest categorisation is expected for these 133 pests or groups and the delivery of the work would be stepwise at regular intervals through the year as detailed below. First priority covers the harmful organisms included in Appendix 1, comprising pests from Annex II Part A Section I and Annex II Part B of Directive 2000/29/EC. The delivery of all pest categorisations for the pests included in Appendix 1 is June 2018. The second priority is the pests included in Appendix 2, comprising the group of *Cicadellidae* (non-EU) known to be vector of Pierce's disease (caused by *Xylella fastidiosa*), the group of *Tephritidae* (non-EU), the group of potato viruses and virus-like organisms, the group of viruses and virus-like organisms of *Cydonia* Mill., *Fragaria* L., *Malus* Mill., *Prunus* L., *Pyrus* L., *Ribes* L., *Rubus* L. and *Vitis* L. and the group of *Margarodes* (non-EU species). The delivery of all pest categorisations for the pests included in Appendix 2 is end 2019. The pests included in Appendix 3 cover pests of Annex I part A section I and all pests categorisations should be delivered by end 2020.

For the above-mentioned groups, each covering a large number of pests, the pest categorisation will be performed for the group and not the individual harmful organisms listed under "such as" notation in the Annexes of the Directive 2000/29/EC. The criteria to be taken particularly under consideration for these cases, is the analysis of host pest combination, investigation of pathways, the damages occurring and the relevant impact.

Finally, as indicated in the text above, all references to 'non-European' should be avoided and replaced by 'non-EU' and refer to all territories with exception of the Union territories as defined in Article 1 point 3 of Regulation (EU) 2016/2031.

¹ Council Directive 2000/29/EC of 8 May 2000 on protective measures against the introduction into the Community of organisms harmful to plants or plant products and against their spread within the Community. OJ L 169/1, 10.7.2000, p. 1–112.

² Regulation (EU) 2016/2031 of the European Parliament of the Council of 26 October 2016 on protective measures against pests of plants. OJ L 317, 23.11.2016, p. 4–104.

³ Regulation (EC) No 178/2002 of the European Parliament and of the Council of 28 January 2002 laying down the general principles and requirements of food law, establishing the European Food Safety Authority and laying down procedures in matters of food safety. OJ L 31/1, 1.2.2002, p. 1–24.

1.1.2.1. Terms of Reference: Appendix 1

List of harmful organisms for which a pest categorisation is requested. The list below follows the annexes of Directive 2000/29/EC.

Annex IIAI

(a) Insects, mites and nematodes, at all stages of their development

<i>Aleurocanthus</i> spp.	<i>Numonia pyrivorella</i> (Matsumura)
<i>Anthonomus bisignifer</i> (Schenkling)	<i>Oligonychus perditus</i> Pritchard and Baker
<i>Anthonomus signatus</i> (Say)	<i>Pissodes</i> spp. (non-EU)
<i>Aschistonyx eppoi</i> Inouye	<i>Scirtothrips aurantii</i> Faure
<i>Carposina niponensis</i> Walsingham	<i>Scirtothrips citri</i> (Moultex)
<i>Enarmonia packardi</i> (Zeller)	<i>Scolytidae</i> spp. (non-EU)
<i>Enarmonia prunivora</i> Walsh	<i>Scrobipalopsis solanivora</i> Povolny
<i>Grapholita inopinata</i> Heinrich	<i>Tachypterellus quadrigibbus</i> Say
<i>Hishomonus phycitis</i>	<i>Toxoptera citricida</i> Kirk.
<i>Leucaspis japonica</i> Ckll.	<i>Unaspis citri</i> Comstock
<i>Listronotus bonariensis</i> (Kuschel)	

(b) Bacteria

Citrus variegated chlorosis	<i>Xanthomonas campestris</i> pv. <i>oryzae</i> (Ishiyama)
<i>Erwinia stewartii</i> (Smith) Dye	Dye and pv. <i>oryzicola</i> (Fang, et al.) Dye

(c) Fungi

<i>Alternaria alternata</i> (Fr.) Keissler (non-EU pathogenic isolates)	<i>Elsinoe</i> spp. Bitanc. and Jenk. Mendes
<i>Anisogramma anomala</i> (Peck) E. Müller	<i>Fusarium oxysporum</i> f. sp. <i>albedinis</i> (Kilian and Maire) Gordon
<i>Apiosporina morbosa</i> (Schwein.) v. Arx	<i>Guignardia piricola</i> (Nosa) Yamamoto
<i>Ceratocystis virescens</i> (Davidson) Moreau	<i>Puccinia pittieriana</i> Hennings
<i>Cercoseptoria pini-densiflorae</i> (Hori and Nambu) Deighton	<i>Stegophora ulmea</i> (Schweinitz: Fries) Sydow & Sydow
<i>Cercospora angolensis</i> Carv. and Mendes	<i>Venturia nashicola</i> Tanaka and Yamamoto

(d) Virus and virus-like organisms

Beet curly top virus (non-EU isolates)	Little cherry pathogen (non- EU isolates)
Black raspberry latent virus	Naturally spreading psorosis
Blight and blight-like	Palm lethal yellowing mycoplasma
Cadang-Cadang viroid	Satsuma dwarf virus
Citrus tristeza virus (non-EU isolates)	Tatter leaf virus
Leprosis	Witches' broom (MLO)

Annex IIB

(a) Insect mites and nematodes, at all stages of their development

<i>Anthonomus grandis</i> (Boh.)	<i>Ips cembrae</i> Heer
<i>Cephalcia lariciphila</i> (Klug)	<i>Ips duplicatus</i> Sahlberg
<i>Dendroctonus micans</i> Kugelan	<i>Ips sexdentatus</i> Börner
<i>Gilpinia hercyniae</i> (Hartig)	<i>Ips typographus</i> Heer
<i>Gonipterus scutellatus</i> Gyll.	<i>Sternochetus mangiferae</i> Fabricius
<i>Ips amitinus</i> Eichhof	

(b) Bacteria

Curtobacterium flaccumfaciens pv. *flaccumfaciens*
(Hedges) Collins and Jones

(c) Fungi

Glomerella gossypii Edgerton

Hypoxyton mammatum (Wahl.) J. Miller

Gremmeniella abietina (Lag.) Morelet

1.1.2.2. Terms of Reference: Appendix 2

List of harmful organisms for which a pest categorisation is requested per group. The list below follows the categorisation included in the annexes of Directive 2000/29/EC.

Annex IAI**(a) Insects, mites and nematodes, at all stages of their development**

Group of Cicadellidae (non-EU) known to be vector of Pierce's disease (caused by *Xylella fastidiosa*), such as:

- | | |
|--|---|
| 1) <i>Carneocephala fulgida</i> Nottingham | 3) <i>Graphocephala atropunctata</i> (Signoret) |
| 2) <i>Draeculacephala minerva</i> Ball | |

Group of Tephritidae (non-EU) such as:

- | | |
|--|---|
| 1) <i>Anastrepha fraterculus</i> (Wiedemann) | 12) <i>Pardalaspis cyanescens</i> Bezzi |
| 2) <i>Anastrepha ludens</i> (Loew) | 13) <i>Pardalaspis quinaria</i> Bezzi |
| 3) <i>Anastrepha obliqua</i> Macquart | 14) <i>Pterandrus rosa</i> (Karsch) |
| 4) <i>Anastrepha suspensa</i> (Loew) | 15) <i>Rhacochlaena japonica</i> Ito |
| 5) <i>Dacus ciliatus</i> Loew | 16) <i>Rhagoletis completa</i> Cresson |
| 6) <i>Dacus curcurbitae</i> Coquillett | 17) <i>Rhagoletis fausta</i> (Osten-Sacken) |
| 7) <i>Dacus dorsalis</i> Hendel | 18) <i>Rhagoletis indifferens</i> Curran |
| 8) <i>Dacus tryoni</i> (Froggatt) | 19) <i>Rhagoletis mendax</i> Curran |
| 9) <i>Dacus tsuneonis</i> Miyake | 20) <i>Rhagoletis pomonella</i> Walsh |
| 10) <i>Dacus zonatus</i> Saund. | 21) <i>Rhagoletis suavis</i> (Loew) |
| 11) <i>Epochra canadensis</i> (Loew) | |

(c) Viruses and virus-like organisms

Group of potato viruses and virus-like organisms such as:

- | | |
|----------------------------------|--|
| 1) Andean potato latent virus | 4) Potato black ringspot virus |
| 2) Andean potato mottle virus | 5) Potato virus T |
| 3) Arracacha virus B, oca strain | 6) non-EU isolates of potato viruses A, M, S, V, X and Y (including Yo, Yn and Yc) and Potato leafroll virus |

Group of viruses and virus-like organisms of *Cydonia* Mill., *Fragaria* L., *Malus* Mill., *Prunus* L., *Pyrus* L., *Ribes* L., *Rubus* L. and *Vitis* L., such as:

- | | |
|--------------------------------------|--|
| 1) Blueberry leaf mottle virus | 8) Peach yellows mycoplasma |
| 2) Cherry rasp leaf virus (American) | 9) Plum line pattern virus (American) |
| 3) Peach mosaic virus (American) | 10) Raspberry leaf curl virus (American) |
| 4) Peach phony rickettsia | 11) Strawberry witches' broom mycoplasma |
| 5) Peach rosette mosaic virus | 12) Non-EU viruses and virus-like organisms of <i>Cydonia</i> Mill., <i>Fragaria</i> L., <i>Malus</i> Mill., <i>Prunus</i> L., <i>Pyrus</i> L., <i>Ribes</i> L., <i>Rubus</i> L. and <i>Vitis</i> L. |
| 6) Peach rosette mycoplasma | |
| 7) Peach X-disease mycoplasma | |

Annex IIAI

(a) Insects, mites and nematodes, at all stages of their development

Group of *Margarodes* (non-EU species) such as:

- | | |
|--|--|
| 1) <i>Margarodes vitis</i> (Phillipi) | 3) <i>Margarodes prieskaensis</i> Jakubski |
| 2) <i>Margarodes vredendalensis</i> de Klerk | |

1.1.2.3. Terms of Reference: Appendix 3

List of harmful organisms for which a pest categorisation is requested. The list below follows the annexes of Directive 2000/29/EC.

Annex IAI

(a) Insects, mites and nematodes, at all stages of their development

<i>Acleris</i> spp. (non-EU)	<i>Longidorus diadecturus</i> Eveleigh and Allen
<i>Amauromyza maculosa</i> (Malloch)	<i>Monochamus</i> spp. (non-EU)
<i>Anomala orientalis</i> Waterhouse	<i>Myndus crudus</i> Van Duzee
<i>Arrhenodes minutus</i> Drury	<i>Nacobbus aberrans</i> (Thorne) Thorne and Allen
<i>Choristoneura</i> spp. (non-EU)	<i>Naupactus leucoloma</i> Boheman
<i>Conotrachelus nenuphar</i> (Herbst)	<i>Premnotrypes</i> spp. (non-EU)
<i>Dendrolimus sibiricus</i> Tschetverikov	<i>Pseudopityophthorus minutissimus</i> (Zimmermann)
<i>Diabrotica barberi</i> Smith and Lawrence	<i>Pseudopityophthorus pruinosus</i> (Eichhoff)
<i>Diabrotica undecimpunctata howardi</i> Barber	<i>Scaphoideus luteolus</i> (Van Duzee)
<i>Diabrotica undecimpunctata undecimpunctata</i> Mannerheim	<i>Spodoptera eridania</i> (Cramer)
<i>Diabrotica virgifera zea</i> Krysan & Smith	<i>Spodoptera frugiperda</i> (Smith)
<i>Diaphorina citri</i> Kuway	<i>Spodoptera litura</i> (Fabricus)
<i>Heliothis zea</i> (Boddie)	<i>Thrips palmi</i> Karny
<i>Hirschmanniella</i> spp., other than <i>Hirschmanniella</i> <i>gracilis</i> (de Man) Luc and Goodey	<i>Xiphinema americanum</i> Cobb <i>sensu lato</i> (non-EU populations)
<i>Liriomyza sativae</i> Blanchard	<i>Xiphinema californicum</i> Lamberti and Bleve-Zacheo

(b) Fungi

<i>Ceratocystis fagacearum</i> (Bretz) Hunt	<i>Mycosphaerella larici-leptolepis</i> Ito et al.
<i>Chrysomyxa arctostaphyli</i> Dietel	<i>Mycosphaerella populorum</i> G. E. Thompson
<i>Cronartium</i> spp. (non-EU)	<i>Phoma andina</i> Turkensteen
<i>Endocronartium</i> spp. (non-EU)	<i>Phyllosticta solitaria</i> Ell. and Ev.
<i>Guignardia laricina</i> (Saw.) Yamamoto and Ito	<i>Septoria lycopersici</i> Speg. var. <i>malagutii</i> Ciccarone and Boerema
<i>Gymnosporangium</i> spp. (non-EU)	<i>Thecaphora solani</i> Barrus
<i>Inonotus weirii</i> (Murril) Kotlaba and Pouzar	<i>Trechispora brinkmannii</i> (Bresad.) Rogers
<i>Melampsora farlowii</i> (Arthur) Davis	

(c) Viruses and virus-like organisms

Tobacco ringspot virus	Pepper mild tigré virus
Tomato ringspot virus	Squash leaf curl virus
Bean golden mosaic virus	Euphorbia mosaic virus
Cowpea mild mottle virus	Florida tomato virus
Lettuce infectious yellows virus	

(d) Parasitic plants

Arceuthobium spp. (non-EU)

Annex I A II

(a) Insects, mites and nematodes, at all stages of their development

Meloidogyne fallax Karssen

Rhizoecus hibisci Kawai and Takagi

Popillia japonica Newman

(b) Bacteria

Clavibacter michiganensis (Smith) Davis et al. ssp. *Ralstonia solanacearum* (Smith) Yabuuchi et al. *sepedonicus* (Spieckermann and Kotthoff) Davis et al.

(c) Fungi

Melampsora medusae Thümen

Synchytrium endobioticum (Schilbersky) Percival

Annex I B

(a) Insects, mites and nematodes, at all stages of their development

Leptinotarsa decemlineata Say

Liriomyza bryoniae (Kaltenbach)

(b) Viruses and virus-like organisms

Beet necrotic yellow vein virus

1.1.3. Interpretation of the Terms of Reference

This opinion provides a list of non-EU phytoplasmas of *Cydonia* Mill., *Fragaria* L., *Malus* Mill., *Prunus* L., *Pyrus* L., *Ribes* L., *Rubus* L. and *Vitis* L. (from now on: 'the host plants'), for which the EFSA Plant Health Panel (from now on: 'the Panel') then conducted a pest categorisation in a separate opinion (EFSA PLH Panel, 2020). This list is based on information collected from databases up to June 2018, as well as information received from EU Member States (MS) during the period February-March 2019.

Non-EU phytoplasmas of the host plants are pests listed in the Appendices to the Terms of Reference (ToR) to be subject to pest categorisation to determine whether they fulfil the criteria of quarantine pests or those of regulated non-quarantine pests for the area of the EU excluding Ceuta, Melilla and the outermost regions of MS referred to in Article 355(1) of the Treaty on the Functioning of the European Union (TFEU), other than Madeira and the Azores.

As a first step towards this goal, the Panel prepared a list of phytoplasmas infecting the host plants. In the process, three groups of phytoplasmas were distinguished:

- (a) non-EU phytoplasmas with presence in the host plants fully supported by literature,
- (b) non-EU phytoplasmas with presence in the host plants not fully supported by literature, and
- (c) phytoplasmas (affecting the host plants) with widespread presence in the EU (known to occur in several MS, frequently reported in the EU, widespread in some MS) or originally described or reported from the EU.

A non-EU phytoplasma is defined by its geographical origin outside of the EU. Therefore, phytoplasmas not reported from the EU and occurring only outside of the EU are considered as non-EU phytoplasmas. Likewise, phytoplasmas occurring outside the EU and having only a limited presence in the EU (reported in only one or few MSs, with restricted distribution) are also considered as non-EU phytoplasmas.

This opinion provides the methodology and results for this classification, thus preparing the ground for the pest categorisation linked to the present mandate (EFSA PLH Panel, 2020). This means that the Panel then performed a pest categorisation for the non-EU phytoplasmas with confirmed ability to infect the host plants. The phytoplasmas with uncertain ability to infect the host plants and the phytoplasmas with significant presence in the EU or originally described or reported from the EU are excluded from further categorisation efforts, unless this will be requested by the risk managers in the future.

In this opinion, to capture the broadest possible range of phytoplasmas, even the poorly characterised ones for which very partial molecular or biological data are available, were considered. In particular, as in some cases there are uncertainties about the 'Ca. P. species definition', related strains will be considered if they infect any of the host plants. Instead, phytoplasma-like diseases of unknown

aetiology or caused by viruses and formerly associated to mycoplasma-like organisms (MLO) or by other graft-transmissible bacteria are not addressed in this opinion.

2. Data and methodologies

2.1. Data

2.1.1. Literature search

The literature considered to generate the list of phytoplasmas infecting the host plants (see Section 1.1.3) and to fill in the extraction tables on their distribution (see Appendixes A–C and Annex A) was obtained from expert knowledge and extensive literature searches performed in Web of Science (WoS, last access June 2018). For each host plant genus, searches in WoS were performed using as keywords: phytoplasma/mycoplasma/witch/spiroplasma combined with the scientific name of the genus OR the common name(s) of the crops. Therefore, for each host plant genus, searches in WoS were performed according to the following strategy:

TS=((Phytoplasma* OR mycoplasma* OR witch* OR spiroplasma*) AND (latin name of the host genus – e.g. *Vitis* – OR common name in English of the crop – e.g. grapevine))

All the references were screened by title, by abstract, and, if needed, by full paper with the specific objective of selecting those providing additional information regarding distribution and host range of the phytoplasmas included in the list or not yet included.

Extensive literature searches in Google Scholar using as keyword the name of a single phytoplasma were also performed for all the phytoplasmas listed in the EPPO Global Database (EPPO GD) (EPPO, 2019), and for all '*Candidatus* Phytoplasma species' described up to 2017 (Naderali et al., 2017).

Information on phytoplasma taxonomy was gathered from either the original reference to species description or IRPCM (International Research Programme on Comparative Mycoplasma) Phytoplasma/Spiroplasma Working Team–Phytoplasma Taxonomy Group (IRPCM, 2004).

Further references and data were obtained from experts, EU National Plant Protection Organisations and from citations within primary references.

2.1.2. Database search

Data on host(s) and distribution of the phytoplasmas were retrieved from the EPPO GD (EPPO, 2019), the Centre for Agriculture and Biosciences International Crop Protection Compendium (CABI, 2019) and relevant publications.

GenBank accessions referring to phytoplasmas were added.

2.2. Methodology

A preliminary list of phytoplasmas infecting the host plants (see Section 1.1.3) was generated by screening for phytoplasma diseases of the host plants present in the EPPO Lists A1 and A2. Further, all phytoplasma diseases listed in the EPPO GD were also screened for their association with the host plants. Then, all phytoplasmas listed in the most recent '*Ca. P. phytoplasma*' species description (Naderali et al., 2017) were screened for their association with the host plants. Finally, the relevant phytoplasmas resulting from the literature search in WoS (as previously described) were included in the list.

The collected information was used to fill an extraction table (Annex A) with data regarding the taxonomy, geographical distribution and host range of each phytoplasma and key references and sources used to obtain that information. Taxonomy, distribution and host range are reported in the table using the following scheme:

- the taxonomy was reported according to '*Ca. P. species*' description, when available. Although phytoplasmas have not yet been cultivated *in vitro*, phylogenetic analyses based on various conserved genes have shown that they represent a distinct, monophyletic clade within the class Mollicutes. Phytoplasmas are therefore accommodated within the '*Candidatus* Phytoplasma' genus. Within this genus, several subtaxa have been described to accommodate organisms sharing less than 97.5% similarity among their 16S rRNA gene sequences. Additional species are described to accommodate organisms that, despite their 16S rRNA gene sequence being > 97.5 % similar to those of other '*Ca. Phytoplasma*' species, are characterised by distinctive biological, phytopathological and genetic properties. Conversely, some organisms, despite their 16S rRNA gene sequence being < 97.5 % similar to that of any other '*Ca. Phytoplasma*' species, are not presently

described as *Candidatus* species, due to their poor overall characterisation (IRPCM, 2004). When a phytoplasma was not classified yet, information on a tentative classification was included based on the original literature source in which the pathogen was reported; to facilitate data retrieval from the literature and available databases, also the 16S rRNA group and subgroups were reported.

- data on distribution and host range of phytoplasmas were first searched in EPPO (2019) and in CABI (2019). Whenever conclusive information was not identified in the two databases or the information retrieved was at odds with expert knowledge, or in the absence of any information, extensive literature searches according to the protocol reported in Section 2.1 were performed.

Because only the non-EU phytoplasmas were subject of further categorisation efforts in the frame of the present mandate, it was decided to have consultation phases with EU MS so that they could provide additional input if necessary. The information provided by EU MS was then considered by the Panel to determine the non-EU phytoplasmas that were further categorised (Section 3.1). The phytoplasmas excluded from this group are referred to here as phytoplasmas excluded from further categorisation in the frame of the present mandate (Section 3.2).

3. Listing of phytoplasmas

3.1. Phytoplasmas considered as non-EU

The phytoplasmas considered as non-EU (Appendix A) belong to two sub-categories:

- Phytoplasmas not known to be present in the EU ('*Ca. P. australiense*', '*Ca. P. hispanicum*', '*Ca. P. pruni*'-related strain (NAGYIII), '*Ca. P. pyri*'-related strain (PYLR) and Buckland valley grapevine yellows phytoplasma)
- Phytoplasmas known to be present outside the EU and with only limited presence (i.e. reported in only one or few MSs or known to have a restricted distribution) in the EU ('*Ca. P. aurantifolia*'-related strains, '*Ca. P. fraxini*', '*Ca. P. phoenicium*', '*Ca. P. trifolii*' and '*Ca. P. ziziphi*').

These phytoplasmas are categorised in EFSA PLH Panel (2020), with the exception of '*Ca. P. phoenicium*', for which a pest risk assessment is already available (EPPO, 2017).

3.2. Phytoplasmas excluded from further categorisation in the frame of the present mandate

The phytoplasmas excluded from further categorisation in the frame of the present mandate are listed in Appendices B and C. The phytoplasmas listed in Appendix B are considered as non-EU, but their ability to infect the host plants was not conclusively supported by the available literature, and belong to two subcategories:

- Phytoplasmas not known to be present in the EU,
- Phytoplasmas known to be present outside the EU and with only limited presence (i.e. reported in only one or few MSs or known to have restricted distribution) in the EU.

Phytoplasmas listed in Appendix C belong to two sub-categories:

- Phytoplasmas originally described or reported from the EU,
- Phytoplasmas known to be present outside the EU, but with a substantial presence also in the EU (known to occur in several MSs, frequently reported in the EU, widespread in some MSs).

3.3. Uncertainties

Uncertainties potentially affecting the current list of non-EU phytoplasmas include:

- The geographic distribution and prevalence of the phytoplasmas.
- The taxonomy and biological status of poorly characterised phytoplasmas.
- The host status of particular plant genera for some phytoplasmas.

4. Conclusions

The Panel was requested by the European Commission to produce a categorisation of 133 harmful organisms or groups listed in annexes of Directive 2000/29/EC. One of the groups for which a categorisation was prepared is non-EU phytoplasmas of *Cydonia*, *Fragaria*, *Malus*, *Prunus*, *Pyrus*,

Ribes, *Rubus* and *Vitis*. As a first step, a systematic approach identified 27 phytoplasmas reported to naturally infect one or more of these genera (Annex A).

Among these phytoplasmas, based on information on distribution and prevalence both inside and outside the EU, the Panel identified 10 non-EU phytoplasmas, known to occur only outside the EU or having only a limited presence in the EU (Appendix A). These phytoplasmas are categorised in EFSA PLH Panel (2020), with the exception of '*Ca. P. phoenicium*', for which a pest risk assessment is already available (EPPO, 2017).

The remaining 17 phytoplasmas (non-EU phytoplasmas, known to occur only outside the EU or having only a limited presence in the EU, whose ability to infect the host species is not fully confirmed by available literature (Appendix B, 3 phytoplasmas), or which have a substantial presence in the EU or are originally described or reported from the EU (Appendix C, 14 phytoplasmas)) were not categorised within the current mandate. However, the European Commission may, at any time, request EFSA to categorise some or all the phytoplasmas excluded from the present exercise.

The main uncertainties of this listing concern the geographic distribution, taxonomy, biology and host range of some phytoplasmas.

References

- Alvizatos AS, 1993. Association of mycoplasma-like organisms with tomato big bud disease in Greece. *Plant Pathology*, 42, 158–162.
- Angelini E, Clair D, Borgo M, Bertaccini A and Boudon-Padieu E, 2001. Flavescence doree in France and Italy - occurrence of closely related phytoplasma isolates and their near relationships to Palatinate grapevine yellows and an alder yellows phytoplasma. *Vitis*, 40, 79–86.
- Arnaud G, Malembic-Maher S, Salar P, Bonnet P, Maixner M, Marcone C, Boudon-Padieu E and Foissac X, 2007. Multilocus sequence typing confirms the close genetic interrelatedness of three distinct flavescence doree phytoplasma strain clusters and group 16SrV phytoplasmas infecting grapevine and alder in Europe. *Applied and Environmental Microbiology*, 73, 4001–4010.
- Avramov Z, Contaldo N, Bertaccini A and Sakalieva D, 2011. First report of stolbur phytoplasmas in *Prunus avium* in Bulgaria. *Bulletin of Insectology*, 64, S71–S72.
- Balakishiyeva G, Qurbanov M, Mammadov A, Bayramov S, Aliyev J and Foissac X, 2011. Detection of '*Candidatus Phytoplasma brasiliense*' in a new geographic region and existence of two genetically distinct populations'. *European Journal of Plant Pathology*, 130, 457–462.
- Bertaccini A, Mittempergher L and Vibio M, 1996. Identification of phytoplasmas associated with a decline of European hackberry (*Celtis australis*). *Annals of Applied Biology*, 128, 245–253.
- Borroto Fernandez EG, Calari A, Hanzer V, Katinger H, Bertaccini A and Laimer M, 2007. Phytoplasma infected plants in Austrian forests: role as a reservoir? *Bulletin of Insectology*, 60, 391.
- Bruni R, Pellati F, Bellardi MG, Benvenuti S, Paltrinieri S, Bertaccini A and Bianchi A, 2005. Herbal drug quality and phytochemical composition of *Hypericum perforatum* L. affected by ash yellows phytoplasma infection. *Journal of Agricultural and Food Chemistry*, 53, 964–968.
- CABI, 2019. Crop Protection Compendium. Available online: <https://www.cabi.org/cpc/overview/> [Accessed: September 2019]
- Cieslinska M, 2011. Detection and characterization of phytoplasmas associated with diseases of *Rubus* spp. in Poland. *Journal of Plant Pathology*, 93, 51–56.
- Constable FE, Whiting JR, Gibb KS and Symons RH, 2002. A new grapevine yellows phytoplasma from the Buckland Valley of Victoria, Australia. *Vitis*, 41, 147–153.
- Constable FE, Gibb KS and Symons RH, 2003. Seasonal distribution of phytoplasmas in Australian grapevines. *Plant Pathology*, 52, 267–276.
- Davies DL, 2000. The occurrence of two phytoplasmas associated with stunted *Rubus* species in the UK. *Plant Pathology*, 49, 86–88.
- Davino S, Calari A, Davino M, Tessitor M, Bertaccini A and Bellardi MG, 2007. Virescence of tenweeks stock associated to phytoplasma infection in Sicily. *Bulletin of Insectology*, 60, 279–280.
- Davis RE, Dally EL, Gundersen DE, Lee IM and Habili N, 1997. '*Candidatus phytoplasma australiense*': a new phytoplasma taxon associated with Australian grapevine yellows. *International Journal of Systematic Bacteriology*, 47, 262–269.
- Davis RE, Zhao Y, Dally EL, Lee IM, Jomantiene R and Douglas SM, 2013. '*Candidatus Phytoplasma pruni*', a novel taxon associated with X-disease of stone fruits, *Prunus* spp.: multilocus characterization based on 16S rRNA, secY, and ribosomal protein genes. *International Journal of Systematic and Evolutionary Microbiology*, 63, 766–776.
- Davis RE, Dally EL, Zhao Y, Lee IM, Wei W, Wolf TK, Beanland L, LeDoux DG, Johnson DA, Fiola JA, Walter-Peterson H, Dami I and Chien M, 2015. Unraveling the etiology of North American Grapevine Yellows (NAGY): novel NAGY phytoplasma sequevars related to '*Candidatus Phytoplasma pruni*'. *Plant Disease*, 99, 1087–1097.

- Davis RE, Harrison NA, Zhao Y, Wei W and Dally EL, 2016. '*Candidatus* Phytoplasma hispanicum', a novel taxon associated with Mexican periwinkle virescence disease of *Catharanthus roseus*. *International Journal of Systematic and Evolutionary Microbiology*, 66, 3463–3467.
- Duduk B, Tian JB, Contaldo N, Fan XP, Paltrinieri S, Chen QF, Zhao QF and Bertaccini A, 2010. Occurrence of phytoplasmas related to stolbur and to '*Candidatus* Phytoplasma japonicum' in woody host plants in China. *Journal of Phytopathology*, 158, 100–104.
- EFSA PLH Panel (EFSA Panel on Plant Health), 2014. Scientific Opinion on pest categorisation of Grapevine Flavescence dorée. *EFSA Journal*, 12, 3851–3881. <https://doi.org/10.2903/j.efsa.2014.3851>
- EFSA PLH Panel (EFSA Panel on Plant Health), Bragard C, Dehnen-Schmutz K, Gonthier P, Jaques Miret JA, Fejer Justesen A, MacLeod A, Magnusson CS, Milonas P, Navas-Cortes JA, Parnell S, Potting R, Reignault PL, Thulke H-H, Van der Werf W, Vicent Civera A, Yuen J, Zappalà L, Bosco D, Chiumenti M, Di Serio F, Galetto L, Marzachi C, Pautasso M and Jacques M-A, 2020. Pest categorisation of the non-EU phytoplasmas of *Cydonia* Mill., *Fragaria* L., *Malus* Mill., *Prunus* L., *Pyrus* L., *Ribes* L., *Rubus* L. and *Vitis* L. *EFSA Journal* 2020;18(1):5929, 100 pp. <https://doi.org/10.2903/j.efsa.2020.5929>
- EPPO (European and Mediterranean Plant Protection Organization), 2017. Pest risk analysis for '*Candidatus* Phytoplasma phoenicium'. EPPO, Paris.
- EPPO (European and Mediterranean Plant Protection Organization), 2019. EPPO Global Database. Available online: <https://gd.eppo.int> [Accessed: September 2019]
- Ertunc F, Orel DC, Bayram S, Paltrinieri S, Bertaccini A, Topkaya S and Soylemezoglu G, 2015. Occurrence and identification of grapevine phytoplasmas in main viticultural regions of Turkey. *Phytoparasitica*, 43, 303–310.
- Fernandez FD, Conci VC, Kirschbaum DS and Conci LR, 2013. Molecular characterization of a phytoplasma of the ash yellows group occurring in strawberry (*Fragaria x ananassa* Duch.) plants in Argentina. *European Journal of Plant Pathology*, 135, 1–4.
- Fernandez FD, Meneguzzi NG, Guzman FA, Kirschbaum DS, Conci VC, Nome CF and Conci LR, 2015. Detection and identification of a novel 16SrXIII subgroup phytoplasma associated with strawberry red leaf disease in Argentina. *International Journal of Systematic and Evolutionary Microbiology*, 65, 2741–2747.
- Ferretti L, Costantini E and Pasquini G, 2015. Identification of 16SrIX-C phytoplasmas in *Argyranthemum frutescens* in Italy. *Phytopathologia Mediterranea*, 54, 21–27.
- Gajardo A, Fiore N, Prodan S, Paltrinieri S, Botti S, Pino AM, Zamorano A, Montealegre J and Bertaccini A, 2009. Phytoplasmas associated with grapevine yellows disease in Chile. *Plant Disease*, 93, 789–796.
- Ghayeb Zamharir M, Paltrinieri S, Hajivand S, Taheri M and Bertaccini A, 2017. Molecular identification of diverse '*Candidatus* Phytoplasma' species associated with grapevine decline in Iran. *Journal of Phytopathology*, 165, 407–413.
- Griffiths HM, Sinclair WA, Smart CD and Davis RE, 1999. The phytoplasma associated with ash yellows and lilac witches'-broom: '*Candidatus* Phytoplasma fraxini'. *International Journal of Systematic Bacteriology*, 49, 1605–1614.
- Harju VA, Skelton AL, Monger WA, Jarvis B and Mumford RA, 2008. Identification of an X-disease (16SrIII) group phytoplasma ('*Candidatus* Phytoplasma pruni') infecting delphiniums in the UK. *Plant Pathology*, 57, 769–769.
- Hashemi-Tameh M, Bahar M and Zirak L, 2014. '*Candidatus* Phytoplasma asteris' and '*Candidatus* Phytoplasma aurantifolia', new phytoplasma species infecting apple trees in Iran. *Journal of Phytopathology*, 162, 472–480.
- Hiruki C and Wang K, 2004. Clover proliferation phytoplasma: '*Candidatus* Phytoplasma trifolii'. *International Journal of Systematic and Evolutionary Microbiology*, 54, 1349–1353.
- IRPCM (International Research Programme on Comparative Mycoplasmaology), 2004. '*Candidatus* Phytoplasma', a taxon for the wall-less, non-helical prokaryotes that colonize plant phloem and insects. *International Journal of Systematic and Evolutionary Microbiology*, 54, 1243–1255.
- Jakovljevic M, Jovic J, Mitrovic M, Krstic O, Kosovac A, Tosevski I and Cvrkovic T, 2015. *Euscelis incisus* (Cicadellidae, Deltocephalinae), a natural vector of 16SrIII-B phytoplasma causing multiple inflorescence disease of *Cirsium arvense*. *Annals of Applied Biology*, 167, 406–419.
- Ježić M, Kontić JK, Preiner D, Maletić E and Curković-Perica M, 2013. Grapevine yellows affecting the Croatian indigenous grapevine cultivar Grk. *Acta Botanica Croatica*, 72, 287–294.
- Ježić M, Poljak I, Idžojtć M, Liber Z and Curković-Perica M, 2016. First report on phytoplasmas infecting wild apples and wild pears in Croatia. *Plant Disease*, 100, 207–208.
- Jomantiene R, Davis RE, Maas J and Dally EL, 1998. Classification of new phytoplasmas associated with diseases of strawberry in Florida, based on analysis of 16S rRNA and ribosomal protein gene operon sequences. *International Journal of Systematic Bacteriology*, 48, 269–277.
- Jomantiene R, Maas JL, Dally EL and Davis RE, 2002. Phytoplasmas in strawberry with fruit phyllody. *Acta Horticulturae*, 567, 635–638.
- Jones P and Arocha Y, 2006. A natural infection of Hebe is associated with an isolate of '*Candidatus* Phytoplasma asteris' causing a yellowing and little-leaf disease in the UK. *Plant Pathology*, 55, 821–821.
- Jones P, Arocha Y, Antesana O, Montilliano E and Franco P, 2005. First report of an isolate of '*Candidatus* Phytoplasma australiense' associated with a yellow leaf roll disease of peach (*Prunus persicae*) in Bolivia. *Plant Pathology*, 54, 558–558.

- Jung HY, Sawayanagi T, Kakizawa S, Nishigawa H, Wei W, Oshima K, Miyata S, Ugaki M, Hibi T and Namba S, 2003. '*Candidatus* Phytoplasma ziziphi', a novel phytoplasma taxon associated with jujube witches'-broom disease. *International Journal of Systematic and Evolutionary Microbiology*, 53, 1037–1041.
- Lee IM, Bertaccini A, Vibio M and Gundersen DE, 1995. Detection of multiple phytoplasmas in perennial fruit-trees with decline symptoms in Italy. *Phytopathology*, 85, 728–735.
- Lee IM, Gundersen-Rindal DE, Davis RE, Bottner KD, Marcone C and Seemüller E, 2004a. '*Candidatus* Phytoplasma asteris', a novel phytoplasma taxon associated with aster yellows and related diseases. *International Journal of Systematic and Evolutionary Microbiology*, 54, 1037–1048.
- Lee IM, Martini M, Marcone C and Zhu SF, 2004b. Classification of phytoplasma strains in the elm yellows group (16SrV) and proposal of '*Candidatus* Phytoplasma ulmi' for the phytoplasma associated with elm yellows. *International Journal of Systematic and Evolutionary Microbiology*, 54, 337–347.
- Li ZN, Bai YB, Liu P, Zhang L and Wu YF, 2014. Occurrence of '*Candidatus* Phytoplasma ziziphi' in apple trees in China. *Forest Pathology*, 44, 417–419.
- Liu SL, Liu HL, Chang SC and Lin CP, 2011. Phytoplasmas of two 16S rDNA groups are with pear decline in Taiwan. *Botanical Studies*, 52, 313–320.
- Maixner M, Reinert W and Darimont H, 2000. Transmission of grapevine yellows by *Oncopsis alni* (Schrank) (Auchenorrhyncha : Macropsinae). *Vitis*, 39, 83–84.
- Malembic-Maher S, Salar P, Filippin L, Carle P, Angelini E and Foissac X, 2011. Genetic diversity of European phytoplasmas of the 16SrV taxonomic group and proposal of '*Candidatus* Phytoplasma rubi'. *International Journal of Systematic and Evolutionary Microbiology*, 61, 2129–2134.
- Marcone C, Ragozzino A and Seemüller E, 1997. Detection and identification of phytoplasmas in yellows-diseased weeds in Italy. *Plant Pathology*, 46, 530–537.
- Marcone C, Ragozzino A, Camele I, Rana GL and Seemüller E, 2001. Updating and extending genetic characterization and classification of phytoplasmas from wild and cultivated plants in southern Italy. *Journal of Plant Pathology*, 83, 133–138.
- Marcone C, Guerra LJ and Uyemoto JK, 2014. Phytoplasma diseases of peach and associated phytoplasma taxa. *Journal of Plant Pathology*, 96, 15–28.
- Mehle N, Mermal S, Vidmar S, Marn MV, Dreo T and Dermastia M, 2018. First report of carrot infection with phytoplasmas in Slovenia. pp. 2–3.
- Montano HG, Davis RE, Dally EL, Hogenhout S, Pimentel JP and Brioso PST, 2001. '*Candidatus* Phytoplasma brasiliense', a new phytoplasma taxon associated with hibiscus witches' broom disease. *International Journal of Systematic and Evolutionary Microbiology*, 51, 1109–1118.
- Morton A, Davies DL, Blomquist CL and Barbara DJ, 2003. Characterization of homologues of the apple proliferation immunodominant membrane protein gene from three related phytoplasmas. *Molecular Plant Pathology*, 4, 109–114.
- Naderali N, Nejat N, Vadamalai G, Davis RE, Wei W, Harrison NA, Kong L, Kadir J, Tani YH and Zhao Y, 2017. '*Candidatus* Phytoplasma wodyetiae', a new taxon associated with yellow decline disease of foxtail palm (*Wodyetia bifurcata*) in Malaysia. *International Journal of Systematic and Evolutionary Microbiology*, 67, 3765–3772.
- Nigro F, Sion V, Antelmi I, Choueiri E, Habib W, Bruno A and Boscia D, 2020. First Report of '*Candidatus* Phytoplasma phoenicium' on almond in Southern Italy. *Plant Disease*, in press., <https://doi.org/10.1094/PDIS-01-19-0157-PDN>
- Nisbet C, Ross S, Monger WA, Hight F and Jeffries C, 2014. First report of '*Candidatus* Phytoplasma asteris' in commercial carrots in the United Kingdom. *New Disease Reports*, 30, 16.
- Paltrinieri S and Bertaccini A, 2007. Detection of phytoplasmas in plantlets grown from different batches of seed-potatoes. *Bulletin of Insectology*, 60, 379–380.
- Paltrinieri S, Botti S, Dal Molin F, Mori N and Fiore N, 2006. Are phytoplasmas involved in a severe peach decline? *Acta Horticulturae*, 713, 421–426.
- Parrella G, Paltrinieri S, Botti S and Bertaccini A, 2008. Molecular identification of phytoplasmas from virescent ranunculus plants and from leafhoppers in southern Italian crops. *Journal of Plant Pathology*, 90, 537–543.
- Pasquini G, Marzachi C, Pollini CP, Faggioli F, Ragozzino A, Bissani R, Vischi A, Barba M, Giunchedi L and Boccardo G, 2000. Molecular identification of phytoplasmas from olive trees in Italy. *Journal of Plant Pathology*, 82, 213–219.
- Pribylova J, Petrzik K and Spak J, 2009. The first detection of '*Candidatus* Phytoplasma trifolii' in *Rhododendron hybridum*. *European Journal of Plant Pathology*, 124, 181–185.
- Quaglino F, Zhao Y, Casati P, Bulgari D, Bianco PA, Wei W and Davis RE, 2013. '*Candidatus* Phytoplasma solani', a novel taxon associated with stolbur-and bois noir-related diseases of plants. *International Journal of Systematic and Evolutionary Microbiology*, 63, 2879–2894.
- Radisek S, Ferant N, Jakse J and Javornik B, 2009. Identification of a phytoplasma from the aster yellows group infecting purple coneflower (*Echinacea purpurea*) in Slovenia. *Plant Pathology*, 58, 392–392.
- Reeder R and Arocha Y, 2008. '*Candidatus* phytoplasma asteris' identified in *Senecio jacobaea* in the United Kingdom. *Plant Pathology*, 57, 769–769.
- Reeder R, Kelly P and Arocha Y, 2010a. First identification of '*Candidatus* Phytoplasma aurantifolia' infecting *Fallopia japonica* in the United Kingdom. *Plant Pathology*, 59, 396–396.

- Reeder R, Kelly PL and Arocha Y, 2010b. '*Candidatus* Phytoplasma asteris' identified in blackberry (*Rubus fruticosus* agg.) in the United Kingdom. *Plant Pathology*, 59, 394–394.
- Sabate J, Lavina A and Batlle A, 2014. First report of '*Candidatus* Phytoplasma pyri' causing peach yellow leaf roll (PYLR) in Spain. *Plant Disease*, 98, 989–990.
- Salehi M, Izadpanah K and Heydarnejad J, 2006. Characterization of a new almond witches' broom phytoplasma in Iran. *Journal of Phytopathology*, 154, 386–391.
- Salehi E, Salehi M, Taghavi SM and Izadpanah K, 2016. First report of a 16SrIX group (pigeon pea witches'-broom) phytoplasma associated with grapevine yellows in Iran. *Journal of Plant Pathology*, 98, 376–376.
- Schneider B and Gibb KS, 1997. Detection of phytoplasmas in declining pears in southern Australia. *Plant Disease*, 81, 254–258.
- Seemüller E and Schneider B, 2004. '*Candidatus* Phytoplasma mali', '*Candidatus* Phytoplasma pyri' and '*Candidatus* phytoplasma prunorum', the causal agents of apple proliferation, pear decline and European stone fruit yellows, respectively. *International Journal of Systematic and Evolutionary Microbiology*, 54, 1217–1226.
- Sharbatkhari M, Bahar M and Ahoonmanesh A, 2008. Detection of the phytoplasma agent of pear decline in Iran, Isfahan province, using nested-PCR. *International Journal of Plant Production*, 2, 167–173.
- Staniulis JB, Davis RE, Jomantiene R, Kalvelyte A and Dally EL, 2000. Single and mixed phytoplasma infections in phyllody- and dwarf-diseased clover plants in Lithuania. *Plant Disease*, 84, 1061–1066.
- Tolu G, Botti S, Garau R, Prota VA, Sechi A, Prota U and Bertaccini A, 2006. Identification of a 16SrII-E phytoplasma in *Calendula arvensis*, *Solanum nigrum*, and *Chenopodium* spp. *Plant Disease*, 90, 325–330.
- Valiunas D, Staniulis J and Davis RE, 2006. '*Candidatus* Phytoplasma fragariae', a novel phytoplasma taxon discovered in yellows diseased strawberry, *Fragaria x ananassa*. *International Journal of Systematic and Evolutionary Microbiology*, 56, 277–281.
- Valiunas D, Jomantiene R and Davis RE, 2007. Phytoplasmas detected in cultivated fruit plants in Lithuania. *Bulletin of Insectology*, 60, 139–140.
- Valiunas D, Jomantiene R, Ivanauskas A, Abraitis R, Staniene G, Zhao Y and Davis RE, 2009. First report of a new phytoplasma subgroup, 16SrIII-T, associated with decline disease affecting sweet and sour cherry trees in Lithuania. *Plant Disease*, 93, 550–550.
- Vellios E and Lioliopoulou F, 2007. Detection and characterization of phytoplasmas infecting tomato plants in Greece. *Bulletin of Insectology*, 60, 157–158.
- Verdin E, Salar P, Danet JL, Choueiri E, Jreijiri F, El Zammam S, Gelie B, Bove JM and Garnier M, 2003. '*Candidatus* Phytoplasma phoenicium' sp nov., a novel phytoplasma associated with an emerging lethal disease of almond trees in Lebanon and Iran. *International Journal of Systematic and Evolutionary Microbiology*, 53, 833–838.
- Wang LP, Hong N, Xu WX, Wang YF, Chofong GN, Martini M, Su J-M and Wang GP, 2014. Identification and molecular characterization of 16 S r V group phytoplasmas with flowering cherry in China. *Forest Pathology*, 44, 166–168.
- Wang JW, Liu QZ, Wei W, Davis RE, Tan Y, Lee IM, Zhu DZ, Wei HR and Zhao Y, 2018. Multilocus genotyping identifies a highly homogeneous phytoplasma lineage associated with sweet cherry virescence disease in China and its carriage by an erythroneurine leafhopper. *Crop Protection*, 106, 13–22.
- White DT, Blackall LL, Scott PT and Walsh KB, 1998. Phylogenetic positions of phytoplasmas associated with dieback, yellow crinkle and mosaic diseases of papaya, and their proposed inclusion in '*Candidatus* Phytoplasma australiense' and a new taxon, '*Candidatus* Phytoplasma australasia'. *International Journal of Systematic Bacteriology*, 48, 941–951.
- Zambon Y, Canel A, Bertaccini A and Contaldo N, 2018. Molecular diversity of phytoplasmas associated with grapevine yellows disease in north-eastern Italy. *Phytopathology*, 108, 206–214.
- Zhu SF, Hadidi A, Gundersen DE, Lee IM and Zhang CL, 1998. Characterization of the phytoplasmas. associated with cherry lethal yellows and jujube witches'-broom diseases in China. *Acta Horticulturae*, 472, 701–714.
- Zirak L, Bahar M and Ahoonmanesh A, 2009. Characterization of phytoplasmas associated with almond diseases in Iran. *Journal of Phytopathology*, 157, 736–741.
- Zirak L, Bahar M and Ahoonmanesh A, 2010a. Characterization of phytoplasmas related to '*Candidatus* Phytoplasma asteris' and peanut WB group associated with sweet cherry diseases in Iran. *Journal of Phytopathology*, 158, 63–65.
- Zirak L, Bahar M and Ahoonmanesh A, 2010b. Molecular characterization of phytoplasmas associated with peach diseases in Iran. *Journal of Phytopathology*, 158, 105–110.
- Zunnoon-Khan S, Arocha-Rosete Y, Scott J, Crosby W, Bertaccini A and Michelutti R, 2010. First report of '*Candidatus* Phytoplasma fraxini' (group 16SrVII phytoplasma) associated with a peach disease in Canada. *Plant Pathology*, 59, 1162–1162.

Abbreviations

Ca. P.	<i>Candidatus</i> Phytoplasma
CrWB	Crotalaria witches'-broom
EPPO	European and Mediterranean Plant Protection Organization
GD	Global Database

IRPCM	International Research Programme on Comparative Mycoplasmaology
KAP	<i>Knautia arvensis</i> phyllody
MLO	mycoplasma-like organisms
MS	Member State
NAGYIII	North American Grapevine Yellows
PCR	polymerase chain reaction
PDTWII	Pear decline Taiwan II
PEY	<i>Picris echioides</i> yellows
PHYPA	<i>Candidatus</i> Phytoplasma australasia
PHYPAS	<i>Candidatus</i> Phytoplasma asteris
PHYPAU	<i>Candidatus</i> Phytoplasma australiense
PHYPBR	<i>Candidatus</i> Phytoplasma brasiliense
PHYPPG	<i>Candidatus</i> Phytoplasma fragariae
PHYPFR	<i>Candidatus</i> Phytoplasma fraxini
PHYPMA	<i>Candidatus</i> Phytoplasma mali
PHYPPH	<i>Candidatus</i> Phytoplasma phoenicium
PHYPPN	<i>Candidatus</i> Phytoplasma pruni
PHYPPR	<i>Candidatus</i> Phytoplasma prunorum
PHYPPY	<i>Candidatus</i> Phytoplasma pyri
PHYPRU	<i>Candidatus</i> Phytoplasma rubi
PHYPSO	<i>Candidatus</i> Phytoplasma solani
PHYPTR	<i>Candidatus</i> Phytoplasma trifolii
PHYPUL	<i>Candidatus</i> Phytoplasma ulmi
PHYPZI	<i>Candidatus</i> Phytoplasma ziziphi
PHYPO1	Tomato big bud
PHYPO7	<i>Candidatus</i> Phytoplasma hispanicum
PHYP19	Clover yellow edge phytoplasma
PHYP39	Sweet potato little leaf
PHYP42	Pigeon pea witches' broom
PHYP45	<i>Knautia phyllody</i> phytoplasma
PHYP64	Grapevine Flavescence dorée phytoplasma
PHYP65	German flavescence dorée phytoplasma
PHYP74	Alder yellows phytoplasma
PLH	Plant Health
PYLR	Peach yellow leaf roll
RFLP	Restriction Fragment Length Polymorphism
SPLL	Sweet potato little leaf
TBB	Tomato big bud
TFEU	Treaty on the Functioning of the European Union
The host plants	<i>Cydonia</i> , <i>Fragaria</i> , <i>Malus</i> , <i>Prunus</i> , <i>Pyrus</i> , <i>Ribes</i> , <i>Rubus</i> and <i>Vitis</i>
ToR	Terms of Reference
WoS	Web of Science

Appendix A – Non-EU phytoplasmas of *Cydonia*, *Fragaria*, *Malus*, *Prunus*, *Pyrus*, *Ribes*, *Rubus* and *Vitis*

ID	Phytoplasma name	Related strain name ⁽¹⁾	Abbreviation (EPPO code)	16S rRNA	<i>Cydonia</i>	<i>Fragaria</i>	<i>Malus</i>	<i>Prunus</i>	<i>Pyrus</i>	<i>Ribes</i>	<i>Rubus</i>	<i>Vitis</i>	Reasoning for considering non-EU	Uncertainties	References
1	<i>Candidatus Phytoplasma aurantifolia</i>	'Ca. P. australasia, pear decline Taiwan II (PDTWII); <i>Crotalaria witches'-broom</i> phytoplasma (CrWB); sweet potato little leaf (SPLL)	PHYPA, PHYPA39	II	–	–	Yes	Yes	Yes	–	–	Yes	Only one report in <i>Fallopia japonica</i> in UK ⁽²⁾ ; Greek report of TBB (tomato big bud) (listed in EPPO) is most probably to be assigned to either 'Ca. P. asteris' or 'Ca. P. solani'; Italian reports refer to few infected individuals	Hashemi-Tameh et al., 2014 (4 infected <i>Malus</i> plants); Ghayeb Zamharir et al., 2017 (4 <i>Vitis</i> plants); Reeder et al., 2010a (only one report from UK, 4 plants detected by nested PCR (polymerase chain reaction) out of 4 tested ones); [PHYPA01]: in Tomato in Greece following EPPO (Alivizatos, 1993; only EM), but Vellios and Lioliopoulou, 2007 identified only 16SrI and XII in tomato in Greece. No record of its presence in Portugal. ⁽²⁾ Absence of PHYPA distribution map	Species description: (White et al., 1998; Liu et al., 2011; IRPCM, 2004) <i>Malus</i> : (Hashemi-Tameh et al., 2014); <i>Prunus</i> : (Zirak et al., 2009, 2010a,b); <i>Vitis</i> : (Constable et al., 2003; Ghayeb Zamharir et al., 2017); <i>Pyrus</i> : (Schneider and Gibb, 1997; Liu et al., 2011); <i>Fallopia japonica</i> in UK (Reeder et al., 2010a); <i>Solanum tuberosum</i> in Italy (Paltrinieri and Bertaccini, 2007); <i>Empoasca decipiens</i> in Italy (Parrella et al., 2008); <i>Calendula arvensis</i> , <i>Solanum nigrum</i> , and <i>Chenopodium</i> spp. In Italy (Tolu et al., 2006); <i>Matthiola incana</i> in Italy (Davino et al., 2007)
2	<i>Candidatus Phytoplasma australiense</i>	–	PHYPAU	XII-B	–	Yes	–	Yes	–	–	Yes	Yes	Not reported to be present in the EU	–	Species description: (Davis et al., 1997); <i>Prunus</i> : (Jones et al., 2005)

ID	Phytoplasma name	Related strain name ⁽¹⁾	Abbreviation (EPPO code)	16S rRNA	Cydonia	Fragaria	Malus	Prunus	Pyrus	Ribes	Rubus	Vitis	Reasoning for considering non-EU	Uncertainties	References
3	<i>Candidatus Phytoplasma fraxini</i>	–	PHYPPFR	VII-A	–	Yes	–	Yes	–	–	–	Yes	Only two reports in Italy	Zambon et al., 2018 (9 plants detected in Italy by nested PCR out of 161 tested ones); Bruni et al., 2005 (<i>Hypericum perforatum</i> in Italy: it is not known on how many plants the phytoplasma was identified)	Species description: (Griffiths et al., 1999); <i>Fragaria</i> : (Fernandez et al., 2013); <i>Prunus</i> : (Zunnoon-Khan et al., 2010); <i>Vitis</i> : (Gajardo et al., 2009; Ghayeb Zamharir et al., 2017; Zambon et al., 2018); Italy: (Bruni et al., 2005; Zambon et al., 2018)
4	<i>Candidatus Phytoplasma hispanicum</i>	–	PHYPP07	XIII	–	Yes	–	–	–	–	–	–	Not reported to be present in the EU	–	Species description: (Davis et al., 2016); <i>Fragaria</i> : (Jomantiene et al., 1998; Fernandez et al., 2015)
5	<i>Candidatus Phytoplasma phoenicium</i>	–	PHYPPH	IX-B, D, F, G ²	–	–	–	Yes	–	–	–	Yes	Only one report in Italy	Ghayeb Zamharir et al., 2017 (3 <i>Vitis</i> plants with probable mixed infections)	Species description: (Verdin et al., 2003; EPPO, 2017); <i>Vitis</i> : (Ghayeb Zamharir et al., 2017); <i>Prunus</i> in Italy: (Nigro et al., 2020)
6	<i>Candidatus Phytoplasma pruni</i>	North American Grapevine Yellows - NAGYIII	–	III	–	–	–	–	–	–	–	Yes	Not reported to be present in the EU	–	Species description: (Davis et al., 2015)
7	<i>Candidatus Phytoplasma pyri</i>	Peach yellow leafroll	–	X	–	–	–	Yes	–	–	–	–	Not reported to be present in the EU	–	Species description: (Morton et al., 2003; Seemüller and Schneider, 2004); <i>Prunus</i> : (Marccone et al., 2014)

ID	Phytoplasma name	Related strain name ⁽¹⁾	Abbreviation (EPPO code)	16S rRNA	Cydonia	Fragaria	Malus	Prunus	Pyrus	Ribes	Rubus	Vitis	Reasoning for considering non-EU	Uncertainties	References
8	<i>Candidatus Phytoplasma trifolii</i>	–	PHYPTR	VI-A	–	Yes	–	Yes	–	–	–	Yes	Reports from the EU MS refer to few infected plants	[PHYPTR]: Pribylova et al., 2009; Borroto Fernandez et al., 2007 (Reports from the EU MS refer to few infected plants, ranging from 1 to 28); Zirak et al., 2010b (only in one <i>Prunus</i> plant out of 91 tested plants)	Species description and <i>Fragaria</i> : (Hiruki and Wang, 2004); <i>Prunus</i> : (Zirak et al., 2010b); Czech Republik: (Pribylova et al., 2009); Austria: (Borroto Fernandez et al., 2007)
9	<i>Candidatus Phytoplasma ziziphi</i>	–	PHYPTZI	V-B	–	–	Yes	Yes	–	–	–	–	One report from Italy is related to mixed infections with 'Ca. P. solani' and 'Ca. P. asteris' and the other one has no further characterization beside PCR and RFLP (restriction fragment length polymorphism) analyses	Paltrinieri et al., 2006 (<i>Prunus</i> in Italy: only a Congress abstracts, no further details); Pasquini et al., 2000 (<i>Olea europea</i> in Italy: phytoplasma detected based only on PCR and RFLP analyses of ribosomal operon)	Species description: (Jung et al., 2003); <i>Malus</i> : (Li et al., 2014); <i>Prunus</i> : (Zhu et al., 1998; Wang et al., 2014, 2018); <i>Prunus</i> in Italy: (Paltrinieri et al., 2006); <i>Olea europea</i> in Italy: (Pasquini et al., 2000)
10	<i>Unclassified</i>	Buckland valley grapevine yellows phytoplasma	–	XXIII	–	–	–	–	–	–	–	Yes	Not reported to be present in the EU	–	Species description: (Constable et al., 2002)

(1): Reference isolate of 'Candidatus Phytoplasma species' is indicated by '-'.
(2): Information provided by MS during commenting phase.

Appendix B – Phytoplasmas of *Cydonia*, *Fragaria*, *Malus*, *Prunus*, *Pyrus*, *Ribes*, *Rubus* and *Vitis* excluded from further categorisation as their presence in the species is not fully supported by available literature

ID	Phytoplasma name	Related strain name ⁽¹⁾	Abbreviation (EPPO code)	16S rRNA	<i>Cydonia</i>	<i>Fragaria</i>	<i>Malus</i>	<i>Prunus</i>	<i>Pyrus</i>	<i>Ribes</i>	<i>Rubus</i>	<i>Vitis</i>	Reasoning for considering non-EU	Uncertainties	References
11	<i>Candidatus Phytoplasma brasiliense</i>	–	PHYPPBR	XV-A	–	–	–	Yes	–	–	–	–	Not reported to be present in the EU. Excluded from further categorisation as its presence in <i>Prunus</i> is not fully supported by available literature	Balakishiyeva et al., 2011 (only in 1 <i>Prunus</i> plant)	Species description: (Montano et al., 2001); <i>Prunus</i> : (Balakishiyeva et al., 2011)
12	<i>Candidatus Phytoplasma pruni</i>	Clover yellow edge, CYE (CYE-C; CYE-L)	PHYPP19	III-B	–	Yes	–	–	–	–	–	–	In the EU reported only in two clover plants and in mixed infections. Excluded from further categorisation as its presence in <i>Fragaria</i> is not fully supported by available literature	Staniulis et al., 2000 (Only in two clover plants and in mixed infections); Jomantiene et al., 2002 (Reported in <i>Fragaria</i> in a unique report of a maximum of 5 tested symptomatic plants)	Species description: (Davis et al., 2013); <i>Fragaria</i> : (Jomantiene et al., 2002); in clover and Lithuania (Staniulis et al., 2000)
13	Unclassified	Pigeon pea witches' broom	PHYPP42	IX-A	–	–	–	–	–	–	–	Yes	Not reported to be present in the EU. Excluded from further categorisation as its presence in <i>Vitis</i> is not fully supported by available literature	Ertunc et al., 2015 (Reported on 1 <i>Vitis</i> plant out of 289 tested plants)	Species description: (Ertunc et al., 2015)

(1): Reference isolate of '*Candidatus* Phytoplasma species' is indicated by '-'.

Appendix C – Phytoplasmas of *Cydonia*, *Fragaria*, *Malus*, *Prunus*, *Pyrus*, *Ribes*, *Rubus* and *Vitis* excluded from further categorisation

ID	Phytoplasma name	Related strain name ⁽¹⁾	Abbreviation (EPPO code)	16S rRNA	EU MS in which the pathogen has been reported	Non-EU European and neighbouring countries	<i>Cydonia</i>	<i>Fragaria</i>	<i>Malus</i>	<i>Prunus</i>	<i>Pyrus</i>	<i>Ribes</i>	<i>Rubus</i>	<i>Vitis</i>	Reasoning for considering as EU	Uncertainties	References
14	<i>Candidatus Phytoplasma asteris</i>	–	PHYPAS	I	Germany, Hungary, Italy (Present widespread); Czech Republic, Spain (Present, restricted distribution); France, Romania (Present, no details); Belgium ⁽²⁾ ; UK ⁽²⁾	Russia (Present, restricted distribution; Belarus (Present, no details)	–	Yes	Yes ⁽²⁾	Yes	Yes	–	Yes	Yes	Reported in the EU (several MS)	–	Species description, <i>Fragaria</i> , <i>Prunus</i> , <i>Pyrus</i> , <i>Vitis</i> : (Lee et al., 2004a) and <i>Rubus</i> (Reeder et al., 2010b); Lithuania (Valiunas et al., 2007); UK (Jones and Arocha, 2006; Reeder and Arocha, 2008; Nisbet et al., 2014), Slovenia: (Mehle et al., 2018; Radisek et al., 2009)
15	<i>Candidatus Phytoplasma fragariae</i>	–	PHYPPFG	XII-E	Slovenia (EPPO report 2018/085); UK (2015/031); Belgium ⁽²⁾		–	Yes	–	–	–	–	–	–	Originally described in the EU	–	Species description: (Valiunas et al., 2006)
16	<i>Candidatus Phytoplasma mali</i>	–	PHYPPMA	X	Czech Republic, Germany, Hungary, Italy, Slovakia, Slovenia (Present widespread); Austria, Belgium, Bulgaria, Croatia, Finland, France, Greece (Present, restricted distribution); Poland, Romania (Present, no details); Netherlands (Present, few occurrences)	Switzerland (Present widespread); Norway (Present, restricted distribution); Albania, Bosnia and Herzegovina, Moldova, Turkey, Ukraine (Present, no details)	–	–	Yes	Yes	Yes	–	–	–	Reported in the EU (several MS)	–	Species description: (Seemüller and Schneider, 2004)

ID	Phytoplasma name	Related strain name ⁽¹⁾	Abbreviation (EPPO code)	16S rRNA	EU MS in which the pathogen has been reported	Non-EU European and neighbouring countries	Cydonia	Fragaria	Malus	Prunus	Pyrus	Ribes	Rubus	Vitis	Reasoning for considering as EU	Uncertainties	References
17	<i>Candidatus Phytoplasma pruni</i>	–	PHYPPN	III-A	Croatia, Italy, Lithuania, Poland, UK	Serbia	–	–	Yes	Yes	Yes	–	Yes	Yes	Reported in the EU (several MS)	PYLR is listed among synonyms of <i>Ca. P. pruni</i> in EPPO (2019), but following Seemüller and Schneider, 2004; and Morton et al., 2003, it is listed here as <i>Ca. P. pyri</i> -related strain	Species description: (Davis et al., 2013); <i>Pyrus</i> in Croatia: (Jezic et al., 2016); <i>Pyrus</i> in Italy: (Lee et al., 1995); <i>Rubus</i> in UK: (Davies, 2000); <i>Rubus</i> in Poland: (Cieslinska, 2011); <i>Vitis</i> in Croatia: (Jezic et al., 2013); <i>Prunus</i> in Lithuania: (Valiunas et al., 2009); <i>Prunus</i> in Italy: (Lee et al., 1995); <i>Celtis australis</i> in Italy: (Bertaccini et al., 1996); <i>Cirsium</i> in Serbia: (Jakovljevic et al., 2015); <i>Delphinium</i> in UK (Harju et al., 2008) ⁽²⁾
18	<i>Candidatus Phytoplasma prunorum</i>	–	PHYPPR	X	Slovenia (Present, widespread); Austria, Bulgaria, Croatia, Czech Republic, France, Germany, Greece, Hungary, Italy, Poland, Romania, Spain, (Present, restricted distribution); Belgium, Slovakia (Present, few details)	Albania, Bosnia and Herzegovina, Switzerland, Turkey, UK (Present, restricted distribution); Belarus, Serbia (Present, no details); Azerbaijan (Present, few details)	–	–	–	Yes	–	–	–	–	Reported in the EU (several MS)	–	Species description: (Seemüller and Schneider, 2004)

ID	Phytoplasma name	Related strain name ⁽¹⁾	Abbreviation (EPPO code)	16S rRNA	EU MS in which the pathogen has been reported	Non-EU European and neighbouring countries	Cydonia	Fragaria	Malus	Prunus	Pyrus	Ribes	Rubus	Vitis	Reasoning for considering as EU	Uncertainties	References
19	<i>Candidatus Phytoplasma pyri</i>	–	PHYPPY	X	Italy, Netherlands (Present widespread); Belgium, Bulgaria, Croatia, Czech Republic, France, Germany, Greece, Poland, Slovakia, Slovenia (Present, restricted distribution); Hungary (Present, no details); Austria, Portugal, Spain (Present, few occurrences)	Switzerland (Present widespread); Norway, Serbia, Turkey (Present, restricted distribution); Albania, Bosnia and Herzegovina, Moldova (Present, no details)	Yes	–	–	Yes	Yes	–	–	–	Reported in the EU (several MS)	–	Species description: (Seemüller and Schneider, 2004); <i>Prunus</i> : (Sabate et al., 2014)
20	<i>Candidatus Phytoplasma rubi</i>	–	PHYPRU	V-E	Denmark, Germany, Italy, Netherlands (Present widespread); Bulgaria, Portugal, Slovakia (Present, no details); UK (Present widespread); Belgium ⁽²⁾ ; Czech Republic ⁽²⁾	Russia (Present, restricted distribution); Norway, Central Russia, Switzerland (Present, no details)	–	–	–	–	–	–	Yes	–	Reported in the EU (several MS)	–	Species description: (Malembic-Maher et al., 2011)

ID	Phytoplasma name	Related strain name ⁽¹⁾	Abbreviation (EPPO code)	16S rRNA	EU MS in which the pathogen has been reported	Non-EU European and neighbouring countries	Cydonia	Fragaria	Malus	Prunus	Pyrus	Ribes	Rubus	Vitis	Reasoning for considering as EU	Uncertainties	References
21	<i>Candidatus Phytoplasma solani</i>	–	PHYPSO	XII-A	Bulgaria, Croatia, France, Germany, Greece, Hungary, Italy, Slovakia, Slovenia, Spain, (Present, restricted distribution); Italy (Present, no details); Austria, Czech Republic, Poland (Present, few occurrences); UK (Transient, under eradication); Belgium (Absent, eradicated) ⁽²⁾ ; Portugal (Present, restricted distribution) ⁽²⁾	Macedonia, Montenegro (Present, widespread); Russia, Serbia, Switzerland, Turkey (Present, restricted distribution); Albania, Armenia, Azerbaijan, Bosnia and Herzegovina, Georgia, Ukraine (Present, no details)	–	Yes	Yes	Yes	–	–	Yes	Yes	Reported in the EU (several MS)	–	Species description: (Quaglino et al., 2013); <i>Malus</i> : (Duduk et al., 2010); <i>Prunus</i> : (Avramov et al., 2011)
22	<i>Candidatus Phytoplasma ulmi</i>	–	PHYFUL	V-A	Croatia (Present, widespread), France, Italy, Slovenia (Present, restricted distribution), Belgium (Present, under eradication) ⁽²⁾ , Czech Republic, Germany (Present, few occurrences), Poland (Transient, under eradication); UK (Absent, eradicated) ⁽²⁾	Serbia (Present, few occurrences)	–	–	–	–	–	–	–	Yes	Reported in the EU (several MS), but not in target species. Reported only once in <i>Vitis</i> (and in 1 plant)	Zambon et al., 2018 (Detected in one <i>Vitis</i> out of 161 tested ones in Italy)	Species description: (Lee et al., 2004b); <i>Vitis</i> : (Zambon et al., 2018)

ID	Phytoplasma name	Related strain name ⁽¹⁾	Abbreviation (EPPO code)	16S rRNA	EU MS in which the pathogen has been reported	Non-EU European and neighbouring countries	Cydonia	Fragaria	Malus	Prunus	Pyrus	Ribes	Rubus	Vitis	Reasoning for considering as EU	Uncertainties	References
23	Unclassified	Grapevine Flavescence dorée phytoplasma	PHYP64	V	Croatia, France, Hungary, Italy, Portugal, Slovenia (Present, restricted distribution); Austria, Spain (Present, few occurrences); Belgium (Absent, no pest record) ⁽²⁾	Serbia, Switzerland (Present, restricted distribution)	-	-	-	-	-	-	-	Yes	Reported in the EU (several MS)	-	Species description: (EFSA PLH Panel, 2014); (Lee et al., 2004b)-
24	Unclassified	German flavescence dorée phytoplasma	PHYP65	V	Germany	-	-	-	-	-	-	-	-	Yes	Originally described in the EU	-	Species description and <i>Vitis</i> : (Angelini et al., 2001)
25	Unclassified	Alder yellows phytoplasma	PHYP74	V-C	France	-	-	-	-	-	-	-	-	Yes	Originally reported in the EU	-	Species description: (Lee et al., 2004b); <i>Vitis</i> (experimental transmission to): (Maixner et al., 2000); <i>Alnus</i> in France: (Arnaud et al., 2007)
26	Unclassified	<i>Knautia phyllody</i> phytoplasma, KAP	PHYP45	IX-C	Italy	-	-	-	Yes	Yes	-	-	-	-	Originally reported in the EU	-	Species description : (Marccone et al., 2001); <i>Knautia arvensis</i> in Italy: (Marccone et al., 1997); <i>Pyrus</i> : (Sharbatkhari et al., 2008); <i>Prunus</i> : (Salehi et al., 2006);
27	Unclassified	Picris echioides yellows, PEY		IX-C	Italy	-	-	-	-	-	-	-	-	Yes	Originally reported in the EU	Salehi et al., 2016 (8 plants)	Species description and <i>Argyranthemum</i> in Italy: (Ferretti et al., 2015); <i>Vitis</i> : (Salehi et al., 2016); <i>Picris echioides</i> in Italy: (Marccone et al., 1997)

(1): Reference isolate of 'Candidatus Phytoplasma species' is indicated by '-'.
 (2): Information provided by MS during commenting phase.

Annex A – List of phytoplasmas considered in the opinion

See excel file in Supplementary Information online.