

# Improving RNAi efficiency for pest control in crop species

Shuo Yan<sup>1</sup>, Binyuan Ren<sup>2</sup>, Bo Zeng<sup>2</sup> & Jie Shen<sup>\*1</sup>

## ABSTRACT

The application of RNAi promotes the development of novel approaches toward plant protection in a sustainable way. Genetically modified crops expressing dsRNA have been developed as commercial products with great potential in insect pest management. Alternatively, some nontransformative approaches, including foliar spray, irrigation and trunk injection, are favorable in actual utilization. In this review, we summarize the recent progress and successful cases of RNAi-based pest management strategy, explore essential implications and possibilities to improve RNAi efficiency by delivery of dsRNA through transformative and nontransformative approaches, and highlight the remaining challenges and important issues related to the application of this technology.

## KEYWORDS

dsRNA delivery • foliar spray • nanoparticle • nontransformative RNAi product • RNAi-based pest management • RNAi efficiency • transformative RNAi product • transplasmidic crop

<sup>1</sup>Department of Entomology, MOA Key Lab of Pest Monitoring & Green Management, College of Plant Protection, China Agricultural University, Beijing, People's Republic China; <sup>2</sup>National Agricultural Technology Extension & Service Center, Beijing, People's Republic China; \*Author for correspondence: [shenjie@cau.edu.cn](mailto:shenjie@cau.edu.cn)

BioTechniques 68: 283–290 (May 2020)  
10.2144/btn-2019-0171

To meet the increasing global demands for food and energy, mankind is facing the biggest challenge: how to increase crop yields in a profitable, efficient and sustainable way. There are several issues constraining agricultural productivity, such as damage by insect pests, diseases and weeds [1]. For instance, insect pests can cause direct and indirect crop damages mainly through loss in yield or quality owing to their adaptive ecological and physiological characteristics, which will be an important constraint to the supply of food in the next 40–50 years [2]. The estimates on potential yield loss of major crops by insect pests reach approximately 18% [3]. Furthermore, climate change can lead to some potential impacts on insect pests. The global yield losses of main crops such as rice, maize and wheat are projected to increase by 10–25% per degree of global mean surface warming [4,5].

Currently, chemical pesticides remain the major approach for suppressing insect pests owing to their well-controlled effect. Unfortunately, the excessive application of chemical pesticides has caused some serious problems threatening the environment and human health [6–8]. Therefore, a great demand for novel and effective alternative approaches has developed in recent years owing to growing consumer awareness and pressure for safer and healthier food. A pest management strategy should be economically, environmentally and farmer friendly. RNAi, first described in *Caenorhabditis elegans*, is known as post-transcriptional gene silencing because exogenous RNAs can induce sequence-specific mRNA degradation [9]. Among the mature biotechnological tools, RNAi has not only provided a novel and powerful reverse genetics tool for identifying gene functions, but also showed a great potential in pest management [10–13].

One promising area is the development of transgenic crops expressing dsRNA against key genes of insect pests, which are now realized by some commercial

products [14–16]. Alternatively, there is also a great demand for nontransformative approaches. For instance, nontransformative products for foliar application, trunk injection, root dipping and seed treatment have their advantages [17–19]. However, there remain some issues hindering the practice and development of transformative and nontransformative RNAi products. When these products were designed for pest management, RNAi efficiency was the biggest technical bottleneck that needed to be overcome before products for pest control could be implemented. In this review, we summarize the successful applications of RNAi-based pest management strategy in crop species, discuss the possibility to further increase RNAi efficiency and propose the remaining challenges and possible solutions.

## RNAi PATHWAYS & MECHANISMS FOR PEST MANAGEMENT

There are three RNAi pathways in insects: the dsRNA/siRNA-mediated siRNA pathway, miRNA-mediated miRNA pathway and piwi-interacting RNA (piRNA)-mediated piRNA pathway [11,20–22]. These pathways play different roles, such as in defense against viruses and transposable elements (transposons) via the siRNA pathway [23,24], regulation of gene expression via the miRNA pathway [25,26] and suppression of germ-line transposon expression via the piRNA pathway [21,27]. The core machinery genes of RNAi pathways may vary among different insect species. Insect pests tend to take up dsRNA rather than siRNA in some coleopteran and dipteran species through clathrin-dependent endocytosis [28–31]; however, siRNA works in some other insect species [32–34]. Therefore, gene silencing is usually triggered by supplying exogenous dsRNA in insects, although some insects are insensitive to RNAi through feeding. However, it is a bit different for fungi and plants, in which siRNAs are usually provided to trigger

RNAi [35–37]. When long dsRNA enters the target cells, it is cleaved by Dcr2 in association with R2D2 in the exo-siRNA pathway or Loqs in the endo-siRNA pathway into siRNA, which is loaded into Ago2 to assemble the RNA-induced silencing complex to degrade the complementary mRNA [38,39].

As shown in Figure 1, when insecticidal dsRNAs are applied in practical production, they need to enter target cells to work. Genetically modified (GM) crops and topical application, known as direct uptake, possess good prospects for wider application, and there have been some successful cases [15,40–42]. However, dsRNA delivery efficiency is often low in topical application, and during conventional transgenesis it is difficult to produce sufficient amounts of stable dsRNA owing to the plant RNAi machinery. However, dsRNAs can also enter the plant vascular system and then undergo uptake by insect pests [43], known as indirect uptake. However, dsRNAs may be restricted to the xylem vessels, and dsRNA delivery inside the plant vascular system is limited [44]. Therefore, some limitations exist when applying this indirect uptake method. Introduction of transplastomic technology and nanotechnology may overcome these current difficulties, and could improve RNAi efficiency, promoting the development and practice of RNAi-based pest management strategies (Figure 1).

## APPLICATION OF TRANSGENIC CROPS FOR PEST MANAGEMENT

### Conventional transgenic crops

GM crops engineered to express Bt toxins have become the most successfully commercialized transgenic crops for pest management [45,46]. Meanwhile, plant-mediated RNAi has developed rapidly in recent years attributed to the great advantage that almost any lethal gene can be targeted in insect pests. The most inspiring breakthrough was the development of GM maize expressing *vATPaseA* dsRNA to efficiently control the western corn rootworm (WCR, *Diabrotica virgifera virgifera*). This pest is susceptible to dsRNA supplied in an artificial diet, which can lead to larval stunting and mortality [15]. Another example targeting a *cytochrome P450* gene encoding a detoxifying enzyme is the Cotton bollworm (*Helicoverpa armigera*), which can resist

gossypol and related sesquiterpene aldehydes that are toxic to many organisms. Mao *et al.* [47] constructed GM plants expressing hairpin RNA (hpRNA) to successfully suppress the *CYP6AE14* expression in cotton bollworm, which reduced the larval tolerance to gossypol. Other successful cases exist in the application of GM crops expressing dsRNAs, such as the GM wheat against aphids [48,49], the GM potato against *Leptinotarsa decemlineata* [50], the GM cotton against *Tetranychus cinnabarinus* [51] and *H. armigera* [52], and the GM tobacco against *Myzus persicae* [53] and *H. armigera* [54].

### Factors determining RNAi efficiency of transgenic crops

Several factors influence RNAi efficiency in insect pests. The success of plant-mediated RNAi for pest management first relies on the stable expression of dsRNA, as GM crops should provide enough dsRNAs to trigger a strong RNAi response. Conventional GM crops use nuclear transformation, and the expressed hpRNAs enter cytoplasm and are usually processed into siRNAs by plant RNAi machinery. RNAi efficiency is clearly dependent on the dsRNA dose, and the desired pest control effect needs to be determined experimentally against various target genes. The length of expressed dsRNA is an important factor affecting RNAi efficiency in some insect species. dsRNAs are taken up by an active process involving the receptor-mediated endocytosis, and insects are more responsive to longer dsRNA. In *Drosophila* S2 cells, dsRNAs of 1000 and 200 bp can induce a significant gene silencing; however, 21 bp siRNAs cannot result in any significant silencing [29]. In WCR, dsRNAs longer than or equal to 60 bp are required for an efficient RNAi, whereas 21 bp siRNAs cannot trigger RNAi [55]. RNAi efficiency is also dependent on insect species that possess different abilities of dsRNA degradation [56]. The activity of dsRNases that can efficiently cleave dsRNA has been identified in several insect species [57–59]. Suppression of specific *dsRNase* genes can lead to the reduction of dsRNA degrading activity and improve RNAi efficiency in *Cylas puncticollis* [60], *Locusta migratoria* [61], *Schistocerca gregaria* [59], *Ostrinia furnacalis* [62] and *L. decemlineata* [63].

### Transplastomic crops with higher RNAi efficiency

Transformation of chloroplast DNA, also referred to as transplastomic crops, overcomes many current difficulties and has a good potential application [64,65]. The high transgene expression levels from chloroplast genome make transplastomic technology an attractive choice in herbicide and insect resistance engineering [66–70]. The greatest advantage in applying chloroplast-expressing dsRNAs is that it permits the accumulation of much higher amounts of stable dsRNA in the chloroplast, and therefore is not cleaved by the plant RNAi machinery [71,72]. In addition, the transplastomic technology provides an environmentally benign method, because plastids are maternally inherited in most crops and, therefore, constrain the pollen-mediated gene flow to decrease the potential environmental risk [64,73,74].

The RNAi efficiencies of nuclear- and chloroplast-transformed potatoes targeting the  $\beta$ -*actin* gene of *L. decemlineata* have been compared [71]. In transplastomic potato, the dsRNAs accumulated to as much as 0.4% of the total cellular RNA, whereas the nuclear-transformed potato produced much less dsRNAs. Meanwhile, the siRNAs specific to the target gene were detected in the beetles feeding on transplastomic potato, but no detectable siRNAs were found in the beetles feeding on nuclear-transformed potato. Reasonably, the transplastomic potato exhibited higher gene silencing and better pest control effects. Similarly, the hpRNAs, targeting the *acetylcholinesterase2* gene of *H. armigera*, were integrated into either the nuclear or chloroplast genome of *Nicotiana benthamiana*. The hpRNAs accumulated in transplastomic *N. benthamiana* to confer a strong resistance to *H. armigera*, whereas the hpRNAs produced by nuclear-transformed crops were cleaved into siRNAs, exhibiting a more modest antifeeding activity [75]. *Chitin synthase*, *cytochrome P450 monooxygenase* and *vATPase* dsRNA expressed via the chloroplast genome decreased the target gene expression and showed a strong resistance to *H. armigera* [76]. These results demonstrate that there is less or no RNAi machinery in chloroplast, and that the dsRNAs produced within chloroplast do not enter the cytoplasm, but can be taken up by the insect midgut cells




Figure 1. Application of transplastomic technology and nanotechnology to improve RNAi efficiency for insect pest management.

to trigger RNAi, making chloroplast a good tool for dsRNA expression.

## APPLICATION OF NONTRANSFORMATIVE DELIVERY FOR PEST MANAGEMENT

### Foliar spray

Sprayable RNAi-based products can be suitable for suppressing pests on stems, foliage or fruits. The dsRNA formulation can be directly sprayed on insect pests, which may penetrate the cuticle to induce lethal effects, and also be sprayed on the crops to feed pests. One famous study exploring the application of sprayable dsRNA formulation was conducted to control *L. decemlineata* [77]. Second instar *L. decemlineata* could not survive to fourth instar on potato plants treated with *actin* dsRNA. The dsRNA was sufficiently stable for at least 4 weeks under greenhouse conditions. However, dsRNAs may be degraded by nuclease in the field, and the stability of dsRNA should be evaluated owing to the complex factors in the environment. Another study was conducted using six siRNAs targeting *acetylcholine esterase* genes of *Plutella xylostella* [33]. The best insecticidal activity with 89% mortality rate was observed when second instar *P. xylostella* were fed with *Brassica* spp. leaves sprayed with siRNAs. Furthermore, the dsRNAs, targeting three functional domains of the *Ostrinia furnacalis methionine-rich storage protein* gene, were sprayed directly on *O. furnacalis* and *H. armigera*, and the dsRNA targeting the C-terminal domain caused high mortality rates in both insect pests [78]. There are also some studies reporting that dsRNA can penetrate insect cuticle to mediate gene silencing. A droplet of 0.5  $\mu$ l dsRNA targeting *TAP1* gene was applied to the dorsal thorax of *Aedes aegypti*, and the mortality rate caused by the combination of three dsRNAs reached 42% at 24 h post-topical application [40]. Two papers from the same study group revealed that the topically applied dsRNA could penetrate the cuticle of *Diaphorina citri*, revealing the dsRNA products can be sprayed directly on some insect species [41,79]. Topical RNAi-mediated gene silencing seems also to work in aphids [80]. However, the penetration ability of dsRNA into the cuticle is different among insects, and the topical application is not fit for all insect pests.

### Irrigation

The delivery of dsRNA via crop roots can trigger RNAi in insect pests, and the irrigation of RNAi-based products seems to be an alternative for suppressing pests feeding/growing in stems and fruits. The longevity of the dsRNAs applied through the root drench seems suitable to develop an area-wide pest suppression approach. The persistence of dsRNA in citrus trees was detectable at least 57 d post-treatment, whereas in psyllids and leafhoppers the detection was 5–8 d post-treatment [81]. One study exploring the applications of dsRNA via crop roots was conducted against the brown planthopper and the Asian corn borer [82]. When *Nilaparvata lugens* fed on rice that had been irrigated with *carboxylesterase* dsRNAs, the mortality rate reached nearly 50% at 5 d post-treatment. Meanwhile, the 5-d mortality rate was more than 45% when *O. furnacalis* was fed on dsRNA-treated maize. A method for RNAi bioassay was developed by feeding dsRNA via plants. The plant stem was detached and inserted into a centrifuge tube containing dsRNA, and the insects were released and reared on the stems and observed for the RNAi effects. This method has already been applied in the evaluation of RNAi effects using the citrus stem for *Toxoptera citricida*, tomato leaf for *Bemisia tabaci*, bean leaf for *Acyrtosiphon pisum* and *Brassica* leaf for *M. persicae* [83–87]. The irrigation is a simple yet practical method to deliver dsRNA; however, dsRNA may be degraded within approximately 2 d after the application to soil, regardless of texture, pH, clay content and other soil differences [88]. Thus, the success of this delivery strategy relies on the advances of formulations to protect dsRNA from degradation.

### Trunk injection

The efficiencies of foliar spray and irrigation are relatively low, and sometimes this method is impractical for trees. Trunk injection is a promising method to deliver agrochemicals in many tree species while reducing the environmental impacts, risk for users and consumer exposure [89,90]. Phloem is considered a preferential channel for the transport of dsRNA/siRNA where it can remain stable for long periods, owing to the RNase-free environment in phloem sap [91,92]. Trunk injection can deliver dsRNAs into the vascular plant systems of

xylem and phloem, and the Arborjet® is available and may be applied to deliver dsRNA [93]. Citrus trees (2.5-m height) and grapevines were treated with dsRNA via root drench and trunk injection, and the dsRNA was taken up into the whole plant system over 3 months to suppress insect pests [81]. The control of some insect pests has been difficult, especially for underground root-feeding pests, and the trunk injection may solve this problem. This strategy may be more effective for sap-sucking pests than for chewing pests feeding largely on leaves [93].

### Improved delivery efficiency of dsRNA by nanoparticles

Nanoparticles are defined as any particle between 1 and 100 nm [94]. In addition to shielding and protecting the dsRNA from environmental nuclease degradation, nanoparticles promote the translocation of dsRNA across the peritrophic membrane, cell membrane and insect cuticle [95–98]. In most cases, the nanoparticle combines with dsRNA into the nanoparticle/dsRNA complex through the electrostatic interactions between the cationic groups in the nanoparticle and the phosphate groups in the dsRNA [99,100]. The complex usually retains a net positive charge that facilitates the interaction with negatively charged cell membrane surface [96]. When the complex is bound to the cell membrane, it can penetrate the cell membrane into the cytoplasm through endocytosis [29–31]. Complexation of the nanoparticle and dsRNA can avoid the degradation within the endocytic vesicles (early endosomes, late endosomes and lysosomes), and the nanoparticle can escape early and late endosomes through a process known as the sponge effect [101–103].

Nanoparticle-mediated dsRNA delivery has the potential to become a more sustainable and eco-friendly pest management method. The first attempt to study nanoparticle-mediated dsRNA delivery was conducted to silence *chitin synthase* genes in *Anopheles gambiae* using chitosan, and the improved RNAi efficiency was observed [104]. Similarly, the nanoparticle-mediated RNAi was then tested in *Spodoptera frugiperda* via a synthetic cationic polymer [105], *A. aegypti* via chitosan, carbon quantum dot and silica [106], *S. exigua* via a



guanylated polymer [107] and *Euschistus heros* and *Blattella germanica* via the liposome [108,109]. He *et al.* [100] designed a cationic core-shell fluorescent nanoparticle to deliver *CHT10* dsRNA through orally feeding and efficiently inhibited the normal development of *O. furnacalis*. A similar study was also performed in *Agrotis ypsilon* [110]. Zheng *et al.* [98] applied a fluorescent nanoparticle to deliver dsRNA to penetrate the aphid cuticle within 1 h. RNAi efficiency reaches 95.4%, and aphid population control effect reaches 80.5%. The transdermal dsRNA delivery system is a benefit for the development of sprayable RNA pesticides, which can be simply applied as chemical pesticides to achieve a high lethal effect. A star polycation was constructed recently as a highly efficient gene and botanical pesticide vector to increase pesticidal activities [97,111,112]. This nanoparticle/dsRNA formulation was sprayed directly on soybean seedlings with *Aphis glycines*, which resulted in a high mortality up to 78.5% [97]. The application of this nanoparticle to deliver pesticide and dsRNA at the same time may be a good option for foliar spray. In addition, nanoparticles can also facilitate the delivery of dsRNA in the *Arabidopsis* plant through the root tip [113], which is beneficial for the development of irrigation and trunk injection.

## CURRENT CHALLENGES IN APPLICATION OF RNAi-BASED PRODUCTS

### High production cost of dsRNA

The large-scale production of dsRNA, with low cost and high efficiency, must be developed for field application, and the dsRNA expression in bacteria seems to be a good alternative. Expression of dsRNA in bacteria strains deficient for RNaseIII is the major method, and the L1440-HT115(DE3) system is the most widely used dsRNA expression system that has been successfully applied in the RNAi of *Mythimna separate* [114], *L. decemlineata* [115] and *Bactrocera dorsalis* [116]. We applied the Scarless Cas9-assisted recombineering system to knock out the *rnc* gene in *Escherichia coli* BL21(DE3) and matched with the RNAi expression vector containing a single T7 promoter to construct a novel dsRNA expression system. The dsRNA expression efficiency of our system was about three-

times that of the L1440-HT115(DE3) system (unpublished data); however, the production efficiency of this system still needs improvement to meet the actual demands.

### Nontarget & off-target effects of siRNA

Nonspecific binding of siRNA may occur within the target and nontarget genomes. The off-target effects may be not a problem in pest management; however, the binding that occurs in nontarget organisms, such as predators and honeybees, may lead to some sublethal effects, which is difficult to predict [17]. The specificity of siRNA for corresponding mRNA is related to sequence homology, and the substantial sequence diversity between two molecules does not preclude gene silencing, which has been confirmed by Baum *et al.* [15]. The *vATPaseA* and *vATPaseE* sequences from *L. decemlineata* and WCR shared 83 and 79% nucleotide-sequence identities, respectively, and the dsRNAs targeting WCR *vATPaseA* and *vATPaseE* also reduced the fitness of *L. decemlineata*. As expected, the *L. decemlineata* dsRNAs appeared more activity than the orthologous WCR dsRNAs. So far, some computational design tools have been developed for the accurate and systemic evaluation of RNAi nontarget and off-target effects, but they should be used with proper bioassays to protect the exposed nontarget organisms.

### Potential RNAi resistance

Insect pests can develop resistance to RNAi-based products through various mechanisms as they do for conventional pesticides, including the mutations of target genes or core RNAi machinery genes, enhanced dsRNA degradation and lower dsRNA uptake [22,96]. Furthermore, the sequence polymorphism of target genes can cause the mismatch between dsRNA and mRNA, which can be potentially selected and lead to the evolution of resistance [117]. The WCR fed on transgenic maize expressing *DvSnf7* dsRNA was proved to exhibit resistance to dsRNA owing to the impaired luminal uptake, and this resistance was not *DvSnf7* dsRNA-specific, as indicated by cross-resistance to all other tested dsRNAs [118]. An RNAi-resistant cell strain of *L. decemlineata* expresses a low-level expression of coleopteran-specific Staufenc, which is required for RNAi and is a potential target for RNAi

resistance [119]. The commercialization of transplastomic crops may give pests a stronger selection pressure than nontransformative RNAi-based products, which may lead to resistance faster.

### Limitations of transgenic crops

The US Environmental Protection Agency and Canadian Food Inspection Agency have approved the application of transgenic crops based on RNAi technology. However, there are several issues to consider about environmental safety, especially foreign gene escape that may lead to some serious ecological consequences [73,120–122]. Although transplastomic crops are regarded as safe, even several magnitudes safer than nuclear-transformed crops attributed to their extranuclear inheritance, we should pay more attention to the horizontal transfer of antibiotic resistant marker genes that may result in some negative effects on humans and microorganisms in the natural environment [123,124]. Furthermore, the lack of proper transformation and especially selection and regeneration protocols to obtain fertile homoplastic crops are major problems in applying transplastomic technology in several major crops [124]. In addition, GM crops still cost more to produce and take a longer time for development [18].

### Potential risk of nanoparticle-mediated dsRNA delivery

The introduction of nanoparticles may bring potential risks for humans and environmental health, including the contamination of water sources and residues on food products [125]. So far, the development of nanopesticides and nanofertilizers has received less or at least delayed attention, and these nanoagrochemicals may be regarded as an intentional diffuse source of engineered nanoparticles in the environment [126]. A tiered approach focusing on key drivers of impact is typically used during their risk assessments, and each titer involves the estimation of a predicted environmental concentration, including the estimated concentration of the active substance in surface water, groundwater and soil [127]. Low cytotoxicity is a vital parameter for ideal dsRNA carrier. A series of fluorescent and nonfluorescent nanoparticles constructed by the group of Shen and Yin showed biocompatibility in ►

*vitro* and *in vivo*, with a much lower cytotoxicity than PEI [97,110,128–130], revealing their safety to some extent. However, studies on the environmental risk of nanoparticle/dsRNA formulation are very limited.

## FUTURE PERSPECTIVE

During the past two decades, RNAi has become an effective tool in functional genomics studies. Fast forward to today, the application of RNAi has helped scientists to find a possible solution to the global problems of agricultural losses attributed to insects and pathogens in a sustainable way. Recent studies reveal that this technology has raised enough attention and received ample funding support [131]. For GM crops expressing dsRNA, transplastomic crops seem to be a preferable strategy to achieve the improved effects. However, they are still considered a GM product in most countries, which requires the crops to undergo a rigorous evaluation before approval, and the extensive regulatory process is constraining the extension of transplastomic technology. The commercialization of SMARTSTAX PRO maize seems to be a good beginning. Furthermore, scientists should develop new chloroplast transformation protocols for major crops to promote the expansion of chloroplast-transformed crop range. For nontransformative RNAi products, the supply of dsRNAs associated with nanoparticles, through foliar spray, irrigation and trunk injection would be a great strategy to improve insecticidal activity, and other delivery methods, especially seed coats, still need to be evaluated. Bacteria-based expression of dsRNA is regarded as the most cost-effective method to produce large batch dsRNA, and some biotech companies are investing in this production method to produce affordable dsRNA for small and large farms [93]. Scientists should also pay more attention to public concerns regarding the specificity of dsRNA, fate of nanoparticle/dsRNA formulation in the environment, effects of RNAi-based products on nontarget organisms, and so on.

## AUTHOR CONTRIBUTIONS

S Yan and J Shen conceived the idea and reviewed the literature. S Yan, J Shen, B Zeng and B Ren revised the manuscript. S Yan wrote the manuscript and contributed to the generation of figures.

## FINANCIAL & COMPETING INTERESTS DISCLOSURE

S Yan is supported by the National Natural Science Foundation of China (grant no. 31900363) and the Beijing Natural Science Foundation (grant no. 6204043). The authors have no other relevant affiliations or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript apart from those disclosed.

No writing assistance was utilized in the production of this manuscript.

## OPEN ACCESS

This work is licensed under the Attribution-NonCommercial-NoDerivatives 4.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>

## REFERENCES

- Chiarappa L, Chiang HC, Smith RF. Plant pests and diseases: assessment of crop losses. *Science* 176(4036), 769–773 (1972).
- Bardner R, Fletcher KE. Insect infestations and their effects on the growth and yield of field crops: a review. *Bull. Entomol. Res.* 64(1), 141–160 (1974).
- Oerke EC. Crop losses to pests. *J. Agr. Sci.* 144(1), 31–43 (2006).
- Guo H, Sun Y, Li Y *et al.* Pea aphid promotes amino acid metabolism both in *Medicago truncatula* and bacteriocytes to favor aphid population growth under elevated CO<sub>2</sub>. *Global Change Biol.* 19(10), 3210–3223 (2013).
- Deutsch CA, Tewksbury JJ, Tigchelaar M *et al.* Increase in crop losses to insect pests in a warming climate. *Science* 361(6405), 916–919 (2018).
- Pimentel D. Amounts of pesticides reaching target pests: environmental impacts and ethics. *J. Agric. Environ. Ethics* 8(1), 17–29 (1995).
- Casida JE, Quistad GB. Golden age of insecticide research: past, present, or future? *Annu. Rev. Entomol.* 43, 1–16 (1998).
- Budzinski H, Couderchet M. Environmental and human health issues related to pesticides: from usage and environmental fate to impact. *Environ. Sci. Pollut. Res.* 25(15), 14277–14279 (2018).
- Fire A, Xu S, Montgomery MK *et al.* Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature* 391(6669), 806–811 (1998).
- Kim YH, Issa MS, Cooper AMW *et al.* RNA interference: applications and advances in insect toxicology and insect pest management. *Pest. Biochem. Phys.* 120, 109–117 (2015).
- Nandety RS, Kuo YW, Nouri S *et al.* Emerging strategies for RNA interference (RNAi) applications in insects. *Bioengineered* 6(1), 8–19 (2015).
- Mamta B, Rajam MV. RNAi technology: a new platform for crop pest control. *Physiol. Mol. Biol. Plants* 23(3), 487–501 (2017).
- Cai Q, He B, Kogel KH *et al.* Cross-kingdom RNA trafficking and environmental RNAi – nature's blueprint for modern crop protection strategies. *Curr. Opin. Microbiol.* 46, 58–64 (2018).
- Gordon KHJ, Waterhouse PM. RNAi for insect-proof plants. *Nat. Biotechnol.* 25(11), 1231–1232 (2007).
- Baum JA, Bogaert T, Clinton W *et al.* Control of coleopteran insect pests through RNA interference. *Nat. Biotechnol.* 25(11), 1322–1326 (2007).
- Head GP, Carroll MW, Evans SP *et al.* Evaluation of SmartStax and SmartStax PRO maize against western corn rootworm and northern corn rootworm: efficacy and resistance management. *Pest Manag. Sci.* 73(9), 1883–1899 (2017).

- Zotti MJ, Smagghe G. RNAi technology for insect management and protection of beneficial insects from diseases: lessons, challenges and risk assessments. *Neotrop. Entomol.* 44(3), 197–213 (2015).
- Cagliari D, Dias NP, Galdeano DM *et al.* Management of pest insects and plant diseases by non-transformative RNAi. *Front. Plant Sci.* 10, 1319 (2019).
- Dubrovina AS, Kiselev KV. Exogenous RNAs for gene regulation and plant resistance. *Int. J. Mol. Sci.* 20(9), 2282 (2019).
- Zhou R, Hotta I, Denli AM *et al.* Comparative analysis of argonaute-dependent small RNA pathways in *Drosophila*. *Mol. Cell* 32(4), 592–599 (2008).
- Han BW, Wang W, Li C *et al.* piRNA-guided transposon cleavage initiates Zucchini-dependent, phased piRNA production. *Science* 348(6236), 817–821 (2015).
- Zhu KY, Palli SR. Mechanisms, applications, and challenges of insect RNA interference. *Annu. Rev. Entomol.* 65, 14.1–14.19 (2020).
- Biryukova I, Ye T. Endogenous siRNAs and piRNAs derived from transposable elements and genes in the malaria vector mosquito *Anopheles gambiae*. *BMC Genomics* 16(1), 278 (2015).
- Swevers L, Liu J, Smagghe G. Defense mechanisms against viral infection in *Drosophila*: RNAi and non-RNAi. *Viruses* 10(5), 230 (2018).
- Yang ML, Wei YY, Jiang F *et al.* MicroRNA-133 inhibits behavioral aggregation by controlling dopamine synthesis in locusts. *PLoS Genet.* 10(2), e1004206 (2014).
- Lucas KJ, Zhao B, Luan J *et al.* Mosquito-specific microRNA-1890 targets the juvenile hormone-regulated serine protease JHA15 in the female mosquito gut. *RNA Biol.* 12(12), 1383–1390 (2015).
- Ninova A, Griffiths-Jones S, Ronshaugen M. Abundant expression of somatic transposon-derived piRNAs throughout *Tribolium castaneum* embryogenesis. *Genome Biol.* 18(1), 184 (2017).
- Feinberg EH, Hunter CP. Transport of dsRNA into cells by the transmembrane protein SID-1. *Science* 301(5639), 1545–1547 (2003).
- Saleh M, Rij RP, Hekele A *et al.* The endocytic pathway mediates cell entry of dsRNA to induce RNAi silencing. *Nat. Cell Biol.* 8(8), 793–802 (2006).
- Wynant N, Santos D, Van Wielendaele P *et al.* Scavenger receptor-mediated endocytosis facilitates RNA interference in the desert locust, *Schistocerca gregaria*. *Insect Mol. Biol.* 23(3), 320–329 (2014).
- Cappelle K, de Oliveira CFR, Van Eynde B *et al.* The involvement of clathrin-mediated endocytosis and teo Sid-1-like transmembrane proteins in double-stranded RNA uptake in the Colorado potato beetle midgut. *Insect Mol. Biol.* 25(3), 315–323 (2016).
- Borgio JF. RNAi mediated gene knockdown in sucking and chewing insect pests. *J. Biopestic.* 3(1), 386–393 (2010).
- Gong L, Chen Y, Hu Z *et al.* Testing insecticidal activity of novel chemically synthesized siRNA against *Plutella xylostella* under laboratory and field conditions. *PLoS ONE* 8(5), e62990 (2013).
- Li H, Jiang W, Zhang Z *et al.* Transcriptome analysis and screening for potential target genes for RNAi-mediated pest control of the beet armyworm, *Spodoptera exigua*. *PLoS ONE* 8(6), e65931 (2013).
- Frizzi A, Huang S. Tapping RNA silencing pathways for plant biotechnology. *Plant Biotechnol. J.* 8(6), 655–677 (2010).
- Koch A, Biedenkopf D, Furch A *et al.* An RNAi-based control of *Fusarium graminearum* infections through spraying of long dsRNAs involves a plant passage and is controlled by the fungal silencing machinery. *PLoS Pathog.* 12(10), e1005901 (2016).
- Wang M, Weiberg A, Lin FM *et al.* Bidirectional cross-kingdom RNAi and fungal uptake of external RNAs confer plant protection. *Nat. Plants* 2(10), 1615 (2016).
- Matranga C, Tomari Y, Shin C *et al.* Passenger-strand cleavage facilitates assembly of siRNA into Ago2-containing RNAi enzyme complex. *Cell* 123(4), 607–620 (2005).
- Marques JT, Kim K, Wu PH *et al.* Loqs and R2D2 act sequentially in the siRNA pathway in *Drosophila*. *Nat. Struct. Mol. Biol.* 17(1), 24–30 (2010).
- Pridgeon JW, Zhao L, Becnel JJ *et al.* Topically applied AaelAP1 double-stranded RNA kills female adults of *Aedes aegypti*. *J. Med. Entomol.* 45(3), 414–420 (2008).
- El-Shesheny I, Hajeri S, El-Hawary I *et al.* Silencing abnormal wing disc gene of the Asian citrus psyllid, *Diaphorina citri* disrupts adult wing development and increases nymph mortality. *PLoS ONE* 8(5), e65392 (2013).



42. Fishilevich E, Vélez AM, Storer NP *et al.* RNAi as a management tool for the western corn rootworm, *Diabrotica virgifera virgifera*. *Pest Manag. Sci.* 72(9), 1652–1663 (2016).
43. Gogoi A, Sarmah N, Kaldis A *et al.* Plant insects and mites uptake double-stranded RNA upon its exogenous application on tomato leaves. *Planta* 246(6), 1233–1241 (2017).
44. Dalakouras A, Jarausch W, Buchholz G *et al.* Delivery of hairpin RNAs and small RNAs into woody and herbaceous plants by trunk injection and petiole absorption. *Front. Plant Sci.* 9, 1253 (2018).
45. James C. A Global overview of biotech (GM) crops: adoption, impact and future prospects. *GM Crops* 1(1), 8–12 (2010).
46. Yan S, Yu J, Han M *et al.* Intercrops can mitigate pollen-mediated gene flow from transgenic cotton while simultaneously reducing pest densities. *Sci. Total Environ.* doi: <https://doi.org/10.1016/j.scitotenv.2019.134855> (2019) (Epub ahead of print).
47. Mao YB, Cai WJ, Wang JW *et al.* Silencing a cotton bollworm P450 monooxygenase gene by plant-mediated RNAi impairs larval tolerance of gossypol. *Nat. Biotechnol.* 25(11), 1307–1313 (2007).
48. Zhao Y, Sui X, Xu L *et al.* Plant-mediated RNAi of grain aphid *CHS1* gene confers common wheat resistance against aphids. *Pest Manag. Sci.* 74(12), 2754–2760 (2018).
49. Hou Q, Xu L, Liu G *et al.* Plant-mediated gene silencing of an essential olfactory-related *Gαq* gene enhances resistance to grain aphid in common wheat in greenhouse and field. *Pest Manag. Sci.* 75(6), 1718–1725 (2019).
50. Guo W, Bai C, Wang Z *et al.* Double-stranded RNAs high-efficiently protect transgenic potato from *Leptinotarsa decemlineata* by disrupting juvenile hormone biosynthesis. *J. Agric. Food Chem.* 66(45), 11990–11999 (2018).
51. Shen GM, Song CG, Ao YQY *et al.* Transgenic cotton expressing CYP392A4 double-stranded RNA decreases the reproductive ability of *Tetranychus cinnabarinus*. *Insect Sci.* 24(4), 559–568 (2017).
52. Han Q, Wang Z, He Y *et al.* Transgenic cotton plants expressing the *HaHR3* gene conferred enhanced resistance to *Helicoverpa armigera* and improved cotton yield. *Int. J. Mol. Sci.* 18(9), E1874 (2017).
53. Mao J, Zeng F. Plant-mediated RNAi of a gap gene-enhanced tobacco tolerance against the *Myzus persicae*. *Transgenic Res.* 23(1), 145–152 (2014).
54. Xiong Y, Zeng H, Zhang Y *et al.* Silencing the *HaHR3* gene by transgenic plant-mediated RNAi to disrupt *Helicoverpa armigera* development. *Int. J. Biol. Sci.* 9(4), 370–381 (2013).
55. Bolognesi R, Ramaseshadri P, Anderson J *et al.* Characterizing the mechanism of action of double-stranded RNA activity against western corn rootworm (*Diabrotica virgifera virgifera* LeConte). *PLoS ONE* 7(10), e47534 (2012).
56. Wang K, Peng Y, Pu J *et al.* Variation in RNAi efficacy among insect species is attributable to dsRNA degradation *in vivo*. *Insect Biochem. Mol. Biol.* 77, 1–9 (2016).
57. Arimatsu Y, Kotani E, Sugimura Y *et al.* Molecular characterization of a cDNA encoding extracellular dsRNase and its expression in the silkworm, *Bombyx mori*. *Insect Biochem. Mol. Biol.* 37(2), 176–183 (2002).
58. Allen ML, Walker III WB. Saliva of *Lygus lineolaris* digests double stranded ribonucleic acids. *J. Insect Physiol.* 58(3), 391–396 (2012).
59. Wynant N, Santos D, Verdonck R *et al.* Identification, functional characterization and phylogenetic analysis of double stranded RNA degrading enzymes present in the gut of the desert locust, *Schistocerca gregaria*. *Insect Biochem. Mol. Biol.* 46, 1–8 (2014).
60. Prentice K, Smaghe G, Gheysen G *et al.* Nuclease activity decreases the RNAi response in the sweet potato weevil *Cylas puncticollis*. *Insect Biochem. Mol. Biol.* 110, 80–89 (2019).
61. Song H, Zhang J, Li D *et al.* A double-stranded RNA degrading enzyme reduces the efficiency of oral RNA interference in migratory locust. *Insect Biochem. Mol. Biol.* 86, 68–80 (2017).
62. Guan RB, Li HC, Fan YJ *et al.* A nuclease specific to lepidopteran insects suppresses RNAi. *J. Biol. Chem.* 293(16), 6011–6021 (2018).
63. Spit J, Philips A, Wynant N *et al.* Knockdown of nuclease activity in the gut enhances RNAi efficiency in the Colorado potato beetle, *Leptinotarsa decemlineata*, but not in the desert locust, *Schistocerca gregaria*. *Insect Biochem. Mol. Biol.* 81, 103–116 (2017).
64. Bock R. Plastid biotechnology: prospects for herbicide and insect resistance, metabolic engineering and molecular farming. *Curr. Opin. Biotechnol.* 18(2), 100–106 (2007).
65. Bock R. Engineering plastid genomes: methods, tools, and applications in basic research and biotechnology. *Annu. Rev. Plant Biol.* 66, 211–241 (2015).
66. Kota M, Daniell H, Varma S *et al.* Overexpression of the *Bacillus thuringiensis* (Bt) Cry1Aa2 protein in chloroplasts confers resistance to plants against susceptible and Bt-resistant insects. *Proc. Natl Acad. Sci. USA* 96(5), 1840–1845 (1999).
67. De Cosa B, Moar W, Lee SB *et al.* Overexpression of the Bt cry2Aa2 operon in chloroplasts leads to formation of insecticidal crystals. *Nat. Biotechnol.* 19(1), 71–74 (2001).
68. Lutz KA, Knapp JE, Maliga P. Expression of bar in the plastid genome confers herbicide resistance. *Plant Physiol.* 125(4), 1585–1591 (2001).
69. Ye GN, Hajdukiewicz TJ, Broyles D *et al.* Plastid-expressed 5-enolpyruvylshikimate-3-phosphate synthase genes provide high level glyphosate tolerance in tobacco. *Plant J.* 25(3), 261–270 (2001).
70. Chen PJ, Senthilkumar R, Jane WN *et al.* Transplastomic *Nicotiana benthamiana* plants expressing multiple defence genes encoding protease inhibitors and chitinase display broad-spectrum resistance against insects, pathogens and abiotic stresses. *Plant Biotechnol. J.* 12(4), 503–515 (2014).
71. Zhang J, Khan SA, Hasse C *et al.* Full crop protection from an insect pest by expression of long double-stranded RNAs in plastids. *Science* 347(6225), 991–994 (2015).
72. Zhang J, Khan SA, Heckel DG *et al.* Next-generation insect-resistant plants: RNAi-mediated crop protection. *Trends Biotechnol.* 35(9), 871–882 (2017).
73. Andow DA, Zwahlen C. Assessing environmental risks of transgenic plants. *Ecol. Lett.* 9(2), 196–214 (2006).
74. Guo L, Zhu J, Sun S *et al.* Biosafety of transgenic crop: research and constraint of potential ecological risk of gene flow. *Crops* 2, 8–14 (2019).
75. Bally J, McIntyre GJ, Doran RL *et al.* In-plant protection against *Helicoverpa armigera* by production of long hpRNA in chloroplasts. *Front. Plant Sci.* 7, 1453 (2016).
76. Jin S, Singh ND, Li L *et al.* Engineered chloroplast dsRNA silences cytochrome p450 monooxygenase, V-ATPase and chitin synthase genes in the insect gut and disrupts *Helicoverpa armigera* larval development and pupation. *Plant Biotechnol. J.* 13(3), 435–446 (2015).
77. San Miguel K, Scott JG. The next generation of insecticides: dsRNA is stable as a foliar-applied insecticide. *Pest Manag. Sci.* 72(4), 801–809 (2016).
78. Zhang H, Li H, Guan R *et al.* Lepidopteran insect species-specific, broad-spectrum, and systemic RNA interference by spraying dsRNA on larvae. *Entomol. Exp. Appl.* 155(3), 218–228 (2015).
79. Killiny N, Hajeri S, Tiwari S *et al.* Double-stranded RNA uptake through topical application, mediates silencing of five CYP4 genes and suppresses insecticide resistance in *Diaphorina citri*. *PLoS ONE* 9(10), 3110536 (2014).
80. Niu J, Yang WJ, Tian Y *et al.* Topical dsRNA delivery induces gene silencing and mortality in the pea aphid. *Pest Manag. Sci.* 75(11), 2873–2881 (2019).
81. Hunter WB, Glick E, Paldi N *et al.* Advances in RNA interference: dsRNA treatment in trees and grapevines for insect pest suppression. *Southwest. Entomol.* 37(1), 85–87 (2012).
82. Li H, Guan R, Guo H *et al.* New insights into an RNAi approach for plant defence against piercing-sucking and stem-borer insect pests. *Plant Cell Environ.* 38(11), 2277–2285 (2015).
83. Luan JB, Ghanim M, Liu SS *et al.* Silencing the ecdysone synthesis and signaling pathway genes disrupts nymphal development in the whitefly. *Insect Biochem. Mol. Biol.* 43(8), 740–746 (2013).
84. Shang F, Ding BY, Xiong Y *et al.* Differential expression of genes in the alate and apterous morphs of the brown citrus aphid, *Toxoptera citricida*. *Sci. Rep.* 6, 32099 (2016).
85. Shang F, Niu JZ, Ding BY *et al.* Vitellogenin and its receptor play essential roles in the development and reproduction of the brown citrus aphid, *Aphis (Toxoptera) citricida*. *Insect Mol. Biol.* 27(2), 221–233 (2018).
86. Shang F, Ding BY, Ye C *et al.* Evaluation of a cuticle protein gene as a potential RNAi target in aphids. *Pest Manag. Sci.* 76(1), 134–140 (2020).
87. Ding BY, Shang F, Zhang Q *et al.* Silencing of two insulin receptor genes disrupts nymph-adult transition of alate brown citrus aphid. *Int. J. Mol. Sci.* 18(2), 357 (2017).
88. Dubelman S, Fischer J, Zapata F *et al.* Environmental fate of double-stranded RNA in agricultural soils. *PLoS ONE* 9(3), e93155 (2014).
89. Wise JC, VanWoerkom AH, Acimovic SG *et al.* Trunk injection: a discriminating delivering system for horticulture crop IPM. *Entomol. Ornithol. Herpetol.* 3(2), 126 (2014).
90. Berger C, Laurent F. Trunk injection of plant protection products to protect trees from pests and diseases. *Crop Prot.* 124, 104831 (2019).
91. Doering-Saad C, Newbury HJ, Bale JS *et al.* Use of aphid stylotomy and RT-PCR for the detection of transporter mRNAs in sieve elements. *J. Exp. Bot.* 53(369), 631–637 (2002).
92. Melnyk CW, Molnar A, Baulcombe DC. Intercellular and systemic movement of RNA silencing signals. *EMBO J.* 30(17), 3553–3563 (2011).
93. Joga MR, Zotti MJ, Smaghe G *et al.* RNAi efficiency, systemic properties, and novel delivery methods for pest insect control: what we know so far. *Front. Physiol.* 7, 553 (2016).
94. Kumar S, Nehra M, Dilbaghi N *et al.* Nano-based smart pesticide formulations: emerging opportunities for agriculture. *J. Control. Release* 294, 131–153 (2019).
95. Shen D, Zhou F, Xu Z *et al.* Systemically interfering with immune response by a fluorescent cationic dendrimer delivered gene suppression. *J. Mater. Chem. B* 2(29), 4653–4659 (2014).
96. Kunte N, McGraw E, Bell S *et al.* Prospects, challenges and current status of RNAi through insect feeding. *Pest Manag. Sci.* 76(1), 26–41 (2020).
97. Yan S, Qian J, Cai C *et al.* Spray method application of transdermal dsRNA delivery system for efficient gene silencing and pest control on soybean aphid *Aphis glycines*. *J. Pest Sci.* 93(1), 449–459 (2020).
98. Zheng Y, Hu Y, Yan S *et al.* A polymer/detergent formulation improves dsRNA penetration through the body wall and RNAi-induced mortality in the soybean aphid *Aphis glycines*. *Pest Manag. Sci.* 75(7), 1993–1999 (2019).
99. Yin M, Feng C, Shen J *et al.* Dual-responsive interaction to detect DNA on template-based fluorescent nanotubes. *Small* 7(12), 1629–1634 (2011).
100. He B, Chu Y, Yin M *et al.* Fluorescent nanoparticle delivered dsRNA toward genetic control of insect pests. *Adv. Mater.* 25(33), 4580–4584 (2013).
101. Benjaminsen RV, Mattheberg MA, Henriksen JR *et al.* The possible “proton sponge” effect of polyethyleneimine (PEI) does not include change in lysosomal pH. *Mol. Ther.* 21(1), 149–157 (2013).
102. Gilleron J, Querbes W, Zeigerer A *et al.* Image-based analysis of lipid nanoparticle-mediated siRNA delivery, intracellular trafficking and endosomal escape. *Nat. Biotechnol.* 31(7), 638–646 (2013).
103. Selby LI, Cortez-Jugo CM, Such GK *et al.* Nanoescapology: progress toward understanding the endosomal escape of polymeric nanoparticles. *Wiley Interdiscip. Rev. Nanomed. Nanobiotechnol.* 9(5), e52523 (2017).
104. Zhang X, Zhang J, Zhu KY. Chitosan/double-stranded RNA nanoparticle-mediated RNA interference to silence chitin synthase genes through larval feeding in the African malaria mosquito (*Anopheles gambiae*). *Insect Mol. Biol.* 19(5), 683–693 (2010).
105. Parsons KH, Mondal MH, McCormick CL *et al.* Guanidinium-functionalized interpolyelectrolyte complexes enabling RNAi in resistant insect pests. *Biomacromolecules* 19(4), 1111–1117 (2018).
106. Das S, Debnath N, Cui Y *et al.* Chitosan, carbon quantum dot, and silica nanoparticle mediated dsRNA delivery for gene silencing in *Aedes aegypti*: a comparative analysis. *ACS Appl. Mater. Interfaces* 7(35), 19530–19535 (2015).
107. Christiaens O, Tardajos MG, Reyna ZLM *et al.* Increased RNAi efficacy in *Spodoptera exigua* via the formulation of dsRNA with guanylated polymers. *Front. Physiol.* 9, 316 (2018).
108. Lin YH, Huang JH, Liu Y *et al.* Oral delivery of dsRNA lipoplexes to German cockroach protects dsRNA from degradation and induces RNAi response. *Pest Manag. Sci.* 73(5), 960–966 (2017).
109. Castellanos NL, Smaghe G, Sharma R *et al.* Liposome encapsulation and EDTA formulation of dsRNA targeting essential genes increase oral RNAi-caused mortality in the neotropical stink bug *Euschistus heros*. *Pest Manag. Sci.* 75(2), 537–548 (2019).
110. Xu Z, He B, Wei W *et al.* Highly water-soluble perylene-diimide-cored poly(amido amine) vector for efficient gene transfection. *J. Mater. Chem. B* 2(20), 3079–3086 (2014).
111. Li J, Qian J, Xu Y *et al.* A facile-synthesized star polycation constructed as a highly efficient gene vector in pest management. *ACS Sustain. Chem. Eng.* 7(6), 6316–6322 (2019).
112. Yan S, Hu Q, Li J *et al.* A star polycation acts as a drug nanocarrier to improve the toxicity and persistence of botanical pesticides. *ACS Sustain. Chem. Eng.* 7(20), 17406–17413 (2019).

113. Jiang L, Ding L, He B *et al.* Systemic gene silencing in plants triggered by fluorescent nanoparticle-delivered double-stranded RNA. *Nanoscale* 6(17), 9965–9969 (2014).
114. Ganbaatar O, Cao B, Zhang Y *et al.* Knockdown of *Mythimna separata chitinase* genes via bacterial expression and oral delivery of RNAi effectors. *BMC Biotechnol.* 17(1), 9 (2017).
115. Zhu F, Xu J, Palli R *et al.* Ingested RNA interference for managing the populations of the Colorado potato beetle, *Leptinotarsa decemlineata*. *Pest Manag. Sci.* 67(2), 175–182 (2011).
116. Li X, Zhang M, Zhang H. RNA interference of four genes in adult *Bactrocera dorsalis* by feeding their dsRNAs. *PLoS ONE* 6(3), e17788 (2011).
117. Scott JG, Michel K, Bartholomay LC *et al.* Towards the elements of successful insect RNAi. *J. Insect Physiol.* 59(12), 1212–1221 (2013).
118. Khajuria C, Ivashuta S, Wiggins E *et al.* Development and characterization of the first dsRNA-resistant insect population from western corn rootworm, *Diabrotica virgifera virgifera* LeConte. *PLoS ONE* 13(5), e0197059 (2018).
119. Yoon JS, Mogilicherla K, Gurusamy D *et al.* Double-stranded RNA binding protein, Staufen, is required for the initiation of RNAi in coleopteran insects. *Proc. Natl Acad. Sci. USA* 115(33), 8334–8339 (2018).
120. Wolf DE, Takebayashi N, Rieseberg LH. Predicting the risk of extinction through hybridization. *Conserv. Biol.* 15(4), 1039–1053 (2001).
121. Chandler S, Dunwell JW. Gene flow, risk assessment and the environmental release of transgenic plants. *Crit. Rev. Plant Sci.* 27(1), 25–49 (2008).
122. Randall BW, Walton DA, Lee DJ *et al.* The risk of pollen-mediated gene flow into a vulnerable eucalypt species. *Forest Ecol. Manag.* 381, 297–304 (2016).
123. Miki B, McHugh S. Selectable marker genes in transgenic plants: applications, alternatives and biosafety. *J. Biotechnol.* 107(3), 193–232 (2004).
124. Wani SH, Sah SK, Sági L *et al.* Transplastomic plants for innovations in agriculture. A review. *Agron. Sustain. Dev.* 35(4), 1391–1430 (2015).
125. Kah M. Nanopesticides and nanofertilizers: emerging contaminants or opportunities for risk mitigation? *Front. Chem.* 3, 64 (2015).
126. Kah M, Beulke S, Tiede K *et al.* Nanopesticides: state of knowledge, environmental fate, and exposure modeling. *Crit. Rev. Environ. Sci. Technol.* 43(16), 1823–1867 (2013).
127. Kookana RS, Boxall ABA, Reeves PT *et al.* Nanopesticides: guiding principles for regulatory evaluation of environmental risks. *J. Agr. Food Chem.* 62(19), 4227–4240 (2014).
128. You S, Cai Q, Zheng Y *et al.* Perylene-cored star-shaped polycations for fluorescent gene vectors and bioimaging. *ACS Appl. Mater. Interfaces* 6(18), 16327–16334 (2014).
129. Liu X, Zheng Y, Zhang S *et al.* Peryleneimide-cored cationic nanocarriers deliver virus DNA to kill insect pests. *Polymer. Chem.* 7(22), 3740–3746 (2016).
130. Gao Z, Huang W, Zheng Y *et al.* Facile synthesis of core-shell magnetic-fluorescent nanoparticles for cell imaging. *RSC Adv.* 6(52), 46226–46230 (2016).
131. Jalaluddin NSM, Othman RY, Harikrishna JA. Global trends in research and commercialization of exogenous and endogenous RNAi technologies for crops. *Crit. Rev. Biotechnol.* 39(1), 67–78 (2019).