



POSTER PRESENTATION

Open Access

Modulation of the immune response in HTLV-1-infections by SM29 antigen in vitro is dependent of IL-10, TGF- β and CTLA-4

Luciane Mota Lima¹, Silvane Braga Santos¹, Ricardo Riccio Oliveira¹, Luciana Santos Cardoso¹, Sergio Costa Oliveira², Alfredo Miranda Góes², Alex Loukas³, Edgar M De Carvalho¹, Maria Ilma Araújo^{1*}

From 16th International Conference on Human Retroviruses: HTLV and Related Viruses
Montreal, Canada. 26-30 June 2013

The HTLV-1 is the causal agent of HTLV-1-associated Myelopathy/Tropical Spastic Paraparesis (HAM/TSP). The immune response in HTLV-1 infection is polarized to the Th1-type, and previous data from our group had demonstrated that the use of *Schistosoma* antigens to cell cultures in HTLV-1 infection resulted in down-modulation of Th1/inflammatory response. The aim of this study was to evaluate whether the modulation of inflammatory response by *S. mansoni* antigens depends upon the presence of IL-10, TGF- β and CTLA-4. The antibodies anti-IL-10, anti-TGF- β and anti-CTLA-4 with or without the *S. mansoni* antigen Sm29 were added to the PBMC cultures of HTLV-1- infected individuals and the levels of cytokines in the supernatants were measure using ELISA sandwich method. Compared to the levels of cytokine in non stimulated cultures (1.073 pg/ml, median values), the levels of IFN-g production (552 pg/ml, 1.002 pg/ml and 548 pg/ml, respectively; p<0.05). On the other hand, the levels of IL-10 were increased by the presence of Sm29 (it passed from 26 pg/ml to 102 pg/ml; p=0.02), however when anti-TGF- β and anti-CTLA-4 were added to the cultures there was a reduction in the levels of IL-10 (19 pg/ml and 23 pg/ml, respectively; p<0.05). The regulation of IFN- γ production by Sm29 antigen in HTLV-1-infected individuals is dependent on IL-10, TGF- β and CTLA-4. Also the levels of IL-10 were affected by the neutralization of TGF- β and CTLA-4.

Financial support

CNPQ (Universal 479417/2008 3), NIH (R01AI079238A).

* Correspondence: marialma.araujo@gmail.com

¹Serviço de Imunologia, Hospital Universitário Professor Edgard Santos, Universidade Federal da Bahia, Salvador-BA, Brazil

Full list of author information is available at the end of the article

Authors' details

¹Serviço de Imunologia, Hospital Universitário Professor Edgard Santos, Universidade Federal da Bahia, Salvador-BA, Brazil. ²Instituto de Ciências Biológicas, Departamento de Bioquímica e Imunologia, Universidade Federal de Minas Gerais, Belo Horizonte-MG, Brazil. ³Division of Infectious Diseases, Queensland Institute of Medical Research, Brisbane, Queensland, Australia.

Published: 7 January 2014

doi:10.1186/1742-4690-11-S1-P68

Cite this article as: Lima et al.: Modulation of the immune response in HTLV-1-infections by SM29 antigen in vitro is dependent of IL-10, TGF- β and CTLA-4. *Retrovirology* 2014 **11**(Suppl 1):P68.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

