W J C C World Journal of Clinical Cases

Submit a Manuscript: https://www.f6publishing.com

World J Clin Cases 2022 August 16; 10(23): 8255-8261

DOI: 10.12998/wjcc.v10.i23.8255

ISSN 2307-8960 (online)

CASE REPORT

Sequential multidisciplinary minimally invasive therapeutic strategy for heart failure caused by four diseases: A case report

Chen-Ze Zhao, Yan Yan, Yong Cui, Ni Zhu, Xue-Yan Ding

Specialty type: Cardiac and cardiovascular systems

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0 Grade B (Very good): B, B Grade C (Good): 0 Grade D (Fair): 0 Grade E (Poor): 0

P-Reviewer: Mydam J, United States; Ong H, Malaysia

Received: December 8, 2021 Peer-review started: December 8, 2021 First decision: June 7, 2022 Revised: June 18, 2022 Accepted: July 11, 2022 Article in press: July 11, 2022 Published online: August 16, 2022



Chen-Ze Zhao, Xue-Yan Ding, Department of Cardiology, Zhejiang Provincial People's Hospital, Hangzhou 310014, Zhejiang Province, China

Yan Yan, Department of Cardiothoracic Surgery, No. 903 Hospital of Chinese People's Liberation Army, Hangzhou 310043, Zhejiang Province, China

Yong Cui, Department of Cardiothoracic Surgery, Zhejiang Provincial People's Hospital, Hangzhou 310014, Zhejiang Province, China

Ni Zhu, Department of Hematology, The First Affiliated Hospital of Zhejiang Chinese Medicine University, Hangzhou 310003, Zhejiang Province, China

Corresponding author: Xue-Yan Ding, MD, Doctor, Department of Cardiology, Zhejiang Provincial People's Hospital, No. 158 Shangtang Road, Hangzhou 310014, Zhejiang Province, China. dingxueyan1213@163.com

Abstract

BACKGROUND

The coexistence with patent ductus arteriosus (PDA), mitral valve prolapse (MVP), atrial fibrillation (AF) and hyperthyroidism is extremely rare and complex. The optimal therapeutic strategy is difficult to develop.

CASE SUMMARY

A 27-year-old female with PDA, MVP, AF and hyperthyroidism presented with severe dyspnea. Given that a one-stage operation for PDA, MVP and AF is high risk, we preferred a sequential multidisciplinary minimally invasive therapeutic strategy. First, PDA transcatheter closure was performed. Hyperthyroidism and heart failure were simultaneously controlled via medical treatment. Video-assisted thoracoscopic mitral valve repair and left atrial appendage occlusion were performed when heart failure was controlled. Under this therapeutic strategy, the patient's sinus rhythm was restored and maintained. Two years after the treatment, the symptoms of heart failure were relieved, and the enlarged heart was reversed.

CONCLUSION

Sequential multidisciplinary therapeutic strategies, which take advantage of both internal medicine and surgical approaches, might be reasonable for this type of disease.

WJCC | https://www.wjgnet.com

Key Words: Patent ductus arteriosus; Mitral valve prolapse; Atrial fibrillation; Heart failure; Sequential multidisciplinary therapeutic strategy; Case report

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: The coexistence of patent ductus arteriosus (PDA), mitral valve prolapse, atrial fibrillation and hyperthyroidism is extremely rare and complex. We proposed a successful sequential multidisciplinary therapeutic strategy for a 27-year-old female suffering from these four diseases in addition to severe heart failure. PDA transcatheter closure, medical treatment and thoracoscopic mitral valve repair were performed sequentially. Two years after the treatment, the symptoms of heart failure were relieved, sinus rhythm was restored, and the enlarged heart was reversed.

Citation: Zhao CZ, Yan Y, Cui Y, Zhu N, Ding XY. Sequential multidisciplinary minimally invasive therapeutic strategy for heart failure caused by four diseases: A case report. World J Clin Cases 2022; 10(23): 8255-8261 URL: https://www.wjgnet.com/2307-8960/full/v10/i23/8255.htm DOI: https://dx.doi.org/10.12998/wjcc.v10.i23.8255

INTRODUCTION

Patent ductus arteriosus (PDA), mitral valve prolapse (MVP), atrial fibrillation (AF) and hyperthyroidism are common diseases, but the combination of these four diseases is extremely rare. Transcatheter closure is the preferred treatment for adult PDA[1]. Catheter ablation is the first-line treatment for nonvalvular AF. However, surgery is the first choice for MVP[2]. Additionally, although simultaneous surgical management for these three diseases may be a choice, the risk of three operations over the same period is high given the uncontrolled hyperthyroidism and severe heart failure. Therefore, the optimal therapeutic strategy is difficult to develop. We herein introduce a successful sequential multidisciplinary minimally invasive therapeutic strategy for treating a 27-year-old female with severe heart failure caused by the combination of these four diseases.

CASE PRESENTATION

Chief complaints

A 27-year-old female was referred to our hospital complaining of palpitations, irritability, edema of the legs and dyspnea for over ten years. She presented with severe shortness of breath and paroxysmal nocturnal dyspnea for 20 d.

History of present illness

The patient developed palpitations, irritability, edema of the legs and dyspnea for over ten years. She was initially diagnosed with hyperthyroidism but failed to achieve a euthyroid state. One month prior, she was diagnosed with PDA, MVP with severe mitral regurgitation (MR) and heart failure [New York Heart Association (NYHA) Classification III] while being treated for hyperthyroidism at a previous hospital. The cardiothoracic surgeons planned to perform one-stage surgery through a standard full median sternotomy when her heart failure had been controlled. She left the hospital and took 7.5 mg qd methimazole, 11.875 mg qd metoprolol, 20 mg qd furosemide and 20 mg qd spironolactone. Twenty days before presentation, she developed severe shortness of breath and paroxysmal nocturnal dyspnea. Given that her heart failure was refractory to medical therapy, she was transferred to our hospital.

History of past illness

The patient had no history of past illness.

Personal and family history

The patient had no personal or family history.

Physical examination

Physical examination showed continuous grade 4/6 murmurs at the left second to third intercostal spaces, systolic grade 3/6 murmurs at the apex with AF, moist rales in both lower lungs and severe edema of both legs.



Laboratory examinations

Laboratory examination showed elevated brain natriuretic peptide (BNP, 890.3 pg/mL) and decreased thyroid hormone (TT3 0.53 μg/L, normal 0.66-1.61 μg/L; TT4 36.3 μg/L, normal 54.4-118.5 μg/L; TSH 0.00 mIU/L, normal 0.34-5.6 mIU/L; FT3 2.45 ng/L, normal 2.14-4.21 ng/L; FT4 5.22 ng/L, normal 5.9-12.5 ng/L).

Imaging examinations

Echocardiography showed generalized cardiac enlargement, 5.1 mm PDA with a continuous shunt (Figure 1A) and anterior mitral leaflet prolapse with severe MR (Figure 2A and B). The estimated pulmonary artery systolic pressure was 68 mmHg, and the estimated right ventricular pressure was 55 mmHg. The electrocardiogram showed AF with a rapid ventricular rate (Figure 3A). Chest computed tomography scan showed pulmonary edema and bilateral pleural effusions.

FINAL DIAGNOSIS

The final diagnosis was PDA, MVP with severe MR, AF, hyperthyroidism, and heart failure (NYHA Classification IV).

TREATMENT

One-stage surgery for PDA, MVP and AF is high risk due to severe refractory heart failure (NYHA Classification IV). Moreover, the patient wanted to avoid median sternotomy given that it results in a large scar on the chest. After a thorough discussion with endocrinologists, cardiologists and cardiothoracic surgeons, a sequential multidisciplinary minimally invasive therapeutic strategy was formulated. First, PDA transcatheter closure was performed. Hyperthyroidism and heart failure were controlled simultaneously via medical treatment. Video-assisted thoracoscopic mitral valve repair and left atrial appendage (LAA) occlusion were performed when her heart failure was controlled.

An 8-10 mm SHSMA PDA duct occluder (Shanghai Shape Memory Alloy Co. Ltd. Shanghai, China) was deployed across the PDA by catheterization through a sheath introduced from the right femoral vein under local anesthesia (Figure 4). Complete closure of the ductus was obtained immediately. As little contrast agent as possible was injected to minimize the impact on the thyroid. After catheterization, the patient was administered sacubitril valsartan sodium (25 mg bid) and a lower dose of methimazole (2.5 mg qd).

Two months later, she exhibited mild exertional dyspnea (NYHA Classification II). Video-assisted thoracoscopic mitral valve repair and LAA occlusion were then performed successfully through a minimal right infra-axillary thoracotomy using femorofemoral extracorporeal circulation. Chordal replacement was used to correct prolapse of the anterior leaflet with three artificial chordaes made of expanded polytetrafluoroethylene sutures (Gore-Tex sutures; W.L. Gore & Associates Inc, Elkton, Md). Annuloplasty ring implantation was performed (32-mm annuloplasty ring, Medtronic Inc., Minnesota, United States). The patient also underwent LAA exclusion using an LAA clip device (PM-LAA-40, Beijing Puhui Biomedical Engineering Co., Ltd, Beijing, China). After weaning from cardiopulmonary bypass, intraoperative transesophageal echocardiography was used to confirm successful LAA occlusion and mitral valve repair (Figure 5). After surgery, the dose of sacubitril valsartan sodium (50 mg bid) was increased.

OUTCOME AND FOLLOW-UP

Sinus rhythm was regained 3 mo after surgery (Figure 3B). Two years after the treatment, the symptoms of heart failure were relieved (NYHA Classification I), and sinus rhythm was maintained. Transthoracic echocardiography showed complete closure of the ductus (Figure 1B) and mind MR (Figure 2D). The enlarged heart gradually shrunk (Table 1). Blood BNP also notably decreased (129.1 vs 890.3 pg/mL).

DISCUSSION

PDA, MVP, AF, and hyperthyroidism are common diseases, but the combination of these four diseases is extremely rare. No previous report has provided a therapeutic strategy for the combination of these four diseases. Minimally invasive and multidisciplinary strategies are the current trends in clinical treatment[3]. Therefore, we developed a sequential multidisciplinary minimally invasive therapeutic strategy based on the pathophysiological characteristics and main treatment methods of these four



Zhao CZ et al. Multidisciplinary therapeutic strategy for heart failure

Table 1 Echocardiographic assessment						
Echocardiographic time	LVEF (%)	LVIDd (mm)	LVIDs (mm)	LA (mm)	RV (mm)	RA (mm)
Pre-PDA closure	57	59	41	50	30	45
Post-PDA closure	69	62	38	49	33	41
Pre-surgery	60	60	40	51	25	46
Post-surgery	51	55	41	39	26	46
Three months after surgery	62	48	32	39	23	44
Six months after surgery	68	49	31	44	23	33
One year after surgery	67	51	32	39	20	34
Two years after surgery	67	52	32	43	20	35

LA: Left atrial anteroposterior diameter; LVEF: Left ventricular ejection fraction; LVIDd: Left ventricular internal diameter in end-diastole; LVIDs: Left ventricular internal diameter in end-systole; PDA: Patent ductus arteriosus; RA: Right atrial diameter; RV: Right ventricular diameter.



DOI: 10.12998/wjcc.v10.i23.8255 Copyright ©The Author(s) 2022.

Figure 1 Patent ductus arteriosus assessment by transthoracic echocardiography. A: A 5.1-mm patent ductus arteriosus before occlusion (arrow); B: No residual shunt 2 years after occlusion (arrow).

> diseases. Hyperthyroidism can lead to AF and high-output heart failure. The long-standing MVP of the patient results in primary mitral regurgitation for many years, during which time the size of the left atrium and ventricle increase and the contractility of the left ventricle decreases. Mitral valve repair is recommended in symptomatic patients with MVP alone and low surgical risk^[2]. PDA results in left-toright shunt and left ventricular volume overload and remodeling, which can also lead to heart failure. Device closure is recommended, as this procedure has excellent technical success and minimal morbidity, supplanting operative ligation in adults^[4]. The treatment of AF includes antiarrhythmic drugs, catheter ablation and surgical ablation. However, AF may be secondary to hyperthyroidism and severe mitral regurgitation in this case. Therefore, MVP and PDA should be treated first after hyperthyroidism is controlled via drugs.

> The traditional approach for treating congenital heart disease with valvular disease is one-stage surgery with median sternotomy. A study showed that one-stage open-heart mitral valve repair, tricuspid annuloplasty, PDA direct closure and radiofrequency-modified maze procedures were successfully performed in a 73-year-old female^[5]. Recently, a one-stage hybrid procedure was performed as a valuable alternative with the advantages of reducing trauma as well as recovery and hospitalization time[6-8]. However, the young female wanted to avoid median sternotomy due to its huge scarring on the chest. However, the patient had a high surgical risk due to severe heart failure. Therefore, one-stage surgery might not represent the ideal therapeutic strategy for this patient.

> Transcatheter closure is the preferred treatment for adult PDA with minimal anesthetic risk and incidence of associated complications^[1]. Moreover, a few reports have confirmed that transcatheter PDA occlusion can reduce the volume load of the left ventricle and functional mitral regurgitation [9, 10]. Closing the PDA is likely to reverse left atrial and ventricular enlargement and will possibly provide symptom relief[11]. Based on the above reasons, transcatheter PDA occlusion combined with antithyroid drug therapy was the optimal choice as the initial treatment with low risk and trauma. As little contrast agent as possible was injected to minimize the impact on the thyroid. After PDA occlusion and optimization of drug therapy for 2 mo, her heart failure improved significantly (NYHA Classi-



WJCC | https://www.wjgnet.com



DOI: 10.12998/wjcc.v10.i23.8255 Copyright ©The Author(s) 2022.

Figure 2 Mitral valve assessment by transthoracic echocardiography. A and B: Echocardiography before surgery showed mitral anterior leaflet prolapse with severe eccentric regurgitation; C and D: Two years after surgery, echocardiography showed mild mitral regurgitation.



DOI: 10.12998/wjcc.v10.i23.8255 Copyright ©The Author(s) 2022.



fication II).

Valve repair combined with maze surgery is the standard treatment for patients with MVP and AF. However, AF may be secondary to hyperthyroidism and severe mitral regurgitation in this case. Accordingly, we did not perform the Maze procedure, which would cause substantial trauma to the left atrium. The minimal right axillary incision was selected for mitral valve repair with satisfactory cosmetic results. The LAA was simultaneously clipped to avoid thrombosis[12]. Consistent with expectations, sinus rhythm was recovered and maintained. Therefore, catheter ablation was not needed, and both the injury and additional financial expenditure of AF ablation to the patient were avoided.

Although multiple hospitalizations are needed, the multidisciplinary sequential minimally invasive therapeutic strategy can significantly reduce the surgical risk, avoid the trauma and complications of median sternotomy, and shorten the recovery time compared with one-stage surgery in this case.

Baishideng® WJCC | https://www.wjgnet.com



DOI: 10.12998/wjcc.v10.i23.8255 Copyright ©The Author(s) 2022.

Figure 4 Transcatheter patent ductus arteriosus occlusion. A: Patent ductus arteriosus (PDA) in the catheterization (arrow); B and C: complete closure of PDA during catheterization and the PDA occluder (arrow).



DOI: 10.12998/wjcc.v10.i23.8255 Copyright ©The Author(s) 2022.

Figure 5 Intraoperative transesophageal echocardiography. A: Left atrial appendage assessment; B: Left atrial appendage occlusion; C: Severe eccentric mitral regurgitation before chordal replacement; D: Decreased mitral regurgitation after chordal replacement.

CONCLUSION

Our successful experience in this patient shows that a sequential multidisciplinary therapeutic strategy, which can take advantage of both internal medicine and surgical approaches, might be reasonable for this type of disease.

FOOTNOTES

Author contributions: Zhao CZ wrote the manuscript; Yan Y and Zhu N revised the manuscript; Cui Y participated in the operation; Ding XY participated in the operation and revised the manuscript; All authors have read and approved the final manuscript.

Supported by National Natural Science Foundation of China, No. 81800342 and 81800138; and Zhejiang Provincial



Raisbideng® WJCC | https://www.wjgnet.com

Natural Science Foundation of China, No. LQ20H020012.

Informed consent statement: Informed written consent was obtained from the patient for publication of this report and any accompanying images.

Conflict-of-interest statement: The authors declare that there are no conflicts of interest related to this report.

CARE Checklist (2016) statement: The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: https://creativecommons.org/Licenses/by-nc/4.0/

Country/Territory of origin: China

ORCID number: Chen-Ze Zhao 0000-0002-6694-2506; Yan Yan 0000-0002-6693-0724; Yong Cui 0000-0002-4703-5038; Ni Zhu 0000-0002-3418-3365; Xue-Yan Ding 0000-0001-7853-3235.

S-Editor: Zhang H L-Editor: A P-Editor: Zhang H

REFERENCES

- Schneider DJ, Moore JW. Patent ductus arteriosus. Circulation 2006; 114: 1873-1882 [PMID: 17060397 DOI: 10.1161/CIRCULATIONAHA.105.592063]
- 2 Vahanian A, Beyersdorf F, Praz F, Milojevic M, Baldus S, Bauersachs J, Capodanno D, Conradi L, De Bonis M, De Paulis R, Delgado V, Freemantle N, Gilard M, Haugaa KH, Jeppsson A, Jüni P, Pierard L, Prendergast BD, Sádaba JR, Tribouilloy C, Wojakowski W; ESC/EACTS Scientific Document Group. 2021 ESC/EACTS Guidelines for the management of valvular heart disease. *Eur Heart J* 2022; 43: 561-632 [PMID: 34453165 DOI: 10.1093/eurheartj/chab395]
- 3 Sugiyama Y, Sasaki M, Kouyama M, Tazaki T, Takahashi S, Nakamitsu A. Current treatment strategies and future perspectives for gastrointestinal stromal tumors. *World J Gastrointest Pathophysiol* 2022; 13: 15-33 [PMID: 35116177 DOI: 10.4291/wjgp.v13.i1.15]
- 4 O'Byrne ML, Smith CL, Gillespie MJ. Device Closure of Patent Ductus Arteriosus in Adults. Can J Cardiol 2020; 36: 795-796 [PMID: 32173055 DOI: 10.1016/j.cjca.2019.10.022]
- 5 Yoshida K, Tobe S, Adachi K, Kawata M. [Adult mitral and tricuspid valve regurgitation due to patent ductus arteriosus combined with atrial fibrillation; report of a case]. *Kyobu Geka* 2004; **57**: 1127-1130 [PMID: 15553030]
- 6 Li Q, Lin K, Gan CP, Feng Y. One-Stage Hybrid Procedure to Treat Aortic Coarctation Complicated by Intracardiac Anomalies in Two Adults. *Ann Thorac Surg* 2015; 100: 2364-2367 [PMID: 26652538 DOI: 10.1016/j.athoracsur.2015.02.130]
- 7 **Guo H**, Wang Y, Wu W, Lai Y. One-Stage Hybrid Procedure for Patent Ductus Arteriosus Combined with Other Cardiac Anomalies in Adults. *Heart Surg Forum* 2018; **21**: E177-E178 [PMID: 29893676 DOI: 10.1532/hsf.1794]
- 8 Li CS, Lu Z, Song XR, Yan ZY. Hybrid procedure for treating adult congenital heart disease with valvular heart disease in two patients. J Cardiothorac Surg 2019; 14: 180 [PMID: 31647015 DOI: 10.1186/s13019-019-1002-z]
- 9 Wang Z, Chen T, Chen L, Qin Y, Zhao X. Safety and Efficacy of Transcatheter Closure of Patent Ductus Arteriosus With Severe Mitral Regurgitation in Adults. *J Invasive Cardiol* 2016; 28: 30-33 [PMID: 26716592]
- 10 Watanabe N, Toyofuku M, Sato T, Shiode N, Masaoka Y, Muraoka Y, Okimoto T, Otsuka M, Tamekiyo H, Mito S, Kawase T, Kagawa Y, Senoo A, Hayashi Y. A case of adult patient ductus arteriosus with congestive heart failure and severe mitral regurgitation. *Cardiovasc Interv Ther* 2011; 26: 278-280 [PMID: 24122597 DOI: 10.1007/s12928-011-0069-9]
- Stout KK, Daniels CJ, Aboulhosn JA, Bozkurt B, Broberg CS, Colman JM, Crumb SR, Dearani JA, Fuller S, Gurvitz M, Khairy P, Landzberg MJ, Saidi A, Valente AM, Van Hare GF. 2018 AHA/ACC Guideline for the Management of Adults With Congenital Heart Disease: Executive Summary: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Circulation* 2019; 139: e637-e697 [PMID: 30586768 DOI: 10.1161/CIR.000000000000602]
- Bedeir K, Warriner S, Kofsky E, Gullett C, Ramlawi B. Left Atrial Appendage Epicardial Clip (AtriClip): Essentials and Post-Procedure Management. J Atr Fibrillation 2019; 11: 2087 [PMID: 31384360 DOI: 10.4022/jafib.2087]

WJCC https://www.wjgnet.com



Published by Baishideng Publishing Group Inc 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA Telephone: +1-925-3991568 E-mail: bpgoffice@wjgnet.com Help Desk: https://www.f6publishing.com/helpdesk https://www.wjgnet.com

