

## RESEARCH ARTICLE

# Procalcitonin as a prognostic marker of patients with acute ischemic stroke

Ling Yan | Shuling Wang | Lanlan Xu | Zhen Zhang | Pu Liao 

Department of Clinical Laboratory,  
Chongqing General Hospital, Chongqing,  
China

**Correspondence**

Pu Liao, Department of Clinical Laboratory,  
Chongqing General Hospital, Chongqing,  
China.  
Email: liaopu2010@126.com

**Abstract**

**Background:** The information on mortality after an acute stroke patient is still limited.

**Objectives:** The aim of this study was to investigate the prognostic potential of procalcitonin (PCT) serum levels in acute ischemic stroke.

**Methods:** A total of 748 patients were enrolled in this study. Prognostic potential of PCT was evaluated by Kaplan-Meier analysis and multivariable Cox hazard regression analyses.

**Results:** Procalcitonin levels were found to be significantly higher in acute ischemic stroke patients who died in 30 days than those who survived. Univariate logistic regression analysis showed that PCT was significantly associated with 30-day mortality, and Cox regression analysis revealed that PCT was a strong predictor of 30-day overall mortality. Kaplan-Meier analysis revealed that overall cumulative 30-day mortality was significantly higher in those with PCT levels >1.5 ng/mL when compared to those with PCT levels <1.5 ng/mL.

**Conclusions:** Procalcitonin is a significant independent prognostic marker of 30-day mortality after the one set of acute ischemic stroke.

## 1 | INTRODUCTION

Ischemic stroke is among the leading cause of mortality and remains serious long-term disability worldwide. There were 2.5 million new stroke cases each year in China,<sup>1</sup> and the incidence of stroke is predicted to rise because of the rapidly aging population. Although subsequent mortality has declined in recent years,<sup>2</sup> a greater understanding of stroke biomarker of subsequent mortality will be required to establish appropriate prevention and treatment strategies.

Systematic inflammation plays a crucial role in the pathogenesis of ischemic stroke and may lead a secondary injury.<sup>3</sup> Many inflammatory biomarkers, such as procalcitonin (PCT) and C-reactive protein (CRP) are also identified as biomarkers of acute ischemic stroke.<sup>4,5</sup> PCT, a prohormone of calcitonin, is produced by C-cells of the thyroid gland. In previous studies, PCT seemed to be a better marker for

the diagnosis of infection than other biomarkers.<sup>6-8</sup> Recent studies showed it may be a prognosis marker of ischemic stroke and may be better than CRP.<sup>9,10</sup> The concentration of PCT was transient increased after severe trauma and seemed proportional to the severity of tissue injury and hypovolemia.<sup>11</sup> However, the relationship between PCT and short-term outcome has not been well examined. In this study, we sought to determine the relationship between PCT and 30-day all-cause mortality after acute ischemic stroke in Chinese patients.

## 2 | MATERIALS AND METHODS

We retrospective study all patients, less than 72 hours after symptom onset, with acute ischemic stroke event from January 2014 to

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. *Journal of Clinical Laboratory Analysis* published by Wiley Periodicals, Inc.

| Characteristic              | Death             | Alive             | P value |
|-----------------------------|-------------------|-------------------|---------|
| Median age ( $\pm$ SD)      | 75.19 $\pm$ 14.27 | 68.81 $\pm$ 13.73 | .572    |
| Male Sex                    | 7 (24.14%)        | 307 (42.70%)      | .047    |
| Comorbidities               |                   |                   |         |
| Hypertension                | 21 (72.41%)       | 492 (68.43%)      | .650    |
| Coronary heart disease      | 7 (24.14%)        | 111 (15.44%)      | .208    |
| Laboratory finding          |                   |                   |         |
| PCT ng/mL                   | 2.03 $\pm$ 2.57   | 0.14 $\pm$ 0.32   | <.001   |
| CRP mg/L                    | 31.75 $\pm$ 69.81 | 22.76 $\pm$ 25.51 | .022    |
| White blood cell            | 3.53 $\pm$ 4.69   | 3.64 $\pm$ 3.50   | .065    |
| Neutrophile granulocyte %   | 85.55 $\pm$ 13.70 | 66.21 $\pm$ 10.63 | <.001   |
| Total bilirubin $\mu$ mol/L | 19.38 $\pm$ 11.99 | 12.7 $\pm$ 5.65   | .019    |
| Triglycerides mmol/L        | 1.02 $\pm$ 0.37   | 1.53 $\pm$ 1.45   | .118    |
| Urea mmol/L                 | 8.35 $\pm$ 5.07   | 5.57 $\pm$ 2.13   | .012    |

**TABLE 1** Baseline characteristics of stroke patients

Abbreviations:CRP, C-reactive protein, PCT, procalcitonin.

December 2018 in a tertiary hospital. Acute ischemic stroke defined according to the World Health Organization criteria.<sup>12</sup> We excluded patients with intracerebral hemorrhage, subarachnoid hemorrhage, systemic infections, transient ischemic attack, inflammation, and incomplete data.

For PCT measurement, the blood sample was collected by venipuncture within 48 hours after hospital admission. Blood samples were centrifuged at 2264 g for 10 minutes. The serum was measured within 2 hours after sample collected. PCT was measured in serum sample (Cobas e601, Roche). All-cause of mortality defined as death occurring after hospital admission. All clinical data were collected from electronic medical records.

All data analyzed using SPSS 22.0. Two groups of any continuous variables were compared using Student's t test. Categorical variables were compared between groups using the chi-square test. Survival analysis by multivariable Cox regression analysis was used to examine the risk factors for 30-day mortality. The survival distribution function was estimated with the Kaplan-Meier method, and a non-parametric log-rank test was used to compare the survival curves among the different groups.  $P < .05$  was considered statistically significant.

### 3 | RESULTS

A total of 748 patients were enrolled in this study. The median age was 68 years, with age ranged between 46 and 101. Demographic data, risk factors, laboratory finding, and 30-day outcome were shown in Table 1. The inflammatory biomarkers including PCT, CRP, and neutrophils percentage in patients who died in 30 days were significantly higher than those who survived. Furthermore, the serum levels of PCT were also higher in patients with unfavorable functional outcome than those in patients with favorable functional outcome  $1.37 \pm 1.81$  ng/mL vs  $0.12 \pm 0.41$  ng/mL.

Death by the 30-day follow-up was associated with total bilirubin, urea, neutrophil percentage, and CRP (Table 2). Hypertension,

coronary heart disease, total bilirubin  $>17 \mu$ mol/L, and urea  $>7.0$  mmol/L were not associated with the mortality.

Multivariate Cox regression analysis was used to analyze the independent risk factors for 30-day mortality. As shown in Table 3, the inflammatory markers, PCT, CRP, and neutrophil percentage, were significant predictors of 30-day overall mortality, with HR of 7.963, 3.289, and 4.438, respectively. Moreover, Kaplan-Meier survival curves showed a trend for higher mortality of patients with PCT  $>1.5$  ng/mL (Figure 1) ( $P < .001$ ).

**TABLE 2** Univariate logistic regression analysis of risk factors for mortality

| Variable                        | HR    | 95% CI       | P value |
|---------------------------------|-------|--------------|---------|
| Sex                             | 2.308 | 0.986-5.403  | .054    |
| Hypertension                    | 1.204 | 0.533-2.717  | .656    |
| Coronary heart disease          | 1.741 | 0.744-4.076  | .201    |
| CRP $>5$ mg/L                   | 3.289 | 1.587-6.813  | .001    |
| PCT $>1.0$ ng/mL                | 2.445 | 1.168-5.119  | .018    |
| PCT $>1.5$ ng/mL                | 7.963 | 3.802-16.678 | <.001   |
| Neutrophile granulocyte $>70\%$ | 4.438 | 1.956-10.020 | <.001   |
| Total bilirubin $>17 \mu$ mol/L | 1.190 | 0.573-2.474  | .641    |
| Urea $>7.0$ mmol/L              | 2.091 | 0.926-4.720  | .076    |


Abbreviations:CRP, C-reactive protein, PCT, procalcitonin.

**TABLE 3** Independent risk factors for mortality in patients with ischemic stroke

| Variable                        | HR (95%CI)           | P value |
|---------------------------------|----------------------|---------|
| CRP $>5$ mg/L                   | 2.242 (1.043-4820)   | .039    |
| PCT $>1.0$ ng/mL                | 1.653 (0.750-3.643)  | .212    |
| PCT $>1.5$ ng/mL                | 5.486 (2.508-12.000) | <.001   |
| Neutrophile granulocyte $>70\%$ | 3.871 (1.707-8.779)  | .001    |

Abbreviations:CRP, C-reactive protein, PCT, procalcitonin.

**FIGURE 1** Kaplan-Meier survival curves according to procalcitonin (PCT) at admission ( $P < .001$ )



## 4 | DISCUSSION

Stroke is the first leading cause of disability and over two-thirds of stroke deaths worldwide occurring in developing countries.<sup>13</sup> Inflammation may play an important role in the progression of stroke, and inflammatory molecules might become both biomarkers and therapeutic targets for stroke management.<sup>14</sup> The infection markers and inflammatory molecules, such as CRP, WBC, and IL-6 were suggested to be biomarkers for outcome of ischemic stroke.<sup>15-17</sup> Compared with these markers, PCT was not a widely diagnostic marker, it is considered to be the best available biomarkers to diagnose infection.<sup>18</sup> In a recent study, PCT was suggested to be an independent risk factor for ischemic stroke.<sup>19</sup> However, the prognostic value of serum PCT in acute ischemic stroke was little to know.<sup>10</sup>

In his study, our results showed that the concentrations of PCT in patients who died in 30 days were significantly higher than those who survived. As shown by multivariate logistic regression PCT level was an independent prognostic marker of 30-day mortality. And Kaplan-Meier analysis indicated PCT levels might help to predict 30-day mortality. These results suggested that the concentration of PCT strongly predicted all-course mortality in patients with acute ischemic stroke in first 30 days. However, hypertension and coronary artery disease, which could increase the incidence of ischemic stroke,<sup>20,21</sup> were not associated with the 30-day mortality of ischemic stroke in our data. Moreover, in our data, PCT levels were also significantly higher in patients with unfavorable functional outcome compared with those in patients with a favorable outcome, which was consistent with previous report.<sup>22</sup>

Procalcitonin was a specific inflammatory biomarker, and systematic inflammation plays a crucial role in the pathogenesis of

ischemic stroke. Thus, the increased PCT level in serum may due to the inflammatory process in acute ischemic stroke. The increased level of PCT was also observed after severe trauma,<sup>11</sup> so we suspected that the increased level of PCT in acute ischemic stroke may also partly because the tissue injury after stroke. In this study, we found that PCT level was associated with the 30-day mortality in Chinese patients, this may indicate that PCT was an independent prognostic marker of short-term mortality after one set of ischemic strokes.

### ORCID

Pu Liao  <https://orcid.org/0000-0002-8884-7059>

### REFERENCES

- Zhang ZG, Wang C, Wang J, et al. Prognostic value of mannose-binding lectin: 90-day outcome in patients with acute ischemic stroke. *Mol Neurobiol.* 2015;51:230-239.
- Rothwell PM, Coull AJ, Giles MF, et al. Change in stroke incidence, mortality, case-fatality, severity, and risk factors in Oxfordshire, UK from 1981 to 2004 (Oxford Vascular Study). *Lancet.* 2004;363:1925-1933.
- Jin R, Yang G, Li G. Inflammatory mechanisms in ischemic stroke: role of inflammatory cells. *J Leukoc Biol.* 2010;87:779-789.
- Tian D, Zhang S, He X, Liu H. Serum procalcitonin as a diagnostic marker in acute ischemic stroke. *Neuroreport.* 2015;26:33-37.
- Luna JM, Moon YP, Liu KM, et al. High-sensitivity C-reactive protein and interleukin-6-dominant inflammation and ischemic stroke risk: the northern Manhattan study. *Stroke.* 2014;45:979-987.
- Massaro KS, Costa SF, Leone C, et al. Procalcitonin (PCT) and C-reactive protein (CRP) as severe systemic infection markers in febrile neutropenic adults. *BMC Infect Dis.* 2007;7:137.
- Jae-Sik J, Sung-Mi J. Diagnostic value of procalcitonin and CRP in critically ill patients admitted with suspected sepsis. *J Dent Anesth Pain Med.* 2015;15:135-140.

8. Harbarth S, Holeckova K, Froidevaux C, et al. Sepsis network. diagnostic value of procalcitonin, interleukin-6, and interleukin-8 in critically ill patients admitted with suspected sepsis. *Am J Respir Crit Care Med.* 2001;164:396-402.
9. Deng WJ, Shen RL, Li M, et al. Relationship between procalcitonin serum levels and functional outcome in stroke patients. *Cell Mol Neurobiol.* 2015;35:355-361.
10. Wang C, Gao L, Zhang ZG, et al. Procalcitonin is a stronger predictor of long-term functional outcome and mortality than high-sensitivity C-reactive protein in patients with ischemic stroke. *Mol Neurobiol.* 2016;53:1509-1517.
11. Mimos O, Benoist JF, Edouard AR, et al. Procalcitonin and C-reactive protein during the early posttraumatic systemic inflammatory response syndrome. *Intensive Care Med.* 1998;24(2):185-188.
12. Hatano S. Experience from a multicentre stroke register: a preliminary report. *Bull World Health Organ.* 1976;54:541-553.
13. Feigin VL. Stroke epidemiology in the developing world. *Lancet.* 2005;365:2160-2161.
14. Laura R, Alba S, Teresa G-B, et al. Inflammatory molecules might become both biomarkers and therapeutic targets for stroke management. *Ther Adv Neurol Dis.* 2018;11:175628641878934.
15. Claudia-Alexandrina I, Mihai V, Marius I, et al. C-reactive protein and T3: new prognostic factors in acute ischemic stroke. *J Stroke Cerebrovasc Dis.* 2018;27(10):2731-2737.
16. Zheng X, Zeng N, Wang A, et al. Prognostic value of white blood cell in acute ischemic stroke patients. *Curr Neurovasc Res.* 2018;15(2):151-157.
17. Bustamante A, Sobrino T, Giralt D, et al. Prognostic value of blood interleukin-6 in the prediction of functional outcome after stroke: a systematic review and meta-analysis. *J Neuroimmunol.* 2014;274:215-224.
18. Aloisio E, Dolci A, Panteghini M. Procalcitonin: between evidence and critical issues. *Clin Chim Acta.* 2019;496:7-12.
19. Katan M, Moon YP, Paik MC, et al. Procalcitonin and midregional proatrial natriuretic peptide as markers of ischemic stroke: the northern Manhattan study. *Stroke.* 2016;47:1714-1719.
20. Huangfu X, Zhu Z, Zhong C, et al. Smoking, hypertension, and their combined effect on ischemic stroke incidence: a prospective study among inner Mongolians in China. *J Stroke Cerebrovasc Dis.* 2017;26:2749-2754.
21. Satue E, Vila-Corcoles A, Ochoa-Gondar O, et al. Incidence and risk conditions of ischemic stroke in older adults. *Acta Neurol Scand.* 2016;134:250-257.
22. Deng WJ, Shen RL, Li M, Teng JF. Relationship between procalcitonin serum levels and functional outcome in stroke patients. *Cell Mol Neurobiol.* 2015;35(3):355-361.

**How to cite this article:** Yan L, Wang S, Xu L, Zhang Z, Liao P. Procalcitonin as a prognostic marker of patients with acute ischemic stroke. *J Clin Lab Anal.* 2020;34:e23301. <https://doi.org/10.1002/jcla.23301>