

Dichlorido[*N*-[2-(diphenylphosphanyl)-benzylidene]-2-(thiophen-2-yl)ethan-amine- κ^2 P,*N*]platinum(II) dichloromethane hemisolvate

Haleden Chiririwa* and Alfred Muller*

Research Center for Synthesis and Catalysis, Department of Chemistry, University of Johannesburg (APK Campus), PO Box 524, Auckland Park, Johannesburg 2006, South Africa

Correspondence e-mail: harrychiririwa@yahoo.com, mullera@uj.ac.za


Received 14 October 2011; accepted 29 December 2011

Key indicators: single-crystal X-ray study; $T = 173$ K; mean $\sigma(\text{C}-\text{C}) = 0.006$ Å; disorder in main residue; R factor = 0.028; wR factor = 0.058; data-to-parameter ratio = 19.8.

The crystal structure of the title compound, $[\text{PtCl}_2(\text{C}_{25}\text{H}_{22}\text{NPS})]\cdot 0.5\text{CH}_2\text{Cl}_2$, was determined to establish the coordination properties of the (phosphanyl)benzylidene-methanamine ligand to platinum. In the unit cell two molecules of *cis*- $[\text{PtCl}_2(\text{C}_{25}\text{H}_{22}\text{NPS})]$ are accompanied by a dichloromethane solvent molecule. The square-planar Pt^{2+} coordination sphere is slightly distorted with the bidentate ligand coordinated *via* the P and the amine N atoms, and the Cl atoms located *cis* at the two remaining coordination sites. Parts of the thiophene ring and the solvate molecule were modeled as disordered with occupancy ratios of 0.55 (2): 0.45 (2) and 0.302 (10):0.198 (10), respectively. Weak $\text{C}-\text{H}\cdots\text{Cl}$ interactions stabilize the crystal packing.

Related literature

For background to related structures, see: Chiririwa *et al.* (2011); Chiririwa & Meijboom (2011*a,b,c*); Ghilardi *et al.* (1992); Sanchez *et al.* (1998, 2001); Coleman *et al.* (2001). For Pt–N and Pt–P bond lengths in similar platinum(II) complexes, see: Ankersmit *et al.* (1996). For background to weak hydrogen-bonding interactions, see Steiner (1996).



Experimental

Crystal data

$[\text{PtCl}_2(\text{C}_{25}\text{H}_{22}\text{NPS})]\cdot 0.5\text{CH}_2\text{Cl}_2$
 $M_r = 707.92$
 Monoclinic, $P2_1/c$
 $a = 9.9635$ (7) Å
 $b = 19.0185$ (14) Å
 $c = 16.0155$ (9) Å
 $\beta = 122.751$ (3)°
 $V = 2552.3$ (3) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 5.97$ mm⁻¹
 $T = 173$ K
 $0.08 \times 0.05 \times 0.03$ mm

Data collection

Bruker APEXII 4K-CCD diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2007)
 $T_{\min} = 0.647$, $T_{\max} = 0.841$
 30158 measured reflections
 6593 independent reflections
 5165 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.052$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.028$
 $wR(F^2) = 0.058$
 $S = 1.01$
 6593 reflections
 333 parameters
 146 restraints
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.84$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.62$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C7}-\text{H7}\cdots\text{Cl2}^i$	0.95	2.71	3.486 (4)	139
$\text{C12}-\text{H12}\cdots\text{Cl1}^{ii}$	0.95	2.71	3.639 (4)	168

Symmetry codes: (i) $x, -y + \frac{3}{2}, z - \frac{1}{2}$; (ii) $x + 1, y, z$.

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT and XPREP (Bruker, 2007); program(s) used to solve structure: SIR97 (Altomare *et al.*, 1999); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: DIAMOND (Brandenburg & Putz, 2005); software used to prepare material for publication: WinGX (Farrugia, 1999).

Financial assistance from the South African National Research Foundation (SA NRF), the Research Fund of the University of Johannesburg and SASOL is gratefully acknowledged.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NR2014).

References

- Altomare, A., Burla, M. C., Camalli, M., Casciarano, G. L., Giacovazzo, C., Guagliardi, A., Moliterni, A. G. G., Polidori, G. & Spagna, R. (1999). *J. Appl. Cryst.* **32**, 115–119.
 Ankersmit, H. A., Loken, B. H., Kooijman, H., Spek, A. L., Vrieze, K. & van Koten, G. (1996). *Inorg. Chim. Acta*, **252**, 141–155.
 Brandenburg, K. & Putz, H. (2005). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
 Bruker (2007). *APEX2, SADABS, SAINT and XPREP*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Chiririwa, H. & Meijboom, R. (2011*a*). *Acta Cryst.* **E67**, m1496.
 Chiririwa, H. & Meijboom, R. (2011*b*). *Acta Cryst.* **E67**, m1497.
 Chiririwa, H. & Meijboom, R. (2011*c*). *Acta Cryst.* **E67**, m1498.
 Chiririwa, H., Meijboom, R. & Omondi, B. (2011). *Acta Cryst.* **E67**, m608–m609.

- Coleman, K. S., Green, M. L. H., Pascu, S. I., Rees, N. H., Cowley, A. R. & Rees, L. H. (2001). *J. Chem. Soc. Dalton Trans.* pp. 3384–3395.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Ghilardi, C. A., Midollini, S., Moneti, S., Orlandini, A. & Scapacci, G. (1992). *J. Chem. Soc. Dalton Trans.* pp. 3371–3376.
- Sanchez, G., Momblona, F., Perez, J. & Lopez, G. (2001). *Transition Met. Chem.* **26**, 100–104.
- Sanchez, G., Serrano, J. L., Ruiz, F. & Lopez, G. (1998). *J. Fluorine Chem.* **91**, 165–169.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Steiner, T. (1996). *Crystallogr. Rev.* **6**, 1–57.

supplementary materials

Acta Cryst. (2012). E68, m116-m117 [doi:10.1107/S1600536811056108]

Dichlorido{*N*-[2-(diphenylphosphanyl)benzylidene]-2-(thiophen-2-yl)ethanamine- κ^2P,N }platinum(II) dichloromethane hemisolvate

H. Chiririwa and A. Muller

Comment

Platinum complexes with (phosphanyl)benzylidene-methanamine ligands have been used as catalysts or catalyst precursors for a variety of organic reactions. Our group and others have recently been interested in these types of complexes and have reported several of these types of complexes (Chiririwa *et al.*, 2011; Chiririwa & Meijboom, 2011*a*, 2011*b*, 2011*c*; Ghilardi *et al.*, 1992; Sanchez *et al.*, (1998, 2001) and Coleman *et al.*, 2001).

The title compound (see Fig. 1) crystallize with a dichloromethane solvate, disordered over an inversion center, for each pair of *cis*-[PtCl₂(C₂₅H₂₁NSP)] molecules. The square planar Pt coordination sphere is slightly distorted (Pt1 displaced -0.0275 (7) Å from the plane formed by Pt1, P1, N1, C11 and Cl2 respectively; r.m.s. deviation of fitted atoms = 0.0423 Å). The distortion is featured most prominently in the N1—Pt1—P1 angle of 86.17 (10)° *versus*. C11—Pt1—Cl2 = 89.08 (4)°, indicating some ring strain induced by the chelation of the bidentate ligand. The average Pt—N and Pt—P bond lengths of 2.030 (3) and 2.2089 (9) Å, respectively are in the range expected for similar platinum(II) complexes (Ankersmit *et al.*(1996)). The initial refinement model of the compound showed some large displacement parameters for the thiophene moiety as well as the dichloromethane solvent. These disorders were elucidated to give an improved model (details can be found under the experimental refinement section). Two weak C—H···Cl interactions (Steiner, 1996) aid in the stabilization of the crystal structure (see Table 1, Fig. 2).

Experimental

To a dry CH₂Cl₂ (10 ml) solution of the precursor [Pt(COD)Cl₂] was added an equimolar amount of (2-(diphenylphosphanyl)benzylidene)(thiophen-2-yl)methanamine in CH₂Cl₂ (10 ml), and stirred at room temperature for 2 hrs. The solvent was reduced and the complex precipitated out on addition of hexane, filtered off, washed with Et₂O (2×5 ml) and dried under vacuum for 4 hrs affording a yellow precipitate in 72% yield. Crystals suitable for X-ray structure determination were obtained by recrystallization from a CH₂Cl₂-hexane mixture at room temperature.

Refinement

All hydrogen atoms were positioned in geometrically idealized positions with C—H = 0.99 Å and 0.95 Å for methylene and aromatic H atoms respectively. All hydrogen atoms were allowed to ride on their parent atoms with $U_{iso}(H) = 1.2U_{eq}$. A disorder refinement model was applied to the thiophene that showed large displacements at C2, C3 and S1. Geometrical (FLAT) restraints were applied to keep the rings C1, C2A/B, C3A/B, C4, S1A/B planar. Ellipsoid displacement (SIMU and DELU) restraints were also applied to the disordered moiety. The occupation parameters of the two disordered thiophene fragments were linked to a free variable so that the two sites add to unity. This showed a distribution of 0.54903:0.45097. The dichloromethane solvate was also refined as disordered on two positions in the asymmetric unit. This resulted in four disordered positions for each dichloromethane at each solvent accessible site in the crystal lattice. The occupancies of these

supplementary materials

sites were linked to a free variable to add to unity and refined to a ratio of 0.60331:0.39669 (based on the asymmetric unit fraction). To keep refinement stable geometrical (DIFX and DANG) restraints were applied to the C—Cl bonds and Cl...Cl distances. All the above restraints were applied with the default standard deviations. The atoms of the solvate molecule was left isotropic due to the extensive nature of the disorder. The highest residual electron density of $0.84 \text{ e.}\text{\AA}^{-3}$ is 0.89 \AA from Pt1 representing no physical meaning.

Figures




Fig. 1. View of title compound showing displacement ellipsoids (drawn at a 30% probability level) and labeling. For clarity: a) hydrogen atoms omitted, b) bonds in part B of the disordered thiophene indicated with dotted lines, and c) minor component as well as symmetry generated disordered parts of dichloromethane created by inversion center omitted.




Fig. 2. Partial packing diagram of title compound to illustrate the weak hydrogen bonding stabilizing crystal packing. All disordered components omitted for clarity.

Dichlorido{*N*-[2-(diphenylphosphanyl)benzylidene]-2-(thiophen-2-yl)ethanamine- κ^2P,N }platinum(II) dichloromethane hemisolvate

Crystal data

[PtCl₂(C₂₅H₂₂NPS)]·0.5CH₂Cl₂

$M_r = 707.92$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 9.9635 (7) \text{ \AA}$

$b = 19.0185 (14) \text{ \AA}$

$c = 16.0155 (9) \text{ \AA}$

$\beta = 122.751 (3)^\circ$

$V = 2552.3 (3) \text{ \AA}^3$

$Z = 4$

$F(000) = 1372$

$D_x = 1.842 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 5962 reflections

$\theta = 2.6\text{--}28.2^\circ$

$\mu = 5.97 \text{ mm}^{-1}$

$T = 173 \text{ K}$

Block, yellow

$0.08 \times 0.05 \times 0.03 \text{ mm}$

Data collection

Bruker APEXII 4K-CCD
diffractometer

graphite

Detector resolution: $8.4 \text{ pixels mm}^{-1}$

6593 independent reflections

5165 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.052$

φ and ω scans $\theta_{\max} = 28.7^\circ$, $\theta_{\min} = 2.1^\circ$
 Absorption correction: multi-scan $h = -13 \rightarrow 13$
 (*SADABS*; Bruker, 2007)
 $T_{\min} = 0.647$, $T_{\max} = 0.841$ $k = -25 \rightarrow 25$
 30158 measured reflections $l = -21 \rightarrow 21$

Refinement

Refinement on F^2 Primary atom site location: structure-invariant direct methods
 Least-squares matrix: full Secondary atom site location: difference Fourier map
 $R[F^2 > 2\sigma(F^2)] = 0.028$ Hydrogen site location: inferred from neighbouring sites
 $wR(F^2) = 0.058$ H-atom parameters constrained
 $S = 1.01$ $w = 1/[\sigma^2(F_o^2) + (0.0216P)^2 + 0.6014P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 6593 reflections $(\Delta/\sigma)_{\max} = 0.002$
 333 parameters $\Delta\rho_{\max} = 0.84 \text{ e } \text{\AA}^{-3}$
 146 restraints $\Delta\rho_{\min} = -0.62 \text{ e } \text{\AA}^{-3}$

Special details

Experimental. The intensity data was collected on a Bruker Apex-II 4 K CCD diffractometer using an exposure time of 80 s/frame. A total of 1315 frames were collected with a frame width of 0.5° covering up to $\theta = 28.72^\circ$ with 99.8% completeness accomplished.

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Pt1	0.557569 (15)	0.701063 (7)	0.420321 (11)	0.02063 (4)	
Cl1	0.38212 (12)	0.79833 (5)	0.36520 (9)	0.0382 (2)	
Cl2	0.57723 (11)	0.70634 (5)	0.56946 (7)	0.0318 (2)	
P1	0.73311 (10)	0.61536 (5)	0.46437 (7)	0.01889 (18)	
N1	0.5342 (4)	0.69787 (16)	0.2864 (2)	0.0278 (7)	
C1	0.1129 (5)	0.6300 (2)	0.1049 (3)	0.0385 (10)	
C2A	0.0513 (10)	0.5796 (8)	0.0305 (9)	0.056 (4)	0.55 (2)
H2A	0.1137	0.5437	0.0264	0.067*	0.55 (2)
C3A	-0.1149 (10)	0.5875 (9)	-0.0387 (9)	0.056 (4)	0.55 (2)
H3A	-0.1758	0.5578	-0.0947	0.067*	0.55 (2)
S1A	-0.0359 (10)	0.6794 (6)	0.0932 (7)	0.060 (2)	0.55 (2)

supplementary materials

C2B	0.0654 (13)	0.6113 (12)	0.0100 (8)	0.069 (5)	0.45 (2)
H2B	0.136	0.5934	-0.0076	0.083*	0.45 (2)
C3B	-0.0994 (13)	0.6213 (13)	-0.0587 (8)	0.060 (5)	0.45 (2)
H3B	-0.1501	0.6136	-0.1281	0.072*	0.45 (2)
S1B	-0.0452 (11)	0.6648 (7)	0.1052 (7)	0.0434 (19)	0.45 (2)
C4	-0.1785 (5)	0.6432 (3)	-0.0158 (3)	0.0511 (12)	
H4A	-0.286	0.6588	-0.0553	0.061*	0.45 (2)
H4B	-0.2912	0.6461	-0.0484	0.061*	0.55 (2)
C5	0.2814 (5)	0.6328 (2)	0.1939 (3)	0.0382 (10)	
H5A	0.3404	0.5912	0.1932	0.046*	
H5B	0.2795	0.6306	0.255	0.046*	
C6	0.3689 (5)	0.6990 (2)	0.1964 (3)	0.0369 (10)	
H6A	0.3728	0.7015	0.136	0.044*	
H6B	0.3115	0.741	0.1979	0.044*	
C7	0.6488 (5)	0.6932 (2)	0.2717 (3)	0.0318 (9)	
H7	0.6203	0.6955	0.2047	0.038*	
C8	0.8184 (5)	0.68468 (19)	0.3477 (3)	0.0272 (8)	
C9	0.8769 (4)	0.64993 (18)	0.4383 (3)	0.0229 (8)	
C10	1.0409 (4)	0.64245 (19)	0.5044 (3)	0.0267 (8)	
H10	1.0809	0.6181	0.5652	0.032*	
C11	1.1464 (5)	0.6705 (2)	0.4817 (3)	0.0345 (10)	
H11	1.2581	0.6662	0.528	0.041*	
C12	1.0910 (5)	0.7042 (2)	0.3935 (4)	0.0422 (11)	
H12	1.1639	0.723	0.3784	0.051*	
C13	0.9267 (5)	0.7108 (2)	0.3254 (3)	0.0365 (10)	
H13	0.8882	0.7332	0.2635	0.044*	
C14	0.8472 (4)	0.58128 (18)	0.5904 (3)	0.0209 (7)	
C15	0.9314 (4)	0.6271 (2)	0.6700 (3)	0.0273 (8)	
H15	0.9295	0.6762	0.6586	0.033*	
C16	1.0187 (5)	0.6010 (2)	0.7665 (3)	0.0341 (9)	
H16	1.0763	0.6323	0.8208	0.041*	
C17	1.0217 (5)	0.5291 (2)	0.7835 (3)	0.0331 (9)	
H17	1.0799	0.5114	0.8495	0.04*	
C18	0.9395 (5)	0.4836 (2)	0.7041 (3)	0.0302 (9)	
H18	0.943	0.4344	0.7156	0.036*	
C19	0.8526 (4)	0.50892 (19)	0.6080 (3)	0.0246 (8)	
H19	0.7964	0.4773	0.5538	0.029*	
C20	0.6463 (4)	0.53857 (17)	0.3865 (3)	0.0201 (7)	
C21	0.7114 (4)	0.50677 (19)	0.3381 (3)	0.0255 (8)	
H21	0.8031	0.5265	0.3432	0.031*	
C22	0.6429 (5)	0.4463 (2)	0.2825 (3)	0.0306 (9)	
H22	0.6885	0.4246	0.25	0.037*	
C23	0.5093 (5)	0.4177 (2)	0.2741 (3)	0.0318 (9)	
H23	0.463	0.3762	0.2361	0.038*	
C24	0.4419 (4)	0.44923 (19)	0.3211 (3)	0.0288 (9)	
H24	0.3494	0.4295	0.315	0.035*	
C25	0.5098 (4)	0.50953 (18)	0.3769 (3)	0.0234 (8)	
H25	0.4633	0.5313	0.4088	0.028*	
C26A	0.485 (3)	0.5266 (10)	-0.0366 (13)	0.070 (6)*	0.302 (10)

H26A	0.5187	0.5656	-0.0625	0.084*	0.302 (10)
H26B	0.3693	0.5203	-0.0843	0.084*	0.302 (10)
Cl3A	0.5086 (19)	0.5549 (5)	0.0677 (8)	0.099 (3)*	0.302 (10)
Cl4A	0.5746 (14)	0.4531 (5)	-0.0434 (8)	0.121 (4)*	0.302 (10)
C26B	0.564 (3)	0.5398 (9)	-0.0142 (17)	0.047 (7)*	0.198 (10)
H26C	0.6572	0.5502	-0.0185	0.057*	0.198 (10)
H26D	0.4676	0.5612	-0.072	0.057*	0.198 (10)
Cl3B	0.591 (3)	0.5697 (8)	0.0906 (11)	0.122 (5)*	0.198 (10)
Cl4B	0.5390 (15)	0.4497 (6)	-0.0105 (12)	0.090 (4)*	0.198 (10)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Pt1	0.01969 (7)	0.01735 (7)	0.02733 (8)	0.00077 (6)	0.01435 (6)	0.00296 (7)
Cl1	0.0329 (5)	0.0267 (5)	0.0624 (7)	0.0114 (4)	0.0307 (5)	0.0171 (5)
Cl2	0.0343 (5)	0.0353 (5)	0.0322 (5)	0.0034 (4)	0.0221 (4)	-0.0041 (4)
P1	0.0200 (4)	0.0179 (4)	0.0209 (5)	-0.0001 (3)	0.0125 (4)	0.0013 (4)
N1	0.0254 (16)	0.0288 (17)	0.0290 (18)	0.0021 (14)	0.0145 (14)	0.0078 (15)
C1	0.030 (2)	0.057 (3)	0.022 (2)	0.0020 (19)	0.0097 (17)	0.0003 (19)
C2A	0.026 (4)	0.097 (9)	0.036 (6)	0.011 (5)	0.011 (4)	-0.024 (6)
C3A	0.027 (4)	0.096 (9)	0.029 (6)	-0.004 (5)	0.005 (4)	-0.022 (6)
S1A	0.048 (2)	0.036 (3)	0.068 (4)	0.0109 (17)	0.012 (2)	-0.009 (2)
C2B	0.033 (5)	0.141 (14)	0.029 (6)	0.006 (8)	0.014 (4)	-0.018 (8)
C3B	0.038 (6)	0.113 (13)	0.020 (5)	-0.002 (7)	0.009 (4)	-0.006 (7)
S1B	0.042 (2)	0.051 (5)	0.033 (2)	0.013 (2)	0.0175 (18)	-0.005 (2)
C4	0.033 (2)	0.059 (3)	0.040 (3)	0.009 (2)	0.006 (2)	-0.002 (2)
C5	0.030 (2)	0.041 (2)	0.028 (2)	-0.0028 (18)	0.0059 (18)	0.0048 (19)
C6	0.030 (2)	0.046 (3)	0.026 (2)	0.0025 (19)	0.0096 (18)	0.013 (2)
C7	0.039 (2)	0.030 (2)	0.032 (2)	-0.0017 (18)	0.0224 (19)	0.0090 (18)
C8	0.032 (2)	0.024 (2)	0.036 (2)	-0.0011 (15)	0.0254 (19)	0.0005 (16)
C9	0.0273 (19)	0.0202 (18)	0.028 (2)	-0.0018 (15)	0.0195 (17)	-0.0026 (16)
C10	0.0253 (19)	0.027 (2)	0.030 (2)	-0.0024 (15)	0.0171 (18)	-0.0036 (17)
C11	0.026 (2)	0.031 (2)	0.051 (3)	-0.0060 (17)	0.024 (2)	-0.010 (2)
C12	0.041 (2)	0.035 (2)	0.071 (3)	-0.004 (2)	0.044 (3)	0.003 (2)
C13	0.045 (3)	0.031 (2)	0.051 (3)	0.0018 (18)	0.038 (2)	0.012 (2)
C14	0.0195 (17)	0.0241 (18)	0.0217 (19)	0.0019 (14)	0.0129 (15)	0.0023 (15)
C15	0.0240 (19)	0.027 (2)	0.026 (2)	-0.0034 (15)	0.0108 (17)	-0.0019 (16)
C16	0.029 (2)	0.041 (2)	0.022 (2)	-0.0067 (18)	0.0081 (17)	-0.0068 (19)
C17	0.029 (2)	0.046 (3)	0.022 (2)	0.0037 (18)	0.0123 (18)	0.0061 (19)
C18	0.033 (2)	0.027 (2)	0.029 (2)	0.0060 (17)	0.0162 (18)	0.0084 (17)
C19	0.029 (2)	0.0231 (19)	0.023 (2)	0.0015 (15)	0.0149 (17)	0.0030 (15)
C20	0.0224 (17)	0.0174 (17)	0.0195 (18)	-0.0007 (14)	0.0107 (15)	0.0001 (14)
C21	0.0245 (19)	0.029 (2)	0.025 (2)	0.0007 (16)	0.0151 (17)	-0.0014 (16)
C22	0.034 (2)	0.033 (2)	0.024 (2)	0.0030 (17)	0.0154 (18)	-0.0066 (17)
C23	0.031 (2)	0.0207 (19)	0.031 (2)	-0.0014 (16)	0.0092 (18)	-0.0039 (17)
C24	0.026 (2)	0.025 (2)	0.031 (2)	-0.0041 (16)	0.0123 (17)	0.0000 (17)
C25	0.0268 (19)	0.0220 (18)	0.027 (2)	0.0008 (15)	0.0179 (16)	0.0008 (16)

supplementary materials

Geometric parameters (Å, °)

Pt1—N1	2.029 (3)	C10—H10	0.95
Pt1—P1	2.2087 (9)	C11—C12	1.367 (6)
Pt1—Cl2	2.2907 (10)	C11—H11	0.95
Pt1—C11	2.3638 (9)	C12—C13	1.397 (6)
P1—C20	1.808 (4)	C12—H12	0.95
P1—C14	1.818 (4)	C13—H13	0.95
P1—C9	1.818 (4)	C14—C15	1.390 (5)
N1—C7	1.288 (5)	C14—C19	1.400 (5)
N1—C6	1.490 (5)	C15—C16	1.391 (5)
C1—C2B	1.373 (11)	C15—H15	0.95
C1—C2A	1.388 (10)	C16—C17	1.391 (6)
C1—C5	1.504 (5)	C16—H16	0.95
C1—S1A	1.678 (9)	C17—C18	1.383 (5)
C1—S1B	1.711 (9)	C17—H17	0.95
C2A—C3A	1.416 (9)	C18—C19	1.381 (5)
C2A—H2A	0.95	C18—H18	0.95
C3A—C4	1.382 (10)	C19—H19	0.95
C3A—H3A	0.95	C20—C21	1.389 (5)
S1A—C4	1.690 (9)	C20—C25	1.397 (5)
C2B—C3B	1.410 (11)	C21—C22	1.387 (5)
C2B—H2B	0.95	C21—H21	0.95
C3B—C4	1.362 (11)	C22—C23	1.376 (5)
C3B—H3B	0.95	C22—H22	0.95
S1B—C4	1.704 (10)	C23—C24	1.386 (5)
C4—H4A	0.95	C23—H23	0.95
C4—H4B	0.95	C24—C25	1.385 (5)
C5—C6	1.519 (5)	C24—H24	0.95
C5—H5A	0.99	C25—H25	0.95
C5—H5B	0.99	C26A—C13A	1.647 (16)
C6—H6A	0.99	C26A—C14A	1.692 (15)
C6—H6B	0.99	C26A—H26A	0.99
C7—C8	1.462 (6)	C26A—H26B	0.99
C7—H7	0.95	C26B—C13B	1.648 (16)
C8—C13	1.397 (5)	C26B—C14B	1.737 (16)
C8—C9	1.403 (5)	C26B—H26C	0.99
C9—C10	1.393 (5)	C26B—H26D	0.99
C10—C11	1.391 (5)		
N1—Pt1—P1	86.20 (9)	C13—C8—C7	116.9 (4)
N1—Pt1—Cl2	178.38 (9)	C9—C8—C7	124.0 (3)
P1—Pt1—Cl2	95.34 (3)	C10—C9—C8	119.6 (3)
N1—Pt1—C11	89.42 (9)	C10—C9—P1	122.3 (3)
P1—Pt1—C11	174.06 (3)	C8—C9—P1	118.1 (3)
Cl2—Pt1—C11	89.08 (4)	C11—C10—C9	120.3 (4)
C20—P1—C14	104.86 (16)	C11—C10—H10	119.9
C20—P1—C9	106.01 (16)	C9—C10—H10	119.9
C14—P1—C9	106.69 (16)	C12—C11—C10	120.7 (4)

C20—P1—Pt1	112.03 (12)	C12—C11—H11	119.7
C14—P1—Pt1	121.94 (11)	C10—C11—H11	119.7
C9—P1—Pt1	104.23 (12)	C11—C12—C13	119.7 (4)
C7—N1—C6	116.7 (3)	C11—C12—H12	120.1
C7—N1—Pt1	126.1 (3)	C13—C12—H12	120.1
C6—N1—Pt1	117.2 (2)	C8—C13—C12	120.6 (4)
C2B—C1—C5	126.7 (6)	C8—C13—H13	119.7
C2A—C1—C5	125.9 (5)	C12—C13—H13	119.7
C2B—C1—S1A	105.5 (6)	C15—C14—C19	119.6 (3)
C2A—C1—S1A	109.7 (5)	C15—C14—P1	120.0 (3)
C5—C1—S1A	123.5 (4)	C19—C14—P1	120.4 (3)
C2B—C1—S1B	109.7 (6)	C14—C15—C16	119.9 (4)
C2A—C1—S1B	106.8 (5)	C14—C15—H15	120
C5—C1—S1B	122.8 (4)	C16—C15—H15	120
C1—C2A—C3A	112.6 (7)	C17—C16—C15	120.2 (4)
C1—C2A—H2A	123.7	C17—C16—H16	119.9
C3A—C2A—H2A	123.7	C15—C16—H16	119.9
C4—C3A—C2A	112.6 (7)	C18—C17—C16	119.7 (4)
C4—C3A—H3A	123.7	C18—C17—H17	120.1
C2A—C3A—H3A	123.7	C16—C17—H17	120.1
C1—S1A—C4	94.9 (5)	C19—C18—C17	120.5 (4)
C1—C2B—C3B	112.8 (8)	C19—C18—H18	119.7
C1—C2B—H2B	123.6	C17—C18—H18	119.7
C3B—C2B—H2B	123.6	C18—C19—C14	120.0 (4)
C4—C3B—C2B	113.3 (8)	C18—C19—H19	120
C4—C3B—H3B	123.4	C14—C19—H19	120
C2B—C3B—H3B	123.4	C21—C20—C25	119.1 (3)
C4—S1B—C1	93.2 (4)	C21—C20—P1	122.9 (3)
C3B—C4—S1A	104.7 (6)	C25—C20—P1	118.0 (3)
C3A—C4—S1A	109.6 (5)	C22—C21—C20	120.2 (3)
C3B—C4—S1B	109.9 (6)	C22—C21—H21	119.9
C3A—C4—S1B	107.4 (5)	C20—C21—H21	119.9
C3B—C4—H4A	120.7	C23—C22—C21	120.3 (4)
C3A—C4—H4A	125.2	C23—C22—H22	119.9
S1A—C4—H4A	125.2	C21—C22—H22	119.9
S1B—C4—H4A	126	C22—C23—C24	120.2 (4)
C3B—C4—H4B	125.1	C22—C23—H23	119.9
C3A—C4—H4B	118.3	C24—C23—H23	119.9
S1A—C4—H4B	128.9	C25—C24—C23	119.8 (4)
S1B—C4—H4B	125	C25—C24—H24	120.1
C1—C5—C6	112.6 (3)	C23—C24—H24	120.1
C1—C5—H5A	109.1	C24—C25—C20	120.4 (3)
C6—C5—H5A	109.1	C24—C25—H25	119.8
C1—C5—H5B	109.1	C20—C25—H25	119.8
C6—C5—H5B	109.1	Cl3A—C26A—Cl4A	123.0 (11)
H5A—C5—H5B	107.8	Cl3A—C26A—H26A	106.6
N1—C6—C5	109.4 (3)	Cl4A—C26A—H26A	106.6
N1—C6—H6A	109.8	Cl3A—C26A—H26B	106.6
C5—C6—H6A	109.8	Cl4A—C26A—H26B	106.6

supplementary materials

N1—C6—H6B	109.8	H26A—C26A—H26B	106.6
C5—C6—H6B	109.8	C13B—C26B—C14B	104.7 (12)
H6A—C6—H6B	108.2	C13B—C26B—H26C	110.8
N1—C7—C8	126.5 (4)	C14B—C26B—H26C	110.8
N1—C7—H7	116.7	C13B—C26B—H26D	110.8
C8—C7—H7	116.7	C14B—C26B—H26D	110.8
C13—C8—C9	119.1 (4)	H26C—C26B—H26D	108.9
N1—Pt1—P1—C20	-59.67 (15)	Pt1—N1—C7—C8	4.2 (6)
C12—Pt1—P1—C20	119.82 (12)	N1—C7—C8—C13	-152.5 (4)
N1—Pt1—P1—C14	174.99 (16)	N1—C7—C8—C9	30.0 (6)
C12—Pt1—P1—C14	-5.52 (14)	C13—C8—C9—C10	0.3 (5)
N1—Pt1—P1—C9	54.51 (15)	C7—C8—C9—C10	177.7 (4)
C12—Pt1—P1—C9	-126.00 (13)	C13—C8—C9—P1	-179.1 (3)
P1—Pt1—N1—C7	-43.9 (3)	C7—C8—C9—P1	-1.7 (5)
C11—Pt1—N1—C7	132.1 (3)	C20—P1—C9—C10	-105.8 (3)
P1—Pt1—N1—C6	134.3 (3)	C14—P1—C9—C10	5.6 (3)
C11—Pt1—N1—C6	-49.8 (3)	Pt1—P1—C9—C10	135.8 (3)
C2B—C1—C2A—C3A	-82.1 (14)	C20—P1—C9—C8	73.6 (3)
C5—C1—C2A—C3A	174.8 (7)	C14—P1—C9—C8	-175.0 (3)
S1A—C1—C2A—C3A	5.3 (10)	Pt1—P1—C9—C8	-44.7 (3)
S1B—C1—C2A—C3A	18.3 (10)	C8—C9—C10—C11	1.3 (5)
C1—C2A—C3A—C4	-0.7 (12)	P1—C9—C10—C11	-179.3 (3)
C2B—C1—S1A—C4	25.5 (9)	C9—C10—C11—C12	-1.6 (6)
C2A—C1—S1A—C4	-6.6 (7)	C10—C11—C12—C13	0.2 (6)
C5—C1—S1A—C4	-176.4 (4)	C9—C8—C13—C12	-1.8 (6)
S1B—C1—S1A—C4	-86 (3)	C7—C8—C13—C12	-179.3 (4)
C2A—C1—C2B—C3B	86.2 (15)	C11—C12—C13—C8	1.5 (6)
C5—C1—C2B—C3B	-173.5 (9)	C20—P1—C14—C15	179.5 (3)
S1A—C1—C2B—C3B	-16.4 (13)	C9—P1—C14—C15	67.3 (3)
S1B—C1—C2B—C3B	-3.8 (12)	Pt1—P1—C14—C15	-52.0 (3)
C1—C2B—C3B—C4	-3.9 (16)	C20—P1—C14—C19	0.4 (3)
C2B—C1—S1B—C4	8.0 (9)	C9—P1—C14—C19	-111.8 (3)
C2A—C1—S1B—C4	-24.5 (8)	Pt1—P1—C14—C19	128.9 (2)
C5—C1—S1B—C4	178.2 (4)	C19—C14—C15—C16	-0.9 (5)
S1A—C1—S1B—C4	81 (2)	P1—C14—C15—C16	180.0 (3)
C2B—C3B—C4—C3A	-81.7 (14)	C14—C15—C16—C17	-0.1 (6)
C2B—C3B—C4—S1A	21.9 (13)	C15—C16—C17—C18	1.0 (6)
C2B—C3B—C4—S1B	9.7 (15)	C16—C17—C18—C19	-1.0 (6)
C2A—C3A—C4—C3B	82.5 (13)	C17—C18—C19—C14	0.1 (6)
C2A—C3A—C4—S1A	-4.1 (11)	C15—C14—C19—C18	0.9 (5)
C2A—C3A—C4—S1B	-17.4 (11)	P1—C14—C19—C18	-180.0 (3)
C1—S1A—C4—C3B	-27.5 (9)	C14—P1—C20—C21	-98.8 (3)
C1—S1A—C4—C3A	6.2 (8)	C9—P1—C20—C21	13.8 (4)
C1—S1A—C4—S1B	88 (2)	Pt1—P1—C20—C21	126.9 (3)
C1—S1B—C4—C3B	-10.1 (10)	C14—P1—C20—C25	79.7 (3)
C1—S1B—C4—C3A	24.2 (8)	C9—P1—C20—C25	-167.6 (3)
C1—S1B—C4—S1A	-78 (2)	Pt1—P1—C20—C25	-54.5 (3)
C2B—C1—C5—C6	79.4 (13)	C25—C20—C21—C22	-1.1 (5)
C2A—C1—C5—C6	118.0 (10)	P1—C20—C21—C22	177.5 (3)

S1A—C1—C5—C6	-73.9 (8)	C20—C21—C22—C23	0.5 (6)
S1B—C1—C5—C6	-89.1 (7)	C21—C22—C23—C24	0.2 (6)
C7—N1—C6—C5	113.9 (4)	C22—C23—C24—C25	-0.3 (6)
Pt1—N1—C6—C5	-64.4 (4)	C23—C24—C25—C20	-0.3 (6)
C1—C5—C6—N1	179.9 (4)	C21—C20—C25—C24	1.0 (5)
C6—N1—C7—C8	-174.0 (4)	P1—C20—C25—C24	-177.6 (3)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C7—H7...C12 ⁱ	0.95	2.71	3.486 (4)	139
C12—H12...C11 ⁱⁱ	0.95	2.71	3.639 (4)	168.

Symmetry codes: (i) $x, -y+3/2, z-1/2$; (ii) $x+1, y, z$.

Fig. 1




Fig. 2

