



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

## Diagnosis of EVALI in the COVID-19 era



Published Online  
October 6, 2020  
[https://doi.org/10.1016/S2213-2600\(20\)30450-1](https://doi.org/10.1016/S2213-2600(20)30450-1)

As of Feb 18, 2020, a total of 2807 cases of e-cigarette, or vaping, product use-associated lung injury (EVALI) and 68 attributed deaths have been reported to the US Centers for Disease Control and Prevention.<sup>1</sup> Diagnostic criteria for EVALI consists of a mixture of non-specific systemic symptoms (eg, fever, chills, and vomiting) and respiratory symptoms (eg, shortness of breath, cough, chest pain, pnoea, and hypoxia), along with detection of lung opacities on imaging. Diagnosis also depends on identifying a history of vaping and careful exclusion of alternative conditions.<sup>2</sup> A urine drug screen positive for tetrahydrocannabinol, although non-specific for the disease, might be helpful in this aspect by identifying marijuana-containing electronic cigarettes and vape use. Patient laboratory findings show elevated inflammatory markers and are suggestive of an active non-specific inflammatory process. Radiological diagnosis of EVALI is often challenging because of a large overlap in its radiological features with other disease processes, particularly viral lung infections. Computer tomography of the chest might provide diagnostic clues, as patients with EVALI often showcase a specific airway-centred acute inhalation injury, presenting as a combination of bilateral symmetric ground glass opacities and lung consolidation with subpleural sparing.<sup>3</sup> In 2019, the bronchoscopy results of 19 patients with EVALI were described in *The Lancet*.<sup>4</sup> The findings suggested that while bronchoalveolar lavage (BAL) outcomes were non-specific, they were helpful in excluding alternative infectious processes via bacterial, fungal, viral, and acid-fast bacilli testing. Qualitative presence of lipid-laden macrophages in BAL was shown to have poor specificity and cannot be suggested as a laboratory marker for EVALI.<sup>5</sup> A more promising alternative might be testing for vitamin E acetate in BAL fluid, which was found in 48 of 51 patients with EVALI in a 2020 study.<sup>6</sup> The test was negative in 52 non-users of electronic cigarettes, and even more importantly, in 18 electronic cigarette users without obvious lung disease. These findings make a strong case for detection of vitamin E acetate in BAL as the laboratory standard for diagnosis of EVALI.<sup>6</sup>

Histological findings in EVALI often present a form of airway-centred chemical pneumonitis with various patterns of acute lung injury, such as acute fibrinous

pneumonitis, diffuse alveolar damage, or organising pneumonia. Foamy macrophages and pneumocyte vacuolisation have been reported in many cases and are considered to be more specific for diagnosis in the clinical context.<sup>7</sup> Considering the presence of vast areas of affected lung parenchyma in most patients, transbronchial lung biopsy with or without endobronchial ultrasound guidance seems to be a reasonable addition to the bronchoscopic investigation.

Despite a legal age requirement of 18 years for purchasing electronic cigarettes, use of vaping products has been increasing among teenagers. As a result, cases of EVALI have been reported in children, who also might present with severe respiratory failure requiring intensive therapy. In one study of 98 patients with EVALI, Layden and colleagues reported 26% were aged younger than 18 years.<sup>2</sup>

The severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) pandemic has made differential diagnosis of EVALI even more challenging due to the high rate of SARS-CoV-2 diagnosis in e-cigarette smokers (5 times more likely among daily users<sup>8</sup>) and large overlap of clinical and radiological features of the two conditions.<sup>9</sup> Adding another level of difficulty is the recent emergence of multisystem inflammatory syndrome in children (MIS-C) with COVID-19. Similar to EVALI, MIS-C is clinically characterised by non-specific systemic symptoms and high concentrations of inflammatory markers. At least 18% of children with MIS-C present with shortness of breath and other respiratory symptoms,<sup>10</sup> not unlike those of EVALI. Unfortunately, BAL characteristics and lung pathology of MIS-C are not well described and diagnosis relies heavily on a history of recent SARS-CoV-2 infection or exposure, which is common for children and young adults residing in COVID-19 hotspots, and more common still in those who use vape products.<sup>8</sup> The overlap of similar clinical features, use of vape products, and probable SARS-CoV-2 exposure make distinguishing the differential diagnosis of EVALI from MIS-C challenging in both children and adults, particularly when MIS-C presents with respiratory symptoms.

In summary, clinical presentation of EVALI is fairly non-specific and differential diagnosis is challenging, particularly in the SARS-CoV-2 pandemic era. Nevertheless, radiological detection of airway-centred

acute inhalation lung injury, along with the presence of vacuolisation of macrophages and pneumocytes, and most importantly the detection of vitamin E acetate in BAL fluid, provide important diagnostic clues for this condition.

We declare no competing interests.

\*Mikhail Kazachkov, Melodi Pirzada  
Mikhail.kazachkov@nyulangone.org

Department of Pediatrics, NYU Langone Health, New York, NY 10016, USA (MK); and Department of Pediatrics, NYU Winthrop Hospital, Mineola, NY, USA (MP)

- 1 Centers for Disease Control and Prevention. Outbreak of lung injury associated with the use of e-cigarette, or vaping products. Feb 25, 2020. [https://www.cdc.gov/tobacco/basic\\_information/e-cigarettes/severe-lung-disease.html](https://www.cdc.gov/tobacco/basic_information/e-cigarettes/severe-lung-disease.html) (accessed Sept 29, 2020).
- 2 Layden JE, Ghinai I, Pray I, et al. Pulmonary illness related to e-cigarette use in Illinois and Wisconsin—final report. *N Engl J Med* 2020; **382**: 903–16.
- 3 Foust AM, Winant AJ, Chu WC, Das KM, Phillips GS, Lee EY. Pediatric SARS, H1N1, MERS, EVALL, and now coronavirus disease (COVID-19) pneumonia: what radiologists need to know. *AJR Am J Roentgenol* 2020; **215**: 736–44.
- 4 Blagev DP, Harris D, Dunn AC, Guidry DW, Grissom CK, Lanspa MJ. Clinical presentation, treatment, and short-term outcomes of lung injury associated with e-cigarettes or vaping: a prospective observational cohort study. *Lancet* 2019; **394**: 2073–83.
- 5 Shields PG, Song MA, Freudenheim JL, et al. Lipid laden macrophages and electronic cigarettes in healthy adults. *EBioMedicine* 2020; **60**: 102982.
- 6 Blount BC, Karwowski MP, Shields PG, et al. Vitamin E acetate in bronchoalveolar-lavage fluid associated with EVALL. *N Engl J Med* 2020; **382**: 697–705.
- 7 Butt YM, Smith ML, Tazelaar HD, et al. Pathology of vaping-associated lung injury. *N Engl J Med* 2019; **381**: 1780–81.
- 8 Gaiha SM, Cheng J, Halpern-Felsher B. Association between youth smoking, electronic cigarette use, and COVID-19. *J Adolesc Health* 2020; **67**: 519–23.
- 9 Callahan SJ, Harris D, Collingridge DS, et al. Diagnosing EVALL in the time of COVID-19. *Chest* 2020; published online June 26. <https://doi.org/10.1016/j.chest.2020.06.029>.
- 10 Ahmed M, Advani S, Moreira A, et al. Multisystem inflammatory syndrome in children: a systematic review. *EClinicalMedicine* 2020; published online Sept 4. <https://doi.org/10.1016/j.eclinm.2020.100527>.



## Dexamethasone in hospitalised patients with COVID-19: addressing uncertainties



Lewis Houghton/Science Photo Library

The impressive results of the RECOVERY trial established that a moderate dose of dexamethasone (6 mg daily for 10 days) reduced mortality in hospitalised patients with COVID-19 and respiratory failure who required therapy with supplemental oxygen or mechanical ventilation.<sup>1</sup> The data also indicated that dexamethasone might increase mortality in hospitalised patients who were not receiving oxygen. This landmark trial and the subsequent practice guidelines from several academic and health organisations recommending dexamethasone use in patients with severe COVID-19 have changed clinical practice for hospitalised patients on supplemental oxygen or mechanical ventilation.<sup>2</sup> These favourable findings are supported by three other trials<sup>3–5</sup> of glucocorticoids for COVID-19, which stopped enrolment in early June, 2020, when the RECOVERY trial results were released. Each of these trials showed some evidence of benefit, although none had completed enrolment. A prospective meta-analysis of these and other trials, totalling 1703 participants (1007 [59%] from the RECOVERY trial), confirmed a reduction in 28-day mortality (summary odds ratio [OR] 0.66, 95% CI 0.53–0.82;  $p < 0.001$ ), with minimal heterogeneity across studies.<sup>6</sup> While confirming beneficial effects of corticosteroids for critically ill hospitalised patients

with COVID-19, some unanswered questions and issues remain that deserve discussion and should be addressed in future research.

Because the design of the largest trial, RECOVERY, was pragmatic, data were scarce in some domains. For example, physicians were able to exclude patients from the trial whom they determined should not be a candidate for treatment with dexamethasone, but reasons for exclusion were not recorded. Thus, we do not know why 1707 patients were unsuitable for randomisation. Patients might have been excluded because of perceived contraindications, including uncontrolled diabetes, acute delirium, underlying malignancy, immunosuppression, or other conditions in which corticosteroids might have harmful effects. Therefore, the benefit–risk profile of corticosteroids across the full spectrum of patients with critical COVID-19 and a range of comorbidities remains uncertain.

A second limitation of the RECOVERY trial is that no data are available on the level of oxygen support. In retrospect, this was an important omission from the database. Such data might have revealed differential benefit or harm with dexamethasone treatment according to level of oxygen supplementation.

Published Online  
October 29, 2020  
[https://doi.org/10.1016/S2213-2600\(20\)30503-8](https://doi.org/10.1016/S2213-2600(20)30503-8)