



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Coxsackie A6 (20%), EV71 (4.8%) and echo 3 (3.4%) were the most commonly detected. Analysis of clinical data showed a significant association between hand-foot-mouth disease (HFMD) and genotypes EV71 (risk ratio, rr13.6; Confidence interval, CI 5.4–33.9) and Coxsackie A6 (rr29.4; CI 17.7–48.8). In 2014, Coxsackie A16 was found in stool samples from one AFP case and EV-D68 was found in respiratory specimens from two AFP cases.

**Conclusion:** This study shows the seasonal fluctuation in circulating EV genotypes from one year to the next, verified in prevalence switches mainly between Coxsackievirus and echovirus genotypes. In addition to the well-known association between HFMD and EV71, we detected an association between Coxsackie A6 and HFMD, which has of late also been observed in other countries. These findings highlight the importance of continued EV surveillance in order to establish the virulence of circulating and upcoming EV genotypes and appropriately guide specific Public Health recommendations.

<http://dx.doi.org/10.1016/j.jcv.2016.08.245>

**Abstract no: 312**

**Presentation at ESCV 2016: Poster 206**

### Different epidemiological characteristics of respiratory virus infections in children and adults



Christiane Prifert\*, Benedikt Weissbrich

*Institute of Virology, University of Wuerzburg, Germany*

Acute respiratory tract infections (RTI) are mostly caused by viruses and are a leading cause of morbidity and mortality especially in young children, the elderly, and in immunocompromised patients. We analyzed epidemiological characteristics of RTI in a university hospital setting in the winter seasons 2015 and 2016.

Diagnostics of respiratory viruses was performed prospectively with the multiplex PCR “FTD Respiratory Pathogens 21” (Fast-track Diagnostics). This kit allows detection of influenza A and B virus (Flu A and B), respiratory syncytial virus (RSV), metapneumovirus (MPV), adenovirus (AdV), coronaviruses (CoV) 229E/OC43/NL63/HKU1, parainfluenzaviruses (PIV) 1–4, rhinovirus, enterovirus, parechovirus and bocavirus (BoV). Results of respiratory virus testing were analyzed of all samples received from the hospital of the University of Würzburg, Germany, during the period January 2015 to April 2016. For data analysis, the study period was subdivided into first season (January 2015 to June 2015) and second season (July 2015 to April 2016).

During the study period, 4136 respiratory samples from 2905 patients at the university hospital Würzburg were tested for the presence of respiratory viruses by multiplex PCR, 2948 in the first season and 1188 in the second season. The median age of the patients was 50.3 years (range 0.01–98.3). Of the 2905 patients 1139 (39.2%) were children. The male versus female ratio was 1.34:1. The overall positivity rate was 37.8% in the first season, 47.1% in the second season, and 40.4% during the whole study period. The positivity rate in children was significantly higher than in adults (61.4% versus 29.5%). Similarly, the rate of double virus detections was significantly higher in children than in adults (17.8% versus 6.5%). Detection of three or more viruses in one sample was only observed in children (4.6%). In both seasons the leading virus detected in respiratory samples was rhinovirus with 27.9% and 25.4%, respectively, of all positive samples. Rhinovirus was followed by Flu A (21.7%), RSV (15.8%), and PIV (12.9%) during the first season and by RSV (18.8%), CoV (17.4%), Flu A (14.7%), and MPV (12.7%) during the second season.

The virus distributions in both seasons were considerably different between children and adults.

In summary, comparison of results of respiratory virus diagnostics in children and adult populations shows substantial differences, which demonstrates the need and usefulness of multiplex PCR for broad spectrum detection of respiratory viruses.

<http://dx.doi.org/10.1016/j.jcv.2016.08.246>

**Abstract no: 313**

**Presentation at ESCV 2016: Poster 207**

### Presence of human bocavirus 1 and other viral co-infections in hospitalized children with lower respiratory tract infection in Latvia



A. Vilmane<sup>1,\*</sup>, S. Rasa<sup>1</sup>, I. Ziemele<sup>2,3</sup>,  
D. Gardovska<sup>2,3</sup>, M. Murovska<sup>1</sup>, Z. Nora-Krukle<sup>1</sup>

<sup>1</sup> A. Kirchenstein Institute of Microbiology and Virology, Riga Stradins University, Riga, Latvia

<sup>2</sup> Children Clinical University Hospital, Riga, Latvia

<sup>3</sup> Department of Paediatrics, Riga Stradins University, Riga, Latvia

**Background:** Acute respiratory tract infection (ARTI), especially lower respiratory tract infection (LRTI), is the common cause of illness and hospitalization in children worldwide. However, in many cases the etiological agent of disease is unknown. The viruses primarily associated with respiratory tract infections in children are respiratory syncytial viruses, influenza viruses, parainfluenza viruses, adenoviruses, coronaviruses, rhinoviruses and enteroviruses. In recent years the role of several new respiratory viruses in respiratory tract diseases have been reported, including human metapneumovirus, four coronaviruses (SARS-CoV, HCoV-NL63, HCoV-HKU1, MERS-CoV) and human bocavirus 1 (HBoV1). The aim of this study was to determine the presence of HBoV1 and 18 other respiratory viruses in nasopharyngeal aspirates (NPAs) from hospitalized children with LRTI in Latvia.

**Material and methods:** Forty four children (28 male and 16 female) aged one to 50 months who were hospitalized in Children's Clinical University Hospital and fulfilled WHO LRTI criteria plus had fever ( $T \geq 38.0^\circ\text{C}$ ) were enrolled in this study. In all cases the etiological agent of the disease was not revealed using standard routine clinical methods. NPAs from all patients were obtained on admission and DNA from NPAs was extracted using phenol-chloroform method. All 44 DNA samples were tested for HBoV1 and 18 other respiratory viruses (influenza viruses A, A-H1, A-H1pdm09, A-H3 and B, respiratory syncytial viruses A and B, adenovirus, enterovirus, parainfluenza viruses 1–4, metapneumovirus, rhinovirus, coronaviruses NL63, 229E and OC43) using multiplex real-time PCR method.

**Results:** Among 44 patients with LRTI, 29 (65.9%) were positive for HBoV1 which was the most frequently detected virus in patients. However, only HBoV1 genomic sequence without any analysed coinfection was detected in two out of 29 (6.9%) patients. Respiratory syncytial virus A was found in 23 out of 44 (52.3%) DNA samples and it was the most common co-infection. Other respiratory viruses detected were: adenovirus in 14 (31.8%), rhinovirus in 9 (20.5%), respiratory syncytial virus B in 7 (15.9%), metapneumovirus in 3 (6.8%), parainfluenza virus 3 in 2 (4.5%), coronavirus 229E in 2 (4.5%), enterovirus in 1, influenza A virus in 1, influenza B virus in 1, coronavirus OC43 in 1 and coronavirus NL63 in 1 patient with LRTI. In 13 cases presence of more than two respiratory pathogens were found and in two cases, none of the tested respiratory viruses were detected.