


Analysis of Quorum-Sensing *Pantoea stewartii* Strain M073a through Whole-Genome Sequencing

Nur Izzati Mohamad,^a Wen-Si Tan,^a Chien-Yi Chang,^{b,c} Kok Keng Tee,^d Wai-Fong Yin,^a  Kok-Gan Chan^a

Division of Genetics and Molecular Biology, Institute of Biological Sciences, Faculty of Science, University of Malaya, Kuala Lumpur, Malaysia^a; Interdisciplinary Computing and Complex BioSystems (ICOS) Research Group, School of Computing Science, Newcastle University, Claremont, Tower, Newcastle upon Tyne, United Kingdom^b; Centre for Bacterial Cell Biology, Medical School, Newcastle University, Richardson Road, Newcastle upon Tyne, United Kingdom^c; Department of Medicine, Faculty of Medicine, Centre of Excellence for Research in AIDS (CERiA), University of Malaya, Kuala Lumpur, Malaysia^d

***Pantoea stewartii* strain M073a is a Gram-negative bacterium isolated from a tropical waterfall. This strain exhibits quorum-sensing activity. Here, the assembly and annotation of its genome are presented.**

Received 7 January 2015 Accepted 13 January 2015 Published 19 February 2015

Citation Mohamad NI, Tan W-S, Chang C-Y, Tee KK, Yin W-F, Chan K-G. 2015. Analysis of quorum-sensing *Pantoea stewartii* strain M073a through whole-genome sequencing. *Genome Announc* 3(1):e00022-15. doi:10.1128/genomeA.00022-15.

Copyright © 2015 Mohamad et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Kok-Gan Chan, kokgan@um.edu.my.

Isolated from a tropical waterfall in Sungai Tua located in the Selangor state of Malaysia, strain M073a belongs to the genus *Pantoea stewartii*, members of which have been reported as pathogens known to cause the Stewart's wilt disease, which infects sweet corn (1). In young seedlings, Stewart's wilt causes wilting, which usually result in death of the seedling. In mature plants, leaf blight can be seen parallel to the veins of the leaves. This blight later forms dark-colored necrotic lesions. In some hybrid varieties of sweet corn only small lesions are formed (1 to 2 cm), which is dependent on the level of resistance (2). Members of *Pantoea* are Gram-negative bacterium, and most of them show quorum-sensing (QS) activity (3–5). It has been reported that *P. stewartii* is endemic in various states of the United States (6). Causing disease that commonly occurs in the winter and late summer, *Pantoea* spp. can be found in various habitats from the western borders to the southern and northern coast, including the Plains (7). Here, we present the draft genome of *Pantoea stewartii* strain M073a for the better understanding of the relationship between its pathogenicity and its QS activity.

Genomic DNA of strain M073a was extracted using the QIAamp DNA minikit (Qiagen, Germany) following the instructions on the kit's manual. Quality of the genomic DNA extracted was evaluated using the NanoDrop spectrophotometer (Thermo Scientific, Waltham, MA, USA) and further quantified with the Qubit 2.0 fluorometer (Life Technologies, Carlsbad, CA, USA). The whole genome of strain M073a was sequenced using an Illumina MiSeq sequencer and generated 1,509,201 filtered reads with 56-fold coverage. The reads were then assembled *de novo* using the CLC Genomic Workbench 5.1 (CLC Bio, Denmark). A total of 39 contigs were yielded, with an N_{50} value of 375,223. The resulting draft genome of *P. stewartii* strain M073a was 4,817,607 bp in length with 53% GC content. Prodigal 2.6 software was used for gene prediction and resulted in 4,298 open reading frames (ORF) (8). The Rapid Annotation using Subsystem Technology (RAST) server was used for the annotation of predicted genes (9) and tRNAscan-SE (version 1.21) (10) was used to predict the number of tRNAs, which was 68. RNA genes were predicted using the

RNAmmer software (11), with which three copies of 5S rRNA genes, one copy of a 16S rRNA gene, and one copy of a 32S rRNA gene were found in this genome analysis.

From the annotated genes of the draft genome, there are two genes that play key roles in the QS mechanism, *luxI* and *luxR*, which are found in contig number 14. This finding is consistent with a study conducted by Tsai and Winans in 2010 which report the discovery of LuxI and LuxR homologues which are in the EsaI/EsaR system in *P. stewartii* (12). Since *P. stewartii* regulates its virulence via QS (1), our work on the whole-genome sequencing of *P. stewartii* strain M073a could serve as a QS model to study the QS relationships in this pathogen.

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession number [JXSF010000000](https://www.ncbi.nlm.nih.gov/nuccore/JXSF010000000). The version described in this paper is the first version, JXSF010000000.

ACKNOWLEDGMENT

This work was supported by High Impact Research Grants from the University of Malaya (UM.C/625/1/HIR/MOHE/CHAN/01, grant no. A-000001-50001 and UM-MOHE HIR Grant UM.C/625/1/HIR/MOHE/CHAN/14/1, no. H-50001-A000027) (awarded to K.-G. Chan), which are gratefully acknowledged.

REFERENCES

- Roper MC. 2011. *Pantoea stewartii* subsp. *stewartii*: lessons learned from a xylem-dwelling pathogen of sweet corn. *Mol Plant Pathol* 12:628–637. <http://dx.doi.org/10.1111/j.1364-3703.2010.00698.x>.
- Pataky JK, du Toit LJ, Kunkel TE, Schmitt RA. 1995. Severe Stewart's wilt in central Illinois in sweet corns hybrids moderately resistant to *Erwinia stewartii*. *Plant Dis* 79:104.
- Yunos NYM, Tan WS, Mohamad NI, Tan PW, Adrian TGS, Yin WF, Chan KG. 2014. Discovery of *Pantoea rodasii* strain nd03 that produces *N*-(3-oxo-hexanoyl)-L-homoserine lactone. *Sensors (Basel)* 14: 9145–9152. <http://dx.doi.org/10.3390/s140509145>.
- Tan WS, Muhamad Yunos NYM, Tan PW, Mohamad NI, Adrian TGS, Yin WF, Chan KG. 2014. *Pantoea* sp. isolated from tropical fresh water exhibiting *N*-acyl homoserine lactone production. *ScientificWorldJournal* 2014:828971. <http://dx.doi.org/10.1155/2014/828971>.
- Hong KW, Gan HM, Low SM, Lee PK, Chong YM, Yin WF, Chan KG.

2012. Draft genome sequence of *Pantoea* sp. strain A4, a *Rafflesia*-associated bacterium that produces *N*-acylhomoserine lactones as quorum-sensing molecules. *J Bacteriol* 194:6610. <http://dx.doi.org/10.1128/JB.01619-12>.
6. Menelas B, Block CC, Esker PD, Nutter FW, Jr. 2006. Quantifying the feeding periods required by corn flea beetles to acquire and transmit *Pantoea stewartii*. *Plant Dis* 90:319–324. <http://dx.doi.org/10.1094/PD-90-0319>.
 7. Reche MHLR, Fiuza LM. 2005. Bacterial diversity in rice-field water in Rio Grande do Sul. *Braz J Microbiol* 36:253–257. <http://dx.doi.org/10.1590/S1517-83822005000300009>.
 8. Hyatt D, Chen GL, Locascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* 11:119. <http://dx.doi.org/10.1186/1471-2105-11-119>.
 9. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formisano K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST Server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <http://dx.doi.org/10.1186/1471-2164-9-75>.
 10. Lowe TM, Eddy SR. 1997. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res* 25:955–964. <http://dx.doi.org/10.1093/nar/25.5.0955>.
 11. Lagesen K, Hallin P, Rødland EA, Staerfeldt HH, Rognes T, Ussery DW. 2007. RNAMmer: consistent and rapid annotation of ribosomal RNA genes. *Nucleic Acids Res* 35:3100–3108. <http://dx.doi.org/10.1093/nar/gkm160>.
 12. Tsai CS, Winans SC. 2010. LuxR-type quorum-sensing regulators that are detached from common scents. *Mol Microbiol* 77:1072–1082. <http://dx.doi.org/10.1111/j.1365-2958.2010.07279.x>.