CORRECTION Open Access

Correction to: The relationship between comorbidity medication adherence and health related quality of life among patients with cancer



Dana Drzayich Antol^{1*}, Adrianne Waldman Casebeer¹, Raya Khoury², Todd Michael², Andrew Renda³, Sari Hopson¹, Aparna Parikh², Alisha Stein¹, Mary Costantino¹, Stephen Stemkowski¹ and Mikele Bunce²

Correction to: J Patient-Rep Outcomes 2, 29 (2018) https://doi.org/10.1186/s41687-018-0057-2

Following publication of the original article [1], some errors were identified in relation to the Morisky Medication Adherence Scale (MMAS-8).

- The legend related to Table 1 should read:

Morisky Medication Adherence Score < 6 was defined as low comorbidity medication adherence.

Morisky Medication Adherence Score = 6-8 was defined as moderate/high medication adherence.

Differences were assessed using a t-test for continuous variables and a chi-square test for categorical variables.

-The legend related to Fig. 2 should read:

Morisky Medication Adherence Score < 6 was defined as low comorbidity medication adherence.

Morisky Medication Adherence Score = 6–8 was defined as moderate/high medication adherence.

Differences were assessed using a t-test for continuous variables and a chi-square test for categorical variables.

-In the reference section, one additional reference should be added after reference 24:Morisky DE, DiMatteo MR. Improving the measurement of self-reported medication nonadherence: Final response. J Clin Epidemio 2011; 64: 258–263. PMID:21144706.

The authors regret any error regarding the Morisky Medication Adherence Scale in the original publication.

The original article can be found online at https://doi.org/10.1186/s41687-018-0057-2.

Full list of author information is available at the end of the article



Author details

¹Louisville, KY, USA. ²South San Francisco, California, USA. ³Humana Inc, Louisville, KY, USA.

Published online: 09 March 2021

Reference

 Antol, D., et al. (2018). The relationship between comorbidity medication adherence and health related quality of life among patients with cancer. J Patient-Rep Outcomes, 2, 29 https://doi.org/10.1186/s41687-018-0057-2.

© The Author(s). 2021 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

^{*} Correspondence: ddrzayichjankus@humana.com ¹Louisville, KY, USA