

POSTER PRESENTATION

Open Access

Expression and characterization of mouse prolactin (mPRL) in CHO dhfr- cells

Miriam Fussae Suzuki*, Aline Barros da Silva, João Ezequiel de Oliveira, Fernanda dos Santos Arthuso, Paolo Bartolini, Carlos Roberto Jorge Soares

From 5th Congress of the Brazilian Biotechnology Society (SBBIOTEC)
Florianópolis, Brazil. 10-14 November 2013

Background

Prolactin (PRL) is a peptide hormone related to innumerable physiological functions such as lactation, reproduction, osmotic and immune regulation [1]. In order to evaluate new drugs many tests are conducted in mouse models. Use of prolactin from different origin in mouse model can result in differences in biological activities [2]. Only 61% sequence homology is observed between mouse PRL (197 amino acid residues, 22.5 kDa) and human PRL (199 amino acid residues, 23 kDa). Authentic mouse PRL, without initial methionine, expressed in periplasmic fluid of *E. coli* has already been characterized and, as known, it does not present the glycosylated isoform (G-PRL) [3]. About 5-30% of PRL from mammalian pituitaries is glycosylated [1]. Physiologically, nonglycosylated and glycosylated PRL presented differences concerning lactogenic activity, binding affinity and mitogenicity. The objective of this work was to produce mPRL in CHO dhfr-cells, with gene amplification based on methotrexate addition, so to obtain the glycosylated and non-glycosylated forms of mPRL directly secreted into serum free media.

Methods

The mPRL gene was obtained by PCR reaction from a pUC57 plasmid introducing the human signal peptide and the XbaI restriction enzyme sequence. After purification, the gene was introduced into the pEDdc expression vector. After DNA sequencing and analysis in 1% agarose of the fragments obtained by digestion with the restriction enzymes XbaI, EcoRI, BamHI, and XhoI confirming the correct DNA sequence, Lipofectamine was used to introduce the plasmid pEDdc-mPRL into CHO dhfr- cells. For selection, cells were cultivated in alpha MEM medium

without nucleosides with 10% dialyzed fetal bovine serum and antibiotics. Dot blot and Western blotting with rabbit anti-mPRL antisera (NIDDK-NIH, USA) were used for analyzing 48 positive clones, methotrexate being used for gene amplification. Cultivation was carried out in serum free media CHO-S-SFM II (Invitrogen, USA) in 10 cm diameter dishes. Purification was based on two chromatographic steps: Ni(II)-based Immobilized Metal Ion Affinity Chromatography (IMAC) and Gel Filtration Chromatography (Sephacryl™ S100-HR) as described previously for human PRL [4,5].

Results and conclusions

Construction of pEDdc-mPRL plasmid was confirmed by DNA sequencing and restriction analysis. After selection and amplification by increasing concentrations of methotrexate (from 20 nM to 150 nM), mPRL was synthesized by adherent CHO cells, secreted into the media. Mouse PRL in its authentic form was collected every day. The best clone expressing 4 µg mPRL/10⁶ cells/mL, which is much higher than bacterial expression (0.1 µg/A₆₀₀/mL) [3], was used for production and purification. For the first time mPRL was expressed in CHO cells, both in nonglycosylated and glycosylated forms, as shown in Western Blotting.

Acknowledgements

This work was supported by FAPESP, São Paulo (Project n. 2007/59540-3) and by CNPq, Brasília (Projects PQ 303839/2008-2, PQ 3000473/2009-5 and DT 310512/2010/7).

Published: 1 October 2014

References

1. Sinha YN: Structural variants of prolactin: occurrence and physiological significance. *Endocr Rev* 1995, **16**:354-369.
2. Utama FE, LeBaron MJ, Neilson LM, Sultan AS, Parlow AF, Wagner KU, Rui H: Human prolactin receptors are insensitive to mouse prolactin: implications for xenotransplant modeling of human breast cancer in mice. *J Endocrinol* 2006, **188**:589-601.

Instituto de Pesquisas Energéticas e Nucleares, IPEN-CNEN/SP, Av. Prof. Lineu Prestes, 2242, São Paulo, SP 05508-000, Brazil

3. Suzuki MF, Arthuso FS, Oliveira JE, Oliveira NAJ, Goulart HR, Capone MVN, Ribela MTCP, Bartolini P, Soares CRJ: **Expression, purification, and characterization of authentic mouse prolactin obtained in *Escherichia coli* periplasmic space.** *Biotechnol Appl Bioc* 2012, **59**:178-185.
4. Soares CR, Morganti L, Miloux B, Lupker JH, Ferrara P, Bartolini P: **High-level synthesis of human prolactin in Chinese-Hamster ovary cells.** *Biotechnol Appl Biochem* 2000, **32**(Pt 2):127-135.
5. Ueda EK, Gout PW, Morganti L: **Ni(II)-based immobilized metal ion affinity chromatography of recombinant human prolactin from periplasmic *Escherichia coli* extracts.** *Journal of chromatography A* 2001, **922**:165-175.

doi:10.1186/1753-6561-8-S4-P4

Cite this article as: Suzuki et al.: Expression and characterization of mouse prolactin (mPRL) in CHO dhfr- cells. *BMC Proceedings* 2014 **8**(Suppl 4):P4.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

