



# BMJ Open Association of trajectory of body mass index with knee pain risk in Japanese middle-aged women in a prospective cohort study: the Japan Nurses' Health Study

Ayumi Ito,<sup>1</sup> Kunihiro Hayashi <sup>1</sup>, Shosuke Suzuki,<sup>2,3</sup> Yuki Ideno,<sup>4</sup> Takumi Kurabayashi,<sup>5</sup> Toru Ogata,<sup>6</sup> Atsushi Seichi,<sup>7</sup> Masami Akai <sup>8</sup>, Tsutomu Iwaya<sup>9</sup>

**To cite:** Ito A, Hayashi K, Suzuki S, *et al.* Association of trajectory of body mass index with knee pain risk in Japanese middle-aged women in a prospective cohort study: the Japan Nurses' Health Study. *BMJ Open* 2020;**10**:e033853. doi:10.1136/bmjopen-2019-033853

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2019-033853>).

Received 18 December 2019  
Revised 27 December 2019  
Accepted 09 January 2020



© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

**Correspondence to**  
Dr Kunihiro Hayashi;  
khayashi@gunma-u.ac.jp

## ABSTRACT

**Objectives** To investigate whether body mass index (BMI) trajectory, lifestyle and reproductive factors are associated with knee pain risk among middle-aged women.

**Design** Prospective study of the Japan Nurses' Health Study (JNHS).

**Setting** The JNHS investigates the health of female nurses in Japan. Biennial follow-up questionnaires are mailed to the participants.

**Participants** The 7434 women aged over 40 years who responded to the 10-year self-administered follow-up questionnaire.

**Primary outcome measure** Self-reported knee pain at the 10-year follow-up was the primary outcome. We analysed BMI (normal or overweight) trajectory data from a baseline survey to the 10-year follow-up survey using group-based trajectory modelling. Exposure measurements were BMI trajectory, BMI at age 18 years, lifestyle variables and reproductive history.

**Results** BMI trajectories from baseline to the 10-year follow-up were divided into four groups: remained normal, remained overweight, gained weight or lost weight. At the 10-year follow-up, 1281 women (17.2%) reported knee pain. Multivariable logistic regression analysis revealed that compared with the remained normal group, multivariable-adjusted ORs (95% CI) of knee pain were 1.93 (1.60 to 2.33) for the remained overweight group, 1.60 (1.23 to 2.08) for the gained weight group and 1.40 (0.88 to 2.21) for the lost weight group. The attributable risk percent (95% CI) of the remained overweight group was 48.1% (37.3% to 57.0%) compared with the reference group of remained normal. Alcohol intake at baseline was significantly associated with knee pain.

**Conclusions** The lost weight group had a lower risk than the remained overweight group and the gained weight group and did not carry statistically significant risks for knee pain. Weight reduction and maintaining a normal BMI in middle age was important for preventing knee pain in women.

## INTRODUCTION

Knee pain, along with lower back pain, is one of the most problematic musculoskeletal

## Strengths and limitations of this study

- The longitudinal cohort study design provides data regarding 10-year trajectory in body mass index.
- The Japan Nurses' Health Study, a large cohort study of female nurses, enabled comprehensive analysis, drawing on rich and accurate information regarding healthcare, reproductive health and lifestyle.
- We selected knee pain as the primary outcome, whereas many studies have investigated risk factors for knee osteoarthritis and insufficiently examined risk factors for knee pain.
- Participants were asked to provide their height and weight via self-assessment; we calculated body mass index on the basis of this information.
- We were unable to obtain information regarding the severity or duration of knee pain or the presence of knee pain at baseline; we therefore excluded women who had been diagnosed with knee osteoarthritis prior to the baseline survey.

disorders, and impedes function and affects movement and daily life.<sup>1-4</sup> Knee pain is caused by the age-related weakness of the knee joint, or excessive load on a knee joint, as with knee osteoarthritis (KOA).<sup>5,6</sup> An example of excessive load on a knee joint is one's own body weight. Furthermore, the load of long-term body weight has a great influence on a knee.<sup>7</sup> Although it is necessary to evaluate changes in body weight over time, when we think of knee pain, few studies have investigated the association of the trajectory of body weight and knee pain. There is a sex difference in the prevalence of knee pain: more women experience knee pain than men.<sup>4,7</sup> Considering this sex difference in the prevalence of knee pain, it is important to examine associations of knee pain with

reproductive and lifestyle factors among women. Moreover, several female reproductive factors were associated with BMI in a prior study,<sup>8</sup> so the association between BMI trajectory and knee pain merits investigation.

KOA is a disease that is typically accompanied by severe knee pain. Many publications have reported risk factors for KOA. Ageing and high BMI are well-known factors; other lifestyle (eg, smoking, alcohol drinking habits and sleep duration) and female reproductive factors are also thought to be associated with KOA.<sup>9–15</sup> Some studies have reported that menopausal status, parity and age at menarche are associated with KOA.<sup>15</sup> However, a systematic review reported no significant associations between female hormonal characteristics and KOA.<sup>16</sup> Risk factors for knee pain in male and female working populations were found to be similar to risk factors for KOA in those same populations.<sup>17</sup>

It has been reported that KOA appears after chronic knee symptoms.<sup>18</sup> We examined knee pain as a primary outcome, because knee pain helps to identify early stage KOA, and pain may appear before the diagnosis of musculoskeletal diseases. We conducted a prospective cohort study on the association between BMI trajectory in middle age, lifestyle, reproductive history and knee pain in middle-aged women.

## METHODS


### Study population

The Japan Nurses' Health Study (JNHS), which started in 2001, is a prospective cohort study to investigate the effects of lifestyle and healthcare practices on the health of female nurses in Japan. Participants were female public health nurses, midwives, registered nurses and assistant nurses aged 25 years or older at the time of registration. The JNHS continues through follow-up surveys conducted once every 2 years by a postal self-administered questionnaire.<sup>19 20</sup> The subjects of the current study included 10 667 women over the age of 40 years at the 10-year follow-up who responded to the 10-year follow-up surveys. We excluded 142 women who had diagnosed KOA on the baseline survey, 575 women who became pregnant at least once during the 10-year follow-up period and 2516 women who had not provided their body weight or height for the calculation of BMI between the baseline survey to the 10 years. The remaining 7434 women were analysed (figure 1).

### Outcomes and covariates

Knee pain on the 10-year follow-up surveys was the primary outcome. Participants were asked if they were currently experiencing knee pain, to which they responded yes or no. We also collected data regarding age, female reproductive factors and menopausal status, and parity on the 10-year follow-up questionnaire.

The other covariate data used from the baseline survey were lifestyle factors including smoking status, alcohol intake and sleep duration, and female reproductive factor including age at menarche and body weight at age 18 years



**Figure 1** Selection of subjects for our analysis from the Japan Nurses' Health Study. Of the 10 667 women over 40 years old who responded to the 10-year follow-up (F10) survey, a total of 7434 were included in our analysis. BMI, body mass index; KOA, knee osteoarthritis.

(included as BMI at age 18 years). To identify the BMI trajectory, weight and height from the baseline survey to the 10-year follow-up survey were used. The participants were listed as either never-smokers or ever-smokers (including former smokers and current smokers). Alcohol intake was defined by intake frequency: non-drinkers were those who never drank or drank one to two times per month, light drinkers were those who drank one to two times per week, moderate drinkers were those who drank three to six times per week and heavy drinkers were those who drank every day. The participants were divided into three groups based on average sleep duration, regardless of shift pattern: short (<6 hours per night), moderate (6–<7 hours per night) and long (≥7 hours per night). For menopausal status, women were classified as premenopausal or postmenopausal; notably, postmenopausal included those who were undetermined (ie, those undergoing hormone replacement therapy and those in whom it could not be determined whether they were postmenopausal due to a transitional period). Age at menarche was categorised by tertiles as early (≤11 years old), moderate (12 years old) or late (≥13 years old). We calculated BMI using the women's self-reported height and weight at age 18 years and between the baseline survey and the 10-year follow-up survey. We converted the continuous BMI variable into either normal or overweight prior to drawing the BMI trajectories. BMI <25 kg/m<sup>2</sup> was regarded as normal, and BMI ≥25 was regarded as overweight for all time points. We defined BMI trajectory as the change in BMI from the baseline survey to the 10-year follow-up survey.

### Statistical analysis

Data analyses were performed using SAS statistical software, V.9.4. Group-based trajectory modelling was used to analyse BMI trajectory for indicating the longitudinal changes using PROC TRAJ in SAS.<sup>21–23</sup> This method is one of finite mixture modelling, where some groups are made around an individual BMI trace.

To examine differences between subjects with and without knee pain,  $\chi^2$  tests were used for nominal variables, and Mann-Whitney U tests were used for ordinal variables. A logistic regression analysis was conducted to explore the associations between knee pain and BMI trajectory, lifestyle and reproductive factors. We selected all variables for logistic regression analysis (full model) regardless of univariate relationships. We calculated multivariable-adjusted ORs and 95% CIs, adjusting for age at the 10-year follow-up survey, and for BMI trajectory, smoking, alcohol intake, sleep duration, menopausal status, parity, age at menarche and BMI at age 18 years. The attributable risk percent of the remained overweight group was estimated using multivariable-adjusted OR for BMI trajectory, when the reference group was the other groups.

For sensitivity analyses, the analysis excluded women who had self-reported having cancer due to the possible influence on BMI, and the analysis with multiple imputation for the missing BMI data from the 2-year to 8-year follow-up survey was conducted to confirm the results of primary analyses.<sup>24</sup>

The results were regarded as statistically significant when two-tailed p values were less than 0.05.

### Ethical consideration

Participants were informed of the purpose and procedures of the JNHS. All participants signed consent sheets when they completed the baseline survey.


### Patient and public involvement

Patients and the public were not involved this study. The results will not be presented at conferences. The announcement of the publication is expected to be sufficient. We send annual newsletters to all study participants; these newsletters include information regarding JNHS papers that were published in the previous year.

**Table 1** Age, BMI trajectories, lifestyle variables and reproductive variables for the population of 7434 women

	N	Percentage		N	Percentage
Age at the 10-year follow-up (years)			Menopausal status		
40s	2521	33.9	Premenopausal	2673	36.0
50s	3166	42.6	Postmenopausal/undetermined	4761	64.0
≥60s	1747	23.5	Parity		
BMI trajectory			0	1605	21.6
Remained normal	6123	82.3	1	907	12.2
Remained overweight	785	10.6	2	2907	39.1
Gained weight	401	5.4	3–8	1947	26.2
Lost weight	125	1.7	Missing	68	0.9
Smoking			Age at menarche		
Never-smoker	5445	73.3	Early (≤11 years)	1598	21.5
Ever-smoker	1936	26.0	Moderate (12 years)	2148	28.9
Missing	53	0.7	Late (≥13 years)	3641	49.0
Alcohol intake			Missing	47	0.6
Non-drinker	4244	57.1	BMI at age 18 years		
Light drinker	1143	15.4	Normal (<25)	6874	92.4
Moderate drinker	1311	17.6	Overweight (≥25)	363	4.9
Heavy drinker	439	5.9	Missing	197	2.7
Missing	297	4.0			
Sleep duration					
Short	1093	14.7			
Moderate	3213	43.2			
Long	2898	39.0			
Missing	230	3.1			

BMI trajectory: BMI from the baseline to the 10-year follow-up survey.  
BMI, body mass index.



**Figure 2** Trajectories of body mass index (BMI) changes. BMI was converted to binary data: BMI  $\geq 25$  as overweight was 1.0, and BMI  $< 25$  as normal was 0.0. Ninety-five per cent CIs are shown as a dashed line. Overall, 82.3% remained normal weight; 10.6% remained overweight; 5.4% gained weight; 1.7% lost weight.

## RESULTS

### Characteristics of the study population

The characteristics of the study population are shown in [table 1](#). The mean (SD) age was 53.5 (7.7) years. The mean BMI at age 18 was 21.1 (2.3), and the mean BMI at the baseline survey was 21.8 (2.8). The percentage of those who were overweight based on BMI at the baseline survey was 11.9%, while 88.1% had a normal BMI. At the 10-year follow-up survey, the mean BMI was 22.2 (3.2), overweight BMI was 16.0% and normal BMI was 84.0%. Regarding the four BMI trajectory groups, there were 6123 (82.3%) women in the remained normal group, 785 (10.6%) in the remained overweight group, 401 (5.4%) in the gained weight group and 125 (1.7%) in the lost weight group ([figure 2](#)). The mean (SD) sleep duration was 6.5 (0.9) hours per night. The mean (SD) age at menarche was 12.6 (1.3) years. The population included 2673 (36.0%) premenopausal women and 4761 (64.0%) postmenopausal women.

### Prevalence of knee pain and musculoskeletal diseases

A total of 1281 (17.2%) women with knee pain at the 10-year follow-up survey were identified in the study population ([table 2](#)). [Table 2](#) shows p values from the  $\chi^2$  tests for nominal variables and from Mann-Whitney U tests for ordinal variables to examine the differences between respondents with and without knee pain.

### Factors associated with knee pain

[Table 2](#) also shows the associations between knee pain and potential risk factors from multivariable logistic regression. Age at the time of the 10-year follow-up survey was significantly associated with risk of knee pain. Multivariable-adjusted ORs (95% CIs) were 1.29 (1.03 to 1.61) and 1.57 (1.21 to 2.04) for women between the age of 50–59 years and age 60 years or older, respectively, compared with women between 40 years and 49 years.

Multivariable-adjusted ORs (95% CIs) were 1.93 (1.60 to 2.33), 1.60 (1.23 to 2.08) and 1.40 (0.88 to 2.21) for remained overweight, gained weight and lost weight groups, respectively, compared with women in the remained normal group. The attributable risk per cent (ie, excess risk fraction) (95% CI) of the remained overweight group was 48.1% (37.3% to 57.0%), 27.5% (–17.0% to 55.1%) and 16.8% (–12.3% to 38.4%) compared with the reference groups of remained normal, lost weight and gained weight groups, respectively. The association between BMI at age 18 years and knee pain was not statistically significant (multivariable-adjusted OR=1.28, 95% CI 0.98 to 1.68).

The multivariable-adjusted OR for experiencing knee pain was 1.37 (95% CI 1.06 to 1.77) for heavy drinker compared with non-drinker. There were no statistically significant associations between knee pain and smoking, sleep duration, menopausal status, parity or age at menarche.

We conducted sensitivity analyses, multivariable-adjusted ORs (95% CIs) excluded women who had self-reported having cancer were 2.09 (1.70 to 2.56), 1.62 (1.22 to 2.15) and 1.40 (0.83 to 2.37) for the remained overweight, gained weight and lost weight groups, respectively, compared with women in the remained normal group. In another sensitivity analysis, the multiple imputation for the missing BMI data from the 2-year to 8-year follow-up survey showed that multivariable-adjusted ORs (95% CIs) were 2.11 (1.78 to 2.49), 1.62 (1.27 to 2.06) and 1.23 (0.93 to 1.62) for the remained overweight, gained weight and lost weight groups, respectively, compared with women in the remained normal group.

According to these sensitivity analyses, we confirmed there were no significant differences in the other resulting ORs of BMI trajectory using logistic regression.



**Table 2** Prevalence and multivariable-adjusted ORs for knee pain on the 10-year follow-up survey

	Knee pain	No knee pain	P value	Multivariate-adjusted ORs (95% CI)
	n=1281	n=6153		
	N (prevalence %)*	N		
<b>Age (years)</b>				
40s	316 (12.5)	2205	<0.0001†	reference
50s	580 (18.3)	2586		<b>1.29 (1.03 to 1.61)</b>
≥60s	385 (22.0)	1362		<b>1.57 (1.21 to 2.04)</b>
<b>BMI trajectory</b>				
Remained normal	939 (15.3)	5184	<0.0001‡	reference
Remained overweight	224 (28.5)	561		<b>1.93 (1.60 to 2.33)</b>
Gained weight	89 (22.2)	312		<b>1.60 (1.23 to 2.08)</b>
Lost weight	29 (23.2)	96		1.40 (0.88 to 2.21)
<b>Smoking</b>				
Never-smoker	902 (16.6)	4543	<b>0.0138‡</b>	reference
Ever-smoker	369 (19.1)	1567		1.14 (0.99 to 1.32)
Missing	10	43		
<b>Alcohol intake</b>				
Non-drinker	719 (16.9)	3525	0.4671†	reference
Light drinker	192 (16.8)	951		1.02 (0.85 to 1.23)
Moderate drinker	212 (16.2)	1099		0.93 (0.77 to 1.10)
Heavy drinker	95 (21.6)	344		<b>1.37 (1.06 to 1.77)</b>
Missing	63	234		
<b>Sleep duration</b>				
Short	217 (19.9)	876	<b>0.0068†</b>	1.12 (0.93 to 1.34)
Moderate	553 (17.2)	2660		reference
Long	463 (16.0)	2435		0.92 (0.79 to 1.06)
Missing	48	182		
<b>Menopausal status</b>				
Premenopausal	337 (12.6)	2336	<0.0001‡	reference
Postmenopausal/undetermined	944 (19.8)	3817		1.24 (0.99 to 1.54)
<b>Parity</b>				
0	263 (16.4)	1342	<b>0.0475†</b>	reference
1	145 (16.0)	762		0.96 (0.76 to 1.22)
2	494 (17.0)	2413		0.90 (0.75 to 1.07)
3–8	363 (18.6)	1584		0.99 (0.82 to 1.20)
Missing	16	52		
<b>Age at menarche</b>				
Early (≤11 years)	276 (17.3)	1322	0.9477†	1.04 (0.86 to 1.25)
Moderate (12 years)	369 (17.2)	1779		reference
Late (≥13 years)	630 (17.3)	3011		1.01 (0.87 to 1.18)
Missing	6	41		
<b>BMI at age 18 years</b>				
Normal (<25)	1158 (16.9)	5716	<0.0001‡	reference
Overweight (≥25)	91 (25.1)	272		1.28 (0.98 to 1.68)
Missing	32	165		

Continued

Table 2 Continued

	Knee pain	No knee pain		Multivariate-adjusted ORs (95% CI)
	n=1281	n=6153		
	N (prevalence %)*	N	P value	

Multivariable-adjusted ORs (95% CI), adjusted for age at the 10-year survey, BMI trajectory, smoking, alcohol intake, sleep duration, menopausal status, parity, age at menarche and BMI at age 18 years.

BMI trajectory: BMI from baseline to the 10-year follow-up survey.

Bold values: statistically significant.

\*Prevalence of knee pain (%).

†Mann-Whitney U test.

‡ $\chi^2$ .

BMI, body mass index.

## DISCUSSION

We found that BMI trajectory was a predictive factor for knee pain. Remaining overweight increased the risk of knee pain. It is known that high BMI is associated with knee pain or KOA,<sup>9</sup> and our results were partially in agreement with prior studies. We found that remaining overweight was associated with a 1.9 times increased risk compared with those who remained at a normal weight. The risk of knee pain was increased approximately 1.6 times when the women who were normal weight became overweight. The women with normal BMI should continue to control her weight in order to maintain the low risk for knee pain. However, if overweight women reduced their weight over 10 years, they were able to decrease the risk of knee pain by 27.5%. Furthermore, remaining overweight group and gained weight group carried statistically significant risks for knee pain; however, lost weight group did not. This BMI trajectory showed the need to reduce the weight and maintain a normal BMI to help prevent knee pain. There are some studies that have described weight change and its influence on knee pain.<sup>9 25 26</sup> Our study showed that a dynamic BMI change was related to knee pain. We found that there was no statistically significant association between BMI at age 18 years and knee pain in middle age, and BMI trajectory in middle age was a much stronger predictor of knee pain compared with BMI at age 18 years, although these results were different from a prior study in men.<sup>27</sup> This phenomenon of knee pain may be caused by loading on the knee joint. The mechanisms of the association between obesity and knee pain constitute a combination of mechanical, structural, metabolic and behavioural changes.<sup>28</sup> Obesity leads to a loss of muscle mass and strength, as well as the accumulation of fat tissue; these factors together contribute to knee joint compression.<sup>29</sup> Conversely, weight loss may improve physical activity level, increase muscle strength and reduce knee pain-related disability.<sup>30</sup> Our findings suggest that major weight gain or the presence of an overweight status for a long period in middle age are probable risk factors for knee pain. In addition, weight loss and maintenance of normal weight may prevent the development of knee pain. Musculoskeletal pain might impact health-related quality of life.<sup>31</sup> Therefore, the continuous control of body weight in middle age may be effective to prevent knee pain and to maintain quality of life.

The present study had several strengths. First, the present study was a longitudinal cohort study, such that a 10-year trajectory in BMI could be examined. Second, because the JNHS comprises a large cohort study of female nurses, we were able to conduct a comprehensive analysis, drawing on rich and accurate information regarding healthcare, reproductive health and lifestyle. Third, we selected knee pain as the primary outcome, whereas many studies have investigated risk factors for KOA and insufficiently examined risk factors for knee pain. Knee pain is a prodromal symptom of KOA; it helps us to identify the risk of KOA. However, the present study had two limitations. First, participants were asked to provide their height and weight, which was a self-assessment. We calculated BMI on the basis of this information. Second, we were unable to obtain information regarding the severity or duration of knee pain or the presence of knee pain at baseline; if we had this information, we could assess the incidence of knee pain from baseline among Japanese female nurses in middle age. However, we excluded women who had been diagnosed with KOA prior to the baseline survey.

The prevalence of knee pain in the present study among Japanese nurses (17.2%) was higher than the prevalence of extremity joint pain previously found among the general population of Japanese women (7.0%).<sup>32</sup> In previous studies involving nurses, the prevalence of knee symptoms (including ache, pain and discomfort) ranged from 10.6% to 24.7% in the USA, Australia, Korea and Estonia.<sup>33–36</sup> Nurses worldwide are clearly in at-risk settings for knee symptoms, and consideration must be given to nurses' knee pain. One of the limitations of the present study was the composition of the study population. Nurses may be exposed to different risk factors, such as manually handling patients, hard physical work and high mental pressure because of their working circumstances.<sup>37</sup> Therefore, our findings regarding prevalence may not be generalisable to the general population. However, there is no reason to suspect that the general population of women would differ in the association between the risk of knee pain and BMI trajectory.

Heavy drinkers had a risk for 10 years later knee pain than non-drinkers in this study, whilst the association between alcohol intake and KOA also had shown in a previous study.<sup>10</sup> There is no convincing explanation about the

mechanism of the association between alcohol drinking habit and knee pain.

In conclusion, the lost weight group had a lower risk than the remained overweight group and the gained weight group. From our longitudinal observational study of the 10-year trajectory in BMI, we found that weight reduction and maintaining a normal BMI in middle age were important for preventing knee pain in women. Further studies are needed to indicate the effectiveness of losing weight and maintaining a normal BMI in the prevention of knee pain.

#### Author affiliations

<sup>1</sup>Unit of International and Community Health Sciences, Graduate School of Health Sciences, Gunma University, Maebashi, Japan

<sup>2</sup>Faculty of Medicine, Gunma University, Maebashi, Japan

<sup>3</sup>Eco-Health Research Group, Iseaki, Japan

<sup>4</sup>Center for Mathematics and Data Science, Gunma University, Maebashi, Japan

<sup>5</sup>Department of Obstetrics & Gynecology, Niigata City General Hospital, Niigata, Japan

<sup>6</sup>Center for Sport Science and Health Promotion, Department of Rehabilitation for the Movement Functions, National Rehabilitation Center for Persons with Disabilities, Tokorozawa, Japan

<sup>7</sup>Department of Orthopaedic Surgery, Mitsui Memorial Hospital, Chiyoda-ku, Japan

<sup>8</sup>Graduate School, International University of Health and Welfare Kohokai Group, Minato-ku, Japan

<sup>9</sup>Nagano University of Health and Medicine, Nagano, Japan

**Acknowledgements** The authors would like to thank Drs Yukiko Miyazaki, Ayumi Kobayashi and Kazue Nagai for their advice.

**Contributors** AI analysed the data and drafted the report. KH designed and initiated the study. AI, KH, SS and YI contributed to the interpretation and discussion of the data and writing of the manuscript. All authors approved the final draft to be published and have agreed to be accountable for all aspects of the work, ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

**Funding** This work was partly supported by a Japanese Society for Musculoskeletal Medicine Research Grant, as well as JSPS KAKENHI Grant Number 18H04069.

**Competing interests** None declared.

**Patient consent for publication** Not required.

**Ethics approval** The JNHS study protocol was approved by the institutional review board of the Gunma University (Approval No. 101, study protocol=JNHS0101).

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** All data relevant to the study are included in the article.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

#### ORCID iDs

Kunihiko Hayashi <http://orcid.org/0000-0003-4648-3936>

Masami Akai <http://orcid.org/0000-0002-9145-7158>

#### REFERENCES

- Alexopoulos EC, Tanagra D, Detorakis I, *et al*. Knee and low back complaints in professional Hospital nurses: occurrence, chronicity, care seeking and absenteeism. *Work* 2011;38:329–35.
- Agaliotis M, Mackey MG, Jan S, *et al*. Burden of reduced work productivity among people with chronic knee pain: a systematic review. *Occup Environ Med* 2014;71:651–9.
- Coggon D, Ntani G, Palmer KT, *et al*. The CUPID (cultural and psychosocial influences on disability) study: methods of data collection and characteristics of study sample. *PLoS One* 2012;7:e39820.
- Andersen RE, Crespo CJ, Ling SM, *et al*. Prevalence of significant knee pain among older Americans: results from the third National health and nutrition examination survey. *J Am Geriatr Soc* 1999;47:1435–8.
- Ding C, Cicuttini F, Blizzard L, *et al*. A longitudinal study of the effect of sex and age on rate of change in knee cartilage volume in adults. *Rheumatology* 2007;46:273–9.
- Mezhov V, Ciccutini FM, Hanna FS, *et al*. Does obesity affect knee cartilage? A systematic review of magnetic resonance imaging data. *Obes Rev* 2014;15:143–57.
- Dawson J, Juszcak E, Thorogood M, *et al*. An investigation of risk factors for symptomatic osteoarthritis of the knee in women using a life course approach. *J Epidemiol Community Health* 2003;57:823–30.
- Newby PK, Dickman PW, Adami H-O, *et al*. Early anthropometric measures and reproductive factors as predictors of body mass index and obesity among older women. *Int J Obes* 2005;29:1084–92.
- Felson DT, Zhang Y, Hannan MT, *et al*. Risk factors for incident radiographic knee osteoarthritis in the elderly: the Framingham study. *Arthritis Rheum* 1997;40:728–33.
- Muthuri SG, Zhang W, Maciewicz RA, *et al*. Beer and wine consumption and risk of knee or hip osteoarthritis: a case control study. *Arthritis Res Ther* 2015;17:23.
- Zhang Y, Zeng C, Wei J, *et al*. Associations of cigarette smoking, betel quid chewing and alcohol consumption with high-sensitivity C-reactive protein in early radiographic knee osteoarthritis: a cross-sectional study. *BMJ Open* 2016;6:e010763.
- Kong L, Wang L, Meng F, *et al*. Association between smoking and risk of knee osteoarthritis: a systematic review and meta-analysis. *Osteoarthritis Cartilage* 2017;25:809–16.
- Jung JH, Seok H, Choi SJ, *et al*. The association between osteoarthritis and sleep duration in Koreans: a nationwide cross-sectional observational study. *Clin Rheumatol* 2018;37:1653–9.
- Taylor-Gjevne RM, Gjevne JA, Nair B, *et al*. Components of sleep quality and sleep fragmentation in rheumatoid arthritis and osteoarthritis. *Musculoskeletal Care* 2011;12:n/a–159.
- Hussain SM, Wang Y, Giles GG, *et al*. Female reproductive and hormonal factors and incidence of primary total knee arthroplasty due to osteoarthritis. *Arthritis Rheumatol* 2018;70:1022–9.
- de Klerk BM, Schiphof D, Groeneweld FPMJ, *et al*. No clear association between female hormonal aspects and osteoarthritis of the hand, hip and knee: a systematic review. *Rheumatology* 2009;48:1160–5.
- Miranda H, Viikari-Juntura E, Martikainen R, *et al*. A prospective study on knee pain and its risk factors. *Osteoarthritis Cartilage* 2002;10:623–30.
- Case R, Thomas E, Clarke E, *et al*. Prodromal symptoms in knee osteoarthritis: a nested case-control study using data from the osteoarthritis initiative. *Osteoarthritis Cartilage* 2015;23:1083–9.
- Hayashi K, Mizunuma H, Fujita T, *et al*. Design of the Japan nurses' health study: a prospective occupational cohort study of women's health in Japan. *Ind Health* 2007;45:679–86.
- Nagai K, Hayashi K, Yasui T, *et al*. Disease history and risk of comorbidity in women's life course: a comprehensive analysis of the Japan nurses' health study baseline survey. *BMJ Open* 2015;5:e006360.
- Nagin DS. Analyzing developmental trajectories: a semiparametric, group-based approach. *Psychol Methods* 1999;4:139–57.
- Jones BL, Nagin DS, ROEDER K. A SAS procedure based on mixture models for estimating developmental trajectories. *Social Methods Res* 2001;29:374–93.
- Jones BL, Nagin DS. Advances in group-based trajectory modeling and an SAS procedure for estimating them. *Social Methods Res* 2007;35:542–71.
- He Y, Zaslavsky AM, Landrum MB, *et al*. Multiple imputation in a large-scale complex survey: a practical guide. *Stat Methods Med Res* 2010;19:653–70.
- Goulston LM, Kiran A, Javaid MK, *et al*. Does obesity predict knee pain over fourteen years in women, independently of radiographic changes? *Arthritis Care Res* 2011;63:1398–406.
- Tanamas SK, Wluka AE, Davies-Tuck M, *et al*. Association of weight gain with incident knee pain, stiffness, and functional difficulties: a longitudinal study. *Arthritis Care Res* 2013;65:34–43.
- Macfarlane GJ, de Silva V, Jones GT. The relationship between body mass index across the life course and knee pain in adulthood: results from the 1958 birth cohort study. *Rheumatology* 2011;50:2251–6.



- 28 Janke EA, Collins A, Kozak AT. Overview of the relationship between pain and obesity: what do we know? where do we go next? *JRRD* 2007;44:245–62.
- 29 Friilander H, Viikari-Juntura E, Heliövaara M, *et al.* Obesity in early adulthood predicts knee pain and walking difficulties among men: a life course study. *Eur J Pain* 2016;20:1278–87.
- 30 Vincent HK, Heywood K, Connelly J, *et al.* Obesity and weight loss in the treatment and prevention of osteoarthritis. *Pm R* 2012;4:S59–67.
- 31 Heo M, Allison DB, Faith MS, *et al.* Obesity and quality of life: mediating effects of pain and comorbidities. *Obes Res* 2003;11:209–16.
- 32 Ministry of Health, Labour and Welfare. Comprehensive survey of living conditions. 2016. Tokyo, Japan: Ministry of health, labour and welfare. Available: [www.mhlw.go.jp/english/database/db-hss/cslc-index.html](http://www.mhlw.go.jp/english/database/db-hss/cslc-index.html) [Accessed 15 Aug 2019].
- 33 Kim H, Dropkin J, Spaeth K, *et al.* Patient handling and musculoskeletal disorders among hospital workers: analysis of 7 years of institutional workers' compensation claims data. *Am J Ind Med* 2012;55:683–90.
- 34 Reed LF, Battistutta D, Young J, *et al.* Prevalence and risk factors for foot and ankle musculoskeletal disorders experienced by nurses. *BMC Musculoskelet Disord* 2014;15:196.
- 35 Kee D, Seo SR. Musculoskeletal disorders among nursing personnel in Korea. *Int J Ind Ergon* 2007;37:207–12.
- 36 Freimann T, Coggon D, Merisalu E, *et al.* Risk factors for musculoskeletal pain amongst nurses in Estonia: a cross-sectional study. *BMC Musculoskelet Disord* 2013;14:334.
- 37 Smith DR, Mihashi M, Adachi Y, *et al.* A detailed analysis of musculoskeletal disorder risk factors among Japanese nurses. *J Safety Res* 2006;37:195–200.