

RESEARCH ARTICLE

Comparative primary paediatric nasal epithelial cell culture differentiation and RSV-induced cytopathogenesis following culture in two commercial media

Lindsay Broadbent¹, Sheerien Manzoor¹, Maria C. Zarcone², Judit Barabas¹, Michael D. Shields^{1,3}, Sejal Saglani², Claire M. Lloyd², Andrew Bush⁴, Adnan Custovic⁵, Peter Ghazal⁶, Mindy Gore², Ben Marsland⁷, Graham Roberts⁸, Jurgan Schwarze⁹, Steve Turner¹⁰, Ultan F. Power^{1*}



1 Wellcome-Wolfson Institute for Experimental Medicine, Queens University Belfast, Belfast, Northern Ireland, United Kingdom, **2** Inflammation, Repair and Development Section, National Heart and Lung Institute, Faculty of Medicine, Imperial College, London, England, United Kingdom, **3** Royal Belfast Hospital for Sick Children, Belfast Health & Social Care Trust, Belfast, Northern Ireland, United Kingdom, **4** Department of Paediatric Respiratory Medicine, Royal Brompton Hospital, London, England, United Kingdom, **5** Department of Paediatrics, Imperial College London, London, England, United Kingdom, **6** Division of Infection and Pathway Medicine, Deanery of Biomedical Sciences, University of Edinburgh Medical School, Edinburgh, Scotland, United Kingdom, **7** Department of Immunology and Pathology, Monash University, Melbourne, Australia, **8** Clinical and Experimental Sciences and Human Development and Health, Faculty of Medicine, University of Southampton, Southampton, England, United Kingdom, **9** Child Life and Health and MRC-Centre for Inflammation Research, Queen's Medical Research Institute, University of Edinburgh, Edinburgh, Scotland, United Kingdom, **10** Child Health, University of Aberdeen, Aberdeen, Scotland, United Kingdom

* u.power@qub.ac.uk

OPEN ACCESS

Citation: Broadbent L, Manzoor S, Zarcone MC, Barabas J, Shields MD, Saglani S, et al. (2020) Comparative primary paediatric nasal epithelial cell culture differentiation and RSV-induced cytopathogenesis following culture in two commercial media. PLoS ONE 15(3): e0228229. <https://doi.org/10.1371/journal.pone.0228229>

Editor: Ralph A. Tripp, University of Georgia, UNITED STATES

Received: January 7, 2020

Accepted: March 5, 2020

Published: March 26, 2020

Peer Review History: PLOS recognizes the benefits of transparency in the peer review process; therefore, we enable the publication of all of the content of peer review and author responses alongside final, published articles. The editorial history of this article is available here: <https://doi.org/10.1371/journal.pone.0228229>

Copyright: © 2020 Broadbent et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: Data available from a public repository run by Queen's University Belfast.

Abstract

The culture of differentiated human airway epithelial cells allows the study of pathogen-host interactions and innate immune responses in a physiologically relevant *in vitro* model. As the use of primary cell culture has gained popularity the availability of the reagents needed to generate these cultures has increased. In this study we assessed two different media, Promocell and PneumaCult, during the differentiation and maintenance of well-differentiated primary nasal epithelial cell cultures (WD-PNECs). We compared and contrasted the consequences of these media on WD-PNEC morphological and physiological characteristics and their responses to respiratory syncytial virus (RSV) infection. We found that cultures generated using PneumaCult resulted in greater total numbers of smaller, tightly packed, pseudostratified cells. However, cultures from both media resulted in similar proportions of ciliated and goblet cells. There were no differences in RSV growth kinetics, although more ciliated cells were infected in the PneumaCult cultures. There was also significantly more IL-29/IFN λ 1 secreted from PneumaCult compared to Promocell cultures following infection. In conclusion, the type of medium used for the differentiation of primary human airway epithelial cells may impact experimental results.

DOI: [10.17034/7c15ce87-a011-47e7-ba55-50e0ccb550a9](https://doi.org/10.1371/journal.pone.0228229).

Funding: MDS, SS, CML, AB, AC, PG, BM, GR, JS, ST and UFP were awarded grant number: 108818/Z/15/B from The Wellcome Trust. <https://wellcome.ac.uk> MDS and UFP were awarded grant number Com/4044/09 from HSC Research & Development (HSC R&D) Division. <https://research.hscni.net>; <https://wellcome.ac.uk/> The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

Introduction

Eukaryotic cell culture is one of the fundamental techniques used by biomedical researchers. Cell culture techniques are routinely used across most disciplines of life science research. Cell culture has advanced dramatically in recent years with the development of differentiated primary epithelial cell cultures[1,2], organoids[3,4] and organ-on-chip systems[5,6]. One of the key aspects of mammalian cell culture is the growth medium. The cell culture medium must provide all of the key nutrients required for cell survival and division, an overview of which is provided by Lodish et al[7]. The choice of cell culture medium is dependent on the cell type in culture and the intended use of the cultures, as components of cell culture medium could affect experimental outcomes. Primary cell culture and the development of more complex cellular models requires highly specialised media to support the growth and differentiation of the cells. This study focused on the culture of air-liquid interface differentiated primary airway epithelial cells and their use in virus-host interaction research.

Advancements in airway epithelial primary cell culture, including the use of growth factors, hormones and the use of an air-liquid interface[8,9], have led to important discoveries in virology and virus-host interactions[10–14]. The main advantage of using well-differentiated primary airway epithelial cell cultures to study respiratory virus-host interactions is the similarity of the cultures to the *in vivo* targets of infection. Well-differentiated primary airway epithelial cell (WD-PAEC) cultures closely mimic the *in vivo* airways, demonstrating pseudostratified morphologies containing ciliated cells, mucus-producing goblet cells and tight junctions[2]. Indeed, we previously demonstrated that WD-PAECs recreate several hallmarks of RSV infection *in vivo*, including RSV infection of ciliated cells but not goblet cells, loss of ciliated cells, increased goblet cell numbers, occasional syncytia, and the secretion of pro-inflammatory chemokines[15].

WD-PAEC cultures derived from patients with specific airway diseases often retain the features of that disease. Cultures derived from cystic fibrosis patients have been used to investigate the potential for personalised treatment[16]. The differentiation of these cultures is essential for measurement of CFTR function. The culture of primary airway epithelial cells has also enhanced the diagnosis of primary ciliary dyskinesia, which is notoriously difficult to diagnose[17].

Initially, the choice of media for the culture of WD-PAECs was limited. However, as the use of these culture systems increases in popularity the availability of specific reagents has also increased. Our laboratory has cultured WD-PAECs for over ten years. Our protocols included the use of Promocell Airway Epithelial Cell Growth Medium to differentiate and maintain the cultures[2]. Using this method, we achieved over 90% success at differentiating primary airway epithelial cell samples derived from paediatric nasal or bronchial brushes in Transwells. However, for a period of ten months we experienced unexplained repeated failure in culture differentiation, and our success rates decreased to ~50%. This led us to assess another primary cell medium, PneumCult-ALI medium, for use in differentiating paediatric primary airway epithelial cells.

In this study, therefore, we evaluated the use of the two media in parallel for the culture and differentiation of airway epithelial cells. We assessed the cultures for the total number of cells, ciliated cells, goblet cells and epithelial integrity. We hypothesised that the choice of differentiation medium would affect the cytopathogenesis and antiviral immune responses of the WD-PNEC cultures to RSV infection.

Materials & methods

Cell lines and viruses

The origin and characterization of the clinical isolate RSV BT2a were previously described [18]. RSV titres in biological samples were determined using HEP-2 cells, as previously described[19].

WD-PNEC cultures

Primary nasal epithelial cells ($n = 3$ donors) were obtained from healthy paediatric patients with full parental consent. The nasal brushes were processed and the monolayer cell cultures were treated as previously described[2]. Cells were passaged twice in Promocell Airway Epithelial Cell Growth Medium (C-21160 Promocell) (supplements added as per the manufacturer's instructions with additional penicillin/streptomycin). When ~90% confluent the cells were seeded onto collagen coated Transwell supports (Corning) at 2×10^4 or 5×10^4 cells per Transwell. Cultures were submerged in modified Promocell Airway Epithelial Cell Growth Medium (see Table 1) supplemented with retinoic acid until fully confluent. After 4–6 days of submersion air-liquid interface (ALI) was initiated by removing the apical medium. This is required to trigger differentiation. Thereafter, half of the Transwell cultures were maintained in Promocell medium and half using PneumaCult-ALI medium supplemented with hydrocortisone and heparin. See Table 1 for constituents of the media, where known. Stemcell Technologies, the producer of PneumaCult, did not disclose the ingredients of the supplements provided with the medium. Medium was replaced with 500 μ L of fresh medium in the basolateral compartment every 2 days. Complete differentiation took at least 21 days. Cultures were only used when hallmarks of excellent differentiation were evident, including extensive apical coverage with beating cilia and obvious mucus production. Trans-epithelial electrical resistance (TEER) was measured using an EVOM2 and ENDOHM 6 mm chamber (World Precision Instruments).

Infection

WD-PNECs were infected apically for 2 h at 37°C with 1.4×10^5 TCID₅₀ RSV BT2a in 50 μ L of DMEM (low glucose, no additives). Cultures were then rinsed 4 times with 250 μ L DMEM (low glucose, no additives). The fourth wash was retained as the 2 hpi time point. At 24 hpi and every 24 h thereafter until 96 hpi apical washes were undertaken and harvested by adding 250 μ L DMEM apically, pipetted up and down gently and aspirated without damaging the cultures, added to cryovials and snap frozen in liquid nitrogen. RSV titres in biological samples were determined by a tissue culture infectious dose 50 (TCID₅₀) assay, as previously described [19].

Immunofluorescence

WD-PNECs were fixed with 200 μ L apically and 500 μ L basolaterally of 4% PFA (v/v in PBS) for 1 h then permeabilised with 0.1% Triton X-100 (v/v in PBS) for 1 h. Cells were blocked

Table 1. Known constituents of promocell and PnemaCult differentiation media.

Promocell Airway Epithelial Cell Growth Medium			PneumaCult-ALI Medium		
Promocell kit supplements	BPE	52 μ g/mL	PneumaCult- ALI x10 supplement	Unknown	Unknown
	hEGF	10 ng/mL			
	Insulin	5 μ g/mL	PneumaCult-ALI maintenance supplement x100	Unknown	Unknown
	Hydrocortisone	0.5 μ g/mL			
	Epinephrine	0.5 μ g/mL			
	Transferrin	10 μ g/mL			
User-optimised supplements	BSA	1.5 μ g/mL	User-optimised supplements	Hydrocortisone	1×10^{-6} M
	Retinoic acid	15 ng/mL		Heparin	4 μ g/mL
	Penicillin	100 U/mL		Penicillin	100 U/mL
	Streptomycin	100 μ g/mL		Streptomycin	100 μ g/mL

<https://doi.org/10.1371/journal.pone.0228229.t001>

with 0.4% BSA (v/v in PBS) for 30 min. Immunofluorescent staining was performed for Muc5Ac (1:100 dilution, mouse monoclonal; Abcam) (goblet cell marker), β -tubulin (1:200 dilution, rabbit polyclonal Cy3 conjugated; Abcam) (ciliated cell marker) and RSV F protein (1:500 dilution, 488 conjugated; Millipore). Cultures were mounted using DAPI mounting medium (Vectashield, Vector Labs) and imaged using a Nikon Eclipse 90i or a Leica SP5 confocal microscope. For ZO-1 images cultures were fixed in 4% PFA for 20 min at room temperature, followed by permeabilization (Permeabilization Buffer set, Ebioscience) and blocking with 2% BSA solution (Sigma). Cells were stained with anti-ZO-1 mouse mAb (ThermoFischer, Alexa Fluor 488). Images were acquired on an inverted laser scanning confocal microscope (SP5, Leica Microsystems).

IFN λ 1/IL-29 ELISA

The concentration of IFN λ 1/IL-29 was measured in basolateral medium from RSV BT2a- or mock-infected cultures at 96 hpi by ELISA (Thermo Fisher Scientific; BMS2049). The manufacturer's instructions were followed.

Microscopy and image analysis

For enumeration of cell types, a minimum of 5 fields were captured per condition/well per patient by UV microscopy (Nikon Eclipse 90i). Differential interference contrast (DIC) microscopy was used to capture bright field images of differentiated cultures. Image analysis was carried out using ImageJ software (<http://rsbweb.nih.gov/ij/>). ImageJ was also used to calculate the diameter of cells. The diameter of >40 cells across 5 fields of view per patient were measured.

Statistical analysis

GraphPad Prism[®] was used to create graphical representations of the data and for statistical analysis. To assess statistical significance results were compared using t tests, except for viral growth kinetics, which were compared by calculating the areas under the curves.

Results

To determine the effect of Promocell or PneumaCult medium on cell proliferation during differentiation, cells were seeded at two different densities on Transwell supports. Trans-epithelial electrical resistance (TEER), a measure of epithelial integrity, was measured in cultures seeded with 5×10^4 cells. There was a trend towards increased TEERs in PneumaCult cultures but this did not reach significance (Fig 1A). Expression of ZO-1, a marker of tight junctions, was clearly evident in cultures differentiated in both media (Fig 1C). Cultures were trypsinised to determine the total cell count (Fig 1B). The seeding density, either 2×10^4 or 5×10^4 cells per Transwell, did not affect the final number of cells in the cultures. PneumaCult medium resulted in ~3-fold higher cell counts following differentiation than Promocell medium.

Cells differentiated in Promocell appeared larger than those in PneumaCult under light and fluorescent microscopy. This was confirmed by imaging the cultures using DIC microscopy and measuring the cell diameters (Fig 2A and 2B). Cells differentiated and maintained in Promocell medium were significantly larger than cells in cultures from the same donors but differentiated using PneumaCult medium. Confocal microscopy revealed that the cells within the PneumaCult cultures appeared more tightly packed (Fig 2C). Orthogonal sections suggest a greater degree of stratification of the cultures differentiated in PneumaCult medium compared to Promocell medium.




Fig 1. Primary paediatric nasal epithelial cells were passed twice then seeded on collagen coated Transwell supports at a seeding density of 2×10^4 or 5×10^4 per Transwell. Cells were maintained in Promocell medium while submerged. Following ALI initiation half of the cultures from each donor were maintained using Promocell and half using Pneumacult. Cultures were differentiated for 21 days. TEER values were measured in the cultures seeded at 5×10^4 ($n = 5$ donors) (A). Cultures were trypsinised and a cell count was performed ($n = 3$ donors) (B). Cultures were fixed in 4% PFA and stained for ZO-1 ($n = 3$), representative images (C). Statistical significance was determined using unpaired t-tests. ** = $p < 0.01$, *** = $p < 0.001$.

<https://doi.org/10.1371/journal.pone.0228229.g001>

The number of total, ciliated and goblet cells in fixed cultures differentiated in Promocell or PneumaCult medium were enumerated from *en face* images following fluorescent microscopy of cultures stained for DAPI (nuclei), β -tubulin (cilia) and Muc5Ac (goblet cells), respectively. Representative images of cultures from both media are presented in Fig 3A. Consistent with the data presented in Fig 1B above, the PneumaCult-maintained cultures demonstrated higher numbers of total cells, as well as ciliated and goblet cells (Fig 3B). In all cases, there was a trend towards increased cell numbers in the PneumaCult cultures, although they did not reach significance. When the proportion of ciliated and goblet cells was calculated, however, there was no difference between Promocell- or PneumaCult-maintained cultures (Promocell: 75.3% ciliated and 5.1% goblet cells; PneumaCult: 75.1% ciliated and 4.2% goblet cells). This was consistent for both seeding densities (data not shown for seeding density 2×10^4) (Fig 3C).

A central theme of research in our laboratory is to study RSV interactions with paediatric airway epithelium. To explore whether the medium used affected RSV growth kinetics or cytopathogenesis, cultures were infected with the low passaged clinical isolate RSV BT2a. The same amount of virus (1.4×10^5 TCID₅₀) was inoculated onto all cultures. At the specified times post infection, apical washes were titrated on HEP-2 cells to determine virus growth kinetics (Fig 4). There was no significant difference in viral growth kinetics between the two initial seeding densities of the cultures or the medium used to differentiate and maintain the cultures. As RSV infects ciliated epithelium and, because of higher ciliated cell numbers, we expected the PneumaCult cultures to reach higher peak viral titres released from them. However, all culture conditions resulted in similar peak viral titres and growth kinetics.

The secretion of IFN λ 1/IL-29, a type-III interferon known to be the main interferon secreted following RSV infection of airway epithelium [20,21], was quantified in the basolateral medium at 96 hpi (Fig 6). There was significantly more IFN λ 1/IL-29 secreted from the PneumaCult compared to the Promocell cultures. This may be due to the larger number of cells




Fig 2. Primary paediatric nasal epithelial cells ($n = 3$ donors) seeded at 5×10^4 per Transwell were differentiated and maintained using either Promocell or Pneumacult medium. Cultures were fixed using 4% paraformaldehyde (PFA) on day 25 post air-liquid interface (ALI) initiation. Images were captured by DIC microscopy at x60 magnification and imageJ was used to determine the diameter of cells (yellow lines) (A). Graphical representation of the average cell diameter in pixels (B). Statistical significance was determined using unpaired t-tests. *** = $p < 0.001$. Cultures were stained for beta-tubulin (red) and DAPI (blue). Z-stacks were obtained using a confocal microscope at x100 magnification (Leica SP5) (C).

<https://doi.org/10.1371/journal.pone.0228229.g002>

present in the PneumaCult cultures. However, there are approximately 3x the number of cells in the PneumaCult compared to the Promocell cultures, yet the IFN λ 1/IL-29 was >6x that secreted from Promocell cultures. The PneumaCult cultures may respond more robustly to infection but further work, investigating different cytokines/chemokines, would be needed to provide further insights into these differences in innate immune responses to RSV infection.

Discussion

In this study we confirmed our hypothesis that the choice of medium affects the resultant cultures. PneumaCult medium resulted in cultures with ~3x more cells than those differentiated




Fig 3. WD-PNEC cultures (n = 3 donors) with an initial seeding density of 5×10^4 per Transwell were differentiated in Promocell or Pneumacult medium. After 21 days cultures were fixed in 4% paraformaldehyde and stained for β -tubulin, a ciliated cell marker; Muc5ac, a goblet cell marker and counterstained for DAPI. Representative images of β -tubulin staining (A). The average number of total, ciliated and goblet cells from 5 fields of view per donor was calculated (B). The percentage of ciliated cells and goblet cells in the culture was calculated (C). Images were acquired using a Nikon Eclipse 90i at x60 magnification.

<https://doi.org/10.1371/journal.pone.0228229.g003>




Fig 4. Primary paediatric nasal epithelial cells (n = 3 donors) seeded on collagen coated Transwell supports at a seeding density of 2×10^4 or 5×10^4 per Transwell. Were maintained in either Promocell or Pneumacult medium for 21 days. Cultures were infected with RSV BT2a 1.4×10^5 TCID₅₀. Apical washes were harvested at 2 and 24 hpi and every 24 h thereafter and titrated on HEP-2 cells to determine virus growth kinetics. RSV-infected cultures were fixed at 96 hpi and the total number of cells and the number of ciliated, goblet and RSV-infected cells were enumerated in *en face* IF images (Fig 5). Following infection, the mean number of cells was significantly different between Promocell- and PneumaCult-differentiated cultures, with a 46% and 37% reduction in mean cell numbers, respectively. All cultures demonstrated a similar loss in ciliated cell numbers following RSV infection, 20% and 23% reduction for Promocell and PneumaCult, respectively. There were significantly more RSV-infected cells in the PneumaCult cultures, consistent with higher numbers of ciliated cells in these cultures compared to the Promocell cultures. However, despite the higher numbers of ciliated cells, as previously mentioned (Fig 3B), and the greater number of RSV-infected cells (Fig 5B), the viral growth kinetics were not significantly different. The percentage of ciliated, goblet and RSV-infected cells in the cultures did not differ significantly as a function of the culture medium used.

<https://doi.org/10.1371/journal.pone.0228229.g004>




Fig 5. WD-PNEC cultures (n = 3 donors) were differentiated in Promocell or Pneumacult medium. After 21 days cultures were infected with RSV BT2a or mock infected. At 96 hpi the cultures were fixed in 4% paraformaldehyde and stained for β -tubulin, a ciliated cell marker; Muc5ac, a goblet cell marker, RSV F and counterstained for DAPI. Representative images of β -tubulin staining (A). The average number of total, ciliated, goblet and RSV infected cells from 5 fields of view per donor was calculated (B). The percentage of ciliated, goblet and RSV infected cells in the culture was calculated (C). Images were acquired using a Nikon Eclipse 90i at x60 magnification. Statistical significance was determined by t-test.

<https://doi.org/10.1371/journal.pone.0228229.g005>

using Promocell medium. Interestingly, despite the differences in total cells counts, the proportions of ciliated cells and goblet cells were similar for both culture conditions. Indeed, the proportion of ciliated cells was consistent with the proportion reported in normal healthy human airway epithelium (50–70%) [22,23]. The limited evidence available suggests that




Fig 6. WD-PNEC cultures (n = 3 donors) were differentiated in Promocell or Pneumacult medium. After 21 days cultures were infected with RSV BT2a or mock infected. Basolateral medium was harvested and snap frozen every 24 hpi. The concentration of IFN λ 1/IL-29 in the basolateral medium at 96 hpi was quantified by ELISA. Statistical significance was determined by t-test. * = p<0.05.

<https://doi.org/10.1371/journal.pone.0228229.g006>

goblet cells represent up to 25% of cells in adult human airway epithelium[24]. The proportions of goblet cells found in our current cultures were considerably lower than this. However, we are unaware of the lower proportion of goblet cells found in normal human paediatric airway epithelium and, therefore, cannot conclude whether or not our cultures are abnormal with respect to goblet cell content. The percent goblet cells that we report here were also lower compared to our previous WD-PNEC cultures derived from newborn and 1-year-old infants [25]. Reasons for these discrepancies remain to be elucidated, although they may be due to nasal epithelial cell donor- or culture-specific factors.

Ciliated cells, mucus and the airway surface liquid (ASL) are key components of the mucociliary escalator, the primary defence mechanism against inhaled pathogens and foreign particulate material. RSV infection disrupts this by having a detrimental effect on the number of ciliated cells[26]. RSV-induced cilia loss was replicated in this WD-PNEC model under both culture conditions, reinforcing the evidence that the WD-PAEC model reliably recreates at least some RSV cytopathogenesis.

Another noticeable impact of the different media was the different cell sizes. There is very little published data available on the size of human nasal epithelial cells *in vivo*. Due to experimental differences, it is difficult to compare cultured nasal epithelial cells. However, previous work from our group demonstrated differentiated nasal epithelial cells of ~12 μm in diameter (data not shown). In the present study cells of ~14 μm and ~25 μm diameters from PneumaCult and Promocell differentiation media, respectively, were evident.

As we are unaware of the precise constituents of the proprietary PneumaCult medium, it is not possible to determine which components, if any, might be responsible for the differential cell count and size between the cultures. We are aware, however, that Promocell medium contains bovine pituitary extract (BPE) (52 $\mu\text{g}/\text{mL}$), while PneumaCult medium apparently does not. BPE contains components which are needed for differentiation of epithelial cells, including growth factors and hormones[27]. As it is derived from animal tissues, the components of BPE can vary between batches. This may explain in part the variable success rates we previously experienced in differentiating WD-PNEC cultures from nasal brushes. However, this remains to be confirmed. BPE-free media, such as PneumaCult, bypass the need for this component by supplementing with a cocktail of hormones and growth factors, allowing for greater reproducibility of the composition of different batches of media.

Although the proportion of ciliated cells is the same in both cultures there was an increase in actual numbers of ciliated cells in the PneumaCult cultures. Ciliated cells are the primary target for RSV infection[15,28]. As such, following RSV infection there were more RSV infected cells in the PneumaCult compared with the Promocell cultures. Both media resulted in cultures that were successfully infected with RSV with very similar viral growth kinetics to that previously reported [2,15]. The difference in cell numbers and, indeed, the difference in the number of RSV+ cells in the cultures did not have a significant impact on apically-released virus titres. This indicates that the number of cells within a culture is not a defining factor in viral growth kinetics. The factors affecting viral growth kinetics are not fully understood. Interestingly, much higher concentrations of IFN λ 1/IL-29 were secreted from RSV-infected PneumaCult cultures, which may be due to the higher cell density within the culture. We previously demonstrated that IFN λ 1/IL-29 was responsible for attenuating RSV growth kinetics in WD-PBECs[21]. The higher IFN λ 1/IL-29 concentrations secreted from the RSV-infected PneumaCult compared to Promocell cultures, therefore, might explain in part the similarities in RSV growth kinetics, despite the higher number of RSV+ cells in the former cultures.

In conclusion, both media tested under these conditions result in WD-PAEC cultures that possess several hallmarks of airway epithelium *in vivo* and resulted in comparable experimental outcomes in several of the parameters assessed. However, our data also indicate that the

choice of medium used to differentiate and maintain primary airway epithelial cell cultures may impact the experimental outcomes and care should be taken in choosing medium for the intended work. However, one should be cognisant of the low donor numbers used in this study. It should also be noted that we did not use either media 'off-the-shelf' and extensive optimisation is often needed to achieve the best culturing conditions. As cell culturing techniques advance and become more sophisticated there will undoubtedly be an increase in reagents created specifically for this purpose. As such, it will be imperative that independent comparisons between different reagents, such as media, are undertaken to ensure reliability of the data generated.

Author Contributions

Conceptualization: Lindsay Broadbent, Maria C. Zarcone, Claire M. Lloyd, Ultan F. Power.

Data curation: Lindsay Broadbent, Maria C. Zarcone.

Formal analysis: Lindsay Broadbent.

Funding acquisition: Michael D. Shields, Sejal Saglani, Claire M. Lloyd, Andrew Bush, Adnan Custovic, Peter Ghazal, Ben Marsland, Graham Roberts, Jurgen Schwarze, Steve Turner, Ultan F. Power.

Investigation: Lindsay Broadbent, Sheerien Manzoor, Judit Barabas.

Methodology: Lindsay Broadbent, Sheerien Manzoor, Sejal Saglani, Claire M. Lloyd, Ultan F. Power.

Project administration: Mindy Gore.

Supervision: Ultan F. Power.

Visualization: Michael D. Shields.

Writing – original draft: Lindsay Broadbent.

Writing – review & editing: Lindsay Broadbent, Sheerien Manzoor, Maria C. Zarcone, Michael D. Shields, Sejal Saglani, Claire M. Lloyd, Andrew Bush, Adnan Custovic, Peter Ghazal, Mindy Gore, Ben Marsland, Graham Roberts, Jurgen Schwarze, Steve Turner, Ultan F. Power.

References

1. Guo-Parke H, Canning P, Douglas I, Villenave R, Heaney LG, Coyle P V, et al. Relative respiratory syncytial virus cytopathogenesis in upper and lower respiratory tract epithelium. *Am J Respir Crit Care Med* [Internet]. 2013 Oct 1 [cited 2014 Jul 30]; 188(7):842–51. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23952745> PMID: 23952745
2. Broadbent L, Villenave R, Guo-Parke H, Douglas I, Shields MD, Power UF. In Vitro Modeling of RSV Infection and Cytopathogenesis in Well-Differentiated Human Primary Airway Epithelial Cells (WD-PAECs). In: *Methods in molecular biology* (Clifton, NJ) [Internet]. 2016 [cited 2016 Sep 5]. p. 119–39. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27464691>
3. Lancaster MA, Knoblich JA. Organogenesis in a dish: Modeling development and disease using organoid technologies. *Science* (80-) [Internet]. 2014 Jul 18 [cited 2019 Oct 18]; 345(6194):1247125–1247125. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25035496>
4. Rossi G, Manfrin A, Lutolf MP. Progress and potential in organoid research. *Nat Rev Genet* [Internet]. 2018 Nov 18 [cited 2019 Oct 18]; 19(11):671–87. Available from: <http://www.nature.com/articles/s41576-018-0051-9> PMID: 30228295
5. Benam KH, Villenave R, Lucchesi C, Varone A, Hubeau C, Lee H-H, et al. Small airway-on-a-chip enables analysis of human lung inflammation and drug responses in vitro. *Nat Methods* [Internet]. 2016

- Feb 21 [cited 2019 Oct 18]; 13(2):151–7. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26689262> PMID: 26689262
6. Nawroth JC, Barrile R, Conegliano D, van Riet S, Hiemstra PS, Villenave R. Stem cell-based Lung-on-Chips: The best of both worlds? *Adv Drug Deliv Rev.* 2018;
 7. Lodish H, Berk A, Zipursky SL, Matsudaira P, Baltimore D, Darnell J. Growth of Animal Cells in Culture. In: *Molecular Cell Biology* [Internet]. 4th ed. W. H. Freeman; 2000 [cited 2019 Aug 26]. <https://www.ncbi.nlm.nih.gov/books/NBK21682/#A1384>
 8. Gruenert DC, Finkbeiner WE, Widdicombe JH. Culture and transformation of human airway epithelial cells. *Am J Physiol Cell Mol Physiol* [Internet]. 1995 Mar [cited 2019 Aug 26]; 268(3):L347–60. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/7900815>
 9. Moller PC, Partridge LR, Cox R, Pellegrini V, Ritchie DG. An in vitro system for the study of tracheal epithelial cells. *Tissue Cell* [Internet]. 1987 Jan [cited 2015 Apr 22]; 19(6):783–91. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/3438910> PMID: 3438910
 10. Stark JM, Huang YT, Carl J, Davis PB. Infection of cultured human tracheal epithelial cells by human parainfluenza virus types 2 and 3. *J Virol Methods* [Internet]. 1991 Jan [cited 2019 Aug 26]; 31(1):31–45. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/1849915> PMID: 1849915
 11. Reiss TF, Gruenert DC, Nadel JA, Jacoby DB. Infection of cultured human airway epithelial cells by influenza A virus. *Life Sci* [Internet]. 1991 [cited 2019 Aug 26]; 49(16):1173–81. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/1895875> PMID: 1895875
 12. Slepshkin VA, Staber PD, Wang G, McCray PB, Davidson BL. Infection of Human Airway Epithelia with H1N1, H2N2, and H3N2 Influenza A Virus Strains. *Mol Ther* [Internet]. 2001 Mar 1 [cited 2019 Aug 26]; 3(3):395–402. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S1525001601902771> PMID: 11273782
 13. Goris K, Uhlenbruck S, Schwegmann-Wessels C, Köhl W, Niedorf F, Stern M, et al. Differential Sensitivity of Differentiated Epithelial Cells to Respiratory Viruses Reveals Different Viral Strategies of Host Infection. *J Virol* [Internet]. 2009 Feb 15 [cited 2019 Aug 26]; 83(4):1962–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/19052091> PMID: 19052091
 14. Villenave R, O'Donoghue D, Thavagnanam S, Touzelet O, Skibinski G, Heaney LG, et al. Differential cytopathogenesis of respiratory syncytial virus prototypic and clinical isolates in primary pediatric bronchial epithelial cells. *Virol J* [Internet]. 2011 Jan [cited 2014 Aug 18]; 8(1):43. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3039598&tool=pmcentrez&rendertype=abstract>
 15. Villenave R, Thavagnanam S, Sarlang S, Parker J, Douglas I, Skibinski G, et al. In vitro modeling of respiratory syncytial virus infection of pediatric bronchial epithelium, the primary target of infection in vivo. *Proc Natl Acad Sci U S A* [Internet]. 2012 Mar 27 [cited 2014 Nov 5]; 109(13):5040–5. Available from: <http://www.pnas.org/content/109/13/5040.short> PMID: 22411804
 16. Awatade NT, Uliyakina I, Farinha CM, Clarke LA, Mendes K, Solé A, et al. Measurements of Functional Responses in Human Primary Lung Cells as a Basis for Personalized Therapy for Cystic Fibrosis. *EBio-Medicine* [Internet]. 2015 Feb [cited 2019 Oct 18]; 2(2):147–53. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26137539> PMID: 26137539
 17. Pifferi M, Montemurro F, Cangiotti AM, Ragazzo V, Di Cicco M, Vinci B, et al. Simplified cell culture method for the diagnosis of atypical primary ciliary dyskinesia. *Thorax* [Internet]. 2009 Dec 1 [cited 2019 Oct 18]; 64(12):1077–81. Available from: <https://thorax.bmj.com/content/64/12/1077> PMID: 19770168
 18. Villenave R, O'Donoghue D, Thavagnanam S, Touzelet O, Skibinski G, Heaney LG, et al. Differential cytopathogenesis of respiratory syncytial virus prototypic and clinical isolates in primary pediatric bronchial epithelial cells. *Virol J.* 2011 Jan; 8(1):43.
 19. Power UF, Plotnicky-Gilquin H, Huss T, Robert A, Trudel M, Stahl S, et al. Induction of protective immunity in rodents by vaccination with a prokaryotically expressed recombinant fusion protein containing a respiratory syncytial virus G protein fragment. *Virology.* 1997 Apr; 230(2):155–66. <https://doi.org/10.1006/viro.1997.8465> PMID: 9143271
 20. Okabayashi T, Kojima T, Masaki T, Yokota S-I, Imaizumi T, Tsutsumi H, et al. Type-III interferon, not type-I, is the predominant interferon induced by respiratory viruses in nasal epithelial cells. *Virus Res* [Internet]. 2011 Sep [cited 2014 Aug 29]; 160(1–2):360–6. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21816185> PMID: 21816185
 21. Villenave R, Broadbent L, Douglas I, Lyons JD, Coyle P V., Teng MN, et al. Induction and Antagonism of Antiviral Responses in Respiratory Syncytial Virus-infected Paediatric Airway Epithelium.
 22. Jafri HS, Chavez-Bueno S, Mejias A, Gomez AM, Rios AM, Nassi SS, et al. Respiratory syncytial virus induces pneumonia, cytokine response, airway obstruction, and chronic inflammatory infiltrates associated with long-term airway hyperresponsiveness in mice. *J Infect Dis* [Internet]. 2004 May 15 [cited 2015 Sep 16]; 189(10):1856–65. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/15122522> PMID: 15122522

23. Schamberger AC, Staab-Weijnitz CA, Mise-Racek N, Eickelberg O. Cigarette smoke alters primary human bronchial epithelial cell differentiation at the air-liquid interface. *Sci Rep* [Internet]. 2015 Jul 2 [cited 2019 Oct 23]; 5(1):8163. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25641363>
24. Rogers DF. The airway goblet cell. *Int J Biochem Cell Biol* [Internet]. 2003 Jan 1 [cited 2019 Oct 25]; 35(1):1–6. Available from: <https://www.sciencedirect.com/science/article/pii/S1357272502000833> PMID: 12467641
25. Groves HE, Guo-Parke H, Broadbent L, Shields MD, Power UF. Characterisation of morphological differences in well-differentiated nasal epithelial cell cultures from preterm and term infants at birth and one-year. *PLoS One* [Internet]. 2018 [cited 2019 Aug 26]; 13(12):e0201328. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/30517096> PMID: 30517096
26. Smith CM, Kulkarni H, Radhakrishnan P, Rutman A, Bankart MJ, Williams G, et al. Ciliary dyskinesia is an early feature of respiratory syncytial virus infection. *Eur Respir J* [Internet]. 2014 Feb 1 [cited 2019 Jun 20]; 43(2):485–96. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23520320> PMID: 23520320
27. Hintze JM, Tchoukalova YD, Sista R, Shah MK, Zhang N, Lott DG. Development of xeno-free epithelial differentiation media for adherent, non-expanded adipose stromal vascular cell cultures. *Biochem Biophys Res Commun* [Internet]. 2018 Sep 18 [cited 2019 Oct 29]; 503(4):3128–33. Available from: <https://www.sciencedirect.com/science/article/pii/S0006291X18317893?via%3Dihub> PMID: 30166060
28. Zhang L, Peeples ME, Boucher RC, Collins PL, Pickles RJ. Respiratory syncytial virus infection of human airway epithelial cells is polarized, specific to ciliated cells, and without obvious cytopathology. *J Virol* [Internet]. 2002 Jun [cited 2018 Aug 20]; 76(11):5654–66. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/11991994>