

- receptor-A expression in the renal inner medullary collecting duct in decompensated heart failure. *J Pharmacol Exp Ther*. 2011;339:203–209.
14. Liu C, Guan J, Kang Y, et al. Inhibition of dehydration-induced water intake by glucocorticoids is associated with activation of hypothalamic natriuretic peptide receptor-a in Rat. *PLoS One*. 2010;5:e15607.
  15. Aguirre JA, Ibarra FR, Barontini M, et al. Effect of glucocorticoids on renal dopamine production. *Eur J Pharmacol*. 1999;370:271–278.
  16. De Matteo R, May CN. Glucocorticoid-induced renal vasodilatation is mediated by a direct renal action involving nitric oxide. *Am J Physiol*. 1997;273:R1972–R1979.
  17. Cope CL, Pearson J. Clinical value of the cortisol secretion rate. *J Clin Pathol*. 1965;18:82–87.
  18. Yancy CW, Jessup M, Bozkurt B, et al. 2013 ACCF/AHA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. 2013;62:e147–e239.
  19. Anand IS, Ferrari R, Kalra GS, et al. Edema of cardiac origin. Studies of body water and sodium, renal function, hemodynamic indexes, and plasma hormones in untreated congestive cardiac failure. *Circulation*. 1989;80:299–305.
  20. Krozowski Z, Li KX, Koyama K, et al. The type I and type II 11beta-hydroxysteroid dehydrogenase enzymes. *J Steroid Biochem Mol Biol*. 1999;69:391–401.
  21. Liu C, Wang N, Meng H, et al. Glucocorticoids for decompensated heart failure. *Cochrane Database Syst Rev*. 2015. In press.
  22. Newman DA. Reversal of intractable cardiac edema with prednisone. *N Y State J Med*. 1959;59:625–633.
  23. Mickerson JN, Swale J. Diuretic effect of steroid therapy in obstinate heart failure. *Br Med J*. 1959;1:876–883.
  24. Bolotin MT. Prednisone in treatment of cardiac decompensation; a new treatment of cardiac pleural effusion. *Am Pract Dig Treat*. 1957;8:1772–1775.
  25. Ashbel SI, Reznik ND, Vladyko NV. Use of methylprednisolone in patients with circulatory failure [in Russian]. *Klin Med (Mosk)*. 1985; 63:111–114.
  26. Damjanic P, Vierhapper H. Permissive action of glucocorticoid substitution therapy on the effects of atrial natriuretic peptide (hANP) in patients with adrenocortical insufficiency. *Exp Clin Endocrinol*. 1990;95:315–321.
  27. De Kloet ER, Vreugdenhil E, Oitzl MS, et al. Brain corticosteroid receptor balance in health and disease. *Endocr Rev*. 1998;19:269–301.
  28. Francis GS, Siegel RM, Goldsmith SR, et al. Acute vasoconstrictor response to intravenous furosemide in patients with chronic congestive heart failure. Activation of the neurohumoral axis. *Ann Intern Med*. 1985; 103:1–6.
  29. Anand IS. Cardiorenal syndrome: a cardiologist's perspective of pathophysiology. *Clin J Am Soc Nephrol*. 2013;8:1800–1807.
  30. Unger T, Badoer E, Gareis C, et al. Atrial natriuretic peptide (ANP) as a neuropeptide: interaction with angiotensin II on volume control and renal sodium handling. *Br J Clin Pharmacol*. 1990;30(suppl 1):83S–88S.
  31. Whitworth JA, Gordon D, Andrews J, et al. The hypertensive effect of synthetic glucocorticoids in man: role of sodium and volume. *J Hypertens*. 1989;7:537–549.
  32. Newton MA, Laragh JH. Effects of glucocorticoid administration on aldosterone excretion and plasma renin in normal subjects, in essential hypertension and in primary aldosteronism. *J Clin Endocrinol Metab*. 1968;28:1014–1022.
  33. Torre-Amione G, Kapadia S, Benedict C, et al. Proinflammatory cytokine levels in patients with depressed left ventricular ejection fraction: a report from the Studies of Left Ventricular Dysfunction (SOLVD). *J Am Coll Cardiol*. 1996;27:1201–1206.
  34. Valgimigli M, Ceconi C, Malagutti P, et al. Tumor necrosis factor-alpha receptor 1 is a major predictor of mortality and new-onset heart failure in patients with acute myocardial infarction: the Cytokine-Activation and Long-Term Prognosis in Myocardial Infarction (C-ALPHA) study. *Circulation*. 2005;111:863–870.
  35. Dunlay SM, Weston SA, Redfield MM, et al. Tumor necrosis factor-alpha and mortality in heart failure: a community study. *Circulation*. 2008;118:625–631.
  36. Amir O, Rogowski O, David M, et al. Circulating interleukin-10: association with higher mortality in systolic heart failure patients with elevated tumor necrosis factor-alpha. *Isr Med Assoc J*. 2010;12:158–162.

## ERRATUM

### Calorie Restriction Attenuates Monocrotaline-induced Pulmonary Arterial Hypertension in Rats: Erratum

In the article that appeared on page 562 of the June 2015 issue, the address given for the corresponding author, Enqing Fu, MD, PhD, was incorrect. The correct address is Department of Respiratory Medicine, Tangdu Hospital, Fourth Military Medical University.

### REFERENCE

Ding M, Lei J, Qu Y, Zhang H, Xin W, Ma F, Liu S, Li Z, Jin F, Fu E. Calorie Restriction Attenuates Monocrotaline-induced Pulmonary Arterial Hypertension in Rats. *J Cardiovasc Pharmacol* 2015;65:562–570.